

# RENDICONTI *del* SEMINARIO MATEMATICO *della* UNIVERSITÀ DI PADOVA

ROSSELLA AGLIARDI

**Fourier integral operators of infinite order  
on  $\mathcal{D}_{L^2}^{\{\sigma\}}$  ( $\mathcal{D}_{L^2}^{\{\sigma\}'}$ ) with an application to a  
certain Cauchy problem**

*Rendiconti del Seminario Matematico della Università di Padova,*  
tome 84 (1990), p. 71-82

[http://www.numdam.org/item?id=RSMUP\\_1990\\_\\_84\\_\\_71\\_0](http://www.numdam.org/item?id=RSMUP_1990__84__71_0)

© Rendiconti del Seminario Matematico della Università di Padova, 1990, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## Fourier Integral Operators of Infinite Order on $\mathcal{D}_L^{(\sigma)}$ ( $\mathcal{D}_L^{(\sigma')}$ ) with an Application to a Certain Cauchy Problem.

ROSSELLA AGLIARDI (\*)

### Introduction.

The aim of this paper is to develop a calculus of Fourier integral operators of infinite order in the spaces  $\mathcal{D}_L^{(\sigma)}$  ( $\mathcal{D}_L^{(\sigma')}$ ) and to apply it to prove some sufficient conditions for a certain Cauchy problem to be well-posed in the above-mentioned spaces.

The calculus we develop here is analogous to the one in [4] in Gevrey classes and in their dual spaces of ultradistributions. As for the spaces  $\mathcal{D}_L^{(\sigma)}$  ( $\mathcal{D}_L^{(\sigma')}$ ) we consider here, we recall that they have been employed many a time in dealing with the Cauchy problem and the propagation of Gevrey singularities. For instance in [15] it is shown that some pseudo-differential and Fourier integral operators of finite order continuously map  $\mathcal{D}_L^{(\sigma)}$  ( $\mathcal{D}_L^{(\sigma')}$ ) to themselves and the same thing is true for the fundamental solution of a hyperbolic equation with constant multiplicities constructed in [14]. Specifically in [14] the hyperbolic equation is reduced to an equivalent system. Therefore at first a fundamental solution is determined for an operator of the form

$$(I) \quad P = \partial_t - i\lambda(t, x, D_x) + a(t, x, D_x)$$

where the symbol of  $\lambda$  is real,  $\lambda$  and  $a$  are continuous in  $t$  with values in some spaces of symbols of Gevrey type  $\sigma$  and of order 1 and  $p$  respectively, for some  $p \in [0, 1]$ . A fundamental solution is found which maps  $\mathcal{D}_L^{(\sigma)}$  ( $\mathcal{D}_L^{(\sigma')}$ ) to itself whenever  $\sigma < 1/p$ . The well-posedness of the Cauchy problem for an operator of the form (I) is well-known when  $\sigma < 1/p$  (see also [11]). A necessary condition for the well-

(\*) Indirizzo dell'A.: Department of Mathematics, University of Bologna (Italy).

This work is dedicated to the memory of Prof. L. Cattabriga.



results of § 1, we construct a parametrix for the Cauchy problem by solving the transport equations, as in [3] and [4]. Under our assumptions it turns out to be a Fourier integral operator of infinite order of the kind examined previously. Then a fundamental solution is determined. Finally we give some results concerning the propagation of Gevrey singularities. I wish to thank Prof. D. Mari and Prof. L. Zanghirati for some suggestions.

**0. Main notation and definitions.**

For  $\xi \in R^n$  we set  $\langle \xi \rangle = \sqrt{1 + \sum_{j=1}^n \xi_j^2}$ . For  $x = (x_1, \dots, x_n) \in R^n$  and  $\alpha = (\alpha_1, \dots, \alpha_n) \in N^n$  we write  $D_x^\alpha = D_{x_1}^{\alpha_1} \dots D_{x_n}^{\alpha_n}$ , where  $D_{x_j} = -i\partial_{x_j}$ . By  $\langle D_x \rangle^N$  we mean a pseudo-differential operator of order  $N$  whose symbol is  $\langle \xi \rangle^N$ .

We recall here the notation concerning symbols of infinite order of Gevrey type that can be found in [3].

We shall say that  $p(x, \xi) \in S_b^{\infty, \sigma, \mu}(R^n \times R_{B_0, B; A}^n)$ , where  $A, B, B_0 \geq 0, \sigma > 1, \mu > 1$ , if  $\forall \varepsilon > 0$  there exists  $C_\varepsilon > 0$  such that:

$$\sup_{x \in R^n} |\partial_\xi^\alpha D_x^\beta p(x, \xi)| \leq C_\varepsilon A^{|\alpha+\beta|} \alpha!^\mu \beta!^\sigma \langle \xi \rangle^{-|\alpha|} \exp [\varepsilon |\xi|^{1/\sigma}]$$

$\forall \alpha, \beta \in N^n, \forall \xi \in R^n, |\xi| \geq B_0 + B|\alpha|^\sigma$ . We shall write  $\tilde{S}_b^{\infty, \sigma, \mu}(R^n \times R_{B_0, B; A}^n)$  for  $S_b^{\infty, \sigma, \mu}(R^n \times R_{B_0, 0; A}^n)$ . We shall denote by  $S_b^{\infty, \sigma, \mu}$  the space

$$\lim_{\substack{A, B_0, B \rightarrow +\infty \\ \rightarrow}} S^{\infty, \sigma, \mu}(R^n \times R).$$

As for formal series of symbols we shall say that  $\sum_{j \geq 0} p_j(x, \xi)$  is in  $FS_b^{\infty, \sigma, \mu}(R^n \times R_{B_0, B; A}^n)$  if  $p_j(x, \xi) \in S_b^{\infty, \sigma, \mu}(R^n \times R_{B_0, B; A}^n)$  and  $\forall \varepsilon > 0 \exists C^\varepsilon > 0$  such that:

$$\sup_{x \in R^n} |\partial_\xi^\alpha D_x^\beta p_j(x, \xi)| \leq C_\varepsilon A^{|\alpha+\beta|} \alpha!^\mu (\beta! j!)^\sigma \langle \xi \rangle^{-|\alpha|-j} \exp [\varepsilon |\xi|^{1/\sigma}]$$

$\forall \xi \in R^n, |\xi| \geq B_0 + B(|\alpha| + j)^\sigma$ .

We shall give the following definition of equivalence of formal series of symbols. We shall write that  $\sum_{j \geq 0} p_j(x, \xi) \sim 0$ , if  $\forall \varepsilon > 0 \exists C_\varepsilon \geq 0$  and  $A, B, B_0 \geq 0$  such that:

$$\sup_{x \in R^n} \left| \partial_\xi^\alpha D_x^\beta \sum_{j < s} p_j(x, x) \right| \leq C_\varepsilon A^{\alpha+\beta+s} \alpha!^\mu (\beta! s!)^\sigma \langle \xi \rangle^{-|\alpha|-s} \exp [\varepsilon |\xi|^{1/\sigma}]$$

$\xi \in R^n, |\xi| \geq B_0 + B(|\alpha| + s)^\sigma$ .

In [15] it is shown how to construct a « true » symbol from a series of formal symbols denoted by  $\sum_{j \geq 0} p_j$ , i.e.  $p(x, \xi) \in S_0^{\infty, \sigma, \mu}$  is found in order to have  $p \sim \sum_{j \geq 0} p_j$ .

We recall also that whenever  $p \sim 0$  in  $FS_0^{\infty, \sigma, \mu}$  then  $\exists A, B_0, C > 0$  and  $h > 0$  such that:

$$\sup_{x \in \mathbb{R}^n} |D_x^\beta p(x, \xi)| \leq CA^{|\beta|} \beta!^\sigma \exp[-h|\xi|^{1/\sigma}] \quad \forall \beta \in \mathbb{N}^n, \quad \forall |\xi| \geq B_0.$$

Now we recall here a few definitions concerning the spaces which are of interest in this work. We refer to [8] and [14] for a more detailed outline. For notational convenience we shall often omit to point out the domain when  $\mathbb{R}^n$  is meant.

Let  $\mathcal{D}_{L^2, \varepsilon}^{(\sigma)} = \{f \in L^2; \exp[\varepsilon \langle \xi \rangle^{1/\sigma}] \cdot (\mathcal{F}f)(\xi) \in L^2\}$  where  $\varepsilon > 0$ ,  $\sigma \geq 1$ . (By  $\mathcal{F}f$  we denote the  $L^2$ -Fourier transform of  $f$ ).

Let  $\mathcal{D}_{L^2, \varepsilon}^{(\sigma)'}$  be the dual space of the Hilbert space  $\mathcal{D}_{L^2, \varepsilon}^{(\sigma)}$  and denote by  $\mathcal{F}$  the transpose of the operator

$$\mathcal{F}: \{v \in L^2; \exp[\varepsilon \langle \xi \rangle^{1/\sigma}] v(\xi) \in L^2\} \rightarrow \mathcal{D}_{L^2, \varepsilon}^{(\sigma)}.$$

By using the notation  $\mathcal{D}_{L^2, \varepsilon}^{(\sigma)}$  for any real number  $\varepsilon$  and by denoting both  $\mathcal{F}f$  and  ${}^t\mathcal{F}f$  by  $\tilde{f}$ , we get

$$\mathcal{D}_{L^2, -\varepsilon}^{(\sigma)} = \mathcal{D}_{L^2, \varepsilon}^{(\sigma)},$$

and we denote the norm in these spaces by

$$\|f\|_{\mathcal{D}_{L^2, \varepsilon}^{(\sigma)}} = \|\exp[\varepsilon \langle \xi \rangle^{1/\sigma}] \tilde{f}(\xi)\|_{L^2(\mathbb{R}_\xi^n)}.$$

Afterwards we define:

$$\begin{aligned} \mathcal{D}_{L^2}^{(\sigma)} &= \lim_{\varepsilon \rightarrow 0^+} \mathcal{D}_{L^2, \varepsilon}^{(\sigma)}, & \mathcal{D}_{L^2}^{(\sigma)} &= \lim_{\varepsilon \rightarrow +\infty} \mathcal{D}_{L^2, \varepsilon}^{(\sigma)}, \\ \mathcal{D}_{L^2}^{(\sigma)'} &= \lim_{\varepsilon \rightarrow 0^+} \mathcal{D}_{L^2, \varepsilon}^{(\sigma)'}, & \mathcal{D}_{L^2}^{(\sigma)'} &= \lim_{\varepsilon \rightarrow +\infty} \mathcal{D}_{L^2, \varepsilon}^{(\sigma)'}. \end{aligned}$$

The following spaces will also be used:

$$\mathcal{S}_{\sigma, \varepsilon} = \{f \in \mathcal{S}; \exp[\varepsilon \langle \xi \rangle^{1/\sigma}] \tilde{f}(\xi) \in \mathcal{S}\}, \quad \varepsilon > 0, \sigma \geq 1.$$

which are Fréchet spaces with the semi-norms:

$$|f|_{\mathcal{S}_{\sigma, \varepsilon}} = \sup_{\xi \in \mathbb{R}^n} \sup_{|\alpha|+k=l} |\langle \xi \rangle^k \partial_{\xi}^{\alpha} (\exp [\varepsilon \langle \xi \rangle^{1/\sigma}] \tilde{f}(\xi))|, \quad l = 0, 1, \dots$$

Finally we shall denote by  $\mathcal{E}_b^{(\sigma)}$  the space of all  $f \in C^{\infty}(\mathbb{R}^n)$  satisfying:

$$\sup_{x \in \mathbb{R}^n} |\partial_x^{\alpha} f(x)| \leq C_0 C_1 \langle \alpha \rangle^{\sigma} \quad \text{for some } C_0, C_1 \geq 0, \quad \sigma \geq 1.$$

**1. Fourier integral operators of infinite order on  $\mathcal{D}_L^{(\sigma)}$  ( $\mathcal{D}_L^{(\sigma)'}$ ).**

In the following pages we shall study operators with amplitude in  $S_b^{\infty, \sigma, \mu}$  and phase  $\varphi \in \mathcal{S}^{1, \sigma, \mu}(\mathbb{R}^n \times \mathbb{R}_{B_0(\varphi), 0; A(\varphi)}^n)$  satisfying:

$$(i) \quad \sum_{|\alpha+\beta| \leq 2} \sup_{\substack{x, \xi \in \mathbb{R}^n \\ |\xi| \geq B_0(\varphi)}} |\partial_{\xi}^{\alpha} D_x^{\beta}(\varphi(x, \xi) - x \cdot \xi)| |\langle \xi \rangle^{1-|\alpha|} \leq \tau \text{ for some } \tau \in [0, 1[$$

(see [9], Def. 1.2 in Cap. 10).

In this paragraph the action of the above-mentioned operators on the spaces  $\mathcal{S}_{\sigma, \varepsilon}$  and  $\mathcal{D}_L^{(\sigma)}$  is investigated. Notice that the operator defined by:

$$(1.1) \quad P_{\varphi} u(x) = \int \exp [i\varphi(x, \xi)] p(x, \xi) \tilde{u}(\xi) p \xi, \quad \forall u \in \mathcal{S}_{\sigma, \varepsilon} \text{ or } \forall u \in \mathcal{D}_L^{(\sigma)}$$

is well-defined. Moreover we have the following

**THEOREM 1.1.** If  $p(x, \xi), \varphi(x, \xi)$  are as above, then  $\forall \varepsilon > 0$  there exists  $\delta_{\varepsilon} > 0$  such that  $\forall \delta \in ]0, \delta_{\varepsilon}]$  the operator  $P_{\varphi}$  defined by (1.1) is a continuous map from  $\mathcal{S}_{\sigma, \varepsilon}$  to  $\mathcal{S}_{\sigma, \delta}$  and from  $\mathcal{D}_L^{(\sigma)}, \varepsilon$  to  $\mathcal{D}_L^{(\sigma)}, \delta$ .

Therefore  $P_{\varphi}: \mathcal{D}_L^{(\sigma)} \rightarrow \mathcal{D}_L^{(\sigma)}$  is a continuous map.

Moreover  $P_{\varphi}$  extends to a continuous operator from  $\mathcal{D}_L^{(\sigma)'}$  to  $\mathcal{D}_L^{(\sigma)'}$ .

Now let us consider  $\sigma$ -regularizing operators.

If  $r(x, \xi)$  has the property:

$$(i) \quad \sup_{x \in \mathbb{R}^n} |D_x^{\beta} r(x, \xi)| \leq C \tilde{A} \langle \beta \rangle^{\sigma} \exp [-h \langle \xi \rangle^{1/\sigma}]$$

with  $C, A, h > 0, \forall \beta \in \mathbb{N}^n, \forall \xi \in \mathbb{R}^n, |\xi|$  sufficiently large, then the operator defined by

$$R_{\varphi} u(x) = \int \exp [i\varphi(x, \xi)] r(x, \xi) \tilde{u}(\xi) p \xi, \quad \forall u \in \mathcal{D}_L^{(\sigma)}$$

extends to an operator defined on  $\mathcal{D}_L^{(\sigma)'}$ .

Furthermore it follows that

$$\sup_{x \in \mathbb{R}^n} |D_x^\beta R_\varphi u(x)| \leq \tilde{C} \tilde{A} |\beta|^\sigma \quad \text{for every } u \in \mathcal{D}_L^{\{\sigma\}'},$$

that is,  $R_\varphi$  maps  $\mathcal{D}_L^{\{\sigma\}'}$  to  $\mathcal{S}_b^{\{\sigma\}}$ .

More precisely we can prove:

**THEOREM 1.2.** If  $r(x, \xi) \in C^\infty(\mathbb{R}^n \times \mathbb{R}^n)$  has the property

$$(ii) \sup_{x \in \mathbb{R}^n} |\partial_\xi^\alpha D_x^\beta r(x, \xi)| \leq C_\alpha A^{|\beta|} |\beta|^\sigma \exp[-h \langle \xi \rangle^{1/\sigma}]$$

for  $|\xi| \geq B_0 + B|\alpha|^\sigma$  and with  $h > 0$ , then the operator  $R_\varphi$  defined on  $\mathcal{D}_L^{\{\sigma\}'}$  by

$$R_\varphi u(x) = \int \exp[i\varphi(x, \xi)] r(x, \xi) \tilde{u}(\xi) d\xi$$

is a continuous map from  $\mathcal{D}_L^{\{\sigma\}'}$  to  $\mathcal{D}_L^{\{\sigma\}}$ .

**THEOREM 1.3** (composition of a pseudo-differential and a Fourier integral operator). Let

$$p_1(x, \xi) \in S_b^{\infty, \sigma, 1}(\mathbb{R}^n \times R_{B_0^1, 0; A^1}^n), \quad p_2(x, \xi) \in S_b^{\infty, \sigma, \mu}(\mathbb{R}^n \times R_{B_0^1, B^1; A^1}^n)$$

and the phase  $\varphi(x, \xi) \in S_b^{\sigma, \mu}(\mathbb{R}^n \times R_{B_0(\varphi), 0; A(\varphi)}^n)$  with property (i).

Let

$$P_1 u(x) = \int \exp[ix \cdot \xi] p_1(x, \xi) \tilde{u}(\xi) d\xi;$$

$$P_{2\varphi} u(x) = \int \exp[i\varphi(x, \xi)] p_2(x, \xi) \tilde{u}(\xi) d\xi$$

for every  $u \in \mathcal{D}_L^{\{\sigma\}}$ .

Then there exists an operator  $Q_\varphi$  defined on  $\mathcal{D}_L^{\{\sigma\}}$  by

$$Q_\varphi u(x) = \int \exp[ib(x, \xi)] q(x, \xi) \tilde{u}(\xi) d\xi$$

and such that  $q(x, \xi) \sim \sum_{j \geq 0} q_j(x, \xi)$ , where

$$q_j(x, \xi) = \sum_{|\gamma|=j} 1/|\gamma| D_x^\gamma \left( (\partial_\xi^\alpha p_1(x, \tilde{\nabla}_x \varphi(x, y, \xi)) p_2(y, \xi) \right) \Big|_{y=x}$$

with

$$\tilde{\nabla}_x \varphi(x, y, \xi) = \int_0^1 \nabla_x \varphi(y + \theta(x - y), \xi) d\theta,$$

and there exists an operator  $R_\varphi$  continuously mapping  $\mathcal{D}_{L^2}^{[\sigma]'}$  to  $\mathcal{D}_{L^2}^{[\sigma]}$ , such that:

$$P_1(x, D_x)(P_{2,\varphi}(x, D_x)u(x)) = Q_\varphi(x, D_x)u(x) + R_\varphi(x, D_x)u(x) \quad \forall u \in \mathcal{D}_{L^2}^{[\sigma]'}$$

*Wave front sets in Gevrey classes.* When  $u \in \mathcal{D}_{L^2}^{[\sigma]'}$  we give the following definition of wave front set.

DEFINITION (see [15]). Let  $u \in \mathcal{D}_{L^2}^{[\sigma]'}$  with  $\sigma > 1$ ; let  $\sigma_1 \geq \sigma$ . We say that a point  $(x_0, \xi_0) \in T^*(R_x^n)$  does not belong to  $WF_{\{\sigma_1\}}(u)$  when there exists a symbol  $a(x, \xi)$  in  $S^{0, \sigma, \sigma}$  with  $a(x_0, \theta\xi_0) \neq 0$  ( $\theta \geq 1$ ) such that  $a(x, D_x)u(x) \in \mathcal{E}_b^{[\sigma]}(R_x^n)$ .

REMARK. In [14] Taniguchi has shown that this definition is equivalent to the one given by Hörmander (1971) in the case where  $u \in \mathcal{E}'$ .

For Fourier integral operators of the kind analyzed above we have the following result:

THEOREM 1.4. If  $p$  and  $\varphi$  satisfy the properties listed previously and moreover  $\varphi(x, \xi)$  is homogeneous in  $\xi$  of degree 1 for  $|\xi|$  sufficiently large, then for every  $\sigma_1 \geq \sigma$  we have:

$$WF_{\{\sigma_1\}}(P_\varphi u) \subset \{(x, \theta \nabla_x \varphi(x, \xi)); \\ \theta \geq 1, (\nabla_\xi \varphi(x, \xi), \xi) \in WF_{\{\sigma_1\}}(u), |\xi| \text{ sufficiently large}\}.$$

## 2. An application of Fourier integral operators of infinite order to the investigation of sufficient conditions for a certain Cauchy problem to be well-posed in $\mathcal{D}_{L^2}^{[\sigma]}(\mathcal{D}_{L^2}^{[\sigma]'})$ .

In this section we consider operators of the form:

$$P = \partial_t - i\lambda(t, x, D_x) + a(t, x, D_x)$$

where  $\lambda$  and  $a$  are pseudo-differential operators whose symbols are in  $\mathcal{C}([0, T]; C^\infty(R^{2n}))$  and satisfy the following properties:

- $\lambda(t, x, \xi)$  is real valued and belongs to  $\mathfrak{B}([0, T]; \tilde{S}^{1,1,1}(R^n \times R^n))$ ;
  - $a(t, x, \xi)$  is in  $\mathfrak{B}([0, T]; \tilde{S}^{p,1,1}(R^n \times R^n))$  with  $p \in [0, 1[$  and verifies:
- (2.i)  $\lim_{\varrho \rightarrow +\infty} \varrho^{-1/\sigma} \operatorname{Re} a(t, x, \varrho\eta) \geq 0$  for every  $t \in [0, T]$  and uniformly with respect to  $(x, \eta) \in R^n \times \mathcal{S}_{n-1}$ , where we assume  $\sigma(2p - 1) < 1$ ,  $\sigma > 1$ .

**REMARK 2.1.** Actually in dealing with the Cauchy problem with initial datum at  $t = s$ ,  $s \in [0, T[$ , it is sufficient to claim:

(2.i')  $\lim_{\rho \rightarrow +\infty} \rho^{-1/\sigma} \int_s^t \operatorname{Re} a(t', x, \nabla_x \varphi(t', s; x, \rho \eta)) dt' > 0$ ,  $\forall t \in [s, T']$ , uniformly with respect to  $(x, \eta) \in R^n \times S_{n-1}$ , where  $\varphi$  and  $T'$  are determined in the following Prop. 2.1.

**REMARK 2.2.** When  $\sigma < 1/p$  the assumption (2.i) is trivially true and it is well-known that the Cauchy problem for  $P$  is well-posed in  $\mathcal{D}_L^{\sigma'}(\mathcal{D}_L^{\sigma'})$ .

We henceforth confine our discussion to the case where  $\sigma \geq 1/p$ . We recall now the following

**PROPOSITION 2.1** (see [4], [14]). If  $\lambda$  is as above, then there exists  $T' > 0$  and there exists a solution  $\varphi(t, s)$  of the eiconal equation

$$(2.1) \quad \begin{cases} \partial_t \varphi(t, s; x, \xi) = \lambda(t, x, \nabla_x \varphi(t, s; x, \xi)) \\ \varphi(s, s; x, \xi) = x \cdot \xi \end{cases}$$

where  $\varphi(x, \xi) - x \cdot \xi \in C^1([0, T']^2; C^\infty(R^{2n})) \cap \mathcal{B}^1([0, T']^2; \tilde{S}^{1,1,1}(R^{2n})) \cap \mathcal{F}(c_0|t-s|)$  for some  $c_0 > 0$ . (For the definition of  $\mathcal{F}(\tau)$  we refer to [9].

**REMARK 2.3.** In the proof of Prop. 2.1 it is shown that if  $T'$  is sufficiently small, a solution  $(p, q)$  of

$$\begin{cases} \dot{q} = -\nabla_\xi \lambda(t, q, p), & \dot{p} = \nabla_x \lambda(t, q, p) \\ (q, p)(t = s) = (y, \eta) \end{cases}$$

can be found with the following properties:

$$q(t, s; y, \eta) - y \in \mathcal{B}^1([0, T']^2; \tilde{S}^{0,1,1}) \cap C^1([0, T']^2; C^\infty(R^{2n}))$$

$$p(t, s; y, \eta) \in \mathcal{B}^1([0, T']^2; \tilde{S}^{1,1,1}) \cap C^1([0, T']^2; C^\infty(R^{2n}))$$

$$|q(t, s; y, \eta) - y| \leq C_1 |t - s|; \quad |p(t, s; y, \eta) - \eta| \leq C_1 |t - s|$$

for some  $C_1 \geq 0$ .

By choosing a suitable  $T'$  it can also be proved that there exists the inverse function  $y = Y(t, s; x, \eta)$  of  $x = q(t, s; y, \eta)$  and

$$|Y(t, s; x, \eta) - x| \leq C_2 |t - s|$$



where

$$g(t, s; x, \xi) = - \sum_{j, k=1, \dots, n} \frac{1}{2} \partial_{\xi_k}^2 \lambda(t, x, \nabla_x \varphi(t, s; x, \xi)) \partial_{x_k}^2 \varphi(t, s; x, \xi) + \\ + a(t, x, \nabla_x \varphi(t, s; x, \xi))$$

and in the case  $h = 1, 2, \dots$

$$(T_h) \left\{ \begin{array}{l} \partial_t e_h(t, s) - \sum_{j=1}^n (\partial_{\xi_j} \lambda)(t, x, \nabla_x \varphi(t, s; x, \xi)) \partial_{x_j} e_h(t, s) + \\ + g(t, s; x, \xi) e_h(t, s) + \\ + \sum_{r=0}^{h-1} \left( -i/\gamma! D_x^r \{ (\partial_{\xi}^2 \lambda)(t, x, \tilde{\nabla}_x \varphi(t, x, z, \xi)) e_r(t, s; z, \xi) \}_{z=x} + \right. \\ \left. + \sum_{|\gamma|=\hbar-r} 1/\gamma! D_x^\gamma \{ (\partial_{\xi}^2 a)(t, x, \tilde{\nabla}_x \varphi(t, x, z, \xi)) e_r(t, s; z, \xi) \}_{z=x} \right) = 0, \\ e_h(s, s) = 0. \end{array} \right.$$

Putting  $\hat{e}_h(t, s; y, \xi) = e_h(t, s; q(t, s; y, \xi), \xi)$ ,  $h = 0, 1, \dots$  and solving the corresponding transport equations, we can prove inductively the following estimate (by applying for instance Lemma 4.2, p. 56 in [5]):

$$(I_h) \quad |\partial_{\xi}^\alpha \partial_y^\beta \hat{e}_h(t, s; y, \xi)| \leq C_\varepsilon \exp[\varepsilon \langle \xi \rangle^{1/\sigma} (t-s)] A^{*|\alpha+\beta|+2h} \cdot \\ \cdot (|\alpha+\beta|+2h)! / h! \langle \xi \rangle^{-|\alpha|-h} \langle \xi \rangle^{(\sigma-1/\sigma)(|\alpha+\beta|+2h)} \sum_{l=\min(|\alpha+\beta|, 1)}^{\alpha+\beta|+3h} \frac{\{C_0(t-s) \langle \xi \rangle^{1/\sigma}\}^l}{l!}$$

which is true for every  $\varepsilon' > 0$ , with suitable positive constants  $A^*, B_0^*, B_1^*, C_0$  and for  $|\xi| \geq B_0^* + B_1^*(|\alpha| + h)^\sigma$ .

From (I<sub>h</sub>) it follows that  $\forall \varepsilon > 0, \forall \mu' > 0$  we have:

$$|\partial_{\xi}^\alpha \partial_y^\beta \hat{e}_h(t, s; y, \xi)| \leq C'_\varepsilon \exp[\varepsilon \langle \xi \rangle^{1/\sigma}] A^{|\alpha+\beta|+h} \langle \xi \rangle^{-|\alpha|-h} (\alpha! \beta!)^{\sigma\mu'+\mu'} h!^{2\sigma\mu'-1+\mu'}$$

Since  $2\sigma\mu' - 1 < \sigma$  (by assumption (2.i)), we take  $\mu' = \sigma + 1 - 2\sigma\mu'$  and conclude with  $\sum_{h \geq 0} \hat{e}_h(t, s) \in FS^{\infty, \sigma, \sigma+1-2\sigma\mu'}$  with respect to  $t$  and  $s$ .

In view of Th. 1.1.21 in [4] it is also  $\sum_{h \geq 0} \hat{e}_h(t, s) \in FS^{\infty, \sigma, \sigma+1-2\sigma\mu'}$  uniformly with respect to  $t, s$ . Thus the theorem is proved.

**REMARK 2.4.** The operator  $R_\varphi$  in Th. 2.1 can be regarded as a pseudo-differential operator, say  $\tilde{R}$ , whose amplitude is  $\tilde{r}(x, \xi) = \exp[i\varphi(x, \xi) - ix \cdot \xi] r(x, \xi)$ , if  $r(x, \xi)$  is the amplitude of  $R$ . Moreover  $\tilde{r}(x, \xi)$  has the same properties as  $r(x, \xi)$ .

Therefore, as we proved in [1], we have the following

LEMMA 2.1. If  $\tilde{R}$  is as in Remark 2.4 then there exists a solution of

$$\tilde{R}(t, s) = -F(t, s) - \int_s^t \tilde{R}(t, \tau) F(\tau, s) d\tau$$

and  $F$  continuously maps  $\mathcal{C}([s, T']; \mathcal{D}_L^{(\sigma)'})$  to  $\mathcal{C}([s, T']; \mathcal{D}_L^{(\sigma)})$ .

THEOREM 2.2. If  $E_\varphi(t, s)$  and  $R_\varphi(t, s)$  are as in Th. 2.1 and  $F(t, s)$  is as in Lemma 2.1, then

$$\tilde{E}_\varphi(t, s) = E_\varphi(t, s) + \int_s^t \tilde{E}_\varphi(t, \tau) F(\tau, s) d\tau$$

is a fundamental solution for the Cauchy problem for  $P$ .

If  $g \in \mathcal{D}_L^{(\sigma)}$ ,  $f \in \mathcal{C}([0, T]; \mathcal{D}_L^{(\sigma)})$  (respectively  $g \in \mathcal{D}_L^{(\sigma)'}$ ,  $f \in \mathcal{C}([0, T]; \mathcal{D}_L^{(\sigma)'})$ ), then for every  $s \in [0, T[$  there exists  $T' \in ]s, T]$  such that  $\forall t \in [s, T']$

$$(2.2) \quad u(t, x) = \tilde{E}_\varphi(t, s)g + \int_s^t \tilde{E}_\varphi(t, \tau) f(\tau, \cdot) d\tau$$

is the solution of the Cauchy problem

$$(C) \quad \begin{cases} Pu(t, \cdot) = f(t, \cdot) & \text{in } [s, T'] \times R^n \\ u(s, \cdot) = g \end{cases}$$

and  $u$  belongs to  $\mathcal{C}^1$  as a function of  $t$  with values in the space  $\mathcal{D}_L^{(\sigma)}$  (respectively  $\mathcal{D}_L^{(\sigma)'}$ ).

Moreover, whenever  $\lambda(t, x, \xi)$  is homogeneous in  $\xi$  for  $|\xi|$  large, then for every  $f$  in  $\mathcal{C}([0, T]; \mathcal{D}_L^{(\sigma)})$  and for every  $g$  in  $\mathcal{D}_L^{(\sigma)'}$ , the solution  $u(t, x)$  of the problem (C) (with  $s \in [0, T[$ ,  $T'$  sufficiently small and  $t \in [s, T']$ ) satisfies

$$(2.2)' \quad WF_{\{\sigma\}}(u(t, \cdot)) \subset \{(q(t, s; y, \eta), \theta p(t, s; y, \eta)); (y, \eta) \in WF_{\{\sigma\}}(g); \\ \theta \geq 1 \text{ and } |\eta| \text{ sufficiently large}\}, \quad \forall t \in [s, T'],$$

where  $q, p$  are solutions to

$$\begin{cases} \dot{q} = -\nabla_\xi \lambda(t, q, p); & \dot{p} = \nabla_x \lambda(t, q, p), \\ (q, p)(t = s) = (y, \eta). \end{cases}$$

PROOF. In view of Th. 2.1 and Lemma 2.1 we have  $P\tilde{E}_\varphi(t, s) = 0$  and  $u(t, x)$  defined in (2.2) satisfies all the claims above. Uniqueness

of the solution follows by a standard argument where the transpose of  $P$  is considered (see [4]).

Finally (2.2)' follows from Th. 1.4 and Th. 2.2.

## REFERENCES

- [1] R. AGLIARDI, *Pseudo-differential operators of infinite order on  $\mathcal{D}'_L(g)$  ( $\mathcal{D}'_L(g')$ ), and applications to the Cauchy problem for some elementary operators*, to appear on Ann. di Mat. Pura e Appl.
- [2] L. CATTABRIGA, *Some remarks on the well-posedness of the Cauchy problem in Gevrey spaces*, in Partial Differential Equations and the Calculus of Variations: *Essays in honour of Ennio De Giorgi*.
- [3] L. CATTABRIGA - D. MARI, *Parametrix of infinite order on Gevrey spaces to the Cauchy problem for hyperbolic operators with one constant multiple characteristics*, Ricerche di Mat., Suppl., **36** (1987), pp. 127-147.
- [4] L. CATTABRIGA - L. ZANGHIRATI, *Fourier integral operators of infinite order on Gevrey spaces-Applications to the Cauchy problem for certain hyperbolic operators*, to appear on Journal of Math. of Kyoto Univ.
- [5] P. HARTMAN, *Ordinary Differential Equations*, John Wiley, 1964.
- [6] S. HASHIMOTO - T. MATSUZAWA - Y. MORIMOTO, *Operateurs pseudo-différentiels et classes de Gevrey*, C.P.D.E., (1983), pp. 1277-1289.
- [7] K. KAJITANI, *Fundamental solution of Cauchy problem for hyperbolic systems and Gevrey classes*, Tsukuba J. Math., **1** (1977), pp. 163-193.
- [8] K. KAJITANI - S. WAKABAYASHI, *Microhyperbolic operators in Gevrey classes*, Publ. RIMS (to appear).
- [9] H. KUMANO-GO, *Pseudo-differential Operators*, M.I.T. Press, 1981.
- [10] S. MISOHATA, *On the Cauchy problem for hyperbolic equations and related problems-micro-local energy methods*, Proc. Taniguchi Intern. Sympos. on Hyperbolic Equations and Related Topics, Kataka, 1984, pp. 193-233.
- [11] S. MIZOHATA, *On the Cauchy Problem*, Science Press, Beijing, 1985.
- [12] Y. MORIMOTO - K. TANIGUCHI, *Propagation of wave front sets of solutions of the Cauchy problem for hyperbolic equations in Gevrey classes*, Osaka J. of Math., **23** (1986), pp. 765-814.
- [13] L. RODINO - L. ZANGHIRATI, *Pseudo-differential operators with multiple characteristics and Gevrey singularities*, C.P.D.E., **11** (1986), pp. 673-711.
- [14] K. TANIGUCHI, *Fourier integral operators in Gevrey classes on  $R^n$  and the fundamental solution for a hyperbolic operator*, Publ. R.I.M.S., **20** (1984), pp. 491-542.
- [15] K. TANIGUCHI, *Pseudo-differential operators acting on ultradistributions*, Math. Japonica, **30** (1985), pp. 719-741.
- [16] L. ZANGHIRATI, *Pseudo-differential operators of infinite order and Gevrey classes*, Ann. Univ. Ferrara - sez. VII - Sc. Mat. (1985), pp. 197-219.

Manoscritto pervenuto in redazione il 29 marzo 1989 e, in forma revisionata, il 19 settembre 1989.