

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

PATRIZIA DONATO

GIOCONDA MOSCARIELLO

**On the homogenization of some nonlinear
problems in perforated domains**

Rendiconti del Seminario Matematico della Università di Padova,
tome 84 (1990), p. 91-108

http://www.numdam.org/item?id=RSMUP_1990__84__91_0

© Rendiconti del Seminario Matematico della Università di Padova, 1990, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

On the Homogenization of Some Nonlinear Problems in Perforated Domains.

PATRIZIA DONATO - GIOCONDA MOSCARIELLO (*)

0. - Introduction.

In this paper we study the homogenization of a class of nonlinear elliptic Neumann problems in perforated domains of \mathbb{R}^n .

Let Ω_ε be a fixed bounded domain Ω from which a set T_ε of holes has been removed. The set T_ε is obtained in the following way: let T a fixed set properly contained in the basic cell Y , let D_ε be the hole homotetic by ratio ε to T . Let us suppose to have a periodic distribution of period εY of D_ε . Then T_ε is the set of the holes of this periodic distribution contained in Ω_ε .

Roughly speaking, let us consider the problem:

$$(\mathfrak{P}_\varepsilon) \quad \begin{cases} -\operatorname{div} a(x/\varepsilon, Du) = f & \text{in } \Omega_\varepsilon, \\ u = 0 & \text{on } \partial\Omega, \\ a(x/\varepsilon, Du) \cdot \nu = \varepsilon g(x, x/\varepsilon) & \text{on } \partial T_\varepsilon, \\ u \in H^{1,p}(\Omega_\varepsilon) & p > 1, \end{cases}$$

where ν denotes the exterior normal with respect to Ω_ε , $f \in L^p(\Omega)$, g is Y -periodic in the second variable and $a(x, \xi)$ is a matrix periodic in x and satisfying suitable coerciveness and growth conditions in ξ .

(*) Indirizzo degli AA.: Dipartimento di Matematica e Applicazioni, Università di Napoli, via Mezzocannone 8, 80134 Napoli.

This work has been performed as part of a National Research Project supported by M.P.I. (40%, 1987).

Our aim is to study the asymptotic behaviour of the solutions u_ε of (P_ε) as $\varepsilon \rightarrow 0$. Indeed, we prove that the « limit » problem of $(\mathcal{P}_\varepsilon)$ is:

$$(\mathcal{P}_0) \quad \begin{cases} -\operatorname{div} b(Du) = f\theta + \mu_\sigma(x) & \text{in } \Omega, \\ u \in H_0^{1,p}(\Omega) & p > 1, \end{cases}$$

with $\theta = |Y \setminus T|/|Y|$ and

$$\mu_\sigma(x) = (1/|Y|) \int_{\partial T} g(x, y) d\sigma_y.$$

Moreover the matrix $b(\xi)$ is given by an explicit formula.

As a consequence, we are able in particular to describe the « homogenized » operator of the problem:

$$\begin{aligned} -\operatorname{div} (|Du|^{p-2} Du) &= f && \text{in } \Omega_\varepsilon, \\ u &= 0 && \text{on } \partial\Omega, \\ (|Du|^{p-2} Du) \cdot \nu &= g(x, x/\varepsilon) && \text{on } \partial T_\varepsilon, \\ u &\in H^{1,p}(\Omega_\varepsilon) && p > 1. \end{aligned}$$

The homogenization theory for linear elliptic operators goes back to De Giorgi-Spagnolo [9], Bensoussan-Lions-Papanicolau [2], Sanchez-Palencia [15].

For Dirichlet nonlinear problems of the type

$$-\operatorname{div} a(x, u, Du) = f$$

some homogenization results for $p = 2$ were first given by Tartar [19] (see also Suquet [16]). By different techniques, for $p > 1$, homogenization results have been recently given in [11].

On the other hand the homogenization of some linear problems in perforated domains has been studied in [6] and in Cioranescu-Saint Jean Paulin [7] by using energy method.

1. - Statement of the problem.

Let Ω be an open set in \mathbb{R}^n , $n \geq 2$, with a smooth boundary $\partial\Omega$. Introduce the representative cell $Y = [0, l_1) \times \dots \times [0, l_n)$ and de-

note by T an open subset of Y , with smooth boundary ∂T , such that $\overline{T} \subset Y$. Denote by $\tau(\varepsilon\overline{T})$ the set of all translated images $\varepsilon(k_1 + \overline{T})$, $k \in \mathbb{Z}^n$, $k_1 = (k_1 l_1, \dots, k_n l_n)$, $\varepsilon > 0$, of $\varepsilon\overline{T}$.

By this way \mathbb{R}^n is periodically perforated by holes of ε -size. We make the following assumption:

The holes $\tau(\varepsilon\overline{T})$ do not intersect the boundary $\partial\Omega$.

This assumption restricts the geometry of the open set Ω (example: Ω is a finite union of rectangles homothetic to the representative cell). Define now the perforated domain:

$$\Omega_\varepsilon = \Omega \cap \{\mathbb{R}^n \setminus \tau(\varepsilon\overline{T})\}.$$

Hence, by the previous assumption it follows that:

$$\partial\Omega_\varepsilon = \partial\Omega \cap \partial T_\varepsilon$$

where T_ε is the subset of $\tau(\varepsilon\overline{T})$ contained in Ω .

The following notations are used in the following:

- 1) $Y^* = Y \setminus \overline{T}$;
- 2) $\theta = |Y^*|/|Y|$;
- 3) $|\omega|$ = the Lebesgue measure of ω (for any measurable set of \mathbb{R}^n);
- 4) $\chi(\omega)$ = the characteristic function of the set ω ;
- 5) \tilde{v} = the zero extension to the whole Ω , for any function v defined on Ω_ε ;
- 6) $\langle f \rangle_E = (1/|E|) \int_E f(x) dx$, for $f \in L^1_{n,loc}(\mathbb{R}^n)$ and $E \subset \mathbb{R}^n$ bounded open set of positive measure.

If $1 < p < +\infty$ and $p' = p/(p-1)$, we shall consider the following spaces:

$$\begin{aligned} H_{\text{per}}^{1,p}(Y) &= \\ &= \{ u(y) \in H^{1,p}(Y) : u \text{ has the same trace on the opposite faces of } Y \}, \\ L_{n, \text{per}}^{p'}(Y) &= \{ q(y) \in L_{n, \text{per}}^{p'}(Y) : \int_Y q \cdot Du dy = 0 \text{ for any } u \in H_{\text{per}}^{1,p}(Y) \}. \end{aligned}$$

Now, let $f \in L^{p'}(\Omega)$ and $g: \Omega \times Y \rightarrow \mathbf{R}$ verifying the following assumptions:

$$(1.1)_1 \quad g(x, \cdot) \text{ is } Y\text{-periodic and measurable for any } x \in \Omega,$$

$$(1.1)_2 \quad g(x, \cdot) \in H^{1-1/p'}(\partial T),$$

$$(1.1)_3 \quad |g(x_1, y) - g(x_2, y)| < \\ < c(1 + |\varphi(y)|) \omega(|x_1 - x_2|) \text{ for any } y \in Y \text{ and } x_1, x_2 \in \Omega,$$

where $\omega(t): [0, +\infty) \rightarrow [0, +\infty)$ is a bounded, concave and continuous function such that $\omega(0) = 0$ and $\varphi(y) \in L^{p'}(\partial T)$.

We shall consider the problem:

$$(\mathcal{P}_\varepsilon) \quad \begin{cases} -\operatorname{div} a(x/\varepsilon, Du) = f & \text{in } \Omega_\varepsilon, \\ u = 0 & \text{on } \partial\Omega, \\ a(x/\varepsilon, Du) \cdot \nu = \varepsilon g_\varepsilon(x) & \text{on } \partial T_\varepsilon, \end{cases}$$

where ν denotes the exterior normal with respect to Ω_ε , $g_\varepsilon(x) = g(x, x/\varepsilon)$ and $a(y, \xi)$ verifies the following structure conditions:

H_1) a is Y -periodic and measurable with respect to y ,

H_2) for any y a.e. in \mathbf{R}^n and any $\xi_1, \xi_2 \in \mathbf{R}^n$ then

if $p > 2$:

$$\text{i) } |a(y, \xi_1) - a(y, \xi_2)| < \beta(1 + |\xi_1| + |\xi_2|)^{p-2} |\xi_1 - \xi_2|$$

$$\text{ii) } (a(y, \xi_1) - a(y, \xi_2), \xi_1 - \xi_2) \geq \alpha |\xi_1 - \xi_2|^p, \quad \alpha > 0$$

or, if $1 < p \leq 2$:

$$\text{j) } |a(y, \xi_1) - a(y, \xi_2)| < \beta |\xi_1 - \xi_2|^{p-1},$$

$$\text{jj) } (a(y, \xi_1) - a(y, \xi_2), \xi_1 - \xi_2) \geq \alpha |\xi_1 - \xi_2|^2 (|\xi_1| + |\xi_2|)^{p-2}, \quad \alpha > 0,$$

H_3) $a(y, 0) \in L_n^{p'}(Y)$.

Set

$$V_\varepsilon^p = \{\varphi \in H^{1,p}(\Omega_\varepsilon): \varphi = 0 \text{ on } \partial\Omega\},$$

a variational solution of problem $(\mathcal{F}_\varepsilon)$ is a function $u_\varepsilon \in V_\varepsilon^p$ such that:

$$(1.2) \quad \int_{\Omega_\varepsilon} a(x/\varepsilon, Du_\varepsilon) \cdot D\varphi \, dx = \int_{\Omega_\varepsilon} f\varphi \, dx + \varepsilon \int_{\partial T_\varepsilon} g_\varepsilon \varphi \, d\sigma$$

for any $\varphi \in V_\varepsilon^p$.

REMARK 1.1. It is well known that, under the above hypotheses, the problem $(\mathcal{F}_\varepsilon)$ has a unique solution $u_\varepsilon \in V_\varepsilon^p$. ■

Finally, let us consider the problem:

$$(1.3) \quad \begin{cases} \int_{Y^*} a(y, Dv(y)) \cdot D\varphi(y) \, dy = 0 & \forall \varphi \in H_{\text{per}}^{1,p}(Y^*), \\ v \in \xi \cdot y + H_{\text{per}}^{1,p}(Y^*), \end{cases}$$

where $\xi \in \mathbb{R}^n$ and

$$\begin{aligned} H_{\text{per}}^{1,p}(Y^*) &= \\ &= \{u(y) \in H^{1,p}(Y^*): u \text{ has the same trace on the opposite faces of } Y\}. \end{aligned}$$

In the § 3 we will prove the convergence of the solutions $u_\varepsilon \in V_\varepsilon^p$ of $(\mathcal{F}_\varepsilon)$ to the solution of the « homogenized » problem:

$$(1.3_0) \quad \begin{cases} -\operatorname{div} b(Du) = f\theta + \mu_\sigma(x) & \text{in } \Omega, \\ u \in H_0^{1,p}(\Omega), \end{cases}$$

where

$$(1.4) \quad \mu_\sigma(x) = (1/|Y|) \int_{\partial T} g(x, y) \, d\sigma_y$$

and, for any $\xi \in \mathbb{R}^n$, if $v(y)$ is the solution of (1.3), b is defined by

$$(1.5) \quad b(\xi) = (1/|Y|) \int_{Y^*} a(y, Dv) \, dy.$$

2. – Preliminary results.

We recall some lemmas about the spaces introduced in § 1 and the existence of a family of extension-operators.

LEMMA 2.1. *If $q(y)$ is an element of $L'_{n, \text{per}}(Y)$, then it can be extended by periodicity to an element of $L'_{n, \text{loc}}(\mathbb{R}^n)$, still denoted by q , such that*

$$\operatorname{div}_x q = 0. \quad \blacksquare$$

LEMMA 2.2. *Let f be an Y -periodic function of $L^p_{\text{loc}}(\mathbb{R}^n)$, $1 \leq p \leq +\infty$ and set*

$$f_\varepsilon(x) = f(x/\varepsilon) \quad \text{for any } x \text{ a.e. in } \mathbb{R}^n,$$

then as $\varepsilon \rightarrow 0$:

$$\begin{aligned} f_\varepsilon &\rightarrow \langle f \rangle && \text{in } w\text{-}L^p_{\text{loc}}(\mathbb{R}^n) \text{ if } p < +\infty \\ f_\varepsilon &\rightarrow \langle f \rangle && \text{in } w^*\text{-}L^\infty(\mathbb{R}^n) \text{ if } p = +\infty. \quad \blacksquare \end{aligned}$$

For a proof of the previous lemmas one may see [16] annexe 2.

LEMMA 2.3 (see [7]). *There exists a linear continuous extension-operator $P \in \mathcal{L}(H^{1,p}(Y^*), H^{1,p}(Y))$ such that:*

$$(2.1) \quad \|D(P\varphi)\|_{L^p_*(\mathbb{R}^*)} \leq \|D\varphi\|_{L^p_*(\mathbb{R}^*)}$$

for any $\varphi \in H^{1,p}(Y^*)$. \blacksquare

LEMMA 2.4 (see [7]). *There exists a family P_ε of linear continuous extension-operators, $P_\varepsilon \in \mathcal{L}(V_\varepsilon^p, H^{1,p}_0(\Omega))$ verifying the following condition:*

$$(2.2) \quad \|D(P_\varepsilon\varphi)\|_{L^p_*(\Omega)} \leq \|D\varphi\|_{L^p_*(\Omega_\varepsilon)}$$

for any $\varphi \in V_\varepsilon^p$, where c is a constant independent of ε . \blacksquare

The previous lemmas are proved in the case $p = 2$ in [7]. The same argument can be used in the general case.

If $\gamma \in W^{1-1/p', p'}(\partial T)$ let us consider the linear form on $H^{1,p}(\Omega)$, $1/p + 1/p' = 1$:

$$\langle \mu_\gamma^\varepsilon, \varphi \rangle = \varepsilon \int_{\partial T_\varepsilon} \gamma(x/\varepsilon) \varphi \, d\sigma$$

and

$$\mu_\gamma = (1/|Y|) \int_{\partial T} \gamma(x) \, d\sigma_\nu.$$

PROPOSITION 2.5 (see [6], [12]). *If $\gamma \in W^{1-1/p', p'}(\partial T)$, $1 < p' < +\infty$, there exists a unique solution $\psi_\gamma \in W^{2,p'}(Y^*)$ of the problem:*

$$(2.3) \quad \begin{cases} -\Delta \psi_\gamma = -(|Y|/|Y^*|) \mu_\gamma & \text{in } Y^*, \\ \partial \psi_\gamma / \partial \nu = \gamma & \text{on } \partial T, \\ \psi_\gamma \text{ } Y\text{-periodic,} \\ \langle \psi_\gamma \rangle_{Y^*} = 0. \end{cases}$$

In particular if γ is a constant function, then $\psi_\gamma \in W^{1,\infty}(Y^)$. ■*

REMARK 2.6. It is easy to verify that the solution of problem (2.3) can be extended by periodicity to $\mathbb{R}^n \setminus \tau(\bar{T})$ and the function

$$\psi_\gamma^\varepsilon(x) = \psi_\gamma^\varepsilon(x/\varepsilon) \quad \text{a.e. } x \in \Omega_\varepsilon$$

verify:

$$(2.4) \quad \begin{cases} -\Delta \psi_\gamma^\varepsilon = -\varepsilon^{-2}(|Y|/|Y^*|) \mu_\gamma & \text{in } \mathbb{R}^n \setminus \tau(\varepsilon \bar{T}), \\ \partial \psi_\gamma^\varepsilon / \partial \nu = \varepsilon^{-1} \gamma(x/\varepsilon) & \text{on } \partial \tau(\varepsilon \bar{T}). \quad \blacksquare \end{cases}$$

Now, we can prove the following lemma that we'll use in the sequel:

LEMMA 2.7. *Let Q be an interval of \mathbb{R}^n , $S_\varepsilon = \tau(\varepsilon \bar{T}) \cap Q$ and $Q_\varepsilon = Q \setminus \bar{S}_\varepsilon$. If $\bar{S}_\varepsilon \cap \partial Q = \emptyset$ then:*

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \int_{\partial S_\varepsilon} \gamma(x/\varepsilon) \varphi_\varepsilon(x) \, d\sigma = \mu_\gamma \int_Q \varphi(x) \, dx$$

for any sequence $\{\varphi_\varepsilon\}$ of $H^{1,p}(\Omega)$ such that $\varphi_\varepsilon \rightarrow \varphi$ in $w\text{-}H^{1,p}(\Omega)$, $1 < p < +\infty$.

PROOF. Let ψ_γ be the solution of (2.3), then by remark 2.6 we have:

$$(2.5) \quad \varepsilon \int_{\partial S_\varepsilon} \gamma(x/\varepsilon) \varphi_\varepsilon(x) d\sigma = \\ = \varepsilon^2 \int_{Q_\varepsilon} D_x \psi_\gamma^\varepsilon \cdot D_x \varphi_\varepsilon dx + (|Y|/|Y^*|) \mu_\gamma \int_{Q_\varepsilon} \varphi_\varepsilon dx - \varepsilon^2 \int_{\partial Q} D_x \psi \varphi_\gamma^\varepsilon \cdot \nu_i d\sigma.$$

Then by lemma 2.2 and by observing that $D_x \psi_\gamma^\varepsilon(x) = (1/\varepsilon) D_\nu \psi_\gamma(x/\varepsilon)$, passing to the limit as $\varepsilon \rightarrow 0$ we get the result. ■

REMARK 2.8. If h is a constant function, by using similar arguments as in the previous lemma, it can be found that

$$\varepsilon \int_{\partial S_\varepsilon} |\varphi_\varepsilon| d\sigma \leq c$$

(with c independent of ε) for any sequence $\{\varphi_\varepsilon\}$ bounded in $H^{1,1}(Q)$. ■

Let $g: \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}$ verifying (1.1)₁ ÷ (1.1)₃ and

$$\langle \mu_\sigma^\varepsilon, \varphi \rangle = \varepsilon \int_{\partial T_\varepsilon} g(x, x/\varepsilon) \varphi d\sigma \quad \forall \varphi \in H_0^{1,p}(\Omega), \quad 1 < p < +\infty.$$

LEMMA 2.9. *The measures $\mu_\sigma^\varepsilon, \mu_\sigma$ are in $H^{-1,p'}(\Omega)$ and*

$$\mu_\sigma^\varepsilon \rightarrow \mu_\sigma \quad \text{in } s\text{-}H^{-1,p'}(\Omega),$$

where μ_σ is given by (1.4).

PROOF. It is enough to prove that

$$(2.6) \quad \langle \mu_\sigma^\varepsilon, \varphi_\varepsilon \rangle \rightarrow \langle \mu_\sigma, \varphi \rangle$$

if $\varphi_\varepsilon, \varphi \in H_0^{1,p}(\Omega)$ and $\varphi_\varepsilon \rightarrow \varphi$ in $w\text{-}H_0^{1,p}(\Omega)$.

Let us consider $\forall \nu \in \mathbb{N}$ a partition of \mathbb{R}^n by intervals $Q_{i\nu}$ of side $2^{-\nu} l_i$. Since the holes T_ε do not intersect $\partial\Omega$, we can assume that $\forall \varepsilon > 0, T_\varepsilon$ does not intersect $\partial Q_{i\nu}$.

Let us denote by $x_{i\nu}$ and $\chi_{i\nu}$ respectively the center and the characteristic function of $Q_{i\nu}$. Set

$$g_\nu(x, y) = \sum_i \chi_{i\nu}(x) g(x_{i\nu}, y).$$

We have

$$\varepsilon \int_{\partial T_\varepsilon} [g(x, x/\varepsilon) - g_\nu(x, x/\varepsilon)] \varphi_\varepsilon d\sigma = \varepsilon \int_{\partial T_\varepsilon} [g(x, x/\varepsilon) - g_\nu(x, x/\varepsilon)] \varphi_\varepsilon d\sigma + \varepsilon \int_{\partial T_\varepsilon} g_\nu(x, x/\varepsilon) \varphi_\varepsilon d\sigma.$$

We observe that by (1.1)₃ $\mu_\nu \in C^0(\bar{\Omega})$, then by lemma 2.7 we have:

$$\begin{aligned} (2.7) \quad \lim_{\nu} \lim_{\varepsilon \rightarrow 0} \varepsilon \int_{\partial T_\varepsilon} g_\nu(x, x/\varepsilon) \varphi_\varepsilon(x) d\sigma &= \lim_{\nu} \lim_{\varepsilon \rightarrow 0} \sum_i \varepsilon \int_{Q_{i\nu} \cap \partial T_\varepsilon} g(x_{i\nu}, x/\varepsilon) \varphi_\varepsilon(x) d\sigma = \\ &= \lim_{\nu} \sum_i \int_{Q_{i\nu} \cap \Omega} \mu_\nu(x_{i\nu}) \varphi(x) dx = \lim_{\nu} \int_{\Omega} \chi_{i\nu} \mu_\nu(x_{i\nu}) \varphi(x) dx = \\ &= \int_{\Omega} \mu_\nu(x) \varphi(x) dx. \end{aligned}$$

On the other hand we have:

$$\begin{aligned} (2.8) \quad \varepsilon \int_{\partial T_\varepsilon} [g(x, x/\varepsilon) - g_\nu(x, x/\varepsilon)] \varphi_\varepsilon(x) d\sigma &\leq \\ &\leq \varepsilon^{1/p'} \left[\sum_i \int_{Q_{i\nu} \cap \partial T_\varepsilon} |g(x, x/\varepsilon) - g(x_{i\nu}, x/\varepsilon)|^{p'} d\sigma \right]^{1/p'} \varepsilon^{1/p} \left[\sum_i \int_{Q_{i\nu} \cap \partial T_\varepsilon} |\varphi_\varepsilon(x)|^p d\sigma \right]^{1/p}. \end{aligned}$$

By remark 2.8 we get

$$(2.9) \quad \varepsilon^{1/p} \left[\sum_i \int_{Q_{i\nu} \cap \partial T_\varepsilon} |\varphi_\varepsilon(x)|^p d\sigma \right]^{1/p} \leq c.$$

Then by (2.8), (2.9) and (1.1)₁ ÷ (1.1)₃:

$$\begin{aligned} \varepsilon \int_{\partial T_\varepsilon} [g(x, x/\varepsilon) - g_\nu(x, x/\varepsilon)] \varphi_\varepsilon(x) d\sigma &\leq c\omega(1/2^\nu) \left[\varepsilon \int_{\partial T_\varepsilon} (1 + |\varphi(x/\varepsilon)|^{p'}) d\sigma \right]^{1/p'} \leq \\ &\leq c'\omega(1/2^\nu) \left[\varepsilon \int_{\partial T} (1 + |\varphi(y)|^{p'}) d\sigma \right]^{1/p'}. \end{aligned}$$

Then passing to the limit as $\varepsilon \rightarrow 0^+$ and $\nu \rightarrow +\infty$ we obtain (2.6). The proof is completed by using standard arguments concerning the duality application between $H_0^{1,p}(\Omega)$ and its dual. ■

We state now some lemmas about the structure properties of b . Using the same arguments of [11] one can prove:

LEMMA 2.10. For any $\xi \in \mathbb{R}^n$

$$|b(\xi)| \leq c(1 + |\xi|)^{p-1}$$

where $c = c(\alpha, \beta, p|Y|, \|a(y, 0)\|_{L_a^{2p'}(x)})$.

Further if v is the solution of problem (1.3) we have:

$$(2.10) \quad \int_{Y^*} |Dv(y)|^p dy \leq c(1 + |\xi|)^p$$

where $c = c(\alpha, \beta, p, |Y|, \|a(y, 0)\|_{L_a^{2p'}(x)})$. ■

LEMMA 2.1. $b(\xi)$ is locally Holder (Lipschitz if $p = 2$). ■

REMARK 2.12. We remark that the limit operator $b(\xi)$, as in the homogenization of Dirichlet problem (see [11]) may not verify the same structure conditions of $a(y, \xi)$.

In some special case, the Holder estimate on $b(\xi)$ can be improved (see [11]). ■

LEMMA 2.13. For $\xi_1, \xi_2 \in \mathbb{R}^n$ we have

$$(2.11) \quad (b(\xi_1) - b(\xi_2), \xi_1 - \xi_2) \geq \alpha |\xi_1 - \xi_2|^p, \quad \text{if } p > 2$$

$$(2.12) \quad (b(\xi_1) - b(\xi_2), \xi_1 - \xi_2) \geq \alpha' |\xi_1 - \xi_2|^2 (1 + |\xi_1| + |\xi_2|)^{p-2},$$

$$\alpha' > 0, \quad \text{if } 1 < p \leq 2.$$

PROOF. Let us denote by v_1 and v_2 the solutions of problem (1.3) defining respectively $b(\xi_1)$ and $b(\xi_2)$.

Let us consider $u_i = v_i - \xi \cdot y$, $i = 1, 2$. Then $u_i(y)$ is an element of $H_{\text{per}}^{1,p}(Y^*)$ and so, by lemma 2.3, we can consider $Pu_i \in H_{\text{per}}^{1,p}(Y)$.

If we extend Pu_i by periodicity, the resulting function (still denoted by Pu_i) is in $H_{\text{loc}}^{1,p}(\mathbb{R}^n)$.

Define

$$w_i^\varepsilon = \varepsilon Pu_i(x/\varepsilon) + \xi_i \cdot x, \quad i = 1, 2,$$

and for $\xi \in \mathbb{R}^n$

$$(2.13) \quad \tilde{a}(y, \xi) = \begin{cases} a(y, \xi) & \text{for } y \in Y^*, \\ 0 & \text{for } y \in \bar{T}. \end{cases}$$

It is easy to verify that

$$(2.14) \quad \begin{cases} w_i^\varepsilon \rightarrow \xi_i \cdot x & \text{in } w-H_{\text{loc}}^{1,p}(\mathbb{R}^n), \\ \tilde{a}(x/\varepsilon, Dw_i^\varepsilon) \rightarrow b(\xi_i) & \text{in } w-L_{n,\text{loc}}^{p'}(\mathbb{R}^n), \\ \operatorname{div} \tilde{a}(x/\varepsilon, Dw_i^\varepsilon) = 0, \end{cases}$$

where the last relation is proved by using lemma 2.1.

If $p \geq 2$ from ii) of H_2 , we get:

$$\alpha \int_{Y^*} \eta |Dw_1^\varepsilon - Dw_2^\varepsilon|^p dx \leq \int_Y \eta (\tilde{a}(x/\varepsilon, Dw_1^\varepsilon) - \tilde{a}(x/\varepsilon, Dw_2^\varepsilon), Dw_1^\varepsilon - Dw_2^\varepsilon) dx$$

where $\eta \in C_0^1(Y^*)$.

Then, passing to the limit as $\varepsilon \rightarrow 0$ and using (2.14), by the compensated compactness result of [12] we get

$$\alpha \int_{Y^*} \eta |\xi_1 - \xi_2|^p \leq \int_{Y^*} \eta (b(\xi_1) - b(\xi_2), \xi_1 - \xi_2) dx.$$

Then from the arbitrariness of η , we deduce (2.11).

If $1 < p \leq 2$, from jj) of H_2 we have:

$$\begin{aligned} \sqrt{\alpha} \int_{Y^*} |Dw_1^\varepsilon - Dw_2^\varepsilon| dx &\leq \left(\int_Y \eta (\tilde{a}(x/\varepsilon, Dw_1^\varepsilon) - \tilde{a}(x/\varepsilon, Dw_2^\varepsilon), Dw_1^\varepsilon - Dw_2^\varepsilon) dx \right)^{\frac{1}{2}} \\ &\quad \cdot \left(\int_Y \eta (|Dw_1^\varepsilon| + |Dw_2^\varepsilon|)^{2-p} dx \right)^{\frac{1}{2}}. \end{aligned}$$

Then, passing to the limit as before, and remarking that by lemma 2.3

$$\int_Y |Dw_i^\varepsilon|^p dx \leq c \int_{Y^*} |Dv_i^\varepsilon|^p dy,$$

we can argue in a similar way to prove (2.12). ■

3. - Homogenization results.

THEOREM 3.1. *If $a(x, \xi)$ verifies the structure conditions $H_1) \div H_3)$ and $g: \Omega \times Y \rightarrow \mathbb{R}$ satisfies (1.1)₁ \div (1.1)₃, then for any $f \in L^{p'}(\Omega)$ the sequence u_ε of the solutions of problem $(\mathcal{P}_\varepsilon)$ verifies:*

$$\begin{aligned} P_\varepsilon u_\varepsilon &\rightarrow u_0 && \text{in } w\text{-}H_0^{1,p}(\Omega), \\ \tilde{a}(x/\varepsilon, Du_\varepsilon) &\rightarrow b(Du_0) && \text{in } w\text{-}L_n^{p'}(\Omega), \end{aligned}$$

where u_0 is the solution of problem (\mathcal{P}_0) .

PROOF. We use the homogenization techniques of nonlinear operators introduced in [10], [11].

Let us denote by $P_\varepsilon u_\varepsilon$ the extension of u_ε given by lemma 2.4. By using lemma 2.9, it is very easy to verify that

$$\|P_\varepsilon u_\varepsilon\|_{H_0^{1,p}(\Omega)} \leq c$$

with c independent of ε .

Then by i) or j) of $H_2)$, we get also:

$$\|\tilde{a}(x/\varepsilon, Du_\varepsilon)\|_{L_n^{p'}(\Omega)} \leq c$$

with c independent of ε .

Hence, up to a subsequence, we have

$$\begin{aligned} P_\varepsilon u_\varepsilon &\rightarrow u_0 && \text{in } w\text{-}H_0^{1,p}(\Omega), \\ \tilde{a}(x/\varepsilon, Du_\varepsilon) &\rightarrow a_0(x) && \text{in } w\text{-}L_n^{p'}(\Omega). \end{aligned}$$

The theorem will be proved if we show that:

$$(3.1) \quad a_0(x) = b(Du_0) \quad \text{a.e. in } \Omega .$$

Indeed, using lemma 2.9 and the fact that

$$(3.2) \quad \chi_{\Omega_\nu} \rightarrow \theta \quad \text{in } w^*-L^\infty(\Omega)$$

we obtain:

$$\int_{\Omega} a_0(x) D\varphi \, dx = \int_{\Omega} \theta f \varphi \, dx + \int_{\Omega} \mu_\nu \varphi \, dx \quad \forall \varphi \in H_0^{1,p}(\Omega) .$$

Let us fix $\nu \in \mathbb{N}$ and denote by $\{Q_{i\nu}\}_i$ a partition of \mathbb{R}^n as in the proof of lemma 2.7. Then we define $I_\nu = \{i: Q_{i\nu} \subset \Omega\}$, $\Omega_\nu = \bigcup_{i \in I_\nu} Q_{i\nu}$. For any i let us consider $\langle Du_0 \rangle_{i\nu} = \langle Du_0 \rangle_{Q_{i\nu}}$. Then if $\chi_{i\nu}$ is the characteristic function of $Q_{i\nu}$, by the continuity of b (see lemma 2.11) we have, if $\nu \rightarrow +\infty$, that:

$$(3.3) \quad \sum_{i \in I_\nu} \chi_{i\nu}(x) b(\langle Du_0 \rangle_{i\nu}) \rightarrow b(Du_0(x)) \quad \text{a.e. in } \Omega .$$

Moreover, from lemma 2.10 we have for any measurable set $E \subset \Omega$:

$$\int_E \left| \sum_{i \in I_\nu} \chi_{i\nu}(x) b(\langle Du_0 \rangle_{i\nu}) \right|^{p'} dx \leq c \int_E \left(1 + \left| \sum_{i \in I_\nu} \chi_{i\nu}(x) \langle Du_0 \rangle_{i\nu} \right|^p \right) dx .$$

So, from the equi-absolute continuity of the integral on the left-hand side and from (3.3) we deduce that:

$$(3.4) \quad \sum_{i \in I_\nu} \chi_{i\nu}(x) b(\langle Du_0 \rangle_{i\nu}) \rightarrow b(Du_0(x)) \quad \text{in } L_n^{p'}(\Omega), \quad \text{as } \nu \rightarrow +\infty .$$

Let $v_{i\nu} \in \langle Du_0 \rangle_{i\nu} \cdot y + H_{\text{per}}^{1,p}(Y^*)$ the solution of problem (1.3) corresponding to $\langle Du_0 \rangle_{i\nu}$:

Then:

$$u_{i\nu} = (v_{i\nu} - \langle Du_0 \rangle_{i\nu} \cdot y) \in H_{\text{per}}^{1,p}(Y^*)$$

and, by lemma 2.3, $Pu_{i\nu} \in H_{\text{per}}^{1,p}(Y)$.

Set

$$w_{i\nu}(y) = Pu_{i\nu}(y) - \langle Du_0 \rangle_{i\nu} \cdot y$$

and

$$w_{i\nu}^\varepsilon(x) = \varepsilon w_{i\nu}(x/\varepsilon),$$

by arguing as in the proof of lemma 2.13 we obtain:

$$(3.5) \quad \begin{cases} w_{i\nu}^\varepsilon \rightarrow \langle Du_0 \rangle_{i\nu} \cdot x & \text{in } w\text{-}H_{\text{loc}}^{1,2}(\mathbb{R}^n), \\ \tilde{a}(x/\varepsilon, Dw_{i\nu}^\varepsilon) \rightarrow b(\langle Du_0 \rangle_{i\nu}) & \text{in } w\text{-}L_{n,\text{loc}}^{p'}(\mathbb{R}^n), \\ \operatorname{div}_x \tilde{a}(x/\varepsilon, Dw_{i\nu}^\varepsilon) = 0. \end{cases}$$

Using the periodicity of $Pu_{i\nu}$ and lemma 2.3, we have:

$$\begin{aligned} \sum_{i \in I_\nu} \int_{Q_{i\nu}} |Dw_{i\nu}^\varepsilon|^p dx &\leq \sum_{i \in I_\nu} 2^{-\nu n} \varepsilon^n (1/\varepsilon + 2^\nu)^n \int_Y |Dw_{i\nu}^\varepsilon(y)|^p dy < \\ &\leq c \sum_{i \in I_\nu} 2^{-\nu n} (1 + \varepsilon^n 2^{\nu n}) \left(\int_{Y^*} |Dv_{i\nu}|^p dy + |\langle Du_0 \rangle_{i\nu}| \right)^p. \end{aligned}$$

The from (2.10), writing the last term as an integral over Ω_ν we have:

$$(3.6) \quad \sum_{i \in I_\nu} \int_{Q_{i\nu}} |Dw_{i\nu}^\varepsilon|^p dx \leq c(1 + \varepsilon^n 2^{\nu n}) \int_\Omega (1 + |Du_0|)^p dx.$$

Let now $\eta \in C_0^1(Q_{i\nu})$, $0 \leq \eta \leq 1$, and extend it by periodicity to the whole \mathbb{R}^n .

Case $p \geq 2$. If $\varphi \in C_n^0(\bar{\Omega})$, set $M_\varphi = \sup_\Omega |\varphi|$, from i) of H_2) we have:

$$\begin{aligned} (3.7) \quad &\left| \int_\Omega \tilde{a}(x/\varepsilon, Du_\varepsilon) \varphi \eta dx - \sum_{i \in I_\nu} \int_{Q_{i\nu}} \tilde{a}(x/\varepsilon, Dw_{i\nu}^\varepsilon) \varphi \eta dx \right| < \\ &\leq c M_\varphi |\Omega \setminus \Omega_\nu|^{1/p} + \left| \sum_{i \in I_\nu} \int_{Q_{i\nu} \cap \Omega_\varepsilon} [a(x/\varepsilon, Du_\varepsilon) - a(x/\varepsilon, Dw_{i\nu}^\varepsilon)] \varphi \eta dx \right| < \\ &\leq c M_\varphi |\Omega \setminus \Omega_\nu|^{1/p} + \sum_{i \in I_\nu} \int_{Q_{i\nu} \cap \Omega_\varepsilon} M_\varphi \eta \{ (|Du_\varepsilon| + |Dw_{i\nu}^\varepsilon|)^{p-2} \cdot |Du_\varepsilon - Dw_{i\nu}^\varepsilon| \} dx < \\ &\leq c M_\varphi |\Omega \setminus \Omega_\nu|^{1/p} + c M_\varphi^{p/(p-1)} \delta^{p/(p-1)} (1 + \varepsilon^n 2^{\nu n}) + \\ &+ \delta^{-p} \sum_{i \in I_\nu} \int_{Q_{i\nu} \cap \Omega_\varepsilon} \eta |Du_\varepsilon - Dw_{i\nu}^\varepsilon|^p dx \end{aligned}$$

where the last inequality is obtained by applying Young inequality with $\delta > 0$ and the estimate (3.5).

On the other hand from ii)

$$\begin{aligned} \sum_{i \in I_\nu} \int_{Q_{i\nu} \cap \Omega_\varepsilon} \eta |Du_\varepsilon - Dw_{i\nu}^\varepsilon|^p dx &\leq \sum_{i \in I_\nu} \int_{Q_{i\nu}} \eta (\tilde{a}(x/\varepsilon, Du_\varepsilon) - \tilde{a}(x/\varepsilon, Dw_{i\nu}^\varepsilon), Du_0 - \langle Du_0 \rangle_{i\nu}) dx + \\ &+ \sum_{i \in I_\nu} \int_{Q_{i\nu}} \eta (\tilde{a}(x/\varepsilon, Du_\varepsilon) - \tilde{a}(x/\varepsilon, Dw_{i\nu}^\varepsilon), (DP_\varepsilon u_\varepsilon - Dw_{i\nu}^\varepsilon) - \\ &- (Du_0 - \langle Du_0 \rangle_{i\nu})) dx. \end{aligned}$$

Then, integrating by parts and using (3.5) and the fact that u_ε is the solution of $(\mathcal{F}_\varepsilon)$, we get:

$$\begin{aligned} (3.8) \quad \sum_{i \in I_\nu} \int_{Q_{i\nu} \cap \Omega_\varepsilon} \eta |Du_\varepsilon - Dw_{i\nu}^\varepsilon|^p dx &\leq \sum_{i \in I_\nu} \int_{Q_{i\nu}} \eta (\tilde{a}(x/\varepsilon, Du_\varepsilon) - \tilde{a}(x/\varepsilon, Dw_{i\nu}^\varepsilon), Du_0 - \langle Du_0 \rangle_{i\nu}) dx + \\ &+ \sum_{i \in I_\nu} \int_{Q_{i\nu}} [\eta f \chi_{\Omega_\varepsilon} - D\eta (\tilde{a}(x/\varepsilon, Du_\varepsilon) - \tilde{a}(x/\varepsilon, Dw_{i\nu}^\varepsilon))] \cdot \\ &\cdot [(P_\varepsilon u_\varepsilon - u_0) - (w_{i\nu}^\varepsilon - \langle Du_0 \rangle_{i\nu} \cdot x)] dx + \\ &+ \langle \mu_\nu^\varepsilon, \boldsymbol{\eta} [(P_\varepsilon u_\varepsilon - u_0) - (w_{i\nu}^\varepsilon - \langle Du_0 \rangle_{i\nu} \cdot x)] \rangle. \end{aligned}$$

Then, passing to the limit as $\varepsilon \rightarrow 0$ in (3.8), by (3.2), (3.5) and lemma 2.9, we obtain

$$\begin{aligned} (3.9) \quad \lim_{\varepsilon \rightarrow 0} \sum_{i \in I_\nu} \int_{Q_{i\nu} \cap \Omega_\varepsilon} \eta |Du_\varepsilon - Dw_{i\nu}^\varepsilon|^p dx &\leq \sum_{i \in I_\nu} \int_{Q_{i\nu}} \eta (a_0(x) - b(\langle Du_0 \rangle_{i\nu}), Du_0 - \langle Du_0 \rangle_{i\nu}) dx. \end{aligned}$$

If we first pass to the limit in (3.7) as $\varepsilon \rightarrow 0$, then $\eta \rightarrow 1, \nu \rightarrow +\infty$, by (3.9) we get:

$$\left| \int_\Omega a_0(x) \varphi dx - \int_\Omega b(Du_0) \varphi dx \right| \leq c M_\varphi^{p/(p-1)} \delta^{p/(p-1)}$$

So, letting $\delta \rightarrow 0$, from the arbitrariness of φ we deduce (3.1).

Case 1 $1 < p < 2$. In this case the proof is very similar to the previous case. Indeed, by using j) of H_2 , we have:

$$\left| \int_{\Omega} \tilde{a}(x/\varepsilon, Du_{\varepsilon}) \varphi \eta \, dx - \sum_{i \in I_r} \int_{Q_{i_r}} \tilde{a}(x/\varepsilon, Dw_{i_r}^{\varepsilon}) \varphi \eta \, dx \right| \leq \\ \leq c M_{\varphi} |\Omega \setminus \Omega_r|^{1/p} + \beta \sum_{i \in I_r} \int_{Q_{i_r} \cap \Omega_{\varepsilon}} |Du_{\varepsilon} - Dw_{i_r}^{\varepsilon}|^{p-1} |\varphi| \eta \, dx .$$

Then, using jj) we can control the last term:

$$\sum_{i \in I_r} \int_{Q_{i_r} \cap \Omega_{\varepsilon}} |Du_{\varepsilon} - Dw_{i_r}^{\varepsilon}|^{p-1} |\varphi| \eta \, dx \leq \\ \leq c \delta^{2/(3-p)} \sum_{i \in I_r} \int_{Q_{i_r} \cap \Omega_{\varepsilon}} (|Du_{\varepsilon}| + |Dw_{i_r}^{\varepsilon}|)^{(2-p)(p-1)/(3-p)} \, dx + \\ + \delta^{-2/(p-1)} \sum_{i \in I_r} \int_{Q_{i_r}} \eta (\tilde{a}(x/\varepsilon, Du_{\varepsilon}) - \tilde{a}(x/\varepsilon, Dw_{i_r}^{\varepsilon}), DP_{\varepsilon} u_{\varepsilon} - Dw_{i_r}^{\varepsilon}) \eta \, dx .$$

Hence, arguing as in the previous case we obtain the result. \blacksquare

An easy consequence of theorem 3.1 is the following result:

COROLLARY 3.2. *Assume that $a(x, \xi)$ verifies the structure conditions $H_1) \div H_3)$ and is homogeneous of degree $p - 1$ with respect to ξ .*

Then for any $f \in L^{p'}(\Omega)$ and $g(x, y)$ verifying (1.1₁) \div (1.1₃) with $\mu_{\varepsilon} \neq 0$, the function $b(\xi)$ is homogeneous of degree $p - 1$ and the sequence v_{ε} of the solutions of the problem:

$$\begin{aligned} - \operatorname{div} a(x/\varepsilon, Dv_{\varepsilon}) &= f && \text{in } \Omega_{\varepsilon} , \\ v_{\varepsilon} &= 0 && \text{on } \partial\Omega , \\ a(x/\varepsilon, Dv_{\varepsilon}) \cdot \nu &= g_{\varepsilon} && \text{on } \partial T_{\varepsilon} , \end{aligned}$$

verifies:

$$\begin{aligned} \varepsilon^{1/(p-1)} P_{\varepsilon} v_{\varepsilon} &\rightarrow v_0 && \text{in } w\text{-}H_0^{1,p}(\Omega) , \\ \varepsilon \tilde{a}(x/\varepsilon, Dv_{\varepsilon}) &\rightarrow b(Dv_0) && \text{in } w\text{-}L_n^{p'}(\Omega) , \end{aligned}$$

where v_0 is the solution of the problem:

$$\begin{aligned} -\operatorname{div} b(Dv_0) &= \mu_0 \quad \text{in } \Omega_\varepsilon, \\ v_0 &\in H_0^{1,p}(\Omega), \end{aligned}$$

with $b(\xi)$ and μ_0 given respectively by (1.4), (1.5) and (1.3).

PROOF. The result follows by applying theorem 3.2 to the sequence $u_\varepsilon = \varepsilon^{1/(p-1)}v_\varepsilon$. ■

REFERENCES

- [1] E. ACERBI - D. PERCIVALE, *Homogenization of noncoercive functionals: periodic materials with soft inclusions*, Appl. Math. Optim., **17** (1988), pp. 91-102.
- [2] A. NENSOUSSAN - J. L. LIONS - G. PAPANICOLAOU, *Asymptotic analysis for periodic structures*, North-Holland, Amsterdam, 1978.
- [3] L. BOCCARDO - F. MURAT, *Homogénéisation de problèmes quasi-linéaires*, Proceeding of the Meeting « Studio di problemi limite dell'Analisi Funzionale », Bressanone 1981, Pitagora, 1982.
- [4] L. CARBONE - C. SBORDONE, *Some properties of Γ -limits of integral functionals*, Ann. Mat. Pura Appl., **122** (1979), pp. 1-60.
- [5] V. CHIADÒ PIAT, *Convergence of minima for non equicoercive functionals and related problems*, preprint S.I.S.S.A., Trieste.
- [6] D. CIORANESCU - P. DONATO, *Homogénéisation du problème de Neumann non homogène dans des ouverts perforés*, to appear on Asymptotic Analysis, **1**, 2 (1988).
- [7] D. CIORANESCU - J. SAINT JEAN PAULIN, *Homogenization in open sets with holes*, J. Math. Anal. Appl., **71** (1979), pp. 590-607.
- [8] P. DONATO, *Una stima per la differenza di H -limiti e qualche applicazione a problemi di omogenizzazione*, Rend. Matematica, **4** (1983), pp. 623-640.
- [9] E. DE GIORGI - S. SPAGNOLO, *Sulla convergenza degli integrali dell'energia per operatori ellittici del secondo ordine*, Boll. Un. Mat. Ital., **8** (1973), pp. 391-411.
- [10] N. FUSCO - G. MOSCARIELLO, *An application of duality to homogenization of integral functionals*, Memorie dell'Acc. dei Lincei, **17**, I (1984), pp. 361-372.
- [11] N. FUSCO - G. MOSCARIELLO, *On the homogenization of quasilinear divergence structure operators*, Ann. Mat. Pura Appl., **4**, 146 (1987), pp. 1-13.

- [12] P. GRISVARD, *Elliptic Problems in Nonsmooth Domains*, Monographs and Studies in Mathematics, **24**, Pitman, London, 1985.
- [13] S. MORTOLA - A. PROFETI. *On the convergence of the minimum points of non equicoercive quadratic functionals*, Comm. Part. Diff. Eqs., **7**, 6 (1982), pp. 645-673.
- [14] F. MURAT, *Compacité par compensation*, Ann. Scuola Norm. Sup. Pisa, **5** (1978), pp. 489-507.
- [15] E. SANCHEZ-PALENCIA, *Nonhomogeneous media and vibration theory*, Lecture Notes in Physics, **127**, Springer-Verlag, Berlin, 1980.
- [16] P. SUQUET, *Plasticité et homogénéisation*, Thèse d'Etat, Univ. de Paris VI, 1982.
- [17] L. TARTAR, *Topics in Nonlinear Analysis*, Publ. Math. Univ. d'Orsay, **13** (1978).
- [18] L. TARTAR, *Homogénéisation et compacité par compensation*, Séminaire Schwartz Exposé, **9** (1978).
- [19] L. TARTAR, *Cours Peccot Collège de France 1977*, partially redacted by F. Murat; *H-convergence*, Séminaire d'Analyse Fonctionnelle et Numérique de l'Univ. d'Alger, 1977/78.

Manoscritto pervenuto in redazione il 25 luglio 1989.