

RENDICONTI  
*del*  
SEMINARIO MATEMATICO  
*della*  
UNIVERSITÀ DI PADOVA

JOHN C. LENNOX

A. MOHAMMADI HASSANABADI

JAMES WIEGOLD

**Some commutativity criteria. - II**

*Rendiconti del Seminario Matematico della Università di Padova*,  
tome 86 (1991), p. 207-211

[http://www.numdam.org/item?id=RSMUP\\_1991\\_\\_86\\_\\_207\\_0](http://www.numdam.org/item?id=RSMUP_1991__86__207_0)

© Rendiconti del Seminario Matematico della Università di Padova, 1991, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## Some Commutativity Criteria. - II

JOHN C. LENNOX - A. MOHAMMADI HASSANABADI  
JAMES WIEGOLD(\*)

In [1] we were concerned with groups  $G$  such that  $XY = YX$  for all  $n$ -sets  $X$  and  $Y$  in  $G$ , the  $P_n$ -groups of that paper. Theorem B stated that all infinite  $P_n$ -groups are abelian, but of course some finite  $P_n$ -groups are non-abelian. Our aim here is to establish the best possible result in this direction:

**THEOREM 1.** (i) Every group in  $P_n$  of order at least  $2n$  is abelian.

(ii) For each  $t$ , every group of order  $t$  is in  $P_n$  whenever  $n > t/2$ .

The same sort of questions can be asked about other algebraic structures than groups. To show how different semigroups are in this context, we prove, with the obvious definitions:

**THEOREM 2.** A semigroup  $S$  with identity is a non-commutative  $P_2$ -semigroup if and only if  $S$  is the disjoint union  $S = A \cup B$ , where

- (i)  $|A| = 2$  and  $A$  is a left zero or right zero semigroup,
- (ii)  $B$  is a commutative subsemigroup containing the identity 1 of  $S$ ,
- (iii)  $xy = yx = x$  for all  $x$  in  $A$ ,  $y$  in  $B$ .

**PROOFS.** To prove Theorem 1, we first establish the simple fact that  $P_n \leq P_{n+1}$  for all  $n$ . Let  $A$  be a group (or indeed a semigroup) in  $P_n$ , and  $X, Y$  any subsets of cardinal  $n + 1$  of  $A$ . Let  $xy$  be any element in the product, with  $x \in X, y \in Y$ . Then  $x$  is in an  $n$ -set  $X_1$  contained in

(\*) Indirizzo degli AA.: J. C. LENNOX and J. WIEGOLD: School of Mathematics, University of Wales, College of Cardiff, Senghennydd Road, Cardiff CF2 4AG, Wales; A. MOHAMMADI HASSANABADI: Department of Mathematics, University of Isfahan, Isfahan, Iran.

$X$ , and similarly  $y$  is contained in an  $n$ -set  $Y_1$  in  $Y$ . Thus  $xy \in X_1 Y_1 = Y_1 X_1 \leq YX$ , so that  $XY \leq YX$ . The converse is obvious, and we have  $XY = YX$ . Because of this, all we need do now is show that groups of order  $2n$  and  $2n + 1$  in  $P_n$  are abelian.

We shall do the case  $|G| = 2n$  as an example. Let  $G$  be a non-abelian group of order  $2n$ , and let  $x_1, x_2$  be non-commuting elements of  $G$ . Further, let  $X = \{x_1, x_2, \dots, x_n\}$  be any  $n$ -set chosen in such a way that  $x_r^{x_1} \neq x_2$  for each  $r = 1, 2, \dots, n$ . The choice is possible since  $|G| \geq 2n$ . Finally, set

$$Y = G \setminus \{1, x_2^{-1}x_1, \dots, x_n^{-1}x_1\} = \{y_1, y_2, \dots, y_n\}, \text{ say.}$$

To establish the theorem, it is enough to show that  $XY \neq YX$ .

Clearly,  $x_1 \notin XY$  by the choice of  $Y$ . We shall show that  $x_1 \in YX$ . We have

$$x_1 \in YX \Leftrightarrow \exists i, j \quad \text{with } y_i = x_1 x_j^{-1},$$

so that

$$\begin{aligned} x_1 \notin YX &\Leftrightarrow \forall i, j: x_1 x_j^{-1} \neq y_i \\ &\Leftrightarrow \forall j: x_1 x_j^{-1} \notin Y \\ &\Leftrightarrow \forall j \exists r: x_1 x_j^{-1} = x_r^{-1} x_1 \\ &\Leftrightarrow \forall j \exists r: x_j = x_r^{x_1}. \end{aligned}$$

Thus

$$x_1 \in XY \Leftrightarrow \exists j \forall r: x_r^{x_1} \neq x_j.$$

However  $x_r^{x_1} \neq x_j$  for all  $j$ , so  $x_1 \in YX$ , as required.

The proof for  $|G| = 2n + 1$  is more-or-less identical: just take  $Y = G \setminus \{1, x_2^{-1}x_1, \dots, x_n^{-1}x_1, x_1\}$  instead.

For part (ii) of the theorem, take  $G$  of order  $t$ ,  $n > t/2$  and two  $n$ -sets  $X, Y$  in  $G$ . Then  $XY = G$  since for all  $g$  in  $G$ ,  $gY^{-1} \cap X \neq \emptyset$  so that  $gy^{-1} = x$  for suitable  $y \in Y$ ,  $x \in X$ , and  $g = xy$ . Similarly,  $G = YX$ , so that  $G \in P_n$ , as required, and this completes the proof of Theorem 1.

As for Theorem 2, it is a matter of routine verification to show that semigroups with the structure indicated in the statement are  $P_2$ -semigroups.

To prove the converse, let  $S$  be a non-commutative  $P_2$ -semigroup with identity 1, and  $a, b$  two non-commuting elements of  $S$ . We show

first that  $\{a, b\}$  is a left or right zero semigroup. Since  $S$  is in  $P_2$ ,

$$\{1, a\}\{1, b\} = \{1, b\}\{1, a\},$$

so

$$\{1, a, b, ab\} = \{1, b, a, ba\}.$$

There are three cases to consider. If  $ab = 1$ , we must have  $ba = a$  or  $ba = b$ , since  $ab \neq ba$ . If  $ba = a$ , then  $b = b(ab) = (ba)b = ab = 1$ , a contradiction; while if  $ba = b$ , we have  $1 = ab = a(ba) = (ab)a = 1 \cdot a = a$ , another contradiction. Thus  $ab \neq 1$ , and, symmetrically,  $ba \neq 1$ .

If  $ab = a$ , then

$$\{1, a, b\} = \{1, a, b, ba\}$$

so that  $ba = b$  since  $ab \neq ba$ . Then

$$a^2 = (ab)a = a(ba) = ab = a,$$

$$b^2 = (ba)b = b(ab) = ba = b,$$

and  $\{a, b\}$  forms a left zero semigroup.

Finally, if  $ab = b$ , the same sort of argument shows that  $A := \{a, b\}$  is a left zero semigroup.

For the remainder of the proof we shall assume without loss that  $A$  is a left zero semigroup, that is,  $a^2 = ab = a$ ,  $b^2 = ba = b$ . We show first that every element  $c$  outside  $A$  must commute with  $a$  or  $b$ . If not,  $\{a, c\}$  and  $\{b, c\}$  are both left or right zero semigroups, and thus there are four cases to consider.

$$1) \ c^2 = c = ca, \ ac = a, \ cb = c, \ bc = b.$$

This is impossible, since

$$\{a, b\}\{1, c\} = \{a, b\},$$

$$\{1, c\}\{a, b\} = \{a, b, c\}.$$

$$2) \ c^2 = c = ca, \ ac = a, \ bc = c, \ cb = b.$$

Here

$$\{a, b\}\{b, c\} = \{a, b, c\},$$

$$\{b, c\}\{a, b\} = \{b, c\}.$$

$$3) \ c^2 = c = ac, \ ca = a, \ bc = b, \ cb = c.$$

Here

$$\begin{aligned}\{a, b\}\{a, c\} &= \{a, b, c\}, \\ \{a, c\}\{a, b\} &= \{a, c\}.\end{aligned}$$

4) Finally in this part of the argument,  $c^2 = c = ac$ ,  $ca = a$ ,  $bc = b$ ,  $cb = b$ . Here

$$\begin{aligned}\{a, b\}\{a, c\} &= \{a, b, c\}, \\ \{a, c\}\{a, b\} &= \{a, c\}.\end{aligned}$$

Thus  $c$  must commute with  $a$  or  $b$ , and it is no loss of generality if we assume that  $ac = ca$ . If  $c$  does not commute with  $b$ , we have two cases to consider, depending on the structure of  $\{b, c\}$ . Recall that  $a^2 = ab = a$ ,  $b^2 = ba = b$ .

$$1) \quad bc = b, \quad c^2 = cb = c.$$

Hence

$$\begin{aligned}\{a, b\}\{a, c\} &= \{a, ac, b\}, \\ \{a, c\}\{a, b\} &= \{a, ac, c\}.\end{aligned}$$

Thus  $b = ac$  and  $c = ac$ , which is a contradiction.

$$2) \quad bc = c = c^2, \quad cb = b.$$

Here

$$\begin{aligned}\{a, b\}\{a, c\} &= \{a, ac, b, c\}, \\ \{a, c\}\{a, b\} &= \{a, ac, b\}.\end{aligned}$$

Thus  $c = ac$  and we have

$$\begin{aligned}\{a, b\}\{b, c\} &= \{a, c, b\}, \\ \{b, c\}\{a, b\} &= \{b, c\}.\end{aligned}$$

which is false.

Hence, thus far we have shown that  $c \notin \{a, b\} \Rightarrow ac = ca$ ,  $bc = cb$ . Consider the following product:

$$\begin{aligned}\{a, b\}\{a, c\} &= \{a, b, ac, bc\}, \\ \{a, c\}\{a, b\} &= \{a, ac, bc\}.\end{aligned}$$

This gives that  $b = bc$ , since the other possibility, *viz.*  $b = ac$ , means

that  $ab = ba$  since  $a$  commutes with  $c$ . Similarly,  $a = ac$ . This completes item (iii).

The next step is to show that  $cd = dc$  for all  $c, d$  outside  $\{a, b\}$ . This is clear, since otherwise  $\{c, d\}$  can play the part of  $\{a, b\}$  in the argument to this point, and we would get  $c = ac = ca = a$ .

The final step is to show that  $c, d \notin \{a, b\} \Rightarrow cd \notin \{a, b\}$ . Suppose, without loss of generality, that  $cd = a$ . Then  $(cd)b = ab = a$ ,  $c(db) = cb = b$ , a contradiction which completes the proof.

We have no idea what happens with  $P_3$ -semigroups with identity, nor with semigroups without identity. The arguments here are likely to be very cumbersome.

#### REFERENCES

- [1] J. C. LENNOX - A. MOHAMMADI HASSANABADI - J. WIEGOLD, *Some commutativity criteria*, Rend. Sem. Mat. Univ. Padova, **84** (1990).

Manoscritto pervenuto in redazione il 7 gennaio 1991.