

The Obstacle Thermistor Problem with Periodic Data

MAURIZIO BADIO (*)

ABSTRACT - The object of this paper is the study of an obstacle thermistor problem with a nonlocal term. Using the classical Lax-Milgram theorem, a result of Lions and a fixed point argument we prove existence of weak periodic solutions. Finally, by means of some a-priori estimates in Campanato's spaces, we obtain the regularity of these solutions.

1. Introduction.

Let Ω be a bounded open set of R^n , $n \geq 1$, with smooth boundary $\partial\Omega$. For given $\omega > 0$, we set $Q := \Omega \times P$, $\Sigma := \partial\Omega \times P$ and $P := \frac{R}{\omega Z}$ denotes the period interval $[0, \omega]$ so the functions defined in Q and Σ are ω -time periodic. In this paper we consider the following parabolic-elliptic system arising from an obstacle evolution thermistor problem

$$(1.1) \quad u_t - \Delta u + \int_{\Omega} G(x, y)u(y, t)dy \geq \sigma(u)|\nabla\varphi|^2 \text{ in } Q,$$

$$(1.2) \quad u(x, t) = 0 \text{ on } \Sigma,$$

$$(1.3) \quad \operatorname{div}(\sigma(u)\nabla\varphi) = 0 \text{ in } Q,$$

$$(1.4) \quad \varphi(x, t) = \varphi_0(x, t) \text{ on } \Sigma.$$

Here Ω is the conductor and the unknowns u and φ are the temperature and the electrical potential in Ω , respectively. The nonlocal term $\int_{\Omega} G(x, y)u(y, t)dy$, with $G(x, y) \geq 0$, describes heat losses to the surrounding Ω

(*) Indirizzo dell'A.: Dipartimento di Matematica "G. Castelnuovo", Università di Roma "La Sapienza", P. A. Moro 2, 00185 Roma, Italy.

E-mail: badio@mat.uniroma1.it

gas while $\sigma(u)$ denotes the temperature dependent electrical conductivity. Finally, the term $\sigma(u)|\nabla\varphi|^2$ represents the Joule heating in Ω . We note that if $\varphi \in L^2(P; W^{1,2}(\Omega)) \cap L^\infty(Q)$, then (1.3) implies

$$\sigma(u)|\nabla\varphi|^2 = \operatorname{div}(\sigma(u)\varphi\nabla\varphi),$$

in the sense of distribution.

Thus, we shall study the problem

$$(1.5) \quad u_t - \Delta u + \int_{\Omega} G(x, y)u(y, t)dy \geq \operatorname{div}(\sigma(u)\varphi\nabla\varphi) \text{ in } Q,$$

$$(1.6) \quad u(x, t) = 0 \text{ on } \Sigma$$

$$(1.7) \quad \operatorname{div}(\sigma(u)\nabla\varphi) = 0 \text{ in } Q,$$

$$(1.8) \quad \varphi(x, t) = \varphi_0(x, t), \text{ on } \Sigma.$$

To solve (1.5)-(1.8) we follow the approach of [1], [2] and introduce the penalized problem

$$(1.9) \quad u_{nt} - \Delta u_n + \left(\int_{\Omega} G(x, y)u_n(y, t)dy \right) I_n(u_n) = \operatorname{div}(\sigma(u_n)\varphi_n\nabla\varphi_n) \text{ in } Q,$$

$$(1.10) \quad u_n(x, t) = 0 \text{ on } \Sigma,$$

$$(1.11) \quad \operatorname{div}(\sigma(u_n)\nabla\varphi_n) = 0 \text{ in } Q,$$

$$(1.12) \quad \varphi_n(x, t) = \varphi_0(x, t) \text{ on } \Sigma,$$

where it is assumed that I_n satisfies

H_n) $I_n \in C(R)$, $0 \leq I_n(s) \leq 1$, for all $s \in R$, $I_n(s) = 0$ if $s \leq 0$ and $I_n(s) \rightarrow H(s)$ in L^2 , where $H(s)$ denotes the Heaviside function.

The plan of the paper is as follows: in Section 2 we give some notations and preliminary results. In Section 3, we consider a linearized version of the elliptic problem (1.7)-(1.8) (see (3.5)-(3.6)) and prove existence and uniqueness of the weak periodic solution φ by means of the classical Lax-Milgram's theorem. Next, in Section 4, we use a result due to Lions [5, Theorem 6.1] to obtain existence and uniqueness of periodic solutions u_n for the penalized problem (1.9)-(1.10). In Section 5, we deduce some uniform estimates for (u_n, φ_n) and utilize a fixed point argument to obtain the

existence of weak periodic solutions (u, φ) of (1.5)-(1.8) by letting $n \rightarrow \infty$. Finally, in Section 6, we establish the Hölder regularity for (u, φ) . This will be done by proving a-priori estimates in Campanato's spaces as in [6], [7].

2. Preliminaries and auxiliary results.

Let us introduce an useful space of ω -periodic functions in which we look for solutions to our problem.

We consider the Hilbert space $V_0 := L^2(P; W_0^{1,2}(\Omega))$ endowed with the norm

$$(2.1) \quad \|v\|_{V_0} := \left(\int_Q |\nabla v(x, t)|^2 dx dt \right)^{1/2}$$

and its dual $V^* := L^2(P; W^{-1,2}(\Omega))$. The duality pairing between V_0 and V^* shall be denoted by $\langle \cdot, \cdot \rangle$. The space V_0 is the closure with respect to the norm (2.1) of $C_0^\infty(\bar{Q})$, the space of periodic functions vanishing near Σ . We recall some notations concerning the Campanato spaces.

NOTATIONS. Let $Q_{t_0, t_1} := \Omega \times (t_0, t_1]$, $0 \leq t_0 < t_1$. A point $(x, t) \in Q_{t_0, t_1}$ shall be denoted by z . Let $B_r(x_0)$ be the ball centered at x_0 with radius r and let $Q_r(z_0)$ be the cylinder $B_r(x_0) \times (t_0 - r^2, t_0]$. Moreover, define $Q_r[z_0, r] := Q_r(z_0) \cap Q_{t_0, t_1}$ and $\Omega[x_0, r] := B_r(x_0) \cap \Omega$.

For $0 \leq \mu < n$, the Campanato space

$$\mathcal{L}^{2,\mu}(\Omega) := \left\{ u \in L^2(\Omega) : \left(\sup_{x_0 \in \bar{\Omega}, r > 0} r^{-\mu} \int_{\Omega[x_0, r]} u^2(x) dx \right)^{1/2} < +\infty \right\}$$

is a Banach space with the equivalent norm

$$\|u\|_{2,\mu,\Omega[x_0, r]} = \left(\sup_{x_0 \in \bar{\Omega}, r > 0} r^{-\mu} \int_{\Omega[x_0, r]} u^2(x) dx \right)^{1/2}.$$

(see [3]).

For $0 \leq \mu < n + 2$, the Campanato space

$$\mathcal{L}^{2,\mu}(Q_{t_0, t_1}) := \left\{ u \in L^2(Q_{t_0, t_1}) : \sup_{z_0 \in Q_{t_0, t_1}, r > 0} \left(r^{-\mu} \int_{Q_r[z_0, r]} u^2(x, t) dx dt \right)^{1/2} < +\infty \right\}.$$

is a Banach space with the equivalent norm

$$\|u\|_{2,\mu,Q_{t_0,t_1}} = \left(\sup_{z_0 \in Q_{t_0,t_1}, r > 0} r^{-\mu} \int_{Q_r[z_0,r]} u^2(x,t) \, dxdt \right)^{1/2}.$$

Furthermore, we need the following result

PROPOSITION A ([7]). The space $\mathcal{L}^{2,n+2+2\mu}(Q_{t_0,t_1})$ is isomorphic topologically and algebraically to $C^{\mu,\mu/2}(\overline{Q_{t_0,t_1}})$, for $\mu \in (0, 1)$.

We shall study problem (1.5)-(1.8) under the following assumptions:

H_σ $\sigma \in C(R)$, $0 < \sigma_* \leq \sigma(s) \leq \sigma^*$ for any $s \in R$;

H_G $G \in C(R^2)$, $G(x, y) \geq 0$, we set $\widehat{G} := \sup_{R^2} G(x, y)$;

H_0 φ_0 is an ω -periodic bounded function with an extension $\widetilde{\varphi}_0$ to Q which satisfies $\widetilde{\varphi}_0 \in L^\infty(P; W^{1,\infty}(\Omega))$.

Moreover, we define the convex set

$$K := \{v \in L^2(P; W_0^{1,2}(\Omega)), v \geq 0 \text{ a.e. in } Q\}.$$

The notion of weak periodic solution for our problem is the following

DEFINITION 2.1. A pair of functions (u, φ) are a weak periodic solution to (1.5)-(1.8) if the following conditions hold

$$u \in K, \quad u_t \in L^2(P; W^{-1,2}(\Omega)) \text{ and } \varphi - \varphi_0 \in V_0,$$

$$(2.2) \quad \int_Q u_t(v-u) \, dxdt + \int_Q \nabla u \nabla(v-u) \, dxdt + \int_Q \int_\Omega G(x,y)u(y,t)(v-u) \, dydxdt \\ \geq - \int_Q \sigma(u)\varphi \nabla \varphi \nabla(v-u) \, dxdt, \text{ for any } v \in K$$

and

$$\int_Q \sigma(u)\nabla \varphi \nabla \xi \, dxdt = 0, \text{ for any } \xi \in V_0.$$

3. The elliptic problem.

In this section we study problem (1.7)-(1.8). Set $v(x, t) := \varphi(x, t) - \varphi_0(x, t)$. Fixed $w \in L^2(Q)$ with $w \geq 0$, we solve the problem

$$(3.1) \quad \operatorname{div}(\sigma(w)\nabla v) = -\operatorname{div}(\sigma(w)\nabla\varphi_0) \text{ in } Q,$$

$$(3.2) \quad v(x, t) = 0 \text{ on } \Sigma,$$

We state the weak formulation of solution to (3.1)-(3.2).

DEFINITION 3.1. A function $v \in V_0$ is called a weak periodic solution of (3.1)-(3.2) if

$$(3.3) \quad \int_Q \sigma(w)\nabla v\nabla\zeta dxdt = - \int_Q \sigma(w)\nabla\varphi_0\nabla\zeta dxdt, \text{ for all } \zeta \in V_0.$$

The existence and uniqueness of the weak solution v shall be obtained by the classical Lax-Milgram theorem.

In agreement with this result, we define the bilinear form

$$a : V_0 \times V_0 \rightarrow R$$

by setting

$$a(v, \zeta) := \int_Q \sigma(w)\nabla v\nabla\zeta dxdt, \text{ for any } \zeta \in V_0$$

We have

PROPOSITION 3.2. If we assume H_σ , then

- i) a is continuous;
- ii) a is coercive.

PROOF. Condition i) is a consequence of Hölder's inequality. In fact

$$\begin{aligned} |a(v, \zeta)| &\leq \sigma^* \left(\int_Q |\nabla v|^2 dxdt \right)^{1/2} \|\zeta\|_{V_0}. \\ &\leq \sigma^* \|v\|_{V_0} \|\zeta\|_{V_0}. \end{aligned}$$

ii) The coercivity of a follows from

$$a(v, v) = \int_Q \sigma(w) |\nabla v|^2 dx dt \geq \sigma_* \|v\|_{V_0}^2$$

which implies

$$\frac{a(v, v)}{\|v\|_{V_0}^2} \geq \sigma_* \|v\|_{V_0} \rightarrow \infty, \text{ as } \|v\|_{V_0} \rightarrow \infty. \quad \square$$

In this way if H_0 holds, problem (3.3) becomes equivalent to the problem

$$(3.4) \quad a(v, \zeta) = \langle G, \zeta \rangle$$

where $G \in V^*$ is the linear functional defined as follows

$$\langle G, \zeta \rangle := - \int_Q \sigma(w) \nabla \varphi_0 \nabla \zeta dx dt, \quad \forall \zeta \in V_0.$$

Now, we can state the main result of this section

THEOREM 3.3. *Let H_σ and H_0 be satisfied. There exists a unique weak periodic solution to (3.4).*

PROOF. By the theorem of Lax-Milgram we easily conclude the existence and uniqueness of the weak periodic solution. \square

Thus, for $w \in L^2(Q)$, $w \geq 0$ we have solved the problem

$$(3.5) \quad \operatorname{div}(\sigma(w) \nabla \varphi) = 0 \text{ in } Q,$$

$$(3.6) \quad \varphi(x, t) = \varphi_0(x, t) \text{ on } \Sigma.$$

4. The parabolic penalized problem.

We consider the problem

$$(4.1) \quad u_{nt} - \Delta u_n + \left(\int_\Omega G(x, y) w(y, t) dy \right) I_n(w) = \operatorname{div}(\sigma(w) \varphi \nabla \varphi) \text{ in } Q,$$

$$(4.2) \quad u_n(x, t) = 0 \text{ on } \Sigma.$$

DEFINITION 4.1. A weak periodic solution of (4.1)-(4.2) corresponding to w , is a function $u_n \in V_0$ such that

$$(4.3) \quad \int_Q u_{nt} \zeta dxdt + \int_Q \nabla u_n \nabla \zeta dxdt + \int_Q \left(\int_\Omega G(x, y) w(y, t) dy \right) I_n(w) \zeta dxdt \\ = - \int_Q \sigma(w) \varphi \nabla \varphi \nabla \zeta dxdt, \text{ for any } \zeta \in V_0.$$

The existence and uniqueness of the periodic solution follows from a Lion's result (see [5, Theorem 6.1]).

For any $u_n \in V_0$ we define the mapping $B : V_0 \rightarrow V^*$ as follows

$$\langle Bu_n, \zeta \rangle := \int_Q \nabla u_n \nabla \zeta dxdt, \text{ for any } \zeta \in V_0.$$

The set

$$D := \{v \in L^2(P; W_0^{1,2}(\Omega)) : v_t \in L^2(P; W^{-1,2}(\Omega))\}$$

is dense in V_0 , because of the density of $C^\infty(\bar{Q}) \subset D$ in V_0 .

Let

$$L : D \rightarrow V^*$$

be the closed skew-adjoint (i.e. $L = -L^*$) linear operator defined by

$$\langle Lu_n, \zeta \rangle := \int_Q u_{nt} \zeta dxdt, \forall \zeta \in V_0.$$

It is known that this operator is maximal monotone (see [4, Lemma 1.1, p. 318]).

The properties of B are contained in the following result.

PROPOSITION 4.2. Assume H_σ , then

- j) B is continuous;
- jj) B is coercive.

PROOF. j) The continuity of B follows from

$$|\langle Bu_n, \zeta \rangle| \leq \|u_n\|_{V_0} \|\zeta\|_{V_0}.$$

jj) By

$$\langle Bu_n, u_n \rangle = \int_Q |\nabla u_n|^2 dx dt = \|u_n\|_{V_0}^2$$

one has

$$\frac{\langle Bu_n, u_n \rangle}{\|u_n\|_{V_0}} = \|u_n\|_{V_0} \rightarrow \infty, \text{ as } \|u_n\|_{V_0} \rightarrow \infty. \quad \square$$

Finally, let $M_n \in V^*$ be the linear functional defined by

$$\langle M_n, \zeta \rangle := \int_Q \left(\int_{\Omega} G(x, y) w(y, t) dy \right) I_n(w) \zeta dx dt - \int_Q \sigma(w) \varphi \nabla \varphi \nabla \zeta dx dt,$$

then, problem (4.3) can be considered in the framework of the abstract problems of the form

$$(4.4) \quad Lu_n + Bu_n = M_n$$

to which we apply [5, Theorem 6.1] to establish the existence and uniqueness of the weak periodic solution.

5. A fixed point argument.

The periodicity of solutions to (1.5)-(1.8) shall be proved utilizing the Schauder fixed point theorem for a suitable operator equation. To this aim, let

$$\Theta : \mathcal{S} \rightarrow \mathcal{S}$$

be the mapping defined by

$$\Theta(w) = u_n$$

and

$$\mathcal{S} := \{v \in L^2(Q) : \|v\|_{L^2(Q)} \leq M\},$$

where u_n is the unique weak periodic solution of (4.1)-(4.2) corresponding to w . The mapping Θ is well-defined. In order to prove its continuity we will prove some crucial estimates and convergences useful to utilize the Schauder fixed point theorem. Let $w_k \in L^2(Q)$ be a non-negative sequence such that $w_k \rightarrow w$ and $\sigma(w_k) \rightarrow \sigma(w)$ strongly in $L^2(Q)$ as $k \rightarrow \infty$. We denote by u_{nk} and φ_k , respectively, the weak periodic

solutions of

$$(5.1) \quad \int_Q u_{nkt} \zeta dxdt + \int_Q \nabla u_{nk} \nabla \zeta dxdt + \int_Q \left(\int_{\Omega} G(x, y) w_k(y, t) dy \right) I_n(w_k) \zeta dxdt = - \int_Q \sigma(w_k) \varphi_k \nabla \varphi_k \nabla \zeta dxdt, \text{ for any } \zeta \in V_0$$

and

$$(5.2) \quad \int_Q \sigma(w_k) \nabla \varphi_k \nabla \zeta dxdt = 0, \text{ for any } \zeta \in V_0.$$

An estimate on $\nabla \varphi_k$ is achieved choosing $\varphi_k - \varphi_0$ as a test function in (5.2). In fact,

$$\begin{aligned} \sigma_* \int_Q |\nabla \varphi_k|^2 dxdt &\leq \int_Q \sigma(w_k) |\nabla \varphi_k|^2 dxdt = \int_Q \sigma(w_k) \nabla \varphi_k \nabla \varphi_0 dxdt \\ &\leq \sigma^* \left(\int_Q |\nabla \varphi_k|^2 dxdt \right)^{1/2} \left(\int_Q |\nabla \varphi_0|^2 dxdt \right)^{1/2} \end{aligned}$$

hence,

$$(5.3) \quad \int_Q |\nabla \varphi_k|^2 dxdt \leq \left(\frac{\sigma^*}{\sigma_*} \right)^2 \|\nabla \varphi_0\|_{L^2(Q)}^2.$$

Moreover, by the weak maximum principle we derive that

$$(5.4) \quad \|\varphi_k\|_{L^\infty(Q)} \leq \|\tilde{\varphi}_0\|_{L^\infty(Q)}.$$

Combining (5.3) and (5.4), one has the usual energy estimate

$$(5.5) \quad \int_Q |\varphi_k(x, t)|^2 dxdt + \int_Q |\nabla \varphi_k(x, t)|^2 dxdt \leq C.$$

In what follows, C stands for a generic positive constant independent of k and n .

Thanks to (5.3), we deduce that φ_k is uniformly bounded in the V norm, with respect to k . Therefore

$$\varphi_k \rightharpoonup \varphi \text{ in } V \text{ as } k \text{ goes to infinity.}$$

Furthermore, one has

LEMMA 5.1. The sequence $\nabla\varphi_k$ converges strongly to $\nabla\varphi$ in $L^2(P; (L^2(\Omega))^n)$.

PROOF. Taking $\varphi_k - \varphi$ as a test function in (5.2), we get

$$\int_Q \sigma(w_k) |\nabla(\varphi_k - \varphi)|^2 dxdt = \int_Q \sigma(w_k) \nabla(\varphi - \varphi_k) \nabla\varphi dxdt$$

from which it follows that

$$\sigma_* \int_Q |\nabla(\varphi_k - \varphi)|^2 dxdt \leq \int_Q \sigma(w_k) \nabla(\varphi - \varphi_k) \nabla\varphi dxdt.$$

The weak convergence of $\sigma(w_k) \nabla(\varphi - \varphi_k)$ to zero as $k \rightarrow \infty$ leads to the conclusion. \square

Choosing u_{nk} as a test function in (5.1), we have

$$\begin{aligned} \int_Q |\nabla u_{nk}|^2 dxdt + \int_Q \left(\int_\Omega G(x, y) w_k(y, t) dy \right) I_n(w_k) u_{nk}(x, t) dxdt \\ = - \int_Q \sigma(w_k) \varphi_k \nabla \varphi_k \nabla u_{nk} dxdt. \end{aligned}$$

Applying Young's and Poincaré's inequalities, one obtains

$$\begin{aligned} & \frac{1}{2} \int_Q |\nabla u_{nk}|^2 dxdt \\ & \leq \frac{\varepsilon}{2} \int_Q |u_{nk}|^2 dxdt + \frac{\widehat{G}^2}{2\varepsilon} \int_Q \left(\int_\Omega w_k(y, t) dy \right)^2 dxdt \\ & \quad + \frac{(\sigma^* \|\tilde{\varphi}_0\|_{L^\infty(Q)})^2}{2} \left(\frac{\sigma^*}{\sigma_*} \right)^2 \|\nabla\varphi_0\|_{L^2(Q)}^2 \\ & \leq \frac{\varepsilon C}{2} \int_Q |\nabla u_{nk}|^2 dxdt + \frac{(\widehat{G}|\Omega|)^2}{2\varepsilon} \int_0^\omega \int_\Omega |w_k(y, t)|^2 dydt \\ & \quad + \frac{(\sigma^* \|\tilde{\varphi}_0\|_{L^\infty(Q)})^2}{2} \left(\frac{\sigma^*}{\sigma_*} \right)^2 \|\nabla\varphi_0\|_{L^2(Q)}^2 \end{aligned}$$

$$\begin{aligned} &\leq \frac{\varepsilon C}{2} \int_Q |\nabla u_{nk}|^2 dxdt + \frac{(\widehat{G}|\Omega|)^2}{2\varepsilon} [\varepsilon + \int_0^\omega \int_\Omega |w(y,t)|^2 dydt \\ &\quad + \frac{(\sigma^* \|\tilde{\varphi}_0\|_{L^\infty(Q)})^2}{2} \left(\frac{\sigma^*}{\sigma_*}\right)^2 \|\nabla \varphi_0\|_{L^2(Q)}^2. \end{aligned}$$

Thus,

$$\int_Q |\nabla u_{nk}(x,t)|^2 dxdt \leq C$$

and we infer the classical energy estimate

$$(5.6) \quad \int_Q |u_{nk}(x,t)|^2 dxdt + \int_Q |\nabla u_{nk}(x,t)|^2 dxdt \leq C.$$

By virtue of (5.5), (5.6) and (5.1), u_{nkt} is uniformly bounded in the V^* norm. Therefore u_{nk} belongs to a bounded set of D i.e.

$$\|u_{nk}\|_D \leq C.$$

Thus, we can select a subsequences, still denoted by u_{nk} , such that

$$u_{nk} \rightharpoonup u_n, \text{ in } D \text{ as } k \rightarrow \infty.$$

A well known result of [4, Lemma 5.1] guarantees that the sequence u_{nk} is precompact in $L^2(Q)$, then

$$u_{nk} \rightarrow u_n \text{ in } L^2(Q) \text{ and a.e. in } Q.$$

LEMMA 5.2. The operator Θ is continuous.

PROOF. From the preceding result, we have

$$u_{nk} \rightarrow u_n \text{ in } L^2(Q) \text{ and a.e. in } Q.$$

$$u_{nkt} \rightharpoonup u_{nt} \text{ in } L^2(P; W^{-1,2}(\Omega))$$

$$\nabla u_{nk} \rightharpoonup \nabla u_n \text{ in } L^2(P; (L^2(\Omega))^n)$$

$$\nabla \varphi_k \rightarrow \nabla \varphi \text{ in } L^2(P; (L^2(\Omega))^n)$$

$$\varphi_k \rightarrow \varphi \text{ in } V \text{ and a.e. in } Q$$

$$w_k \rightarrow w \text{ in } L^2(Q)$$

$$\sigma(w_k) \rightarrow \sigma(w) \text{ in } L^2(Q)$$

$$I_n(w_k) \rightarrow I_n(w) \text{ in } L^2(Q).$$

These convergences enable us to conclude that $\Theta(w_k) = u_{nk}$ converges strongly to $\Theta(w) = u_n$ in $L^2(Q)$. \square

LEMMA 5.3. There exists a constant $M > 0$ such that

$$\|\Theta(w)\|_{L^2(Q)} \leq M, \text{ for all } w \in L^2(Q).$$

PROOF. The assertion is obtained as above. \square

Since $\Theta(L^2(Q)) \subset D$ and the embedding $D \hookrightarrow L^2(Q)$ is compact, Θ is a compact operator from $L^2(Q)$ into itself. \square

Our main result, is given in the next statement.

THEOREM 5.4. If $H_\sigma) - H_0)$ are fulfilled, there exists at least one weak periodic solution to (1.5)-(1.8).

PROOF. As a consequence of Lemmas 5.2 and 5.3 the mapping Θ is both continuous and compact. Hence, by the Schauder fixed point theorem, Θ has a fixed points which corresponds to a weak periodic solutions to (1.9)-(1.12). \square

As far as previously proved, we can conclude that

$$\begin{aligned} u_{nt} &\rightharpoonup u_t \text{ in } L^2(P; W^{-1,2}(\Omega)) \\ u_n &\rightarrow u \text{ in } L^2(Q) \text{ and a.e. in } Q. \\ \nabla u_n &\rightharpoonup \nabla u \text{ in } L^2(P; (L^2(\Omega))^n) \\ \nabla \varphi_n &\rightharpoonup \nabla \varphi \text{ in } L^2(P; (L^2(\Omega))^n) \\ \varphi_n &\rightarrow \varphi \text{ in } V \text{ and a.e. in } Q \\ w_n &\rightarrow w \text{ in } L^2(Q) \\ \sigma(w_n) &\rightarrow \sigma(w) \text{ in } L^2(Q) \\ I_n(u_n) &\rightarrow \rho \text{ weak-}^* \text{ in } L^\infty(Q), \text{ with } 0 \leq \rho \leq 1. \end{aligned}$$

These u_n are nonnegative. In fact, for $\zeta = u_n^-$, we have

$$\begin{aligned} (5.7) \quad \int_Q u_{nt} u_n^- dxdt &+ \int_Q \nabla u_n \nabla u_n^- dxdt + \int_Q \int_\Omega G(x, y) u_n(y, t) I_n(u_n) u_n^-(x, t) dy dxdt \\ &= \int_Q \sigma(u_n) |\nabla \varphi_n|^2 u_n^- dxdt, \forall \zeta \in V_0 \end{aligned}$$

from which it follows that

$$-\int_Q |\nabla u_n|^2 dxdt = \int_Q \sigma(u_n) |\nabla \varphi_n|^2 u_n^- dxdt.$$

This implies the nonnegativity of u_n . Since $u_n \geq 0$, so is u .

Passing to the limit in

$$\begin{aligned} (5.8) \quad \int_Q u_n \zeta dxdt + \int_Q \nabla u_n \nabla \zeta dxdt + \int_Q \int_\Omega G(x, y) u_n(y, t) I_n(u_n) \zeta(x, t) dy dxdt \\ = - \int_Q \sigma(u_n) \varphi_n \nabla \varphi_n \nabla \zeta dxdt \end{aligned}$$

we have

$$\begin{aligned} (5.9) \quad \int_Q u_t \zeta dxdt + \int_Q \nabla u \nabla \zeta dxdt + \int_Q \int_\Omega G(x, y) u(y, t) \rho(x, t) \zeta(x, t) dy dxdt \\ = - \int_Q \sigma(u) \varphi \nabla \varphi \zeta dxdt, \forall \zeta \in V_0. \end{aligned}$$

We observe that $\rho(x_0, t_0) = 1$ if $u(x_0, t_0) > 0$ thus $\rho(x, t)(v - u) \leq (v - u)$ for all $v \geq 0$. If we replace ζ with $v - u$ in (5.9) with $v \in K$, then (2.2) is satisfied. □

6. Regularity.

This part of paper is devoted to the regularity of weak periodic solutions (u, φ) to problem (1.5)-(1.8). We will prove the Hölder continuity of (u, φ) by means of some a priori estimates in the Campanato spaces.

LEMMA 6.1. Let φ_n be the weak periodic solution of (1.11)-(1.12). If $0 \leq \mu_0 < n - 2 + 2\delta_0, \delta_0 \in (0, 1)$ one has

$$(6.1) \quad \|\nabla \varphi_n(t)\|_{2, \mu_0, \Omega} \leq c(\|\varphi_0(t)\|_{2, \mu_0, \Omega} + \|\nabla \varphi_0(t)\|_{2, (\mu_0 - 2)^+, \Omega}),$$

for a.e. $t \in P$.

PROOF. From a result of [6] we get

$$\|\nabla \varphi_n(t)\|_{2, \mu_0, \Omega} \leq c(\|\varphi_0(t)\|_{2, \mu_0, \Omega} + \|\nabla \varphi_0(t)\|_{2, (\mu_0 - 2)^+, \Omega} + \|\varphi_n(t)\|_{H^1(\Omega)})$$

a.e. t . Since $\varphi_0(t), \nabla\varphi_0(t) \in L^\infty(\Omega)$ for a.e. t and the fact that $L^\infty(\Omega)$ is a multiplier for $\mathcal{L}^{2,\mu_1}(\Omega)$ for $\mu_1 < n$, by the inclusion $\mathcal{L}^{2,(\mu_0-2)^+}(\Omega) \hookrightarrow L^2(\Omega)$ and (5.3) we infer that

$$\|\nabla\varphi_n(t)\|_{L^2(\Omega)} \leq \left(\frac{\sigma^*}{\sigma_*}\right) \|\nabla\varphi_0(t)\|_{2,(\mu_0-2)^+, \Omega}$$

for a.e. t , which implies

$$\|\varphi_n(t)\|_{H^1(\Omega)} \leq c\|\varphi_0(t)\|_{2,\mu_0,\Omega} + \left(\frac{\sigma^*}{\sigma_*}\right) \|\nabla\varphi_0(t)\|_{2,(\mu_0-2)^+, \Omega}.$$

This completes the proof of Lemma 6.1. \square

PROPOSITION 6.2. The weak periodic solutions (u, φ) of (1.5)-(1.8) belong to the space $C^{\alpha,\alpha/2}(\overline{Q})$.

PROOF. From (6.1), one has

$$\|\nabla\varphi_n(t)\|_{2,\mu_0,\Omega} \leq C$$

for a.e. t . The definition of Campanato's spaces gives us

$$(6.2) \quad r^{-(\mu_0+2)} \int_{Q[z_0,r]} |\nabla\varphi_n(x,t)|^2 dx dt \leq C$$

for all $z_0 \in Q$ and $r > 0$.

Being $\sigma(u_n)\varphi_n \in L^\infty(Q)$ and $L^\infty(Q)$ a multiplier of $\mathcal{L}^{2,\mu}(Q)$ if $0 \leq \mu \leq n + 2\delta_0$, then $\sigma(u_n)\varphi_n \nabla\varphi_n \in \mathcal{L}^{2,\mu}(Q)$ and the inequality (6.2) implies that

$$\|\nabla\varphi_n\|_{2,\mu}^2 \leq C.$$

By the comparison principle, we have that the nonnegative weak periodic solution ψ of problem

$$(6.3) \quad \begin{aligned} \psi_t - \Delta\psi &= \operatorname{div}(\sigma(u_n)\varphi_n \nabla\varphi_n) \text{ in } Q, \\ \psi &= 0 \text{ on } \Sigma, \end{aligned}$$

is a supersolution for (5.2), so that

$$(6.4) \quad 0 \leq u_n \leq \psi.$$

According to a result in [7, Theorem 1], we get

$$\|\nabla\psi\|_{2,\mu} \leq C$$

for $0 < \mu \leq n + 2\alpha$ and ψ belongs to $\mathcal{L}^{2,\mu+2}(Q)$ (see [7, Lemma 2.6]). Because of Proposition A, one has that $\psi \in C^{\alpha,\alpha/2}(\bar{Q})$ and consequently

$$\|\psi\|_{L^\infty(Q)} \leq C.$$

The boundedness of ψ implies that $u_n \in L^\infty(Q)$. Theorem 1 in [7] allows to conclude that

$$(6.5) \quad \|\nabla u_n\|_{2,\mu} \leq c(\|\sigma(u_n)\varphi_n \nabla \varphi_n\|_{2,\mu} + \|I_n(u_n)\|_{2,(\mu-2)^+} + \|u_n\|_V)$$

and the embedding of $L^\infty(Q) \hookrightarrow \mathcal{L}^{2,\mu}(Q)$ yields

$$\left\| \int_{\Omega} I_n(u_n)(G(x, y)u_n(y, t)dy) \right\|_{2,(\mu-2)^+} \leq c \left\| \int_{\Omega} G(x, y)u_n(y, t)dy \right\|_{L^\infty(Q)} \leq C.$$

Thus, if $0 < \mu \leq n + 2\alpha$ by (6.5), we obtain

$$\|\nabla u_n\|_{2,\mu} \leq C$$

and $u_n \in \mathcal{L}^{2,\mu+2}(Q)$, (see [7, Lemma 2.6]). Since we showed that $u_n \rightarrow u$ and $\varphi_n \rightarrow \varphi$ a.e. in Q with u_n, φ_n in $L^\infty(Q)$, we conclude that $u, \varphi \in \mathcal{L}^{2,\mu}(Q)$ for $\mu = n + 2 + 2\alpha$. Finally, by Proposition A $u, \varphi \in C^{\alpha,\alpha/2}(\bar{Q})$ for $\alpha = \frac{\mu - n}{2}$. \square

REFERENCES

- [1] W. ALLEGRETTO - Y. LIN - S. MA, *Hölder continuous solutions of an obstacle thermistor problem*, Discrete and Continuous Dynamic Systems-Series B, vol. 4, n. 4 (2004), pp. 983–997.
- [2] W. ALLEGRETTO - Y. LIN - S. MA, *Existence and long time behaviour of solutions to obstacle thermistor equations*, Discrete and Continuous Dynamic Systems-Series B, vol. 8, n. 8 (2002), pp. 757–780.
- [3] M. GIAQUINTA, *Multiple integrals in the calculus and nonlinear elliptic systems*, Annals of Mathematics Studies, Princeton, New Jersey 1983.
- [4] J. L. LIONS, *Quelques méthodes de résolution des problèmes aux limites non-linéaires*, Dunod, Paris 1968.
- [5] J. L. LIONS - E. MAGENES, *Non-homogeneous boundary value problems and applications*, vol. I, Springer Verlag Berlin Heidelberg New York 1972.
- [6] H. XIE, $\mathcal{L}^{2,\mu}(\Omega)$ estimate to the mixed boundary value problem for second order elliptic equations and its applications in the thermistor problem, Nonlinear Anal., 24 (1995), pp. 9–27.
- [7] H. YIN, $\mathcal{L}^{2,\mu}(Q)$ -estimates for parabolic equations and applications, J. Partial Diff. Eqn., 10 (1997), pp. 31–44.

