

Astérisque

NIGEL J. HITCHIN

The Yang-Mills equations and the topology of 4-manifolds

Astérisque, tome 105-106 (1983), Séminaire Bourbaki,
exp. n° 606, p. 167-178

http://www.numdam.org/item?id=SB_1982-1983__25__167_0

© Société mathématique de France, 1983, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

THE YANG-MILLS EQUATIONS
AND THE TOPOLOGY OF 4-MANIFOLDS

[after Simon K. Donaldson]

by Nigel J. HITCHIN

§ 1. The result

(1.1) THEOREM (S.K. Donaldson [8]).— *Let X be a compact, smooth, simply connected, oriented 4-manifold such that the intersection form Q on $H^2(X, \mathbb{Z})$ is positive definite. Then there exists an integral basis for $H^2(X, \mathbb{Z})$ such that*

$$Q(u, u) = u_1^2 + u_2^2 + \dots + u_r^2 .$$

This theorem should be contrasted with

(1.2) THEOREM (M.H. Freedman [9]).— *Let Q be any unimodular quadratic form over \mathbb{Z} . Then there exists a compact, simply connected, topological 4-manifold X such that Q is equivalent to the intersection form on $H^2(X, \mathbb{Z})$.*

There are sufficient examples of definite unimodular forms (see [17]) to see that Donaldson's theorem imposes strong restrictions on smooth 4-manifolds.

(1.3) Proof of Theorem (1.1)

Let $r = \text{rank } H^2(X, \mathbb{Z})$ and $2n = \# \{u \in H^2(X, \mathbb{Z}) \mid Q(u, u) = 1\}$. The proof consists of constructing (as in § 3-§ 7) an oriented cobordism between X and n copies of $\mathbb{C}P^2$. Let p of these have the canonical orientation of the complex structure and $q = n - p$ the opposite orientation. Then

(i) By the cobordism invariance of signature,

$$r = \text{Sign } X = (p - q) \text{Sign } \mathbb{C}P^2 = p - q \leq n .$$

(ii) Let $\{\pm x_1, \pm x_2, \dots, \pm x_n\} = \{u \in H^2(X, \mathbb{Z}) \mid Q(u, u) = 1\}$, then $Q(x_i, x_j) \in \mathbb{Z}$ but by the Cauchy-Schwarz inequality $|Q(x_i, x_j)| < 1$ if $i \neq j$. Hence $\{x_1, \dots, x_n\}$ is orthonormal and $n \leq r$.

(iii) From (i) and (ii) $n = r$ and $\{x_1, \dots, x_n\}$ is an orthonormal basis for $H^2(X, \mathbb{R})$. Thus for $u \in H^2(X, \mathbb{Z})$, $u = \sum_{i=1}^n Q(u, x_i) x_i = \sum_{i=1}^n u_i x_i$ with $u_i \in \mathbb{Z}$ and $\{x_1, \dots, x_n\}$ is a basis for $H^2(X, \mathbb{Z})$. Hence $Q(u, u) = \sum_{i=1}^n u_i^2$.

§ 2. Background

(2.1) Let X be an oriented riemannian 4-manifold. A 2-form $\alpha \in \Omega^2$ is said to be *self-dual* (resp. *anti-self-dual*) if $*\alpha = \alpha$ (resp. $*\alpha = -\alpha$) where $*$: $\Omega^2 \rightarrow \Omega^2$ denotes the Hodge star operator.

Let G be a compact Lie group and P a principal G -bundle over X . A connection A on P has curvature $F(A) \in \Omega^2(\mathfrak{g})$ where \mathfrak{g} denotes the vector bundle associated to P by the adjoint representation. For any bundle V associated to P a connection A defines a differential operator $d_A : \Omega^p(V) \rightarrow \Omega^{p+1}(V)$. The metric on X defines the formal adjoint $d_A^* : \Omega^{p+1}(V) \rightarrow \Omega^p(V)$. The *Bianchi identity*, satisfied by all connections, is $d_A^* F(A) = 0$. The *Yang-Mills equations* are $d_A^* F(A) = 0$.

A connection A on P is said to be *self-dual* if $F(A) = *F(A)$. In this case $d_A^* F(A) = *d_A *F(A) = *d_A F(A) = 0$ by the Bianchi identity, so a self-dual connection automatically satisfies the Yang-Mills equations.

The Yang-Mills equations describe the critical points for the Yang-Mills functional (or action).

$$\|F(A)\|_{L^2}^2 = \int_X |F(A)|^2 d\mu .$$

The self-dual connections give the absolute minimum for compact X which, if $G = SU(2)$, may be expressed via the Chern-Weil theorem as $-8\pi^2 c_2(P)$ where $c_2(P)$ is the 2nd Chern class of the associated rank 2 vector bundle.

The Yang-Mills functional and Yang-Mills equations are invariant under (i) conformal changes of the metric on X (ii) automorphisms of the principal bundle P ("gauge transformations").

(2.2) The initial mathematical development of the study of self-dual connections, motivated by the interest of mathematical physicists, concentrated on the case $X = S^4$ and an explicit description of all solutions was possible [2] using the twistor approach of R. Penrose and R.S. Ward [6] which converted the problem into one of holomorphic bundles on \mathbb{CP}^3 .

More recently the self-duality equations have been studied on more general 4-manifolds. There are three major lines of thought which have spurred this progress :

(2.3) If X is a Kähler manifold, the space of anti-self-dual 2-forms $\Omega_0^{2,1}$, the space of primitive 2-forms of type (1,1). A vector bundle with an anti-self-dual connection is then automatically endowed with a holomorphic structure (see [3]) and is moreover *stable* in the sense of Mumford and Takemoto (see [8], [11]). Converse results have been conjectured and in some cases proved ([13], [8]).

(2.4) The analysis of self-dual connections has been pushed forward by the funda-

mental results of K.K. Uhlenbeck ([20], [21]). Amongst these is the following *removable singularity theorem* : If A is an $SU(2)$ connection (on the trivial bundle) over the punctured ball $B^4 \setminus \{0\}$, self-dual with respect to some smooth riemannian metric on B^4 and with finite action ; then there is a bundle automorphism $g : B^4 \setminus \{0\} \rightarrow SU(2)$ such that $g(A)$ extends smoothly over B^4 .

(2.5) The existence of self-dual connections is assured under very general circumstances by a theorem of C.H. Taubes [19] : Let X be a compact, oriented, riemannian 4-manifold with positive definite intersection form Q , and let P be a principal $SU(2)$ bundle over X with $c_2(P) \leq 0$. Then P admits an irreducible self-dual connection. Taubes' construction makes use of an implicit function theorem which involves L^p estimates on curvature. It should be noted that anti-self-dual harmonic 2-forms may certainly obstruct the existence of self-dual connections, as can be seen by considering $\mathbb{C}P^2$ with opposite orientation. There are no stable rank 2 bundles on $\mathbb{C}P^2$ with $c_2(P) = 1$ [16] and hence by (2.3) no anti-self-dual connections. Taubes' hypotheses and result are the starting point for Donaldson's theorem.

(2.6) As an example of a self-dual connection, take $X = \mathbb{R}^4$ and $G = SU(2)$. Then in terms of a quaternionic coordinate $x \in \mathbb{H} \cong \mathbb{R}^4$ and using the isomorphism $SU(2) \cong \text{Im } \mathbb{H}$ the 1-instanton [7] solution of the self-duality equations is given by

$$A_\lambda = \text{Im} \left(\frac{x d\bar{x}}{\lambda^2 + |x|^2} \right) \quad \text{with} \quad F(A_\lambda) = \frac{\lambda^2 dx \wedge d\bar{x}}{(\lambda^2 + |x|^2)^2}$$

and action $8\pi^2$.

(2.7) PROPOSITION.— Let A be a self-dual $SU(2)$ connection on \mathbb{R}^4 with action $8\pi^2$. Then up to a gauge transformation and a translation of \mathbb{R}^4 , A is equal to A_λ for some $\lambda \in \mathbb{R}$.

Proof.— By conformal invariance and stereographic projection A is defined on $S^4 \setminus \{x\}$, and by the removable singularity theorem is defined on a bundle $P \rightarrow S^4$. Now use [3] § 9 or [2] or [6].

§ 3. The moduli space

(3.1) The cobordism in the proof of (1.1) is modelled on a moduli space of self-dual connections whose general structure is described next.

Let X be as in Theorem (1.1), and given a riemannian metric. Let P be a principal $SU(2)$ bundle over X with $c_2(P) = -1$. Using the covariant derivative of a fixed smooth connection A_0 on P , one may define Sobolev spaces $L^p_q(V)$ of sections of any associated vector bundle V .

Let \mathcal{C} denote the affine space of connections on P differing from A_0 by

an element of $L^2_3(\Omega^1(g))$, and let \mathcal{G} denote the group of L^2_4 sections of $PX_{\text{Ad}}G$ ($\subset \text{End } V$ for some faithful representation). Then \mathcal{G} is a Banach Lie group of gauge transformations acting smoothly on \mathcal{A} by $g(A) = A - (d_A g)g^{-1}$. Let \mathcal{B} denote the quotient space with projection $p : \mathcal{A} \rightarrow \mathcal{B}$, and $p(A) = [A]$.

(3.2) Recall that a connection on P is *reducible* if its holonomy group is a proper subgroup of $SU(2)$. Since X is simply-connected and P is topologically non-trivial, the only possible reduction is to $U(1) \subset SU(2)$. Let $\Gamma_A \subset \mathcal{G}$ denote the subgroup of covariant constant sections with respect to the connection A . Then A is reducible iff $\Gamma_A \cong U(1)$. The equivalence classes of irreducible connections form an open subset $\mathcal{B}^* \subset \mathcal{B}$.

(3.3) PROPOSITION.— (i) \mathcal{B} is a Hausdorff space in the quotient topology.

(ii) \mathcal{B}^* is a Banach manifold with charts constructed from the slices

$T_{A,\epsilon} = \{A+a \mid d_A^* a = 0, \|a\|_{L^2_3} < \epsilon\}$ of the action of \mathcal{G} .

(iii) $p : p^{-1}(\mathcal{B}^*) \rightarrow \mathcal{B}^*$ is a principal $\mathcal{G}/\pm 1$ bundle with a connection defined by the slices.

(iv) If A is reducible, Γ_A acts on $T_{A,\epsilon}$ and the map $T_{A,\epsilon}/\Gamma_A \rightarrow \mathcal{B}$ is a homeomorphism to a neighbourhood of $[A] \in \mathcal{B}$, smooth away from the fixed point set.

Proof.— Standard methods (see [3], [12], [14]) using Banach space inverse and implicit function theorems.

(3.4) Let $\mathcal{M} \subset \mathcal{B}$ denote the subspace of equivalence classes of *self-dual* connections on P . \mathcal{M} is the *moduli space*. If $A \in \mathcal{A}$ is reduced to a connection on a principal $U(1)$ bundle $Q \subset P$, then (since $\pi_1(X) = 0$) its equivalence class is determined by its curvature $F(A) \in \Omega^2$. If A is self-dual, $F(A)$ is a self-dual closed 2-form, hence harmonic. By Hodge theory $F(A)$ is determined by its cohomology class $2\pi ic_1(Q)$. The reduction to $U(1)$ is well-defined modulo the Weyl group, so $[A] \in \mathcal{M}$ is determined by $\pm c_1(Q)$. Since $c_2(P) = -c_1(Q)^2 = -1$ there are n distinguished points in \mathcal{M} representing the reducible self-dual connections, where $2n = \# \{u \in H^2(X, \mathbb{Z}) \mid Q(u, u) = 1\}$. From (2.5) there are also irreducible connections.

(3.5) If A is a self-dual connection on P , then there exists an elliptic complex [3]

$$\Omega^0(g) \xrightarrow{d_A} \Omega^1(g) \xrightarrow{d_A^-} \Omega^2(g)$$

where d_A^- is the projection of d_A onto the anti-self-dual 2-forms. Let H_A^p ($0 \leq p \leq 2$) denote the associated harmonic spaces, then by the Atiyah-Singer index theorem (see [3])

$$-\sum_{p=0}^2 (-1)^p \dim H_A^p = 8|c_2(P)| - \frac{3}{2}(\chi(X) - \text{Sign}(X)) = 5.$$

(3.6) PROPOSITION.— Let A be a self-dual connection on P .

Then there exists a neighbourhood U of $0 \in H_A^1$ and a smooth map $\phi : U \rightarrow H_A^2$ such that :

(i) if A is irreducible, a neighbourhood of $[A] \in \mathcal{M}$ is diffeomorphic to $\phi^{-1}(0) \subseteq H_A^1$.

(ii) if A is reducible, a neighbourhood of $[A] \in \mathcal{M}$ is diffeomorphic to $\phi^{-1}(0)/\Gamma_A$.

Proof.— The connection $A+a$ is self-dual iff

$$\Phi(A+a) = F_-(A+a) = d_A^- a + \frac{1}{2}[a, a] = 0 \in L_2^2(\Omega_2^2(g)) .$$

Restricted to a slice $T_{A, \epsilon}$ the derivative $D\Phi_A$ of Φ at A is the Fredholm operator $d_A^- : \text{Ker } d_A^*(\subseteq L_3^2(\Omega^1(g))) \rightarrow L_2^2(\Omega_2^2(g))$, and so Φ is a Fredholm map ([1], [18]). After a local diffeomorphism Φ may be represented as $\Phi(x) = (D\Phi_A)_x + \phi(x)$. The argument is analogous to the methods applied to moduli of complex structures [10].

(3.7) As a consequence of (3.5) and (3.6), if A is irreducible and $H_A^2 = 0$, then \mathcal{M} is a smooth 5-manifold in a neighbourhood of $[A]$. A particular case when this holds for all irreducible A is when the underlying metric on X is self-dual with positive scalar curvature (see [3]). Note that if A is reducible, Γ_A acts on H_A^1 by complex multiplication ($b_1(X) = 0$) so that if $H_A^2 = 0$, $H_A^1/\Gamma_A \cong \mathbb{C}^3/S^1$ from the index theorem and $\dim H_A^0 = \dim \Gamma_A = 1$.

§ 4. A key result

(4.1) An important tool in understanding the global structure of the moduli space is the following : (see also [15]).

(4.2) PROPOSITION.— Let $\tilde{A}_i \in \mathcal{A}$ be a sequence of self-dual connections on P . Then there is a subsequence such that either :

(i) each \tilde{A}_i is gauge equivalent to $A_i \in \mathcal{A}$ converging in C^∞ to a self-dual connection A_∞ on P , and hence $[\tilde{A}_i] \rightarrow [A_\infty] \in \mathcal{M}$.

or

(ii) there is a point $x \in X$ and trivializations ρ_i of $P|_K$ on the complement K of any geodesic ball about x such that $\rho_i^* \tilde{A}_i \rightarrow \mathfrak{S}$ (the trivial flat connection) in $C^\infty(K)$.

Proof.— The proof uses two lemmas :

(4.3) Lemma.— Given $L, C > 0$ let $\{f_i\}$ be a sequence of integrable functions on X with $f_i \geq 0$ and $\int_X f_i d\mu \leq L$. Then there exists a subsequence, a finite set $\{x_1, \dots, x_\rho\} \subset X$ and a countable collection $\{B_\alpha\}$ of geodesic balls in X such that the half-sized balls cover $X \setminus \{x_1, \dots, x_\rho\}$ and for each α , $\limsup \int_{B_\alpha} f_i d\mu < C$.

Proof.— Elementary : the x_i 's are characterized by the property that each lies in

no ball with $\limsup \int_B f_i d\mu \leq \frac{1}{2}C$.

(4.4) Lemma.— Let h_i be a sequence of metrics on B^4 , sufficiently close to the Euclidean metric, and converging in $C^\infty(\bar{B}^4)$ to h_∞ . Let \tilde{A}_i be a sequence of connections on the trivial bundle over B^4 with \tilde{A}_i self-dual with respect to h_i . Then there is a constant C (independent of h_i and \tilde{A}_i) such that if $\int_{B^4} |F(\tilde{A}_i)|^2 d\mu \leq C$, there is a subsequence such that A_i (gauge equivalent to \tilde{A}_i) converges in $C^\infty(\frac{1}{2}\bar{B}^4)$ to A_∞ , a connection which is self-dual with respect to h_∞ .

Proof.— Consequence of ([21] Theorem (1.3)).

(4.5) To obtain (4.2) first consider a geodesic coordinate system χ on a geodesic ball $B \subset X$ of radius r . Thus χ defines a diffeomorphism $\chi : B_r^4 \rightarrow B$ from the euclidean ball of radius r to B . Pulling back the metric h , and putting it on the Euclidean unit ball by dilation gives a metric

$$h_r = \chi^*h(rx) = r^2(\delta_{ij} + r^2O(|y|^2))dy_i dy_j.$$

Choose r small enough that the metric $r^{-2}h_r$ on B^4 satisfies the condition for (4.4). By conformal invariance each \tilde{A}_i is self-dual with respect to h_r .

Now in Lemma (4.3) take the constant C from (4.4), $f_i = |F(\tilde{A}_i)|^2$ and $L = 8\pi^2$. Thus from (4.4) on each ball $\frac{1}{2}\bar{B}_\alpha$ some subsequence converges (after gauge transformations) to $A_\infty(\alpha)$. By a diagonal argument the convergence may be achieved simultaneously for all α .

The gauge transformations introduced in the above process give rise to connection matrices $A_i(\alpha) \rightarrow A_\infty(\alpha)$ in $C^\infty(\frac{1}{2}B_\alpha)$ and transition functions $g_i(\alpha, \beta) : \frac{1}{2}B_\alpha \cap \frac{1}{2}B_\beta \rightarrow SU(2)$ satisfying :

$$(4.6) \quad A_i(\alpha) = -dg_i(\alpha, \beta)g_i(\alpha, \beta)^{-1} + g_i(\alpha, \beta)A_i(\beta)g_i(\alpha, \beta)^{-1}.$$

The compactness of $SU(2)$ gives a uniform bound to dg_i in (4.6) and so one can find a uniformly convergent subsequence. Repeatedly applying (4.6) gives convergence in C^∞ , and using a diagonal argument one obtains a subsequence

$$(A_i(\alpha), g_i(\alpha, \beta)) \rightarrow (A_\infty(\alpha), g_\infty(\alpha, \beta))$$

for all (α, β) simultaneously. This represents a self-dual connection on a bundle Q over $X \setminus \{x_1, \dots, x_\ell\}$. Furthermore, if $K \subset X \setminus \{x_1, \dots, x_\ell\}$ is compact then by induction on the number of balls $\frac{1}{2}B_\alpha$ covering K (see [21] Sect. 3) one obtains isomorphisms $\rho_i : Q|_K \rightarrow P|_K$ such that $\rho_i^* : A_i \rightarrow A_\infty$ in $C^\infty(K)$.

(4.7) Let $B_j^!$ be a small punctured ball centred on x_j ($1 \leq j \leq \ell$). Since $\int_{B_j^!} |F(\tilde{A}_i)|^2 d\mu \leq 8\pi^2$, by Fatou's lemma $\int_{B_j^!} |F(A_\infty)|^2 d\mu \leq 8\pi^2$. Hence by the removable singularity theorem (2.4) the connection A_∞ and bundle Q extend over X . By the definition of x_j , $\lim \int_{B_j^!} |F(\tilde{A}_i)|^2 d\mu > \frac{1}{2}C$ for all balls $B_j^!$ hence for a sufficiently small ball

$$\int_{B_j^!} |F(A_\infty)|^2 d\mu < \liminf \int_{B_j^!} |F(\tilde{A}_1)|^2 d\mu .$$

(4.8) On the other hand, since all connections are self-dual these integrands are Chern forms. They may therefore be evaluated mod. $8\pi^2\mathbb{Z}$ by boundary integrals (Chern-Simons invariants). Hence by uniform convergence on the boundary ∂B_j ,

$$\int_{B_j^!} |F(A_\infty)|^2 d\mu = \liminf \int_{B_j^!} |F(\tilde{A}_1)|^2 d\mu \quad \text{mod. } 8\pi^2\mathbb{Z} .$$

(4.9) However, since $\int_{B_j^!} |F(A_\infty)|^2 d\mu \geq 0$ and $\int_{B_j^!} |F(\tilde{A}_1)|^2 d\mu \leq 8\pi^2$ the only possibilities from (4.7) and (4.8) are :

- (i) $\ell = 0$ or
- (ii) $\liminf \int_{B_j^!} |F(\tilde{A}_1)|^2 d\mu = 8\pi^2$ and $\int_X |F(A_\infty)|^2 d\mu < 8\pi^2$ and hence Q is trivial and A_∞ flat. Thus Proposition (4.2) follows.

(4.10) The proposition shows that a self-dual connection on P can only degenerate by having its curvature concentrate in the neighbourhood of a point. An example is the instanton A_λ in (2.6) as $\lambda \rightarrow 0$.

§ 5. The boundary of \mathcal{M}

(5.1) Let $\beta : \mathbb{R} \rightarrow \mathbb{R}$ be a bump function approximating and dominated by $\chi_{[-1,1]}$ and set $R_A(x, s) = \int_X \beta(d(x, y)/s) |F(A)|^2 d\mu_y$, where $d(x, y)$ is the geodesic distance in X . Then define

$$(5.2) \quad \lambda(A) = K^{-1} \min \{s \mid \exists x \text{ with } R_A(x, s) = 4\pi^2\}$$

where K is chosen so that $\lambda(A_1) = 1$ for the instanton A_1 . Donaldson introduces this convenient but ad hoc function as a measure of the concentration of curvature : if β is replaced by $\chi_{[-1,1]}$ then $\lambda(A)$ becomes the radius of the smallest ball containing half the action. In any case a ball of radius $\lambda(A)$ contains more than half the action and hence any sequence $[A_i] \in \mathcal{M}$ without convergent subsequences has $\lambda(A_i) \rightarrow 0$ from (4.2). It is thus a measure of the distance from the boundary.

(5.3) PROPOSITION.— *There exists $\lambda_0 > 0$ such that if A is a self-dual connection on P with $\lambda(A) < \lambda_0$, then the minimum in (5.2) is attained at a unique point $x(A) \in X$.*

Proof.— Take a small geodesic ball of radius r centred on a minimum x for A , and pull back the metric and connection as in (4.5) to the Euclidean ball of radius $r/\lambda(A)$. For each sequence of connections with $\lambda(A_i) \rightarrow 0$, the pulled-back connections \hat{A}_i satisfy $\lambda(\hat{A}_i) = 1$ by construction and applying (4.4) and (4.2) there is a subsequence converging to a self-dual connection on \mathbb{R}^4 . From the classification (2.7) and normalization this is the instanton A_1 . Since $\lambda(\hat{A}_1) = 1$, from

(4.2) every subsequence converges and since the limit is unique, $\hat{A}_i \rightarrow A_1$ as $\lambda(A_i) \rightarrow 0$. Now the function R_{A_1} has a unique non-degenerate minimum so for sufficiently small $\lambda(A)$, so will $R_{\hat{A}}$. Any two minima for A must however be separated by a distance of at most $2\lambda(A)$, since the ball of radius $\lambda(A)$ about each contains more than half the action, thus a unique minimum for $R_{\hat{A}}$ implies a unique one for R_A .

Note how the connectedness of the moduli space for R^4 is essential for this argument.

(5.4) Let $\mathcal{M}_{\lambda_0} = \{[A] \in \mathcal{M} \mid \lambda(A) < \lambda_0\}$, and define $p : \mathcal{M}_{\lambda_0} \rightarrow X \times (0, \lambda_0)$ by $p(A) = (x(A), \lambda(A))$.

- (5.5) PROPOSITION.— (i) \mathcal{M}_{λ_0} is compact.
 (ii) \mathcal{M}_{λ_0} is a smooth manifold.
 (iii) p is a smooth covering map.

Proof.— (i) Immediate from Proposition (4.2).

(ii) As $\lambda(A) \rightarrow 0$, $[A] \rightarrow \mathfrak{G}$ in $C^\infty(X \setminus B(x(A), r))$ from (4.2). Then using an argument of Taubes [19], $H_A^2 = 0$. The result follows from (3.6).

(iii) p is smooth because the minimum of R_A is non-degenerate, and proper by (4.2). Thus one only needs to check that the derivative of p is an isomorphism. Taubes' implicit function theorem provides an inverse.

(5.6) PROPOSITION.— p is a diffeomorphism.

Proof.— This is the most technical part of Donaldson's proof, and involves delicate curvature estimates. The idea is to show that any two self-dual connections A, B with $x(A) = x(B)$ and $\lambda(A) = \lambda(B)$ sufficiently small may be joined by a short path in \mathcal{M} (see [8]).

§ 6. Perturbation of \mathcal{M}

(6.1) If $H_A^2 = 0$ for all self-dual connections then \mathcal{M} is a smooth manifold except at the $n(Q)$ points corresponding to the reducible connections. This may not be true in general and there may be a subset $K \subset \mathcal{M}$ (compact from (5.5)) for which $H_A^2 \neq 0$. A perturbation of \mathcal{M} is then necessary to obtain a manifold.

(6.2) Perturbation around the reducible connections is dealt with in a straightforward manner: the finite-dimensional map $\phi(x)$ in the decomposition $\Phi(x) = (D\Phi_A)x + \phi(x)$ is modified by a nearby map with surjective derivative. Then, as in (3.6) a neighbourhood of $[A]$ is diffeomorphic to \mathbb{C}^3/S^1 — a cone on $\mathbb{C}P^2$. One may assume, then, that $K \subset \mathcal{M} \cap \mathcal{B}^*$.

(6.3) The group $\mathcal{G}/\pm 1$ acts on the Banach spaces $L_3^2(\Omega_-^2(g))$ and $L_2^2(\Omega_-^2(g))$ and

associated to the principal $\mathcal{G}/\pm 1$ bundle $p^{-1}(\mathcal{B}^*)$ over \mathcal{B}^* one obtains vector bundles $\mathcal{L}^3 \subset \mathcal{L}^2$ with norms and connections. There is a canonical section $\Phi = F_-(A)$ of \mathcal{L}^2 and one seeks perturbations $\sigma \in C^\infty(\mathcal{B}^*, \mathcal{L}^3)$, such that $\Phi + \sigma$ vanishes non-degenerately.

(6.4) PROPOSITION.— *There exists $\sigma \in C^\infty(\mathcal{B}^*, \mathcal{L}^3)$, supported in a neighbourhood of K , such that $(\Phi + \sigma)^{-1}(0)$ is a smooth 5-manifold.*

Proof.— Covering K with a finite number of slices $T_{A, \varepsilon}$ and shrinking, take open sets U_1, U_2 with $K \subset U_1$ and $\bar{U}_1 \subset U_2$, and let σ be a bounded section of \mathcal{L}^3 supported in U_2 . Then $\hat{K} = \{[A] \in \bar{U}_1 \mid \|(\Phi + \sigma)(A)\|_{L^2_3} \leq \varepsilon\}$ is compact. This follows from the fact that U_2 is covered by a finite number of slices and on each one $\Phi(A) = d^-_A a + \frac{1}{2}[a, a] + \sigma(A)$ with $d^*_A a = 0$ and $\|a\|_{L^2_3} < \varepsilon$. Thus L^2_3 bounds on $\sigma(A)$, a and $(\Phi + \sigma)(A)$ give an L^2_3 bound on $(d^-_A + d^*_A)a$ and so by ellipticity an L^2_4 bound on a . Since $L^2_4 \subset L^2_3$ is compact the statement follows. Thus if $\Phi + \sigma$ vanishes non-degenerately in \bar{U}_1 , so do nearby sections $\Phi + \sigma'$ in the topology of uniform convergence of σ and its derivative on compact sets.

The space of such non-degenerate perturbations is also dense: at each point take a slice on which there is a decomposition $\Phi + \sigma = L + \phi$ where L is linear and ϕ finite dimensional. By compactness, take a finite subcovering and modify $\Phi + \sigma$ by subtracting a regular value of ϕ , extended by a bump function. By Sard's theorem such perturbations can be made arbitrarily close in L^2_3 norm to $\Phi + \sigma$.

The section Φ itself vanishes non-degenerately outside \bar{U}_1 . By the density argument choose a perturbation σ sufficiently small that $\Phi + \sigma$ (by the openness argument on $U_2 \setminus \bar{U}_1$) vanishes non-degenerately on $\overline{U_2 \setminus \bar{U}_1}$. Then $\Phi + \sigma$ is non-degenerate everywhere. Let $\mathcal{M}^\sigma = (\Phi + \sigma)^{-1}(0)$, a 5-manifold with n quotient singularities \mathbb{E}^3/S^1 and boundary X .

§ 7. Orientability of \mathcal{M}^σ

(7.1) On the manifold $\mathcal{M}^\sigma \cap \mathcal{B}^*$ one must consider the Stiefel-Whitney class $w_1(\text{Ker } \nabla(\Phi + \sigma))$. The singular points can be avoided by using the gauge transformations $\mathcal{G}_0 \subset \mathcal{G}$ which are the identity at a fixed point $x_0 \in X$. These then act freely on \mathcal{B} to give quotient $\hat{\mathcal{B}} \xrightarrow{\pi} \mathcal{B}$. Over \mathcal{B}^* , π gives a principal $SO(3)$ bundle, so $T\mathcal{M}^\sigma \cap \mathcal{B}^*$ is orientable iff its pull back to $\pi^{-1}(\mathcal{M}^\sigma \cap \mathcal{B}^*)$ is.

(7.2) The vector bundle $\text{Ker } \nabla(\Phi + \sigma)$ restricted to any compact subset $Y \subset \pi^{-1}(\mathcal{M}^\sigma \cap \mathcal{B}^*)$ defines an element of $KO(Y)$. This is the *index class* [5] of the family of Fredholm operators $d^*_A + d^-_A + (\nabla\sigma)A$, which by considering the deformation $d^*_A + d^-_A + t(\nabla\sigma)A$, $0 \leq t \leq 1$, is independent of σ . Since w_1 factors

through KO , the orientability can be decided by considering $\text{ind}(d_A^* + d_A^-) \in KO(Y)$ where Y is a loop. Since this is now defined for all equivalence classes of connections, the loop may be deformed in $\hat{\mathcal{B}}$.

(7.3) If $SU(2)$ is embedded in $SU(3)$ in the standard way, the Lie algebra bundle $\tilde{\mathcal{G}}$ of the associated $SU(3)$ connection \tilde{A} splits as $\tilde{\mathcal{G}} = \mathfrak{g} \oplus \mathbb{R} \oplus V$ where V is a complex rank 2 bundle and \mathbb{R} a trivial bundle, all preserved by the connection. Thus $w_1(\text{ind}(d_A^* + d_A^-)) = w_1(\text{ind}(d_A^* + d_A^-))$ and so the loop may be deformed in the space $\hat{\mathcal{B}}_3$ of equivalence classes of $SU(3)$ connections.

(7.4) PROPOSITION.— $\pi_1(\hat{\mathcal{B}}_3) = 0$.

Proof.— Since the group \mathcal{G}_0 of $SU(3)$ gauge transformations preserving x_0 acts freely, $\pi_1(\hat{\mathcal{B}}_3) \cong \pi_0(\mathcal{G}_0)$. The principal bundle P is trivial on the complement of a point and in particular on the 2-skeleton of X . Since $\pi_2(SU(3)) = 0$ any element of \mathcal{G}_0 can be deformed to one which is the identity on the 2-skeleton. Collapsing the 2-skeleton of X gives a sphere S^4 . The homotopy type of \mathcal{G}_0 on S^4 is independent of $c_2(P)$ (see [4]), so the question reduces to the trivial bundle. But $\pi_4(SU(3)) = 0$, so \mathcal{G}_0 is connected.

Thus $\mathcal{M}^G \cap \mathcal{B}^*$ is an oriented 5-manifold which, putting in the boundaries of the quotient singularities provides the cobordism of Theorem (1.1).

§ 8. Examples

(8.1) Let $X = S^4$, with the canonical metric. Then any self-dual connection on P is gauge equivalent to f^*A where $f : S^4 \rightarrow \mathbb{H}P^1$ is a conformal map and A is the canonical connection on the quaternionic Hopf bundle. Since isometries of $\mathbb{H}P^1$ preserve A , the moduli space is $SO(5,1)/SO(5) \cong$ hyperbolic 5-space. This is the ball B^5 with boundary S^4 . There are many ways of proving this ([2], [3], [6]).

(8.2) Let $X = \mathbb{C}P^2$ with its canonical metric. In the non-compact component of \mathcal{M} , any connection is gauge equivalent to f^*A where $f : \mathbb{C}P^2 \rightarrow \mathbb{H}P^2$ is equivalent under the action of $SU(3)$ on $\mathbb{C}P^2$ to a map of the form

$$(z, w) \mapsto \left(z, \frac{az + w}{\sqrt{1 - a^2}} \right) \quad a \in [0, 1)$$

in affine coordinates. When $a = 0$ this is the standard embedding $\mathbb{C}P^2 \subset \mathbb{H}P^2$ and gives the reducible connection. The moduli space is a cone on $\mathbb{C}P^2$ where a is essentially the distance from the vertex. This was proved by Donaldson (unpublished) using the algebraic geometry of the flag manifold F_3 , and the Penrose/Ward approach.

§ 9. References

- [1] ABRAHAM R., ROBBIN J. - *Transversal mappings and flows*, Benjamin, New York (1967).
- [2] ATIYAH M.F., HITCHIN N.J., DRINFELD V.G., MANIN, YU I. - *Construction of instantons*, *Phys. Lett.* 65A(1978), 185.
- [3] ATIYAH M.F., HITCHIN N.J., SINGER I.M. - *Self-duality in four-dimensional Riemannian geometry*, *Proc. R. Soc. (Lond.)* A362(1978), 425-461.
- [4] ATIYAH M.F., JONES J.D.S. - *Topological aspects of Yang-Mills theory*, *Commun. Math. Phys.* 61(1978), 97-118.
- [5] ATIYAH M.F., SINGER I.M. - *The index of elliptic operators IV*, *Ann. of Math.* 93(1971), 119-138.
- [6] ATIYAH M.F., WARD R.S. - *Instantons and algebraic geometry*, *Commun. Math. Phys.* 55(1977), 117-124.
- [7] BELAVIN A.A., POLYAKOV A.M., SCHWARTZ A.S., TYUPKIN Y.S. - *Pseudoparticle solutions of the Yang-Mills equations*, *Phys. Lett.* 59B(1975), 85.
- [8] DONALDSON S.K. - D. Phil. thesis (Oxford) (1982).
- [9] FREEDMAN M.H. - *The topology of four-dimensional manifolds*, *J. Differential Geometry* 17(1982), 357-453.
- [10] KURANISHI M. - *A new proof for the existence of locally complete families of complex structures*, *Proc. Conf. on Complex Analysis*, Minneapolis, Springer-Verlag, New York, (1964), 142-156.
- [11] LÜBKE M. - *Stability of Einstein-Hermitian Vector bundles*, Preprint (Bayreuth).
- [12] MITTER P.K., VIALLET C.M. - *On the bundle of connections and the gauge orbit manifold in Yang-Mills theory*, *Commun. Math. Phys.* 79(1981), 457-472.
- [13] NARASIMHAN M.S., SESHADRI C.S. - *Stable and unitary vector bundles on a compact Riemann surface*, *Ann. of Math.* 82(1965), 540-567.
- [14] PARKER T.H. - *Gauge theories on four-dimensional Riemannian manifolds*, *Commun. Math. Phys.* 85(1982), 563-602.
- [15] SEDLACEK S. - *A direct method for minimizing the Yang-Mills functional over 4-manifolds*, *Commun. Math. Phys.* 86(1982), 515-527.
- [16] SCHWARZENBERGER R.L.E. - *Vector bundles on the projective plane*, *Proc. Lond. Math. Soc.* 11(1961), 623-640.
- [17] SERRE J.-P. - *A course in arithmetic*, Springer-Verlag, New York, (1973).
- [18] SMALE S. - *An infinite dimensional version of Sard's theorem*, *Amer. J. Math.* 87(1965), 861-866.
- [19] TAUBES C.H. - *Self-dual Yang-Mills connections on non-self-dual 4-manifolds*, *J. Differential Geometry* 17(1982), 139-170.

N. J. HITCHIN

- [20] UHLENBECK K.K. - *Removable singularities in Yang-Mills fields*, Commun. Math. Phys. 83(1982), 11-30.
- [21] UHLENBECK K.K. - *Connections with L^p bounds on curvature*, Commun. Math. Phys. 83(1982), 31-42.

Nigel J. HITCHIN
St. Catherine's College
GB - OXFORD OX1 3UJ