

SÉMINAIRE DE PROBABILITÉS (STRASBOURG)

NATHALIE EISENBAUM

Exponential inequalities for Bessel processes

Séminaire de probabilités (Strasbourg), tome 34 (2000), p. 146-150

http://www.numdam.org/item?id=SPS_2000__34__146_0

© Springer-Verlag, Berlin Heidelberg New York, 2000, tous droits réservés.

L'accès aux archives du séminaire de probabilités (Strasbourg) (<http://portail.mathdoc.fr/SemProba/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

EXPONENTIAL INEQUALITIES FOR BESSEL PROCESSES

Nathalie Eisenbaum

Laboratoire de Probabilités, Tour 56, 3^{ème} étage, 4 Place Jussieu
75252 Paris Cedex 05.

Abstract : Let $R_d^*(t)$ be the supremum at time t of a Bessel process with dimension d . For T a stopping time, Burkholder has compared the expectation of $\left(\frac{R_d^*(T)}{\sqrt{d}}\right)^p$ and $(\sqrt{T})^p$ for $p > 0$. Replacing the function x^p by exponential functions, we obtain some variant of his results.

I - Introduction and notations

Let $(B_s)_{s \geq 0}$ be a linear Brownian motion starting from zero. Let R_d be a positive process such that R_d^2 is a solution of the equation :

$$X_t = 2 \int_0^t \sqrt{|X_s|} dB_s + dt, \quad d > 0;$$

i.e. : R_d is a Bessel process of dimension d starting from 0.

We set : $R_d^*(t) = \sup_{0 \leq s \leq t} R_d(s)$.

Let T be a stopping time with respect to the natural filtration of B .

Burkholder [B] established for any $p > 0$, that $\left(E\left(\frac{R_d^*(T)}{\sqrt{d}}\right)^p / E(\sqrt{T})^p\right)_{d \in \mathbb{N}^*}$ converges to 1, uniformly in T , as d tends to ∞ .

Refinements of this convergence have since been proved by Davis [D].

Making use of Poincaré's Lemma, Yor ([Y], p.55) could prove adequate modifications of these results for other times than stopping times.

A consequence of Burkholder's result is that $\left(E\left(\frac{R_d^*(T)}{d^{1/2}}\right)^p / E(T^{p/2})\right)_{d \in \mathbb{N}^*}$ is uniformly bounded in d and T . In this paper we consider the following question : What happens if the moderate function x^p is replaced by an exponential function ? For example, is there a function F such that the

sequence $(E(\exp(\frac{R_d^*(T)}{d^{1/2}})) / E(F(\sqrt{T})))_{d \geq 0}$ is uniformly bounded in d and T , or even converging as d tends to ∞ .

We will see that the answer is affirmative for the uniform boundedness question and negative for the convergence question.

II - Exponential inequalities for Bessel processes

Theorem 1 :

(i) *There exist two strictly positive constants c and β such that for any $\lambda > 0$ and any d in \mathbb{N}^* , we have for any stopping time T :*

$$E(\exp\{\lambda(\frac{R_d^*(T)}{d^{1/2}})\}) \leq c E(\exp\{\frac{\beta^2}{2} \lambda^2 T\})$$

Moreover β can be taken equal to $2\sqrt{e}$.

(ii) *For any p in $(0,2)$ there exist two strictly positive constants b_p and β_p such that for any $\lambda > 0$ and any d in \mathbb{N}^* , we have for any stopping time T :*

$$E(\exp\{\lambda(\frac{R_d^*(T)}{d^{1/2}})^p\}) \leq b_p E(\exp\{\beta_p \lambda^{\frac{2}{2-p}} T^{\frac{p}{2-p}}\})$$

Moreover β_p can be taken equal to $(\frac{1}{p} - \frac{1}{2}) p^{\frac{2}{2-p}} (4e)^{\frac{p}{2-p}}$.

The proof of Theorem 1 is based on the following result.

Theorem 2. *There exists a strictly positive constant c such that for every stopping time T , every d in \mathbb{N}^* and every $p > 0$, we have :*

$$(1)_p \quad E(R_d^*(T))^p \leq c (2\sqrt{e})^p d^{p/2} E((B_1^*)^p) E(T^{p/2})$$

Proof of Theorem 2 : Jacka and Yor have proved (see [J-Y] section 4) that there exists a constant a_p such that for all stopping times T with respect to the natural filtration of B and every d in \mathbb{N}^* :

$$E(R_d^*(T))^p \leq a_p d^{p/2} E(T^{p/2})$$

with $a_p \leq 2e 2^p (p + \frac{1}{2})^{p/2}$ when $p \geq 2$.

Thus, we are looking for $\beta > 0$ such that there exists $c > 0$ with :

$$c \cdot \beta^p E(B_1^*)^p \geq a_p$$

Since for any $p > 0$: $E(|B_1|^{2p}) = \frac{2^p}{\pi^{1/2}} \Gamma(p + \frac{1}{2})$,

Stirling's asymptotic formula gives the following equivalency :

$$E(|B_1|^{2p}) \underset{p \rightarrow \infty}{\sim} 2^{p+\frac{1}{2}} (p + \frac{1}{2})^p \exp\{-(p + \frac{1}{2})\}$$

Hence, for any $\beta \geq 2\sqrt{e}$, there exists a constant $c > 0$ such that for every $p > 0$

$$c \cdot \beta^p E(B_1^*)^p \geq a_p. \quad \square$$

We note that (1)_p can be rewritten as follows :

$$(2)_p \quad E(R_d^*(T))^p \leq c (2\sqrt{e})^p d^{p/2} E(\tilde{B}_1^*)^p T^{p/2}$$

where \tilde{B} is an independent copy of B .

Summing the inequalities (2)_{np} , n running through \mathbb{N} and p being a fixed value in $(0, 2)$, we obtain the following result :

$$E(\exp\{\lambda (R_d^*(T))^p\}) \leq c E(\exp\{\lambda (2\sqrt{e})^p d^{p/2} (\tilde{B}_1^*)^p T^{p/2}\})$$

We then use the following majorizations already established in [D-E] :

$$E(\exp\{\lambda B_1^*\}) \leq 4 \exp\left(\frac{\lambda^2}{2}\right) \quad \text{and}$$

$$E(\exp\{\lambda (B_1^*)^p\}) \leq b_p \exp\left\{\left(\frac{1}{p} - \frac{1}{2}\right)(\lambda p)^{\frac{2}{2-p}}\right\}$$

where b_p is a strictly positive constant.

to obtain Theorem 1.

We can write similar relations for exponential functions vanishing at zero. As an example, we have the following theorem .

Theorem 3 : *There exist two strictly positive constants c and β such that*

for any $\lambda > 0$ and any d in \mathbb{N}^* , we have for any stopping time T :

$$E\left(\cosh\left\{\lambda\left(\frac{R_d^*(T)}{d^{1/2}}\right)\right\} - 1\right) \leq c E\left(\exp\left\{\frac{\beta^2}{2} \lambda^2 T\right\} - 1\right)$$

Moreover: $\beta \leq 2\sqrt{e}$.

Proof : By summing the inequalities $(2)_{2n}$, n running through \mathbb{N}^* , we obtain :

$$E\left(\cosh\left\{\lambda\left(\frac{R_d^*(T)}{d^{1/2}}\right)\right\} - 1\right) \leq c E\left(\cosh\{\lambda\beta \tilde{B}_1^* \cdot T^{1/2}\} - 1\right)$$

We then note that : $E\left(\cosh\{\lambda B_1^*\} - 1\right) \leq 2\left(\exp\left(\frac{\lambda^2}{2}\right) - 1\right)$.

Remark : In view of the results of Burkholder [B] and Davis [D] it is natural to look for a function F such that, for example,

$E\left(\exp\left(\frac{R_d^*(T)}{d^{1/2}}\right)\right) / E(F(T^{1/2}))$ would converge to 1, uniformly in T , when d tends to infinity.

Assuming such a function F exists, we would obtain that for a given $\epsilon > 0$, if d is big enough, for every stopping time T :

$$(*) \quad (1-\epsilon) E(F(T^{1/2})) \leq E\left(\exp\left(\frac{R_d^*(T)}{d^{1/2}}\right)\right).$$

We would also have : $\lim_{d \rightarrow \infty} E\left(\exp\left(\frac{R_d^*(t)}{d^{1/2}}\right)\right) = F(t^{1/2})$, for $t > 0$.

But we know that for every p in \mathbb{N}^* , $E\left(\frac{R_d^*(t)}{d^{1/2}}\right)^p$ converges to $t^{p/2}$.

Consequently for every n in \mathbb{N}^* :

$$\lim_{d \rightarrow \infty} E\left(\exp\left(\frac{R_d^*(t)}{d^{1/2}}\right)\right) \geq \lim_{d \rightarrow \infty} \sum_{p=0}^n \frac{1}{p!} E\left(\frac{R_d^*(t)}{d^{1/2}}\right)^p = \sum_{p=0}^n \frac{t^{p/2}}{p!}.$$

We finally obtain : $F(t^{1/2}) \geq \exp(t^{1/2})$, for every $t > 0$.

Hence the inequality (*) implies :

$$(1-\epsilon) E(\exp(T^{1/2})) \leq E\left(\exp\left(\frac{R_d^*(T)}{d^{1/2}}\right)\right).$$

In [J-Y] Jacka and Yor have proved that such an inequality can not hold

when $d=1$. Since their argument is exclusively based on the scaling property of the Brownian motion, we can easily extend their result to any $d>1$.

In conclusion, there is no function F verifying such hypothesis.

Moreover, we see thanks to the same kind of argument, that there is not

even a function F such that $E(\exp(-\frac{R_d^*(T)}{d^{1/2}}))/E(F(T^{1/2}))$ would be uniformly minorized by a strictly positive constant.

References

- [B] D.L. BURKHOLDER "Exit times of Brownian motion, and Hardy spaces" *Advances in Math.* 26, 182-205 (1977).
- [D] B. DAVIS "On stopping times for n- dimensional Brownian motion" *Annals of Proba.* 6 ,651-659,(1978).
- [D-E] V.H. DE LA PENA and N.EISENBAUM "Exponential Burkholder Davis Gundy inequalities" *Bull.Lond.Math.Soc.* 29 ,239-242 (1996).
- [J-Y] S.D. JACKA and M. YOR "Inequalities for non-moderate functions of a pair of stochastic processes".*Proc.London Math.Soc.* (3) 67,649-672(1993).
- [Y] M. YOR "Some aspects of Brownian motion - II - Some recent martingale problems" *Lecture in Math.(Zürich) Birkhäuser*(1997).