SÉMINAIRE DE THÉORIE SPECTRALE ET GÉOMÉTRIE

WERNER BALLMANN

On the Dirichlet problem at infinity for manifolds of non-positive curvature (résumé)

Séminaire de Théorie spectrale et géométrie, tome 7 (1988-1989), p. 33-34

http://www.numdam.org/item?id=TSG_1988-1989__7__33_0

© Séminaire de Théorie spectrale et géométrie (Chambéry-Grenoble), 1988-1989, tous droits réservés.

L'accès aux archives de la revue « Séminaire de Théorie spectrale et géométrie » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

ON THE DIRICHLET PROBLEM AT INFINITY FOR MANIFOLDS OF NON-POSITIVE CURVATURE (résumé)

by *Werner BALLMANN*

A complete Riemannian manifold M^n does not admit non-constant harmonic functions in L_p for $1 < p < \infty$, see [Y 2]. As for $p = \infty$, this also holds if the sectional curvature of M is non-negative [Y 1]. The case of non-positive curvature is discussed below. We assume that M is simply connected. Then M is homeomorphic to the unit disc and has a natural compactification by a sphere at infinity, $M(\infty)$.

Consider first the case of a symmetric space $M = G/K$ of non-compact type. Then the space of bounded harmonic functions on M was described by Fürstenberg [F]. It is naturally isomorphic to the space of bounded measurable functions on G/P , where $P \subset G$ is minimal parabolic. Note that $\dim(G/P) = n - \text{rank}(M) \leq n - 1$.

Next consider the Dirichlet problem at infinity : given a continuous function f on $M(\infty)$, is there a harmonic function on M which extends continuously to f at $M(\infty)$. If M is a symmetric space of non-compact type, the Dirichlet problem at infinity (for arbitrary f) can be solved iff $\text{rank}(M) = 1$. This is quite clear from the result of Fürstenberg mentioned above. More generally, if the curvature K of M is strictly negative, $-a^2 \leq K \leq -b^2 < 0$, then the Dirichlet problem at infinity is solvable, as was shown by Anderson [Ad] and Sullivan [S] (compare also [K 1]).

Suppose now that M admits a discrete group of isometries F such that M/F is compact. If M is not a Riemannian product, then M is a symmetric space of non-compact type and $\text{rank}(M) > 1$, or M admits a geodesic which has no non-zero perpendicular parallel Jacobi field [B 2, BS]. In the latter case we say that M is a space of rank one. Such spaces were studied in [B 1], and in many respects they resemble spaces of negative curvature.

THEOREM [B 3]. — *Suppose M admits a discrete group of isometries F such that M/F is compact. If M has rank one, then the Dirichlet problem at infinity is solvable.*

In the proof it is shown that Brownian motion starting at a point $p \in M$ converges at $M(\infty)$ and that the hitting measure tends weakly to the Dirac measure at $z \in M(\infty)$ if p approaches z .

In the case of strictly negative curvature, more detailed results on harmonic functions were obtained by Anderson and Schoen [AS]. They showed, among others, that $M(\infty)$ is naturally isomorphic to the Martin boundary of M . Most of their results have been reproved (and partly extended) by other means by Ancona [Ac] and Kifer [K 2].

References

- [Ac] ANCONA A. — *Negatively curved manifolds, elliptic operators, and the Martin boundary*, Ann. of Math., **125** (1987), 495–536.
- [Ad] ANDERSON M. — *The Dirichlet problem at infinity for manifolds of negative curvature*, J. Differential Geom., **18** (1983), 701–721.
- [AS] ANDERSON M, SCHOEN R. — *Positive harmonic functions on complete manifolds of negative curvature*, Ann. of Math., **121** (1985), 429–461.
- [B 1] BALLMANN W. — *Axial isometries of manifolds of non-positive curvature*, Math. Ann., **259** (1982), 131–144.
- [B 2] BALLMANN W. — *Nonpositively curved manifolds of higher rank*, Ann. of Math., **122** (1985), 597–609.
- [B 3] BALLMANN W. — *On the Dirichlet problem at infinity for manifolds of nonpositive curvature*, Forum Math., **1** (1989), 201–213.
- [BS] BURNS K., SPATZIER R. — *Manifolds of nonpositive curvature and their buildings*, Publications Math. IHES, **65** (1987), 35–59.
- [F] FÜRSTENBERG H. — *A Poisson for semi-simple Lie groups*, Ann. of Math., **77** (1963), 335–386.
- [K 1] KIFER Y. — *Brownian motion and harmonic functions on manifolds of negative curvature*, Theory Probab. Appl., **21** (1976), 81–95.
- [K 2] KIFER Y. — *Brownian motion and positive harmonic functions on complete manifolds of non-positive curvature*, Pitman Research Notes in Mathematics, **150** (1986), 132–178.
- [S] SULLIVAN D. — *The Dirichlet problem at infinity for a negatively curved manifold*, J. Differential Geom., **18** (1983), 723–732.
- [Y 1] YAU S.T. — *Harmonic functions on complete Riemannian manifolds*, Comm. Pure Appl. Math., **28** (1975), 201–228.
- [Y 2] YAU S.T. — *Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry*, Ind. Univ. Math. J., **25** (1976), 659–670.

Werner BALLMANN
 Universität Bonn
 Math. Institut
 5300 BONN (R.D.A.)