

Positive solutions for the p -Laplacian involving critical and supercritical nonlinearities with zeros

Leonelo Iturriaga^{a,*}, Sebastián Lorca^{a,2}, Eugenio Massa^{b,3}

^a Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica, Chile

^b Departamento de Matemática, Instituto de Ciências Matemáticas e de Computação, Universidade de São Paulo, Campus de São Carlos, Caixa Postal 668, 13560-970, São Carlos SP, Brazil

Received 28 September 2009; received in revised form 4 November 2009; accepted 4 November 2009

Available online 10 November 2009

Abstract

In this paper we show the existence of multiple solutions to a class of quasilinear elliptic equations when the continuous nonlinearity has a positive zero and it satisfies a p -linear condition only at zero. In particular, our approach allows us to consider superlinear, critical and supercritical nonlinearities.

© 2009 Elsevier Masson SAS. All rights reserved.

1. Introduction

In this paper, we look for positive $C^1(\overline{\Omega})$ weak solutions of the problem

$$\begin{cases} -\Delta_p u = \lambda f(u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (P_\lambda)$$

where Ω is a convex bounded domain in \mathbb{R}^N with smooth boundary, $N > p > 1$, λ is a positive parameter and f satisfies $f(0) = f(1) = 0$, $f(x) > 0$ for any $x \notin \{0; 1\}$; we will show the existence of at least two positive solutions for λ large, without restrictions on the growth of the nonlinearity at infinity.

It is known from [17,18] that if the domain Ω is star-shaped and the nonlinearity is $|u|^{r-2}u$ with r greater or equal to the critical exponent $p^* = pN/(N-p)$, then no nontrivial solution exists. A solution could be recovered either by considering more topologically complex domains, or by perturbing the nonlinearity. In this second direction several authors considered nonlinearities with any growth at infinity but which behave like $|u|^{q-2}u$ with $q \in (p, p^*)$ near zero; for instance, [3,16,9] assume this type of condition, then they truncate the nonlinearity and look for estimates on the possible solutions. These estimates allow to prove that the solutions are below the truncation point for suitable values of λ , and then a solution of the original problem is obtained.

* Corresponding author.

E-mail addresses: leonelo.iturriaga@gmail.com (L. Iturriaga), slorca@uta.cl (S. Lorca), eug.mass@gmail.com (E. Massa).

¹ Partially supported by FONDECYT No. 11080203 and Convenio de desempeño UTA MECESUP 2.

² Partially supported by FONDECYT No. 1080500.

³ The author was partially supported by Fapesp and CNPq Brazil.

Problems with a nonlinearity which is nonnegative but has a zero at a positive value were first considered in [14] for the Laplacian operator, and two solutions were obtained, through topological degree arguments, in the subcritical case. The existence and behavior of a solution below the zero of f are studied in many works (see for instance [7] and references therein), and it can be proved that this solution converges to 1 (the positive zero of the nonlinearity), when $\lambda \rightarrow \infty$. The existence of a solution whose maximum is above 1 is more delicate and usually requires some hypotheses on the growth of f at infinity. In [11], we considered the p -Laplacian operator and we allowed f to depend also on the variable $x \in \Omega$, but only in the subcritical case: two positive solutions were obtained for λ above the first eigenvalue of the asymptotical problem at the origin, and it was proved that both solutions converge at least pointwise to 1 when $\lambda \rightarrow \infty$.

This behavior suggests that also for this problem, truncation procedures like those in [3,16,9] could be used to prove the existence of two solutions when considering critical or supercritical nonlinearities. However, the pointwise convergence is not enough to guarantee a suitable control on the L^∞ norm of the solutions.

In this paper we suppose that Ω is convex and that f is independent of $x \in \Omega$, in order to use suitable monotonicity results (such as [2,6]) which imply a better knowledge of the geometry of the solutions, and then allow to estimate the L^∞ norm when $\lambda \rightarrow \infty$ and finally obtain the existence result for critical or supercritical nonlinearities. Even in the subcritical case, our result gives new information with respect to [11] in the sense that we may substitute global hypotheses on f with (much weaker) local ones (see Remark 1.1).

We remand to [11] for a further discussion on the literature related to (P_λ) .

1.1. Statement of the results

We will consider the following hypotheses on f .

- (F₁) $f : [0, +\infty) \rightarrow [0, +\infty)$ is a continuous function which is locally Lipschitz continuous in $(0, \infty)$, $f(0) = f(1) = 0$ and $f(x) > 0$ for $x \notin \{0; 1\}$.
 (F₂) $\liminf_{s \rightarrow 0^+} \frac{f(s)}{s^{p-1}} \geq 1$.
 (F₃) There exist $\gamma > 0$ and $\sigma \in (p-1, p_*-1)$ such that

$$\lim_{t \rightarrow 1} \frac{f(t)}{|t-1|^\sigma} = \gamma,$$

where p_* denotes the Serrin's exponent given by $p_* = \frac{(N-1)p}{N-p}$.

- (F₄) There exist $k > 0$ and $T > 1$ such that the map $t \mapsto f(t) + kt^{p-1}$ is increasing for $t \in [0, T]$.

Our result is the following

Theorem 1.1. *Assume that Ω is a convex smooth domain. Then, under the hypotheses (F₁) through (F₄), there exists $\lambda^* > 0$ such that the problem (P_λ) has at least two C^1 weak positive solutions $u_{1,\lambda}$, $u_{2,\lambda}$, for $\lambda > \lambda^*$.*

Moreover, these solutions satisfy $\|u_{1,\lambda}\|_\infty \rightarrow 1^-$ and $\|u_{2,\lambda}\|_\infty \rightarrow 1^+$, when $\lambda \rightarrow \infty$.

A simple example of a function f satisfying the four assumptions of the preceding theorem is $f(u) = u^{p-1}e^u|1-u|^\sigma$ where $\sigma \in (p-1, p_*-1)$.

1.2. Some comments on the problem

In [11], two solutions for the problem (P_λ) were encountered by mainly variational techniques, the first solution being a local minimum (which could also be obtained via sub- and supersolutions), while the second one was obtained via mountain pass. It was also proved, using a combination of a Liouville-type theorem, a priori estimates and the blow-up technique, that both solutions tend pointwise to 1.

For the above result of pointwise convergence, a blow-up argument centered at an arbitrary fixed point in Ω and a new Liouville-type theorem in \mathbb{R}^N (see Lemma 2.2) were combined. A stronger result could be achieved if one centers the blow-up at the maximum point of the solution; however, in that case we did not know if the maximum of

the solutions stayed far from the boundary or not, so that the limiting problem could be in a half-space instead of \mathbb{R}^N , and Liouville-type theorems in the half-space are not available for the kind of nonlinearity that we are considering.

Since we are aiming to treat also supercritical nonlinearities, variational techniques cannot be used directly here. For this reason we perform a truncation of the nonlinearity and we look for solutions below the truncation point. As observed above, in order to obtain an estimate on the L^∞ norm, when applying the blow-up argument, we need to be sure that the maximum point of the solutions stays far from the boundary, so that the limiting problem is defined in the whole of \mathbb{R}^N . This will be obtained by assuming the convexity of Ω and using [6].

However, the results in [6] hold for locally Lipschitz and strictly positive nonlinearities. The first condition imposes a restriction on p and N (actually, assumption (F_3) is not possible for a Lipschitz function if $p < 2$ and N is large, see Remark 1.1) while, due to the second condition, we will need to solve first an auxiliary problem, where a perturbation is added which makes the nonlinearity strictly positive for $u > 0$. A first solution for the perturbed problem is obtained via sub- and supersolutions, and a second one by using topological degree (see Propositions 3.3–3.4).

Finally, the first solution in Theorem 1.1 is just the same as in [11], while the second one is obtained as the limit of the solutions of the perturbed problem; since we need to distinguish these two solutions, it is crucial to know that one lies below 1 and the other does not: for this reason, instead of the mountain pass theorem used in [11], we obtain the second solution for the perturbed problem by a degree approach, which has the advantage to furnish the information that its maximum is greater than the supersolution available.

We conclude this Introduction with some remarks on the hypotheses.

Remark 1.1.

- Hypothesis (F_2) is classical in order to have a subsolution for λ above the first eigenvalue of the operator. On the other hand, the constant function 1 is always a supersolution for (P_λ) , but not for the perturbed problem: hypothesis (F_3) will be used to obtain a family of supersolutions, in particular a supersolution strictly below 1 and one strictly above: this will help to distinguish the two solutions when taking limit.
- Hypothesis (F_3) is required also to obtain inequality (2.1) for the truncated nonlinearity, which is necessary for applying the Liouville-type theorem from [11] (Lemma 2.2). We also remark that, in fact, (F_3) avoids the formation of the so-called flat core (a solution which coincides with 1 in a whole open set, see [12,20] for instance); actually, if this phenomenon could occur, then it would become difficult to separate the two solutions and obtain the multiplicity result.
- Hypothesis (F_4) is a standard condition required in order to apply the sub- and supersolution method and comparison principles. We remark that in [11] we had to impose hypothesis (F_4) with $T = \infty$ and condition (2.1) had to be imposed directly; here these hypotheses are replaced by the local conditions (F_3) and (F_4) , since it will be possible to verify the global conditions when performing the truncation of the nonlinearity.
- We observe that in fact Theorem 1.1 is meaningful only for $p > 4/3$, moreover, if $p \in (4/3, 2)$ we have an upper bound for the dimension N : actually, hypothesis (F_3) is possible for a Lipschitz function only if the Serrin’s exponent $p_* > 2$, which implies, for $p < 2$, that $p < N < p/(2 - p)$, and this cannot be satisfied if $p \leq 4/3$.

2. Preliminaries

We will denote by λ_1 the first eigenvalue of $(-\Delta_p)$ in Ω and by ϕ_1 the first eigenfunction, which can be chosen positive in Ω .

By hypothesis (F_3) there exist $R > 1$ and $\gamma' > 0$ such that $f(t) \geq \gamma'|t - 1|^\sigma$ for $t \in [1, R]$; without loss of generality we may suppose that $R \leq T$ from hypothesis (F_4) . Then we truncate f as follows

$$f_R(t) = \begin{cases} f(t^+), & t \leq R, \\ \frac{f(R)}{R^\sigma} t^\sigma, & t \geq R, \end{cases}$$

where $t^+ = \max\{0, t\}$. With this definition, f_R has a power growth at infinity with exponent below the Serrin’s exponent and satisfies the following properties:

$$f_R(t) \geq \gamma''|t - 1|^\sigma \quad \text{for } t \geq 1, \tag{2.1}$$

if $\gamma'' = \min\{\gamma', \frac{f(R)}{R^\sigma}\} > 0$ and

$$\text{the map } t \mapsto f_R(t) + kt^{p-1} \text{ is increasing for } t \in [0, \infty], \quad (2.2)$$

where k is as in hypothesis (F_4) .

We consider then the auxiliary problem

$$\begin{cases} -\Delta_p u = \lambda f_R(u) + \tau(u^+)^{p-1} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (Q_{\lambda, \tau})$$

where τ is a nonnegative parameter.

We remark that, by the strong maximum principle (see [23]), the nontrivial solutions of the problem $(Q_{\lambda, \tau})$ are positive and, by hypothesis (F_1) and since $\sigma < p_* - 1$, they are in $C^{1, \alpha}(\overline{\Omega})$ for some $\alpha \in (0, 1)$ (see [8]); moreover, since $f_R \geq 0$, $(Q_{\lambda, \tau})$ has no positive solution if $\tau > \lambda_1$.

The following lemma is a consequence of the results in [6], and will be used in our argument.

Lemma 2.1. *Under the hypotheses (F_1) and (F_3) , if Ω is convex, there exists $\delta_\Omega > 0$ which depends only on Ω (but not on f , R , τ and λ) with the following property: for any $C^1(\overline{\Omega})$ weak solution u of $(Q_{\lambda, \tau})$ with $\tau > 0$, there exists a point $x \in \Omega$ such that $\text{dist}(x, \partial\Omega) > \delta_\Omega$ and $u(x) = \|u\|_\infty$.*

Our purpose will be to obtain a solution for $(Q_{\lambda, 0})$ as the limit of solutions of $(Q_{\lambda, \tau})$ with $\tau > 0$, so that the conclusion of Lemma 2.1 holds also for this solution. Then it will be possible to prove that, for large λ , this is also a solution for (P_λ) ; for this we will make use of the following result, which (using the fact that f_R was constructed satisfying (2.1) and with power growth with exponent below $p_* - 1$) is a consequence of the Liouville-type theorem proved in [11].

Lemma 2.2. *Under hypotheses (F_1) , (F_2) and (F_3) , any C^1 weak solution of the problem*

$$\begin{cases} -\Delta_p w = f_R(w) & \text{in } \mathbb{R}^N, \\ w \geq 0 \end{cases} \quad (2.3)$$

is either $w \equiv 0$ or $w \equiv 1$.

Remark 2.1. We observe that hypothesis (F_2) is weaker than the one appearing in [11], but the extension to include this case is straightforward.

3. Proofs

Our first step will be to derive some a priori estimates for the solutions of $(Q_{\lambda, \tau})$; we remark that this result holds also for $\tau = 0$.

Lemma 3.1. *Under hypotheses (F_1) and (F_2) , we have*

(1) *given $\tilde{\lambda} > 0$, there exists a constant $D_{\tilde{\lambda}}$ such that, if $u \in C^1(\overline{\Omega})$ is a weak solution of the problem $(Q_{\lambda, \tau})$ with $\lambda > \tilde{\lambda}$ and $\tau \geq 0$ then*

$$\|u\|_\infty \leq D_{\tilde{\lambda}};$$

(2) *given $\lambda > 0$, there exist constants $C_\lambda > 0$ and $\alpha \in (0, 1)$ such that one has also the estimate*

$$\|u\|_{C^{1, \alpha}(\overline{\Omega})} \leq C_\lambda. \quad (3.4)$$

Proof. Suppose, for sake of contradiction, that there exists a sequence $\{(u_n, \lambda_n, \tau_n)\}_{n \in \mathbb{N}}$ with u_n being a positive C^1 -solution of (Q_{λ_n, τ_n}) , such that $S_n := \max_{\overline{\Omega}} u_n = u_n(x_n) \xrightarrow{n \rightarrow \infty} \infty$, where $\{x_n\} \subset \Omega$ is a sequence of points where the maximum is attained. We remark that since we are not supposing $\tau > 0$ at this point, this sequence may not be bounded away from the boundary.

Let now $\delta_n = \text{dist}(x_n, \partial\Omega)$ and define $w_n(y) = S_n^{-1}u_n(A_n y + x_n)$, where A_n will be fixed later; then w_n satisfies

$$-\Delta_p w_n(y) = \lambda_n \frac{A_n^p}{S_n^{p-1}} f_R(S_n w_n(y)) + \tau_n A_n^p w_n(y) \quad \text{in } B(0, \delta_n A_n^{-1}) \tag{3.5}$$

and $w_n(0) = \max w_n = 1$.

We choose $A_n^p = \lambda_n^{-1} S_n^{p-1-\sigma} f(R)^{-1} R^\sigma$: since $S_n \rightarrow \infty$, $\lambda_n > \tilde{\lambda}$ and $\tau_n \leq \lambda_1$ (since no positive solution of $(Q_{\lambda,\tau})$ exists for $\tau > \lambda_1$), we conclude that $A_n \rightarrow 0$ and $\tau_n A_n^p \rightarrow 0$. Then, the right hand side of (3.5) becomes $\frac{R^\sigma f_R(S_n w_n)}{f(R) S_n^\sigma} + o(1)$ and then by the continuity of f and the definition of f_R it is bounded. This allows us to apply the regularity theorem in [22] for the p -Laplacian operator, indeed, if Ω_n is the rescaled domain then, according to whether the limit of δ_n/A_n is infinity or not, Ω_n tends to \mathbb{R}^N or to a half-space; fixed an open subset $\tilde{\Omega}$ such that $\tilde{\Omega} \subseteq \Omega_n$ for n large, for any compact set $\Omega' \subseteq \tilde{\Omega}$ one obtains $\alpha \in (0, 1)$ and $C > 0$ such that, since w_n is also uniformly bounded in L^∞ , the estimate $\|w_n\|_{C^{1,\alpha}(\Omega')} \leq C$ holds.

Then, up to a subsequence, $w_n \rightarrow w$ in the C^1 norm in compact sets, where w is a C^1 -function defined on \mathbb{R}^N or on a half-space.

Finally, taking the limit in (3.5), we have that w satisfies, in the weak sense, the following:

$$\begin{cases} -\Delta_p w = w^\sigma, \\ w > 0, \\ w(0) = \max w = 1; \end{cases}$$

this contradicts the Liouville-type theorem in [19, Corollary II] in the case of \mathbb{R}^N and in [15] for the half-space.

This contradiction proves that $\|u\|_\infty \leq C$ for any solution of the problem $(Q_{\lambda,\tau})$ with $\lambda > \tilde{\lambda}$ and $\tau \geq 0$, that is, the item (1) of Lemma 3.1, and also for any solution with a given λ and $\tau \geq 0$. In this second case, using the regularity theorem in [13], one obtains the uniform bound also for the $C^{1,\alpha}$ norm, as claimed in the item (2). \square

Now we look for a family of supersolutions: for this purpose, let $e \in W_0^{1,p}(\Omega)$ be the solution of

$$\begin{cases} -\Delta_p e = 1 & \text{in } \Omega, \\ e = 0 & \text{on } \partial\Omega \end{cases}$$

and $n := \|e\|_\infty$.

Lemma 3.2. *Under hypothesis (F_3) , for any $\lambda > 0$ there exist $\tau_\lambda^*, \delta_\lambda > 0$ such that $v_\xi = 1 + \xi + \frac{\delta_\lambda}{4n}e$ is a supersolution for $(Q_{\lambda,\tau})$ for any $\xi \in [-\delta_\lambda, \delta_\lambda/2]$ and $\tau \in [0, \tau_\lambda^*)$. Moreover, we may choose δ_λ as a nonincreasing function of λ .*

Proof. Fixed $\lambda > 0$, by the hypothesis (F_3) we have that

$$\lim_{t \rightarrow 1} \frac{\lambda f_R(t)}{|t-1|^{p-1}} = 0,$$

and then there exists $\delta > 0$ such that $\lambda f_R(t) < (\frac{|t-1|}{8n})^{p-1} < (\frac{\delta}{8n})^{p-1}$ for $|t-1| \leq \delta$. Since this estimate still holds for lower values of λ we deduce that δ may be chosen as a nonincreasing function of λ .

If $\tau^* > 0$ is such that $\tau u^{p-1} < (\frac{\delta}{8n})^{p-1}$ for $\tau \in [0, \tau^*)$, $u \in (0, 1 + \delta]$, then

$$\lambda f_R(u) + \tau u^{p-1} < \left(\frac{\delta}{4n}\right)^{p-1} \quad \text{for } \tau \in [0, \tau^*), \quad u \in [1 - \delta, 1 + \delta].$$

If we define $v_\xi = 1 + \xi + \frac{\delta}{4n}e$, we have that $v_\xi \in [1 - \delta, 1 + \delta]$ provided $\tau \in [0, \tau^*)$, $\xi \in [-\delta, \delta/2]$ and then

$$-\Delta_p v_\xi = \left(\frac{\delta}{4n}\right)^{p-1} > \lambda f_R(v_\xi) + \tau v_\xi^{p-1},$$

which proves that v_ξ is a supersolution. \square

Now we prove the existence of a first solution for $(Q_{\lambda,\tau})$ via the sub- and supersolution method: for this we need hypothesis (F_4) .

Proposition 3.3. *If hypotheses (F₁)–(F₄) hold, then the problem (Q_{λ,τ}) has a positive solution u_{1,λ,τ} < 1 for λ > λ₁ and 0 ≤ τ < τ_λ^{*}.*

Moreover, the following property holds: given λ̄ > λ₁ there exists ε > 0 such that εφ₁ ≤ u_{1,λ,τ} < 1 for any λ > λ̄ and τ ∈ [0, τ_λ^{}).*

Proof. In a standard way, using (F₂), we may find a ε > 0 (as small as desired) such that λf_R(t) > λ₁t^{p-1} for any t ∈ (0, max{εφ₁}) and any λ > λ̄ > λ₁; then εφ₁ is a subsolution for the problem (Q_{λ,τ}) for any τ ≥ 0 and λ > λ̄.

For τ ∈ [0, τ_λ^{*}), we have the supersolution v_{-δ_λ} < 1 from Lemma 3.2; since δ_λ is not increasing in λ, we may choose ε such that εφ₁ < v_{-δ_λ}/2 for any λ > λ̄. Then the sub- and supersolutions method gives a solution u_{1,λ,τ} with the claimed properties. □

Now, we work with τ > 0 and we show that a second solution exists: we will apply a topological degree argument, adapting a result obtained, for p = 2, by de Figueiredo and Lions in [4], see also [10] for the general case.

Proposition 3.4. *In the same hypotheses as Proposition 3.3, if λ > λ₁ and τ₀ ∈ (0, τ_λ^{*}), then (P_{λ,τ₀}) has a second positive solution u_{2,λ,τ₀}. Moreover ||u_{2,λ,τ₀}||_∞ > 1.*

Proof. Let us fix λ > λ₁ and denote by X the Banach space of C¹-functions on Ω̄ which are 0 on ∂Ω, endowed with the usual C¹-norm. Also, we will write u ≪ v to say that u < v in Ω and ∂u/∂ν > ∂v/∂ν on ∂Ω, where ν denotes the unitary outward normal to ∂Ω. Let k be as in (2.2) and K_τ : X → X be defined as follows: K_τv = u, where u is the unique solution of the Dirichlet problem

$$\begin{cases} -\Delta_p u + \lambda k u^{p-1} = \lambda f_R(v) + (\lambda k + \tau)v^{p-1} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega; \end{cases} \quad (3.6)$$

the mapping K_τ so defined is compact.

We consider the bounded open set

$$\mathcal{O} = \{u \in X : \|u\|_X < C_\lambda + B_\lambda + 1, u \gg \varepsilon\phi_1\},$$

where C_λ, B_λ > 0 will be chosen below (see in (3.7) and (3.9), respectively) and ε > 0 is as in the proof of Proposition 3.3, so that εφ₁ < 1 and it is a strict subsolution for all problems (Q_{λ,τ}), τ ≥ 0 (in particular λ₁(εφ₁)^{p-1} < λf_R(εφ₁)).

We need that 0 ∉ (I - K_τ)(∂O) (i.e., no solution of (Q_{λ,τ}) lies on ∂O), so that the degree deg(I - K_τ, O, 0) will be well defined and independent of τ. To obtain this we get C_λ from Lemma 3.1 part (2), so that

$$\|u\|_X \leq C_\lambda \quad (3.7)$$

for all possible solutions of (Q_{λ,τ}) with τ ≥ 0.

Then, we claim that any solution u of (Q_{λ,τ}) such that u ≥ εφ₁ in Ω satisfies u ≫ εφ₁ (and then it is not on ∂O). Actually, we have

$$\begin{cases} -\Delta_p u + \lambda k u^{p-1} = \lambda f_R(u) + (\lambda k + \tau)u^{p-1}, \\ -\Delta_p(\varepsilon\phi_1) + \lambda k(\varepsilon\phi_1)^{p-1} = \lambda_1(\varepsilon\phi_1)^{p-1} + (\lambda k + \tau)(\varepsilon\phi_1)^{p-1}, \end{cases} \quad (3.8)$$

by hypothesis (F₄) and since u ≥ εφ₁, we have λf_R(u) + (λk + τ)u^{p-1} ≥ λf_R(εφ₁) + (λk + τ)(εφ₁)^{p-1}, and then a strict inequality holds between the (continuous) right hand sides of (3.8). Thus, using the comparison result in [5], the claim is proved.

By the above computations, we obtain that

$$\deg(I - K_\tau, \mathcal{O}, 0) = 0 \quad \text{for any } \tau > 0,$$

since (Q_{λ,τ}) has no solutions for τ > λ₁.

At this point we fix τ = τ₀, we consider the supersolution a := v_{ξ=0} > 1 from Lemma 3.2, and we assume that no solution of (Q_{λ,τ₀}) touches it, otherwise such a solution would satisfy the claim and the proposition would be true. Using the L[∞] estimate in [1] and then [13] we obtain that we may choose the constant B_λ > 0 such that

$$\|K_{\tau_0} v\|_X \leq B_\lambda, \quad \forall v \in X : 0 \leq v \leq a; \quad (3.9)$$

we consider the open subset of \mathcal{O}

$$\mathcal{O}' = \{u \in \mathcal{O}: u < a \text{ in } \Omega\}$$

and we claim that $\text{deg}(I - K_{\tau_0}, \mathcal{O}', 0) = 1$.

Observe that K_{τ_0} maps $\overline{\mathcal{O}'}$ into $\overline{\mathcal{O}'}$. Indeed, if $v \in \overline{\mathcal{O}'}$, then $\|K_{\tau_0}v\|_X \leq B_\lambda v$ by (3.9), and if we consider $u = K_{\tau_0}v$ we have

$$\begin{cases} -\Delta_p a + \lambda k a^{p-1} \geq \lambda f_R(a) + (\lambda k + \tau_0) a^{p-1}, \\ -\Delta_p u + \lambda k u^{p-1} = \lambda f_R(v) + (\lambda k + \tau_0) v^{p-1}, \\ -\Delta_p(\varepsilon\phi_1) + \lambda k(\varepsilon\phi_1)^{p-1} = \lambda_1(\varepsilon\phi_1)^{p-1} + (\lambda k + \tau_0)(\varepsilon\phi_1)^{p-1}, \end{cases} \tag{3.10}$$

then, since $\varepsilon\phi_1 \leq v \leq a$, the comparison principle in [21] implies that $\varepsilon\phi_1 \leq K_{\tau_0}v \leq a$.

Now, let $u_0 \in \mathcal{O}'$ and consider the constant mapping $C : \overline{\mathcal{O}'} \rightarrow \overline{\mathcal{O}'}$ defined by $C(u) = u_0$: one obtains that $I - \mu K_{\tau_0}(v) - (1 - \mu)u_0$, $\mu \in [0, 1]$, is a homotopy between $I - K_{\tau_0}$ and $I - C$ in $\overline{\mathcal{O}'}$ without zeros on $\partial\mathcal{O}'$: in fact, if $v \in \partial\mathcal{O}'$ then (since \mathcal{O}' is convex) $\mu K_{\tau_0}(v) + (1 - \mu)u_0 \in \mathcal{O}'$ for $\mu \neq 1$, and then it is different from v , while for $\mu = 1$ we have $v \neq K_{\tau_0}(v)$ since we are assuming that no solution touches a .

Hence $\text{deg}(I - K_{\tau_0}, \mathcal{O}', 0) = \text{deg}(I - C, \mathcal{O}', 0) = 1$, as we claimed.

Then, applying the excision property, it follows that $\text{deg}(I - K_{\tau_0}, \mathcal{O} \setminus \overline{\mathcal{O}'}, 0) = -1$, so (Q_{λ, τ_0}) has a solution $u_2 \in \mathcal{O} \setminus \overline{\mathcal{O}'}$; in particular, $u_2(x_0) > a(x_0) > 1$ in some point $x_0 \in \Omega$, since otherwise it would be on $\partial\mathcal{O}'$, and then u_2 is distinct from u_{1, λ, τ_0} from Proposition 3.3. \square

Now, we will obtain a solution for $(Q_{\lambda, 0})$ as the limit of the solutions obtained in the previous proposition; as a result, such solution inherits the property in Lemma 2.1.

Lemma 3.5. *In the same hypotheses as Propositions 3.3–3.4, if moreover Ω is convex, then given $\lambda > \lambda_1$, there exists a solution $u_{2, \lambda, 0}$ for the problem $(Q_{\lambda, 0})$, which satisfies $\|u_{2, \lambda, 0}\|_\infty \geq 1$.*

Moreover, there exists $x \in \Omega$ such that $d := \text{dist}(x, \partial\Omega) \geq \delta_\Omega$ and $u_{2, \lambda, 0}(x) = \|u_{2, \lambda, 0}\|_\infty$.

Proof. Given $\lambda > \lambda_1$ we will consider a sequence $\tau_n \rightarrow 0$ and we will focus on the solution $u_n := u_{2, \lambda, \tau_n}$ from Proposition 3.4, so that we know that $\|u_n\|_\infty > 1$, and that, by Lemma 2.1, there exists $x_n \in \Omega$ such that $d_n := \text{dist}(x_n, \partial\Omega) > \delta_\Omega$ and $u_n(x_n) = \|u_n\|_\infty$.

By Lemma 3.1 point (2), we have a uniform bound for $\|u_n\|_{C^{1, \alpha}(\overline{\Omega})}$ for some $\alpha \in (0, 1)$. Then, up to a subsequence, $u_n \rightarrow u$ in $C^1(\overline{\Omega})$, where u is a nonnegative weak solution of $(Q_{\lambda, 0})$.

From $\|u_n\|_\infty > 1$ we obtain $\|u\|_\infty \geq 1$, thus u is nontrivial and then positive. Finally, up to a subsequence, $x_n \rightarrow x \in \Omega$ with $\text{dist}(x, \partial\Omega) \geq \delta_\Omega$ and taking limit $u(x) = \|u\|_\infty$. \square

The following lemma will show that, for λ large, the solution from Lemma 3.5 is a solution also for the supercritical problem (P_λ) .

Lemma 3.6. *The solutions $u_{2, \lambda, 0}$ from Lemma 3.5 satisfy $\|u_{2, \lambda, 0}\|_\infty \rightarrow 1$ when $\lambda \rightarrow \infty$.*

In particular, there exists λ^ such that if $\lambda > \lambda^*$ then $\|u_{2, \lambda, 0}\|_\infty \leq R$.*

Proof. Given $\eta > 1$, suppose by contradiction that there exists a sequence $\lambda_n \rightarrow \infty$ such that the corresponding solutions $u_n := u_{2, \lambda_n, 0}$ satisfy $\|u_n\|_\infty > \eta$, in particular there exists a sequence $x_n \in \Omega$ such that $d_n := \text{dist}(x_n, \partial\Omega) \geq \delta_\Omega$ and $u_n(x_n) = \|u_n\|_\infty > \eta$.

Letting $w_n(x) = u_n(x_n + \lambda_n^{-\frac{1}{p}}x)$ we see that w_n satisfies

$$-\Delta_p w_n(x) = f_R(w_n) \quad \text{in } B(0, d_n \lambda_n^{1/p}) \tag{3.11}$$

and $w_n(0) = u_n(x_n)$.

As in the proof of point (2) in Lemma 3.1, we obtain (since w_n is bounded in L^∞ by the point (1) in the same lemma) also a uniform bound in the $C^{1, \alpha}$ norm in compact sets, for some $\alpha \in (0, 1)$; then, up to a subsequence,

$w_n \rightarrow w$ in the C^1 norm in compact sets, where now w is a C^1 function defined in \mathbb{R}^N , since $d_n \lambda_n^{1/p} \rightarrow \infty$. Thus, w is a weak solution of the problem

$$\begin{cases} -\Delta_p w = f_R(w) & \text{in } \mathbb{R}^N, \\ w \geq 0. \end{cases} \quad (3.12)$$

According to Lemma 2.2 we conclude that either $w \equiv 0$ or $w \equiv 1$.

This contradicts the fact that $w_n(0) = u_n(x_n) > \eta > 1$, and then the lemma is proved. \square

We are now in a position to prove our main result.

Proof of Theorem 1.1. The first solution is $u_{1,\lambda,0}$ from Proposition 3.3, and satisfies $\|u_{1,\lambda,0}\|_\infty < 1$. By Lemma 3.6 we see that, for λ large, the solutions $u_{2,\lambda,0}$ from Lemma 3.5 satisfy $1 \leq \|u_{2,\lambda,0}\|_\infty < R$, and then are solutions of the supercritical problem (P_λ) . Therefore we have obtained the existence of a second solution.

We have already proved that $\|u_{2,\lambda,0}\|_\infty \rightarrow 1$ when $\lambda \rightarrow \infty$.

By hypotheses (F_1) and (F_2) , if t_λ is the largest real such that $\lambda f(t) > \lambda_1 t^{p-1}$ for $t \in (0, t_\lambda)$, then $t_\lambda \rightarrow 1$ when $\lambda \rightarrow \infty$. Since no positive solution of (P_λ) may exist below t_λ , we deduce that also $\|u_{1,\lambda,0}\|_\infty \rightarrow 1$ when $\lambda \rightarrow \infty$. \square

Acknowledgements

This work was started while the third author was visiting the Instituto de Alta Investigación, Universidad de Tarapacá, Chile, and then concluded while the first author was visiting the ICMC-USP/BRAZIL. Both authors thank the faculties and staff of ICMC-USP and Instituto de Alta Investigación for their kind hospitality.

References

- [1] A. Anane, Etude des valeurs propres et de la résonance pour l'opérateur p -Laplacien, PhD thesis, Université Libre de Bruxelles, 1987.
- [2] H. Berestycki, L. Nirenberg, On the method of moving planes and the sliding method, *Bol. Soc. Brasil. Mat. (N.S.)* 22 (1) (1991) 1–37. MR MR1159383 (93a:35048).
- [3] S. Chen, S. Li, On a nonlinear elliptic eigenvalue problem, *J. Math. Anal. Appl.* 307 (2) (2005) 691–698. MR MR2142453 (2006a:35225).
- [4] D.G. de Figueiredo, P.-L. Lions, On pairs of positive solutions for a class of semilinear elliptic problems, *Indiana Univ. Math. J.* 34 (3) (1985) 591–606. MR MR794577 (86j:35058).
- [5] D.G. de Figueiredo, J.-P. Gossez, P. Ubilla, Local “superlinearity” and “sublinearity” for the p -Laplacian, *J. Funct. Anal.* 257 (3) (2009) 721–752. MR 35J60 (47J30 58E05).
- [6] L. Damascelli, B. Sciunzi, Regularity, monotonicity and symmetry of positive solutions of m -Laplace equations, *J. Differential Equations* 206 (2) (2004) 483–515. MR MR2096703 (2005h:35116).
- [7] J. García-Melián, J. Sabina de Lis, Stationary profiles of degenerate problems when a parameter is large, *Differential Integral Equations* 13 (10–12) (2000) 1201–1232. MR MR1785705 (2001h:35059).
- [8] M. Guedda, L. Véron, Quasilinear elliptic equations involving critical Sobolev exponents, *Nonlinear Anal.* 13 (8) (1989) 879–902. MR MR1009077 (90h:35100).
- [9] L. Iturriaga, S. Lorca, M. Montenegro, Existence of solutions to quasilinear elliptic equations with singular weights, *Adv. Nonlinear Stud.*, in press.
- [10] L. Iturriaga, S. Lorca, J. Sánchez, Existence and multiplicity results for the p -Laplacian with a p -gradient term, *NoDEA Nonlinear Differential Equations Appl.* 15 (6) (2008) 729–743. MR MR2465780.
- [11] L. Iturriaga, E. Massa, J. Sanchez, P. Ubilla, Positive solutions for the p -Laplacian with a nonlinear term with zeros, *J. Differential Equations* 248 (2) (2010) 309–327.
- [12] S. Kamin, L. Véron, Flat core properties associated to the p -Laplace operator, *Proc. Amer. Math. Soc.* 118 (4) (1993) 1079–1085. MR MR1139470 (93j:35071).
- [13] G.M. Lieberman, Boundary regularity for solutions of degenerate elliptic equations, *Nonlinear Anal.* 12 (11) (1988) 1203–1219. MR MR969499 (90a:35098).
- [14] P.-L. Lions, On the existence of positive solutions of semilinear elliptic equations, *SIAM Rev.* 24 (4) (1982) 441–467. MR MR678562 (84a:35093).
- [15] S. Lorca, Nonexistence of positive solution for quasilinear elliptic problems in the half-space, *J. Inequal. Appl.* (2007), Art. ID 65126, 4 pp. MR MR2291650 (2007m:35073).
- [16] S. Lorca, P. Ubilla, Partial differential equations involving subcritical, critical and supercritical nonlinearities, *Nonlinear Anal.* 56 (1) (2004) 119–131. MR MR2031438 (2004m:35087).
- [17] S.I. Pohožaev, On the eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$, *Dokl. Akad. Nauk SSSR* 165 (1965) 36–39. MR MR0192184 (33 #411).
- [18] P. Pucci, J. Serrin, A general variational identity, *Indiana Univ. Math. J.* 35 (3) (1986) 681–703. MR MR855181 (88b:35072).

- [19] J. Serrin, H. Zou, Cauchy–Liouville and universal boundedness theorems for quasilinear elliptic equations and inequalities, *Acta Math.* 189 (1) (2002) 79–142. MR MR1946918 (2003j:35107).
- [20] S. Takeuchi, Partial flat core properties associated to the p -Laplace operator, in: *Dynamical Systems and Differential Equations. Proceedings of the 6th AIMS International Conference*, *Discrete Contin. Dyn. Syst. (Suppl.)* (2007) 965–973. MR MR2409934.
- [21] P. Tolksdorf, On the Dirichlet problem for quasilinear equations in domains with conical boundary points, *Comm. Partial Differential Equations* 8 (7) (1983) 773–817. MR MR700735 (85g:35053).
- [22] P. Tolksdorf, Regularity for a more general class of quasilinear elliptic equations, *J. Differential Equations* 51 (1) (1984) 126–150. MR MR727034 (85g:35047).
- [23] J.L. Vázquez, A strong maximum principle for some quasilinear elliptic equations, *Appl. Math. Optim.* 12 (3) (1984) 191–202. MR MR768629 (86m:35018).