



Dini estimates for nonlocal fully nonlinear elliptic equations

Hongjie Dong^{*,1}, Hong Zhang²

Division of Applied Mathematics, Brown University, 182 George Street, Providence, RI 02912, USA

Received 26 December 2016; received in revised form 13 September 2017; accepted 26 September 2017
Available online 2 October 2017

Abstract

We obtain Dini type estimates for a class of concave fully nonlinear nonlocal elliptic equations of order $\sigma \in (0, 2)$ with rough and non-symmetric kernels. The proof is based on a novel application of Campanato's approach and a refined $C^{\sigma+\alpha}$ estimate in [9].
© 2017 Elsevier Masson SAS. All rights reserved.

Keywords: Nonlocal fully nonlinear elliptic equations; Dini type estimates; A priori estimates

1. Introduction and main results

The paper is a continuation of our previous work [9], where we studied Schauder estimates for concave fully nonlinear nonlocal elliptic and parabolic equations. In particular, when the kernels are translation invariant and the data are merely bounded and measurable, we proved the C^σ estimate, which is very different from the classical theory for second-order elliptic and parabolic equations. In this paper, we consider concave fully nonlinear nonlocal elliptic equations with Dini continuous coefficients and nonhomogeneous terms, and establish a C^σ estimate under these assumptions.

The study of classical elliptic equations with Dini continuous coefficients and data has a long history. Burch [4] first considered divergence type linear elliptic equations with Dini continuous coefficients and data, and estimated the modulus of continuity of the derivatives of solutions. The corresponding result for concave fully nonlinear elliptic equations was obtained by Kovats [15], which generalized a previous result by Safonov [25]. Wang [29] studied linear non-divergence type elliptic and parabolic equations with Dini continuous coefficients and data, and gave a simple proof to estimate the modulus of continuity of the second-order derivatives of solutions. See, also [19,28,1,12,20,18], and the references therein.

Recently, there is extensive work on the regularity theory for nonlocal elliptic and parabolic equations. For example, C^α estimates, $C^{1,\alpha}$ estimates, Evans–Krylov type theorem, and Schauder estimates were established in the past

* Corresponding author.

E-mail addresses: Hongjie_Dong@brown.edu (H. Dong), Hong_Zhang@brown.edu (H. Zhang).

¹ H. Dong was partially supported by the NSF under agreements DMS-1056737 and DMS-1600593.

² H. Zhang was partially supported by the NSF under agreement DMS-1056737.

decade. See, for instance, [5,6,10,11,13,7,8,17,16,27,23], and the references therein. In particular, Mou [23] investigated a class of concave fully nonlinear nonlocal elliptic equations with smooth symmetric kernels, and obtained the C^σ estimate under a slightly stronger assumption than the usual Dini continuity on the coefficients and data. The author implemented a recursive Evans–Krylov theorem, which was first studied by Jin and Xiong [17], as well as a perturbation type argument. In this paper, by using a novel perturbation type argument, we relax the regularity assumption to simply Dini continuity and also remove the symmetry and smoothness assumptions on the kernels.

To be more specific, we are interested in fully nonlinear nonlocal elliptic equations in the form

$$\inf_{\beta \in \mathcal{A}} (L_\beta u + f_\beta) = 0, \tag{1.1}$$

where \mathcal{A} is an index set and for each $\beta \in \mathcal{A}$,

$$L_\beta u = \int_{\mathbb{R}^d} \delta u(x, y) K_\beta(x, y) dy,$$

$$\delta u(x, y) = \begin{cases} u(x + y) - u(x) - y \cdot Du(x) & \text{for } \sigma \in (1, 2), \\ u(x + y) - u(x) - y \cdot Du(x) \chi_{B_1} & \text{for } \sigma = 1, \\ u(x + y) - u(x) & \text{for } \sigma \in (0, 1), \end{cases}$$

and

$$K_\beta(x, y) = a_\beta(x, y)|y|^{-d-\sigma}.$$

This type of nonlocal operators was first investigated by Komatsu [14], Mikulevičius and Pragarauskas [21,22], and later by Dong and Kim [10,11], and Schwab and Silvestre [26], to name a few.

We assume that $a(\cdot, \cdot) \in [(2 - \sigma)\lambda, (2 - \sigma)\Lambda]$ for some ellipticity constants $0 < \lambda \leq \Lambda$, and is merely measurable with respect to the y variable. When $\sigma = 1$, we additionally assume that

$$\int_{S_r} y K_\beta(x, y) ds = 0, \tag{1.2}$$

for any $r > 0$, where S_r is the sphere of radius r centered at the origin. We say that a function f is Dini continuous if its modulus of continuity ω_f is a Dini function, i.e.,

$$\int_0^1 \omega_f(r)/r dr < \infty.$$

The following theorem is our main result, in which for simplicity we assume $u \in C^{\sigma^+}(B_1)$, which means that $u \in C^{\sigma+\varepsilon}(B_1)$ for some arbitrary $\varepsilon > 0$. This condition is only needed for $L_\beta u$ to be well defined, and may be replaced by other weaker conditions.

Theorem 1.1. *Let $\sigma \in (0, 2)$, $0 < \lambda \leq \Lambda < \infty$, and \mathcal{A} be an index set. Assume for each $\beta \in \mathcal{A}$, K_β satisfies (1.2) when $\sigma = 1$, and*

$$\begin{aligned} |a_\beta(x, y) - a_\beta(x', y)| &\leq (2 - \sigma)\Lambda\omega_a(|x - x'|), \\ |f_\beta(x) - f_\beta(x')| &\leq \omega_f(|x - x'|), \quad \sup_{\beta \in \mathcal{A}} \|f_\beta\|_{L_\infty(B_1)} < \infty, \end{aligned}$$

where ω_a and ω_f are Dini functions. Suppose $u \in C^{\sigma^+}(B_1)$ is a solution of (1.1) in B_1 and is Dini continuous in \mathbb{R}^d . Then we have the a priori estimate

$$[u]_{\sigma; B_{1/2}} \leq C \|u\|_{L_\infty} + C \sup_{\beta} \|f_\beta\|_{L_\infty(B_1)} + C \sum_{j=1}^\infty (\omega_u(2^{-j}) + \omega_f(2^{-j})) \tag{1.3}$$

where $C > 0$ is a constant depending only on $d, \sigma, \lambda, \Lambda$, and ω_a , and is uniformly bounded as $\sigma \rightarrow 2$. Moreover, when $\sigma \neq 1$, we have

$$\sup_{x_0 \in B_{1/2}} [u]_{\sigma; B_r(x_0)} \rightarrow 0 \quad \text{as } r \rightarrow 0$$

with a decay rate depending only on $d, \sigma, \lambda, \Lambda, \omega_a, \omega_f, \omega_u$, and $\sup_{\beta \in \mathcal{A}} \|f_\beta\|_{L^\infty(B_1)}$. When $\sigma = 1$, Du is uniformly continuous in $B_{1/2}$ with a modulus of continuity controlled by the quantities before.

Remark 1.2. By a careful inspection of the proofs below, one can see that the estimates above in fact only depend on $d, \sigma, \lambda, \Lambda, \sup_{\beta \in \mathcal{A}} \|f_\beta\|_{L^\infty(B_1)}$, the modulus continuity ω_f of f_β in B_1 , $\omega_a(r), \omega_u(r)$ for $r \in (0, 1)$, and $\|u\|_{L_{1,w}}$, where the weight $w = w(x)$ is equal to $(1 + |x|)^{-d-\sigma}$. In particular, u does not need to be globally bounded in \mathbb{R}^d .

Remark 1.3. In the literature, it was also common to define δu as

$$\delta u(x, y) = u(x + y) - u(x) - y \cdot Du(x) \chi_{B_1}$$

even for $\sigma \in (1, 2)$. See, for instance, [2]. In this case, the corresponding operators can be written as

$$\tilde{L}_\beta u = L_\beta u - b \cdot \nabla u,$$

where

$$b_\beta = b_\beta(x) = \int_{\mathbb{R}^d \setminus B_1} y K_\beta(x, y) dy.$$

It is easily seen that b_β is Dini continuous with a modulus of continuity ω_b depending only on d, Λ , and ω_a . Since $\sigma > 1$, $b \cdot \nabla u$ is a lower-order term. Then the same result follows by using the interpolation and iteration arguments as in the proof of Theorem 1.1.

Remark 1.4. By keeping track of the constants in the proofs below, in the symmetric case, if $\sigma \in [\sigma_0, 2)$ for some $\sigma_0 \in (0, 1)$, then the constant C in (1.3) depends on σ_0 , not σ . In the non-symmetric case, if $0 < \sigma_0 \leq \sigma \leq \sigma_1 < 1$ (or $1 < \sigma_2 \leq \sigma < 2$), then the constant C depends on σ_0 and σ_1 (or σ_2), not σ . In particular, C does not blow up as σ approaches 2.

Roughly speaking, the proof can be divided into two steps: We first show that Theorem 1.1 holds when the equation is satisfied in the whole space; Then we implement a localization argument to treat the general case. In Step one, our proof is based on a refined $C^{\sigma+\alpha}$ estimate in our previous paper [9] and a new perturbation type argument, as the standard perturbation techniques do not seem to work here. The novelty of this method is that instead of estimating C^σ semi-norm of the solution, we construct and bound certain semi-norms of the solution, see Lemmas 2.1 and 2.2. When $\sigma < 1$, such semi-norm is defined as a series of lower-order Hölder semi-norms of u . This is in the spirit of Campanato’s approach first developed in [3]. Heuristically, in order for the nonlocal operator to be well defined, the solution needs to be smoother than C^σ . To resolve this problem, we divide the integral domain into annuli, which allows us to use a lower-order semi-norm to estimate the integral in each annulus. The series of lower-order semi-norms, which turns out to be slightly stronger than the C^σ semi-norm, further implies that

$$[u]_{\sigma; B_r(x_0)} \rightarrow 0 \quad \text{as } r \rightarrow 0$$

uniformly in x_0 . In particular, when $\sigma = 1$ we are able to estimate the modulus of continuity of the gradient of solutions. The proof of the case when $\sigma \geq 1$ is more difficult than that of the case when $\sigma < 1$. This is mainly due to the fact that the series of lower-order Hölder semi-norms of the solution itself is no longer sufficient to estimate the C^σ norm. Therefore, we need to subtract a polynomial from the solution in the construction of the semi-norm. Similar idea was also used in the literature to derive Cordes–Nirenberg type estimates. See [24]. In some sense, the polynomial should be taken to minimize the series. It turns out that when $\sigma > 1$, up to a constant we can choose the polynomial to be the first-order Taylor’s expansion of the solution. The case $\sigma = 1$ is particularly challenging since the polynomial needs to be selected carefully, for which an additional mollification argument is applied.

The organization of this paper is as follows. In the next section, we introduce some notation and preliminary results that are necessary in the proof of our main theorem. Some of these results might be of independent interest. In section 3, we first prove a global version of Theorem 1.1 and then localize the result to obtain Theorem 1.1.

2. Preliminaries

For $\alpha \in (0, 2)$ and a domain $\Omega \subset \mathbb{R}^d$, we define the Lipschitz–Zygmund semi-norm by

$$[u]_{\Lambda^\alpha(\Omega)} = \sup_{\substack{x, y, z \in \Omega \\ x \neq z, x+z=2y}} |x - y|^{-\alpha} |u(x) + u(z) - 2u(y)|.$$

For $\alpha > 0$, denote $[u]_{\alpha; \Omega}$ to be the usual α -Hölder semi-norm of u in Ω , and the α -Hölder norm is defined by

$$\|u\|_{\alpha; \Omega} := [u]_{\alpha; \Omega} + \|u\|_{L^\infty(\Omega)}.$$

We use the convention $[\cdot]_\alpha = [\cdot]_{\alpha; \mathbb{R}^d}$, etc.

We will frequently use the following identity

$$\begin{aligned} & 2^j (u(x + 2^{-j} \ell) - u(x)) - (u(x + \ell) - u(x)) \\ &= \sum_{k=1}^j 2^{k-1} (2u(x + 2^{-k} \ell) - u(x + 2^{-k+1} \ell) - u(x)), \end{aligned} \tag{2.1}$$

which holds for any $\ell \in \mathbb{R}^d$ and nonnegative integer j .

Denote \mathcal{P}_1 to be the set of first-order polynomials of x .

Lemma 2.1. *Let $\alpha \in (0, \sigma)$ be a constant.*

(i) *When $\sigma \in (0, 1)$, we have*

$$[u]_\sigma \leq C \sup_{r>0} \sup_{x_0 \in \mathbb{R}^d} r^{\alpha-\sigma} [u]_{\Lambda^\alpha(B_r(x_0))} \leq C \sup_{r>0} \sup_{x_0 \in \mathbb{R}^d} r^{\alpha-\sigma} [u]_{\alpha; B_r(x_0)}, \tag{2.2}$$

where $C > 0$ is a constant depending only on d, α , and σ .

(ii) *When $\sigma \in (1, 2)$, we have*

$$[u]_\sigma \leq C \sup_{r>0} \sup_{x_0 \in \mathbb{R}^d} r^{\alpha-\sigma} [u]_{\Lambda^\alpha(B_r(x_0))} \leq C \sup_{r>0} \sup_{x_0 \in \mathbb{R}^d} r^{\alpha-\sigma} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_r(x_0)}, \tag{2.3}$$

where $C > 0$ is a constant depending only on d, α , and σ .

(iii) *When $\sigma = 1$, we have*

$$\begin{aligned} \|Du\|_{L^\infty} &\leq C \sum_{k=0}^\infty \sup_{x_0 \in \mathbb{R}^d} 2^{-k(\alpha-1)} [u]_{\Lambda^\alpha(B_{2^{-k}}(x_0))} + C \sup_{\substack{x, x' \in \mathbb{R}^d \\ |x-x'|=1}} |u(x) - u(x')| \\ &\leq C \sum_{k=0}^\infty \sup_{x_0 \in \mathbb{R}^d} 2^{-k(\alpha-1)} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-k}}(x_0)} + C \sup_{\substack{x, x' \in \mathbb{R}^d \\ |x-x'|=1}} |u(x) - u(x')|, \end{aligned} \tag{2.4}$$

where $C > 0$ is a constant depending only on d and α . Moreover, we can estimate the modulus of continuity of Du by the remainder of the summation on the right-hand side of (2.4).

Proof. First we consider the case when $\sigma \in (0, 1)$. Let $x, x' \in \mathbb{R}^d$ be two different points. Denote $h = |x - x'|$. Since

$$u(x') - u(x) = \frac{1}{2}(u(2x' - x) - u(x)) - \frac{1}{2}(u(2x' - x) - 2u(x') + u(x)),$$

we get

$$\begin{aligned} & h^{-\sigma} |u(x') - u(x)| \\ &\leq 2^{\sigma-1} (2h)^{-\sigma} (u(2x' - x) - u(x)) + h^{-\sigma} |u(2x' - x) - 2u(x') + u(x)| \\ &\leq 2^{\sigma-1} (2h)^{-\sigma} (u(2x' - x) - u(x)) + \sup_{x \in \mathbb{R}^d} h^{\alpha-\sigma} [u]_{\Lambda^\alpha(B_h(x))}. \end{aligned}$$

Taking the supremum with respect to x and x' on both sides, we get

$$[u]_\sigma \leq 2^{\sigma-1}[u]_\sigma + \sup_{x \in \mathbb{R}^d} h^{\alpha-\sigma} [u]_{\Lambda^\alpha(B_h(x))},$$

which together with the triangle inequality gives (2.2).

For $\sigma \in (1, 2)$, let $\ell \in \mathbb{R}^d$ be a unit vector and $\varepsilon \in (0, 1/16)$ be a small constant to be specified later. For any two distinct points $x, x' \in \mathbb{R}^d$, we denote $h = |x - x'|$. By the triangle inequality,

$$h^{1-\sigma} |D_\ell u(x) - D_\ell u(x')| \leq I_1 + I_2 + I_3, \tag{2.5}$$

where

$$\begin{aligned} I_1 &= h^{1-\sigma} |D_\ell u(x) - (\varepsilon h)^{-1}(u(x + \varepsilon h \ell) - u(x))|, \\ I_2 &= h^{1-\sigma} |D_\ell u(x') - (\varepsilon h)^{-1}(u(x' + \varepsilon h \ell) - u(x'))|, \\ I_3 &= h^{1-\sigma} (\varepsilon h)^{-1} |(u(x + \varepsilon h \ell) - u(x)) - (u(x' + \varepsilon h \ell) - u(x'))|. \end{aligned}$$

By the mean value theorem,

$$I_1 + I_2 \leq 2\varepsilon^{\sigma-1} [Du]_\sigma. \tag{2.6}$$

Now we choose and fix a ε sufficiently small depending only on σ such that $2\varepsilon^{\sigma-1} \leq 1/2$. Using the triangle inequality, we have

$$I_3 \leq Ch^{-\sigma} (|u(x + \varepsilon h \ell) + u(x') - 2u(\bar{x})| + |u(x' + \varepsilon h \ell) + u(x) - 2u(\bar{x})|),$$

where $\bar{x} = (x + \varepsilon h \ell + x')/2$. Thus,

$$I_3 \leq Ch^{\alpha-\sigma} [u]_{\Lambda^\alpha(B_h(\bar{x}))}. \tag{2.7}$$

Combining (2.5), (2.6), and (2.7), we get (2.3) as before.

Finally, we treat the case when $\sigma = 1$. It follows from (2.1) that

$$2^j |u(x + 2^{-j} \ell) - u(x)| \leq 2|u(x + \ell) - u(x)| + \sum_{k=1}^j 2^{-k(\alpha-1)} [u]_{\Lambda^\alpha(B_{2^{-k}}(x+2^{-k}\ell))}.$$

Taking $j \rightarrow \infty$, we obtain the desired inequality. For the continuity estimate, let $\ell \in \mathbb{R}^d$ be a unit vector. Assume that $|x - x'| \in [2^{-i-1}, 2^{-i}]$ for some positive integer i . From (2.1), for any $j \geq i + 1$,

$$\begin{aligned} &2^j (u(x + 2^{-j} \ell) - u(x)) - 2^i (u(x + 2^{-i} \ell) - u(x)) \\ &= \sum_{k=i+1}^j 2^{k-1} (2u(x + 2^{-k} \ell) - u(x + 2^{-k+1} \ell) - u(x)) \end{aligned}$$

and a similar identity holds with x' in place of x . Then we have

$$\begin{aligned} |D_\ell u(x) - D_\ell u(x')| &= \lim_{j \rightarrow \infty} \left| 2^j (u(x + 2^{-j} \ell) - u(x)) - 2^j (u(x' + 2^{-j} \ell) - u(x')) \right| \\ &\leq \left| 2^i (u(x + 2^{-i} \ell) - u(x)) - 2^i (u(x' + 2^{-i} \ell) - u(x')) \right| \\ &\quad + \sum_{k=i+1}^{\infty} \sup_{x_0 \in \mathbb{R}^d} 2^{-k(\alpha-1)} [u]_{\Lambda^\alpha(B_{2^{-k}}(x_0))}. \end{aligned}$$

By the triangle inequality, the first term on the right-hand side is bounded by

$$2^i |u(x + 2^{-i} \ell) - 2u(\bar{x}) + u(x')| + 2^i |u(x' + 2^{-i} \ell) - 2u(\bar{x}) + u(x)|$$

with $\bar{x} = (x + 2^{-i} \ell + x')/2$, which is further bounded by

$$2^{1+i(1-\alpha)} [u]_{\Lambda^\alpha(B_{2^{-i}}(\bar{x}))}.$$

Therefore,

$$|D_\ell u(x) - D_\ell u(y)| \leq C \sum_{k=i}^\infty \sup_{x_0 \in \mathbb{R}^d} 2^{-k(\alpha-1)} [u]_{\Lambda^\alpha(B_{2^{-k}}(x_0))},$$

which converges to 0 as $i \rightarrow \infty$ uniformly with respect to ℓ . The lemma is proved. \square

The following lemma will be used to estimate the error term in the freezing coefficient argument.

Lemma 2.2. *Let $\alpha \in (0, 1)$ and $\sigma \in (1, 2)$ be constants. Then for any $u \in C^1$, we have*

$$\sum_{k=0}^\infty 2^{k(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u - P_{x_0}u]_{\alpha; B_{2^{-k}}(x_0)} \leq C \sum_{k=0}^\infty 2^{k(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-k}}(x_0)} \tag{2.8}$$

and

$$\sum_{k=0}^\infty 2^{k\sigma} \sup_{x_0 \in \mathbb{R}^d} \|u - P_{x_0}u\|_{L^\infty(B_{2^{-k}}(x_0))} \leq C \sum_{k=0}^\infty 2^{k(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\Lambda^\alpha(B_{2^{-k}}(x_0))}, \tag{2.9}$$

where $P_{x_0}u$ is the first-order Taylor expansion of u at x_0 , and $C > 0$ is a constant depending only on d, α , and σ .

Proof. Denote

$$b_k := 2^{k(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-k}}(x_0)}.$$

Then for any $x_0 \in \mathbb{R}^d$ and each $k = 0, 1, \dots$, there exists $p_k \in \mathcal{P}_1$ such that

$$[u - p_k]_{\alpha; B_{2^{-k}}(x_0)} \leq 2b_k 2^{-k(\sigma-\alpha)}.$$

By the triangle inequality, for $k \geq 1$ we have

$$[p_{k-1} - p_k]_{\alpha; B_{2^{-k}}(x_0)} \leq 2b_k 2^{-k(\sigma-\alpha)} + 2b_{k-1} 2^{-(k-1)(\sigma-\alpha)}. \tag{2.10}$$

It is easily seen that

$$[p_{k-1} - p_k]_{\alpha; B_{2^{-k}}(x_0)} = |\nabla p_{k-1} - \nabla p_k| 2^{-(k-1)(1-\alpha)},$$

which together with (2.10) implies that

$$|\nabla p_{k-1} - \nabla p_k| \leq C(b_k + b_{k-1}) 2^{-k(\sigma-1)}. \tag{2.11}$$

Since $\sum_0^k b_k < \infty$, from (2.11) we see that $\{\nabla p_k\}$ is a Cauchy sequence in \mathbb{R}^d . Let $q = q(x_0) \in \mathbb{R}^d$ be its limit, which clearly satisfies for each $k \geq 0$,

$$|q - \nabla p_k| \leq C \sum_{j=k}^\infty 2^{-j(\sigma-1)} b_j.$$

By the triangle inequality, we get

$$\begin{aligned} [u - q \cdot x]_{\alpha; B_{2^{-k}}(x_0)} &\leq [u - p_k]_{\alpha; B_{2^{-k}}(x_0)} + [p_k - q \cdot x]_{\alpha; B_{2^{-k}}(x_0)} \\ &\leq C 2^{-k(1-\alpha)} \sum_{j=k}^\infty 2^{-j(\sigma-1)} b_j \leq C 2^{-k(\sigma-\alpha)}, \end{aligned} \tag{2.12}$$

which implies that

$$\|u - u(x_0) - q \cdot (x - x_0)\|_{L^\infty(B_{2^{-k}}(x_0))} \leq C 2^{-k\sigma},$$

and thus $q = \nabla u(x_0)$. It then follows (2.12) that

$$\begin{aligned} \sum_{k=0}^{\infty} 2^{k(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u - P_{x_0} u]_{\alpha; B_{2^{-k}}(x_0)} &\leq C \sum_{k=0}^{\infty} 2^{k(\sigma-1)} \sum_{j=k}^{\infty} 2^{-j(\sigma-1)} b_j \\ &= C \sum_{j=0}^{\infty} 2^{-j(\sigma-1)} b_j \sum_{k=0}^j 2^{k(\sigma-1)} \leq C \sum_{j=0}^{\infty} b_j. \end{aligned}$$

This completes the proof of (2.8).

Next we show (2.9). For any $x \in B_{2^{-k}}$, it follows from (2.1) that for $j \geq 1$,

$$\begin{aligned} u(x) - u(0) - 2^j(u(2^{-j}x) - u(0)) \\ = \sum_{i=0}^{j-1} 2^i(u(2^{-i}x) + u(0) - 2u(2^{-i-1}x)). \end{aligned}$$

Sending $j \rightarrow \infty$, we obtain

$$\begin{aligned} |u(x) - u(0) - x \cdot \nabla u(0)| &\leq \sum_{i=0}^{\infty} 2^i |u(2^{-i}x) + u(0) - 2u(2^{-i-1}x)| \\ &\leq 2^{-\alpha} \sum_{i=0}^{\infty} 2^{i-(i+k)\alpha} [u]_{\Lambda^\alpha(B_{2^{-(k+i)}})} = 2^{-\alpha} \sum_{i=k}^{\infty} 2^{i-k-i\alpha} [u]_{\Lambda^\alpha(B_{2^{-i}})}, \end{aligned}$$

where we shifted the index in the last equality. Therefore, by shifting the coordinates and sum in k , we have

$$\begin{aligned} \sum_{k=0}^{\infty} 2^{k\sigma} \sup_{x_0 \in \mathbb{R}^d} \|u - P_{x_0} u\|_{L^\infty(B_{2^{-k}}(x_0))} \\ \leq C \sum_{k=0}^{\infty} 2^{k(\sigma-1)} \sum_{i=k}^{\infty} 2^{i(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\Lambda^\alpha(B_{2^{-i}}(x_0))} \\ = C \sum_{i=0}^{\infty} 2^{i(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\Lambda^\alpha(B_{2^{-i}}(x_0))} \sum_{k=0}^i 2^{k(\sigma-1)} \\ \leq C \sum_{i=0}^{\infty} 2^{i(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\Lambda^\alpha(B_{2^{-i}}(x_0))}, \end{aligned}$$

where we switched the order of the summations in the second equality and in the last inequality we used the condition that $\sigma > 1$. The lemma is proved. \square

Let $\zeta \in C_0^\infty(B_1)$ be a nonnegative radial function with unit integral. For $R > 0$, we define the mollification of a function u by

$$u^{(R)}(x) = \int_{\mathbb{R}^d} u(x - Ry)\zeta(y) dy.$$

The next lemmas will be used in the estimate of M_j in Proposition 3.1 when $\sigma = 1$.

Lemma 2.3. *Let $\beta \in (0, 1]$, $\alpha \in (0, 1 + \beta)$, and $0 < R \leq R_1 < \infty$. Then for any $u \in \Lambda^\alpha(B_{2R_1})$, we have*

$$[Du^{(R)}]_{\beta; B_{R_1}} \leq C(d, \beta, \alpha) R^{\alpha-1-\beta} [u]_{\Lambda^\alpha(B_{2R_1})}. \tag{2.13}$$

Proof. We begin by estimating $\|D_\ell^2 u\|_{0; B_{R_1}}$ for a fixed unit vector $\ell \in \mathbb{R}^d$. Because $D_\ell^2 \zeta$ is even with respect to x and has zero integral, using integration by parts we have for any $x \in B_{R_1}$,

$$\begin{aligned}
 |D_\ell^2 u^{(R)}(x)| &= R^{-2} \left| \int_{\mathbb{R}^d} u(x - Ry) D_\ell^2 \zeta(y) dy \right| \\
 &= \frac{R^{-2}}{2} \left| \int_{\mathbb{R}^d} (u(x - Ry) + u(x + Ry) - 2u(x)) D_\ell^2 \zeta(y) dy \right| \\
 &\leq CR^{\alpha-2} [u]_{\Lambda^\alpha(B_{2R_1})} \int_{\mathbb{R}^d} |y|^\alpha D_\ell^2 \zeta(y) dy \leq CR^{\alpha-2} [u]_{\Lambda^\alpha(B_{2R_1})}.
 \end{aligned}$$

Using the identity, $2D_{ij}u = 2D_\ell^2 u - D_i^2 u - D_j^2 u$, where $\ell = (e_i + e_j)/\sqrt{2}$, we obtain the desired inequality (2.13) when $\beta = 1$.

Next we consider the case when $\beta \in (0, 1)$. We follow the proof of Lemma 2.1. Let $\ell \in \mathbb{R}^d$ be a unit vector, and $\varepsilon \in (0, 1/16)$ be a small constant to be specified later. For any two distinct points $x, x' \in B_{R_1}$, let $h = |x - x'| (< 2R_1)$. It is easily seen that there exist two points $y \in B_{\varepsilon h}(x) \cap B_{R_1}$ and $y' \in B_{\varepsilon h}(x') \cap B_{R_1}$ such that

$$y + \varepsilon h \ell \in B_{\varepsilon h}(x) \cap B_{R_1}, \quad y' + \varepsilon h \ell \in B_{\varepsilon h}(x') \cap B_{R_1}.$$

By the triangle inequality,

$$h^{-\beta} |D_\ell u^{(R)}(x) - D_\ell u^{(R)}(x')| \leq I_1 + I_2 + I_3,$$

where

$$\begin{aligned}
 I_1 &= h^{-\beta} |D_\ell u^{(R)}(x) - (\varepsilon h)^{-1} (u^{(R)}(y + \varepsilon h \ell) - u^{(R)}(y))|, \\
 I_2 &= h^{-\beta} |D_\ell u^{(R)}(x') - (\varepsilon h)^{-1} (u^{(R)}(y' + \varepsilon h \ell) - u^{(R)}(y'))|, \\
 I_3 &= h^{-\beta} (\varepsilon h)^{-1} |(u^{(R)}(y + \varepsilon h \ell) - u^{(R)}(y)) - (u^{(R)}(y' + \varepsilon h \ell) - u^{(R)}(y'))|.
 \end{aligned}$$

By the mean value theorem,

$$I_1 + I_2 \leq 2\varepsilon^\beta [Du^{(R)}]_{\beta; B_{R_1}}. \tag{2.14}$$

Now we choose ε depending only on d and β such that $2\varepsilon^\beta \leq 1/2$. To estimate I_3 , we consider two cases. If $h > R$, by the triangle inequality, we have

$$\begin{aligned}
 I_3 &\leq Ch^{-1-\beta} (|u^{(R)}(y + \varepsilon h \ell) + u^{(R)}(y') - 2u^{(R)}(\bar{y})| \\
 &\quad + |u^{(R)}(y' + \varepsilon h \ell) + u^{(R)}(y) - 2u^{(R)}(\bar{y})|),
 \end{aligned}$$

where $\bar{y} = (y + \varepsilon h \ell + y')/2$. Then by the Minkowski inequality,

$$I_3 \leq Ch^{\alpha-1-\beta} [u^{(R)}]_{\Lambda^\alpha(B_{R_1})} \leq CR^{\alpha-1-\beta} [u]_{\Lambda^\alpha(B_{2R_1})}. \tag{2.15}$$

On the other hand, if $h \in (0, R)$, by the mean value theorem and (2.13) with $\beta = 1$,

$$I_3 \leq Ch^{1-\beta} [Du^{(R)}]_{1; B_{R_1}} \leq Ch^{1-\beta} R^{\alpha-2} [u]_{\Lambda^\alpha(B_{2R_1})} \leq CR^{\alpha-1-\beta} [u]_{\Lambda^\alpha(B_{2R_1})}. \tag{2.16}$$

Combining (2.14), (2.15), and (2.16), we obtain

$$h^{-\beta} |D_\ell u^{(R)}(x) - D_\ell u^{(R)}(x')| \leq \frac{1}{2} [Du^{(R)}]_{\beta; B_{R_1}} + CR^{\alpha-1-\beta} [u]_{\Lambda^\alpha(B_{2R_1})}.$$

Taking the supremum of the left-hand side above with respect to unit vector $\ell \in \mathbb{R}^d$ and $x, x' \in B_{R_1}$, we immediately get (2.13). The lemma is proved. \square

Lemma 2.4. *Let $\alpha \in (0, 1)$, $\beta \in (0, 1)$, and $R > 0$ be constants. Let $p = p(x)$ be the first-order Taylor expansion of $u^{(R)}$ at the origin and $\tilde{u} = u - p$. Then for any integer $j \geq 0$, we have*

$$\|\tilde{u}\|_{L^\infty(B_{2^{j+1}R})} \leq C2^{j(1+\beta)} R^\alpha [u]_{\Lambda^\alpha(B_{2^{j+2}R})}, \tag{2.17}$$

$$\sup_{\substack{x, x' \in B_{2^j R} \\ 0 < |x - x'| < 2R}} \frac{|\tilde{u}(x) - \tilde{u}(x')|}{|x - x'|^\alpha} \leq C2^{j\beta} [u]_{\Lambda^\alpha(B_{2^{j+2}R})}, \tag{2.18}$$

where $C > 0$ is a constant depending only on d, β , and α .

Proof. Since $\zeta \in C_0^\infty(B_1)$ is radial and has unit integral, we have for any $x \in B_{2^{j+1}R}$,

$$\begin{aligned} & |u^{(R)}(x) - u(x)| \\ &= \left| \frac{1}{2} \int_{\mathbb{R}^d} (u(x + Ry) - u(x - Ry) - 2u(x)) \zeta(y) dy \right| \leq CR^\alpha [u]_{\Lambda^\alpha(B_{2^{j+2}R})}. \end{aligned} \tag{2.19}$$

By the mean value theorem and Lemma 2.3, for any $x \in B_{2^{j+1}R}$,

$$|u^{(R)}(x) - p(x)| \leq C(2^{j+1}R)^{1+\beta} [u^{(R)}]_{1+\beta; B_{2^{j+1}R}} \leq C2^{j(1+\beta)} R^\alpha [u]_{\Lambda^\alpha(B_{2^{j+2}R})},$$

which together with (2.19) implies (2.17). Next we show (2.18). For any two distinct points $x, x' \in B_{2^j R}$ satisfying $0 < |x - x'| < 2R$, denote $h = |x - x'| (< 2R)$. Let k be the largest nonnegative integer such that $2^k(x' - x) + x \in B_{2^{j+1}R}$. Clearly,

$$2^k h \in (2^{j-1}R, 2^{j+2}R). \tag{2.20}$$

It follows from (2.1) that

$$\begin{aligned} \tilde{u}(x') - \tilde{u}(x) &= 2^{-k} (\tilde{u}(2^k(x' - x) + x) - \tilde{u}(x)) \\ &+ \sum_{i=0}^{k-1} 2^{-i-1} (2\tilde{u}(2^i(x' - x) + x) - \tilde{u}(x) - \tilde{u}(2^{i+1}(x' - x) + x)). \end{aligned} \tag{2.21}$$

By (2.20), (2.21), and (2.17), we obtain

$$\begin{aligned} h^{-\alpha} |\tilde{u}(x') - \tilde{u}(x)| &\leq 2^{-k+1} h^{-\alpha} \|\tilde{u}\|_{L^\infty(B_{2^{j+1}R})} + C[u]_{\Lambda^\alpha(B_{2^{j+1}R})} \\ &\leq C2^{-j} R^{-1} h^{1-\alpha} \cdot 2^{j(1+\beta)} R^\alpha [u]_{\Lambda^\alpha(B_{2^{j+2}R})} + C[u]_{\Lambda^\alpha(B_{2^{j+1}R})} \\ &\leq C2^{j\beta} [u]_{\Lambda^\alpha(B_{2^{j+2}R})}, \end{aligned}$$

where we used $h < 2R$ in the last inequality. The lemma is proved. \square

3. Proofs

The following proposition is a further refinement of [9, Corollary 4.6].

Proposition 3.1. *Let $\sigma \in (0, 2)$ and $0 < \lambda \leq \Lambda$. Assume that for any $\beta \in \mathcal{A}$, K_β only depends on y . There is a constant $\hat{\alpha} \in (0, 1)$ depending on d, σ, λ , and Λ so that the following holds. Let $\alpha \in (0, \hat{\alpha})$. Suppose $u \in C^{\sigma+\alpha}(B_1) \cap C^\alpha(\mathbb{R}^d)$ is a solution of*

$$\inf_{\beta \in \mathcal{A}} (L_\beta u + f_\beta) = 0 \quad \text{in } B_1.$$

Then,

$$[u]_{\alpha+\sigma; B_{1/2}} \leq C \sum_{j=1}^{\infty} 2^{-j\sigma} M_j + C \sup_{\beta} [f_\beta]_{\alpha; B_1},$$

where

$$M_j = \sup_{x, x' \in B_{2^j}, 0 < |x-x'| < 2} \frac{|u(x) - u(x')|}{|x - x'|^\alpha}$$

and $C > 0$ depends only on $d, \lambda, \Lambda, \alpha,$ and $\sigma,$ and is uniformly bounded as $\sigma \rightarrow 2.$

Proof. This follows from the proof of [9, Corollary 4.6] by observing that in the estimate of $[h_\beta]_{\alpha; B_1},$ the term $[u]_{\alpha; B_{2^j}}$ can be replaced by $M_j.$ Moreover, by replacing u by $u - u(0),$ we see that

$$\|u\|_{\alpha; B_2} \leq C[u]_{\alpha; B_2}.$$

The lemma is proved. \square

Proposition 3.2. *Suppose that (1.1) is satisfied in $\mathbb{R}^d.$ Then under the conditions of Theorem 1.1, we have*

$$[u]_\sigma \leq C\|u\|_{L_\infty} + C \sum_{k=1}^\infty \omega_f(2^{-k}), \tag{3.1}$$

where $C > 0$ is a constant depending only on $d, \lambda, \Lambda, \omega_\alpha,$ and $\sigma,$ and is uniformly bounded as $\sigma \rightarrow 2.$

Proof. Case 1: $\sigma \in (0, 1).$ For $k \in \mathbb{N},$ let v be the solution of

$$\begin{cases} \inf_{\beta \in \mathcal{A}} (L_\beta(0)v + f_\beta(0)) = 0 & \text{in } B_{2^{-k}} \\ v = u & \text{in } B_{2^{-k}}^c \end{cases}, \tag{3.2}$$

where $L_\beta(0)$ is the operator with kernel $K_\beta(0, y).$ Then by Proposition 3.1 with scaling, we have

$$\begin{aligned} [v]_{\alpha+\sigma; B_{2^{-k-1}}} &\leq C \sum_{j=1}^\infty 2^{(k-j)\sigma} M_j + C2^{k\sigma} [v]_{\alpha; B_{2^{-k}}} \\ &\leq C \sum_{j=1}^k 2^{(k-j)\sigma} M_j + C[u]_\alpha + C2^{k\sigma} [v]_{\alpha; B_{2^{-k}}}, \end{aligned} \tag{3.3}$$

where $\alpha \in (0, \hat{\alpha})$ satisfying $\sigma + \alpha < 1$ and

$$M_j = \sup_{x, x' \in B_{2^{j-k}}, 0 < |x-x'| < 2^{-k+1}} \frac{|u(x) - u(x')|}{|x - x'|^\alpha}.$$

Let $k_0, k_1 \geq 1$ be integers to be specified. From (3.3), we get

$$[v]_{\alpha; B_{2^{-k-k_0}}} \leq C2^{-(k+k_0)\sigma} \sum_{j=1}^k 2^{(k-j)\sigma} M_j + C2^{-(k+k_0)\sigma} [u]_\alpha + C2^{-k_0\sigma} [v]_{\alpha; B_{2^{-k}}}. \tag{3.4}$$

Next, $w := u - v$ satisfies

$$\begin{cases} \mathcal{M}^+ w \geq -C_k & \text{in } B_{2^{-k}}, \\ \mathcal{M}^- w \leq C_k & \text{in } B_{2^{-k}}, \\ w = 0 & \text{in } B_{2^{-k}}^c, \end{cases} \tag{3.5}$$

where

$$C_k = \sup_{\beta \in \mathcal{A}} \|f_\beta - f_\beta(0) + (L_\beta - L_\beta(0))u\|_{L_\infty(B_{2^{-k}})}.$$

It is easily seen that

$$\begin{aligned}
 C_k &\leq \omega_f(2^{-k}) + C\omega_a(2^{-k}) \int_{\mathbb{R}^d} |u(x+y) - u(x)| |y|^{-d-\sigma} dy \\
 &\leq \omega_f(2^{-k}) + C\omega_a(2^{-k}) \left(\sup_{x_0 \in B_{2^{-k}}} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} [u]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L^\infty} \right).
 \end{aligned}$$

Then by the Hölder estimate [9, Lemma 2.5], we have

$$\begin{aligned}
 [w]_{\alpha; B_{2^{-k}}} &\leq C2^{-k(\sigma-\alpha)} C_k \\
 &\leq C2^{-k(\sigma-\alpha)} \left(\omega_f(2^{-k}) + \omega_a(2^{-k}) \left(\sup_{x_0 \in B_{2^{-k}}} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} [u]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L^\infty} \right) \right).
 \end{aligned} \tag{3.6}$$

Combining (3.4) and (3.6) yields

$$\begin{aligned}
 &2^{(k+k_0)(\sigma-\alpha)} [u]_{\alpha; B_{2^{-k-k_0}}} \\
 &\leq C2^{-(k+k_0)\alpha} \sum_{j=1}^k 2^{(k-j)\sigma} [u]_{\alpha; B_{2^{j-k}}} + C2^{-(k+k_0)\alpha} [u]_{\alpha} + C2^{-k_0\alpha} 2^{k(\sigma-\alpha)} [u]_{\alpha; B_{2^{-k}}} \\
 &\quad + C2^{k_0(\sigma-\alpha)} \left(\omega_f(2^{-k}) + \omega_a(2^{-k}) \left(\sup_{x_0 \in B_{2^{-k}}} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} [u]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L^\infty} \right) \right).
 \end{aligned} \tag{3.7}$$

Shifting the coordinates, from (3.7) we get

$$\begin{aligned}
 &2^{(k+k_0)(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-k-k_0}}(x_0)} \\
 &\leq C2^{-(k+k_0)\alpha} \sup_{x_0 \in \mathbb{R}^d} \sum_{j=1}^k 2^{(k-j)\sigma} [u]_{\alpha; B_{2^{j-k}}(x_0)} + C2^{-(k+k_0)\alpha} [u]_{\alpha} \\
 &\quad + C2^{-k_0\alpha} 2^{k(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-k}}(x_0)} + C2^{k_0(\sigma-\alpha)} \left(\omega_f(2^{-k}) \right. \\
 &\quad \left. + \omega_a(2^{-k}) \left(\sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} [u]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L^\infty} \right) \right).
 \end{aligned} \tag{3.8}$$

We take the summation of (3.8) in $k = k_1, k_1 + 1, \dots$ to obtain

$$\begin{aligned}
 &\sum_{k=k_1}^{\infty} 2^{(k+k_0)(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-k-k_0}}(x_0)} \\
 &\leq C \sum_{k=k_1}^{\infty} 2^{-(k+k_0)\alpha} \left(\sup_{x_0 \in \mathbb{R}^d} \sum_{j=1}^k 2^{(k-j)\sigma} [u]_{\alpha; B_{2^{j-k}}(x_0)} \right) + C2^{-(k_1+k_0)\alpha} [u]_{\alpha} \\
 &\quad + C2^{-k_0\alpha} \sum_{k=k_1}^{\infty} 2^{k(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-k}}(x_0)} + C2^{k_0(\sigma-\alpha)} \sum_{k=k_1}^{\infty} \left(\omega_f(2^{-k}) \right. \\
 &\quad \left. + \omega_a(2^{-k}) \left(\sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L^\infty} \right) \right),
 \end{aligned}$$

which by switching the order of summations is further bounded by

$$\begin{aligned}
 & C2^{-k_0\alpha} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-j}}(x_0)} \\
 & + C2^{-(k_1+k_0)\alpha} [u]_{\alpha} + C2^{k_0(\sigma-\alpha)} \sum_{k=k_1}^{\infty} \omega_f(2^{-k}) \\
 & + C2^{k_0(\sigma-\alpha)} \sum_{k=k_1}^{\infty} \omega_a(2^{-k}) \cdot \left(\sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L_{\infty}} \right).
 \end{aligned}$$

The bound above together with the obvious inequality

$$\sum_{j=0}^{k_1+k_0-1} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-j}}(x_0)} \leq C2^{(k_1+k_0)(\sigma-\alpha)} [u]_{\alpha},$$

implies that

$$\begin{aligned}
 & \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-j}}(x_0)} \leq C2^{-k_0\alpha} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-j}}(x_0)} \\
 & + C2^{(k_1+k_0)(\sigma-\alpha)} [u]_{\alpha} + C2^{k_0(\sigma-\alpha)} \sum_{k=k_1}^{\infty} \omega_f(2^{-k}) \\
 & + C2^{k_0(\sigma-\alpha)} \sum_{k=k_1}^{\infty} \omega_a(2^{-k}) \cdot \left(\sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-j}}(x_0)} + C\|u\|_{L_{\infty}} \right).
 \end{aligned}$$

By first choosing k_0 sufficiently large and then k_1 sufficiently large, we get

$$\sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u]_{\alpha; B_{2^{-j}}(x_0)} \leq C\|u\|_{\alpha} + C \sum_{k=1}^{\infty} \omega_f(2^{-k}),$$

which together with Lemma 2.1 (i) and the interpolation inequality gives (3.1).

Case 2: $\sigma \in (1, 2)$. For $k \in \mathbb{N}$, let v_M be the solution of

$$\begin{cases} \inf_{\beta \in \mathcal{A}} (L_{\beta}(0)v_M + f_{\beta}(0)) = 0 & \text{in } B_{2^{-k}} \\ v_M = g_M & \text{in } B_{2^{-k}}^c \end{cases},$$

where $M \geq 2\|u - p_0\|_{L_{\infty}(B_{2^{-k}})}$ is a constant to be specified later,

$$g_M = \max(\min(u - p_0, M), -M),$$

and p_0 is the first-order Taylor’s expansion of u at the origin.

By Proposition 3.1, instead of (3.3), we have

$$\begin{aligned}
 [v_M]_{\alpha+\sigma; B_{2^{-k-1}}} & \leq C \sum_{j=0}^{\infty} 2^{(k-j)\sigma} M_j + C2^{k\sigma} [v_M]_{\alpha; B_{2^{-k}}} \\
 & \leq C \sum_{j=0}^k 2^{(k-j)\sigma} M_j + C\|Du\|_{L_{\infty}} + C2^{k\sigma} [v_M]_{\alpha; B_{2^{-k}}},
 \end{aligned} \tag{3.9}$$

where $\alpha \in (0, \hat{\alpha})$ and

$$M_j = \sup_{x, x' \in B_{2^{-j-k}}, 0 < |x-x'| < 2^{-k+1}} \frac{|u(x) - p_0(x) - u(x') + p_0(x')|}{|x - x'|^{\alpha}}.$$

From (3.9) and the mean value formula,

$$\begin{aligned} \|v_M - p_1\|_{L_\infty(B_{2^{-k-k_0}})} &\leq C2^{-(k+k_0)(\sigma+\alpha)} \sum_{j=0}^k 2^{(k-j)\sigma} M_j \\ &+ C2^{-(k+k_0)(\sigma+\alpha)} \|Du\|_{L_\infty} + C2^{-k\alpha-k_0(\sigma+\alpha)} [v_M]_{\alpha; B_{2^{-k}}}, \end{aligned}$$

where p_1 is the first-order Taylor’s expansion of v_M at the origin. The above inequality, (3.9), and the interpolation inequality imply

$$\begin{aligned} [v_M - p_1]_{\alpha; B_{2^{-k-k_0}}} &\leq C2^{-(k+k_0)\sigma} \sum_{j=0}^k 2^{(k-j)\sigma} M_j \\ &+ C2^{-(k+k_0)\sigma} \|Du\|_{L_\infty} + C2^{-k_0\sigma} [v_M]_{\alpha; B_{2^{-k}}}. \end{aligned} \tag{3.10}$$

Next $w_M := g_M - v_M$ satisfies

$$\begin{cases} \mathcal{M}^+ w_M \geq h_M - C_k & \text{in } B_{2^{-k}}, \\ \mathcal{M}^- w_M \leq \hat{h}_M + C_k & \text{in } B_{2^{-k}}, \\ w_M = 0 & \text{in } B_{2^{-k}}^c, \end{cases}$$

where

$$h_M := \mathcal{M}^-(g_M - (u - p_0)), \quad \hat{h}_M := \mathcal{M}^+(g_M - (u - p_0)).$$

By the dominated convergence theorem, it is easy to see that

$$\|h_M\|_{L_\infty(B_{2^{-k}})}, \|\hat{h}_M\|_{L_\infty(B_{2^{-k}})} \rightarrow 0 \quad \text{as } M \rightarrow \infty.$$

By the same argument as in the previous case,

$$C_k \leq \omega_f(2^{-k}) + C\omega_a(2^{-k}) \left(\sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^\infty 2^{j(\sigma-\alpha)} [u - P_{x_0}u]_{\alpha; B_{2^{-j}}(x_0)} + \|Du\|_{L_\infty} \right).$$

Thus similar to (3.6), choosing M sufficiently large so that

$$\|h_M\|_{L_\infty(B_{2^{-k}})}, \|\hat{h}_M\|_{L_\infty(B_{2^{-k}})} \leq C_k/2,$$

we have

$$\begin{aligned} [w_M]_{\alpha; B_{2^{-k}}} &\leq C2^{-k(\sigma-\alpha)} \left(\omega_f(2^{-k}) + \omega_a(2^{-k}) \|Du\|_{L_\infty} \right. \\ &\quad \left. + \omega_a(2^{-k}) \sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^\infty 2^{j(\sigma-\alpha)} [u - P_{x_0}u]_{\alpha; B_{2^{-j}}(x_0)} \right). \end{aligned} \tag{3.11}$$

Combining (3.10) and (3.11), similar to (3.8), we obtain

$$\begin{aligned} &2^{(k+k_0)(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-k-k_0}}(x_0)} \\ &\leq C2^{-(k+k_0)\alpha} \sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^k 2^{(k-j)\sigma} [u - P_{x_0}u]_{\alpha; B_{2^{j-k}}(x_0)} + C2^{-(k+k_0)\alpha} \|Du\|_{L_\infty} \\ &\quad + C2^{-k_0\alpha} 2^{k(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u - P_{x_0}]_{\alpha; B_{2^{-k}}(x_0)} + C2^{k_0(\sigma-\alpha)} \left(\omega_f(2^{-k}) \right. \\ &\quad \left. + \omega_a(2^{-k}) \left(\sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^\infty 2^{j(\sigma-\alpha)} [u - P_{x_0}u]_{\alpha; B_{2^{-j}}(x_0)} + \|Du\|_{L_\infty} \right) \right). \end{aligned} \tag{3.12}$$

Using (3.12), as before we get

$$\begin{aligned}
 & \sum_{k=k_1}^{\infty} 2^{(k+k_0)(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-k-k_0}}(x_0)} \\
 & \leq C 2^{-k_0\alpha} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u - P_{x_0} u]_{\alpha; B_{2^{-j}}(x_0)} \\
 & \quad + C 2^{-(k_1+k_0)\alpha} \|u\|_1 + C 2^{k_0(\sigma-\alpha)} \sum_{k=k_1}^{\infty} \omega_f(2^{-k}) \\
 & \quad + C 2^{k_0(\sigma-\alpha)} \sum_{k=k_1}^{\infty} \omega_{\alpha}(2^{-k}) \cdot \sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} [u - P_{x_0} u]_{\alpha; B_{2^{-j}}(x_0)}, \tag{3.13}
 \end{aligned}$$

and

$$\begin{aligned}
 & \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} \\
 & \leq C 2^{-k_0\alpha} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} [u - P_{x_0} u]_{\alpha; B_{2^{-j}}(x_0)} \\
 & \quad + C 2^{(k_1+k_0)(\sigma-\alpha)} \|u\|_1 + C 2^{k_0(\sigma-\alpha)} \sum_{k=k_1}^{\infty} \omega_f(2^{-k}) \\
 & \quad + C 2^{k_0(\sigma-\alpha)} \sum_{k=k_1}^{\infty} \omega_{\alpha}(2^{-k}) \cdot \sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} [u - P_{x_0} u]_{\alpha; B_{2^{-j}}(x_0)}.
 \end{aligned}$$

By choosing k_0 and k_1 sufficiently large and applying Lemma 2.2, we obtain

$$\sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} \leq C \|u\|_1 + C \sum_{k=1}^{\infty} \omega_f(2^{-k}). \tag{3.14}$$

Finally, by Lemma 2.1 (ii) and the interpolation inequality, we get (3.1).

Case 3: $\sigma = 1$. We proceed as in the previous case, but instead take p_0 to be the first-order Taylor’s expansion of the mollification $u^{(2^{-k})}$ at the origin. We also assume that the solution v to (3.2) exists without carrying out another approximation argument. By Proposition 3.1 and Lemma 2.4 with $\beta = \alpha/2$,

$$\begin{aligned}
 [v]_{\alpha+1; B_{2^{-k-1}}} & \leq C \sum_{j=0}^{\infty} 2^{k-j} M_j + C 2^k [v]_{\alpha; B_{2^{-k}}} \\
 & \leq C \sum_{j=0}^{\infty} 2^{k-j+j\alpha/2} [u]_{\Lambda^{\alpha}(B_{2^{j-k}})} + C 2^k [v]_{\alpha; B_{2^{-k}}} \\
 & \leq C \sum_{j=0}^k 2^{k-j+j\alpha/2} [u]_{\Lambda^{\alpha}(B_{2^{j-k}})} + C 2^{k\alpha/2} [u]_{\alpha} + C 2^k [v]_{\alpha; B_{2^{-k}}}. \tag{3.15}
 \end{aligned}$$

From (3.15) and the interpolation inequality, we obtain

$$\begin{aligned}
 & [v - p_1]_{\alpha; B_{2^{-k-k_0}}} \\
 & \leq C 2^{-(k+k_0)} \sum_{j=0}^k 2^{k-j+j\alpha/2} [u]_{\Lambda^{\alpha}(B_{2^{j-k}})} + C 2^{-(k+k_0)+k\alpha/2} [u]_{\alpha} + C 2^{-k_0} [v]_{\alpha; B_{2^{-k}}}
 \end{aligned}$$

$$\begin{aligned} &\leq C2^{-(k+k_0)} \sum_{j=0}^k 2^{k-j+j\alpha/2} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{j-k}}} \\ &\quad + C2^{-(k+k_0)+k\alpha/2} [u]_{\alpha} + C2^{-k_0} [v]_{\alpha; B_{2^{-k}}}, \end{aligned} \tag{3.16}$$

where p_1 is the first-order Taylor’s expansion of v at the origin. Next $w := u - p_0 - v$ satisfies (3.5), where by the cancelation property (1.2),

$$C_k \leq \omega_f(2^{-k}) + C\omega_a(2^{-k}) \left(\sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^{\infty} 2^{j(1-\alpha)} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L_{\infty}} \right).$$

Therefore, similar to (3.6), we have

$$\begin{aligned} [w]_{\alpha; B_{2^{-k}}} &\leq C2^{-k(1-\alpha)} \left(\omega_f(2^{-k}) \right. \\ &\quad \left. + \omega_a(2^{-k}) \left(\sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^{\infty} 2^{j(1-\alpha)} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L_{\infty}} \right) \right). \end{aligned} \tag{3.17}$$

Notice that from (2.18) and the triangle inequality,

$$\begin{aligned} [v]_{\alpha; B_{2^{-k}}} &\leq [w]_{\alpha; B_{2^{-k}}} + [u - p_0]_{\alpha; B_{2^{-k}}} \\ &\leq [w]_{\alpha; B_{2^{-k}}} + C[u]_{\Lambda^{\alpha}(B_{2^{-k+2}})} \leq [w]_{\alpha; B_{2^{-k}}} + C \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-k+2}}}. \end{aligned}$$

Similar to (3.8), combining (3.16), (3.17), and the inequality above, we obtain

$$\begin{aligned} &2^{(k+k_0)(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-k-k_0}}(x_0)} \\ &\leq C2^{-(k+k_0)\alpha} \sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^k 2^{k-j+j\alpha/2} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{j-k}}(x_0)} + C2^{-(k/2+k_0)\alpha} [u]_{\alpha} \\ &\quad + C2^{-k_0\alpha+(k-2)(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-k+2}}(x_0)} + C2^{k_0(1-\alpha)} \left(\omega_f(2^{-k}) \right. \\ &\quad \left. + \omega_a(2^{-k}) \left(\sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^{\infty} 2^{j(1-\alpha)} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L_{\infty}} \right) \right), \end{aligned}$$

which by summing in $k = k_1, k_1 + 1, \dots$ implies that

$$\begin{aligned} &\sum_{k=k_1}^{\infty} 2^{(k+k_0)(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-k-k_0}}(x_0)} \\ &\leq C2^{-k_0\alpha} \sum_{j=0}^{\infty} 2^{j(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} \\ &\quad + C2^{-(k/2+k_0)\alpha} [u]_{\alpha} + C2^{k_0(1-\alpha)} \sum_{k=k_1}^{\infty} \omega_f(2^{-k}) + C2^{k_0(1-\alpha)} \sum_{k=k_1}^{\infty} \omega_a(2^{-k}) \\ &\quad \cdot \left(\sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^{\infty} 2^{j(1-\alpha)} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L_{\infty}} \right), \end{aligned}$$

where for the first term on the right-hand side, we switched the order of summations to get

$$\begin{aligned} & \sum_{k=k_1}^{\infty} 2^{-(k+k_0)\alpha} \sup_{x_0 \in \mathbb{R}^d} \sum_{j=0}^k 2^{k-j+j\alpha/2} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{j-k}}(x_0)} \\ & \leq \sum_{k=0}^{\infty} 2^{-(k+k_0)\alpha} \sum_{j=0}^k 2^{j+(k-j)\alpha/2} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} \\ & = 2^{-k_0\alpha} \sum_{j=0}^{\infty} 2^{j(1-\alpha/2)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} \sum_{k=j}^{\infty} 2^{-k\alpha/2} \\ & \leq C 2^{-k_0\alpha} \sum_{j=0}^{\infty} 2^{j(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)}. \end{aligned}$$

Therefore,

$$\begin{aligned} & \sum_{j=0}^{\infty} 2^{j(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} \\ & \leq C 2^{-k_0\alpha} \sum_{j=0}^{\infty} 2^{j(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} \\ & \quad + C 2^{(k_1+k_0)(1-\alpha)} [u]_{\alpha} + C 2^{k_0(1-\alpha)} \sum_{k=k_1}^{\infty} \omega_f(2^{-k}) + C 2^{k_0(1-\alpha)} \sum_{k=k_1}^{\infty} \omega_a(2^{-k}) \\ & \quad \cdot \left(\sum_{j=0}^{\infty} 2^{j(1-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [u - p]_{\alpha; B_{2^{-j}}(x_0)} + \|u\|_{L_{\infty}} \right). \end{aligned}$$

Finally, to get (3.1) it suffices to choose k_0 and k_1 sufficiently large and apply Lemma 2.1 (iii). \square

Next we employ a localization argument as in [9].

Proof of Theorem 1.1. Since the proof of the case when $\sigma \in (0, 1)$ is almost the same as $\sigma \in (1, 2)$ and actually simpler, we only present the latter and sketch the proof of the case when $\sigma = 1$ in the end.

The case when $\sigma \in (1, 2)$. We divide the proof into three steps.

Step 1. For $k = 1, 2, \dots$, denote $B^k := B_{1-2^{-k}}$. Let $\eta_k \in C_0^{\infty}(B^{k+1})$ be a sequence of nonnegative smooth cutoff functions satisfying $\eta \equiv 1$ in B^k , $|\eta| \leq 1$ in B^{k+1} , and $\|D^i \eta_k\|_{L_{\infty}} \leq C 2^{ki}$ for each $i \geq 0$. Set $v_k := u \eta_k \in C^{\sigma+}$. A simple calculation reveals that

$$\inf_{\beta \in \mathcal{A}} (L_{\beta} v_k - h_{k\beta} + \eta_k f_{\beta}) = 0 \quad \text{in } \mathbb{R}^d,$$

where

$$h_{k\beta} = h_{k\beta}(x) = \int_{\mathbb{R}^d} \frac{\xi_k(x, y) a_{\beta}(x, y)}{|y|^{d+\sigma}} dy$$

and

$$\xi_k(x, y) = u(x + y)(\eta_k(x + y) - \eta_k(x)) - y \cdot D\eta_k(x)u(x).$$

Obviously, $\eta_k f_{\beta}$ is a Dini continuous function in \mathbb{R}^d and

$$\begin{aligned} & |\eta_k(x) f_{\beta}(x) - \eta_k(x') f_{\beta}(x')| \\ & \leq \|\eta_k\|_{L_{\infty}} \omega_f(|x - x'|) + \|f_{\beta}\|_{L_{\infty}(B_1)} \|D\eta_k\|_{L_{\infty}} |x - x'| \\ & \leq \omega_f(|x - x'|) + C 2^k \|f_{\beta}\|_{L_{\infty}(B_1)} |x - x'|, \end{aligned}$$

where C only depends on d .

Step 2. We first estimate the L_∞ norm of $h_{k\beta}$. By the fundamental theorem of calculus,

$$\xi_k(x, y) = y \cdot \int_0^1 u(x + y) D\eta_k(x + ty) - u(x) D\eta_k(x) dt.$$

For $|y| \geq 2^{-k-3}$, $|\xi_k(x, y)| \leq C2^k |y| \|u\|_{L_\infty}$. For $|y| < 2^{-k-3}$, we can further write

$$\xi_k(x, y) = y \cdot \int_0^1 (u(x + y) - u(x)) D\eta_k(x + ty) + u(x) (D\eta_k(x + ty) - D\eta_k(x)) dt,$$

where the second term on the right-hand side is bounded by $C2^{2k}|y|^2|u(x)|$. To estimate the first term, we consider two cases: when $|x| \geq 1 - 2^{-k-2}$, because $|y| < 2^{-k-3}$, $\xi_k(x, y) \equiv 0$; when $|x| < 1 - 2^{-k-2}$, we have

$$\left| y \cdot \int_0^1 (u(x + y) - u(x)) D\eta_k(x + ty) dt \right| \leq C2^k |y|^2 \|Du\|_{L_\infty(B^{k+3})}.$$

Hence for $|y| < 2^{-k-3}$,

$$|\xi_k(x, y)| \leq C|y|^2 (2^{2k}|u(x)| + 2^k \|Du\|_{L_\infty(B^{k+3})}).$$

Combining with the case when $|y| > 2^{-k-3}$, we see that

$$\|h_{k\beta}\|_{L_\infty} \leq C2^{\sigma k} (\|u\|_{L_\infty} + \|Du\|_{L_\infty(B^{k+3})}). \tag{3.18}$$

Next we estimate the modulus of continuity of $h_{k\beta}$. By the triangle inequality,

$$\begin{aligned} & |h_{k\beta}(x) - h_{k\beta}(x')| \\ & \leq \int_{\mathbb{R}^d} \frac{|(\xi_k(x, y) - \xi_k(x', y))a_\beta(x, y)|}{|y|^{d+\sigma}} + \frac{|\xi_k(x', y)(a_\beta(x, y) - a_\beta(x', y))|}{|y|^{d+\sigma}} dy \\ & := \text{I} + \text{II}. \end{aligned} \tag{3.19}$$

Similar to (3.18), by the estimates of $|\xi_k(x, y)|$ above, we have

$$\text{II} \leq C2^{\sigma k} (\|u\|_{L_\infty} + \|Du\|_{L_\infty(B^{k+3})}) \omega_a(|x - x'|), \tag{3.20}$$

where C depends on d, σ , and Λ , and is uniformly bounded as $\sigma \rightarrow 2$. For I , by the fundamental theorem of calculus,

$$\begin{aligned} \xi_k(x, y) - \xi_k(x', y) &= y \cdot \int_0^1 \left(u(x + y) D\eta_k(x + ty) - u(x) D\eta_k(x) \right. \\ & \quad \left. - u(x' + y) D\eta_k(x' + ty) + u(x') D\eta_k(x') \right) dt. \end{aligned}$$

When $|y| \geq 2^{-k-3}$, similar to the estimate of $\xi_k(x, y)$, it follows that

$$|\xi_k(x, y) - \xi_k(x', y)| \leq C|y| (2^k \omega_u(|x - x'|) + 2^{2k} \|u\|_{L_\infty} |x - x'|). \tag{3.21}$$

The case when $|y| < 2^{-k-3}$ is a bit more delicate. First, by the fundamental theorem of calculus,

$$\begin{aligned} & |\xi_k(x, y) - \xi_k(x', y)| \\ & \leq |y| \int_0^1 |(u(x + y) - u(x)) D\eta_k(x + ty) - (u(x' + y) - u(x')) D\eta_k(x' + ty)| dt \\ & \quad + |y|^2 \int_0^1 \int_0^1 |u(x) D^2\eta_k(x + tsy) - u(x') D^2\eta_k(x' + tsy)| dt ds := \text{III} + \text{IV}. \end{aligned}$$

It is easily seen that

$$IV \leq C|y|^2(2^{2k}\omega_u(|x - x'|) + 2^{3k}\|u\|_{L^\infty}|x - x'|).$$

Next we bound III by considering four cases. When $x, x' \in (B^{k+2})^c$, we have $III \equiv 0$. When $x, x' \in B^{k+2}$,

$$\begin{aligned} III &\leq |y|^2 \int_0^1 \int_0^1 |Du(x + sy)D\eta_k(x + ty) - Du(x' + sy)D\eta_k(x' + ty)| ds dt \\ &\leq C|y|^2(2^k[u]_{1+\alpha; B^{k+3}}|x - x'|^\alpha + 2^{2k}\|Du\|_{L^\infty(B^{k+3})}|x - x'|), \end{aligned}$$

where we choose $\alpha = \frac{\sigma-1}{2}$. When $x \in B^{k+2}$ and $x' \in (B^{k+2})^c$,

$$\begin{aligned} III &= |y| \int_0^1 |(u(x + y) - u(x))D\eta_k(x + ty)| dt \\ &\leq |y|^2 \int_0^1 \int_0^1 |Du(x + sy)(D\eta_k(x + ty) - D\eta_k(x' + ty))| ds dt \\ &\leq C|y|^2 2^{2k}\|Du\|_{L^\infty(B^{k+3})}|x - x'|. \end{aligned}$$

The last case is similar. In conclusion, we obtain

$$III \leq C|y|^2(2^k[u]_{1+\alpha; B^{k+3}}|x - x'|^\alpha + 2^{2k}\|Du\|_{L^\infty(B^{k+3})}|x - x'|).$$

Combining the estimates of III, IV, and (3.21), we obtain

$$\begin{aligned} I &\leq C2^{k(\sigma+1)}(\omega_u(|x - x'|) + [u]_{1+\alpha; B^{k+3}}|x - x'|^\alpha \\ &\quad + (\|Du\|_{L^\infty(B^{k+3})} + \|u\|_{L^\infty})|x - x'|). \end{aligned} \tag{3.22}$$

By combining (3.19), (3.20), and (3.22), we obtain

$$|h_{k\beta}(x) - h_{k\beta}(x')| \leq \omega_h(|x - x'|),$$

where

$$\begin{aligned} \omega_h(r) &:= C2^{\sigma k}(\|u\|_{L^\infty} + \|Du\|_{L^\infty(B^{k+3})})\omega_a(r) \\ &\quad + C2^{k(\sigma+1)}(\omega_u(r) + [u]_{1+\alpha; B^{k+3}}r^\alpha + (\|Du\|_{L^\infty(B^{k+3})} + \|u\|_{L^\infty})r) \end{aligned} \tag{3.23}$$

is a Dini function.

Step 3. We apply Proposition 3.2 to v_k to obtain

$$\begin{aligned} [v_k]_\sigma &\leq C\|v_k\|_{L^\infty} + C \sum_{j=1}^\infty (\omega_h(2^{-j}) + \omega_f(2^{-j})) + C2^k \sup_\beta \|f_\beta\|_{L^\infty(B_1)} \\ &\leq C\|v_k\|_{L^\infty} + C2^{k(\sigma+1)}([u]_{1+\alpha; B^{k+3}} + \|Du\|_{L^\infty(B^{k+3})} + \|u\|_{L^\infty}) \\ &\quad + C \sum_{j=1}^\infty (2^{k(\sigma+1)}\omega_u(2^{-j}) + \omega_f(2^{-j})) + C2^k \sup_\beta \|f_\beta\|_{L^\infty(B_1)}, \end{aligned}$$

where C depends on $d, \lambda, \Lambda, \sigma$, and ω_a , but independent of k . Since $\eta_k \equiv 1$ in B^k , it follows that

$$\begin{aligned} [u]_{\sigma; B^k} &\leq C2^{k(\sigma+1)}\|u\|_{L^\infty} + C2^{k(\sigma+1)}([u]_{1+\alpha; B^{k+3}} + \|Du\|_{L^\infty(B^{k+3})}) \\ &\quad + C_0 \sum_{j=1}^\infty (2^{k(\sigma+1)}\omega_u(2^{-j}) + \omega_f(2^{-j})) + C2^k \sup_\beta \|f_\beta\|_{L^\infty(B_1)}. \end{aligned} \tag{3.24}$$

By the interpolation inequality, for any $\varepsilon \in (0, 1)$,

$$[u]_{1+\alpha; B^{k+3}} + \|Du\|_{L_\infty(B^{k+3})} \leq \varepsilon [u]_{\sigma; B^{k+3}} + C\varepsilon^{-\frac{1+\alpha}{\sigma-(1+\alpha)}} \|u\|_{L_\infty}. \tag{3.25}$$

Recall that $\alpha = \frac{\sigma-1}{2}$ and denote

$$N := \frac{1 + \alpha}{\sigma - (1 + \alpha)} = \frac{\sigma + 1}{\sigma - 1} (> 3).$$

Combining (3.24) and (3.25) with $\varepsilon = C_0^{-1}2^{-3k-12N-1}$, we obtain

$$[u]_{\sigma; B^k} \leq C2^{3k+(3k+12N)N} \|u\|_{L_\infty} + 2^{-12N-1} [u]_{\sigma; B^{k+3}} + C2^k \sup_{\beta} \|f_{\beta}\|_{L_\infty(B_1)} + C \sum_{j=1}^{\infty} (2^{3k} \omega_u(2^{-j}) + \omega_f(2^{-j})).$$

Then we multiply 2^{-4kN} to both sides of the inequality above and get

$$2^{-4kN} [u]_{\sigma; B^k} \leq C2^{3k-kN} \|u\|_{L_\infty} + 2^{-4N(k+3)-1} [u]_{\sigma; B^{k+3}} + C2^{-4kN+k} \sup_{\beta} \|f_{\beta}\|_{L_\infty(B_1)} + C2^{-kN} \sum_{j=1}^{\infty} (\omega_u(2^{-j}) + \omega_f(2^{-j})).$$

We sum up the both sides of the inequality above and obtain

$$\sum_{k=1}^{\infty} 2^{-4kN} [u]_{\sigma; B^k} \leq C \sum_{k=1}^{\infty} 2^{3k-kN} \|u\|_{L_\infty} + \frac{1}{2} \sum_{k=4}^{\infty} 2^{-4kN} [u]_{\sigma; B^k} + C \sum_{k=1}^{\infty} 2^{-4kN+k} \sup_{\beta} \|f_{\beta}\|_{L_\infty(B_1)} + C \sum_{j=1}^{\infty} (\omega_u(2^{-j}) + \omega_f(2^{-j})),$$

which further implies that

$$\sum_{k=1}^{\infty} 2^{-4kN} [u]_{\sigma; B^k} \leq C \|u\|_{L_\infty} + C \sup_{\beta} \|f_{\beta}\|_{L_\infty(B_1)} + C \sum_{j=1}^{\infty} (\omega_u(2^{-j}) + \omega_f(2^{-j})),$$

where C depends on $d, \lambda, \Lambda, \sigma$, and ω_a . In particular, when $k = 4$, we deduce

$$[u]_{\sigma; B^4} \leq C \|u\|_{L_\infty} + C \sup_{\beta} \|f_{\beta}\|_{L_\infty(B_1)} + C \sum_{j=1}^{\infty} (\omega_u(2^{-j}) + \omega_f(2^{-j})), \tag{3.26}$$

which apparently implies (1.3).

Finally, since $\|v_1\|_1$ is bounded by the right-hand side (3.26), from (3.14), we see that

$$\sum_{j=0}^{\infty} 2^{j(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [v_1 - p]_{\alpha; B_{2^{-j}}(x_0)} \leq C.$$

This and (3.13) with u replaced by v_1 and f_{β} replaced by $\eta_1 f_{\beta} - h_1 \beta$ give

$$\begin{aligned} & \sum_{j=k_1}^{\infty} 2^{(j+k_0)(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [v_1 - p]_{\alpha; B_{2^{-j-k_0}}(x_0)} \\ & \leq C2^{-k_0\alpha} + C2^{k_0(\sigma-\alpha)} \sum_{j=k_1}^{\infty} (\omega_f(2^{-j}) + \omega_a(2^{-j}) + \omega_u(2^{-j}) + 2^{-j\alpha}). \end{aligned}$$

Here we also used Lemma 2.2 and (3.23) with $k = 1$. Therefore, for any small $\varepsilon > 0$, we can find k_0 sufficiently large then k_1 sufficiently large, depending only on $C, \sigma, \alpha, \omega_f, \omega_a, \omega_f$, and ω_u , such that

$$\sum_{j=k_1}^{\infty} 2^{(j+k_0)(\sigma-\alpha)} \sup_{x_0 \in \mathbb{R}^d} \inf_{p \in \mathcal{P}_1} [v_1 - p]_{\alpha; B_{2^{-j-k_0}}(x_0)} < \varepsilon,$$

which, together with the fact that $v_1 = u$ in $B_{1/2}$ and the proof of Lemma 2.1 (ii), indicates that

$$\sup_{x_0 \in B_{1/2}} [u]_{\sigma; B_r(x_0)} \rightarrow 0 \quad \text{as } r \rightarrow 0$$

with a decay rate depending only on $d, \lambda, \Lambda, \omega_a, \omega_f, \omega_u, \sup_{\beta \in \mathcal{A}} \|f_\beta\|_{L_\infty(B_1)}$, and σ . Hence, the proof of the case when $\sigma \in (1, 2)$ is completed.

The case when $\sigma = 1$. The proof is very similar to the case when $\sigma \in (1, 2)$ and we only provide a sketch here. We use the same notation as in the previous case

$$h_{k\beta}(x) = \int_{\mathbb{R}^d} \frac{\xi_k(x, y)a_\beta(x, y)}{|y|^{d+1}} dy,$$

where

$$\xi_k(x, y) := u(x + y)(\eta_k(x + y) - \eta_k(x)) - u(x)y \cdot D\eta_k(x)\chi_{B_1}.$$

It is easy to see that when $|y| \geq 2^{-k-3}$,

$$|\xi_k(x, y)| \leq C2^k|y|\|u\|_{L_\infty}.$$

On the other hand, when $|y| < 2^{-k-3}$,

$$\begin{aligned} |\xi_k(x, y)| &\leq |y| \int_0^1 |u(x + y)D\eta_k(x + ty) - u(x)D\eta_k(x)| dt \\ &\leq C2^k|y|w_u(|y|) + C2^{2k}|y|^2|u(x)|. \end{aligned}$$

Therefore,

$$\|h_{k\beta}\|_{L_\infty} \leq C2^k \left(\|u\|_{L_\infty} + \int_0^1 \frac{w_u(r)}{r} dr \right).$$

Next we estimate the modulus of continuity of $h_{k\beta}$ and proceed as in the case when $\sigma \in (1, 2)$. Indeed, it is easily seen that

$$\Pi \leq C2^k \left(\|u\|_{L_\infty} + \int_0^1 \frac{\omega_u(r)}{r} dr \right) \omega_a(|x - x'|).$$

To estimate I, we write

$$\begin{aligned} \xi_k(x, y) - \xi_k(x', y) &= u(x + y)(\eta_k(x + y) - \eta_k(x)) - u(x)y \cdot D\eta_k(x)\chi_{B_1} \\ &\quad - u(x' + y)(\eta_k(x' + y) - \eta_k(x')) + u(x')y \cdot D\eta_k(x')\chi_{B_1}. \end{aligned}$$

Obviously, when $|y| \geq 2^{-k-3}$

$$|\xi_k(x, y) - \xi_k(x', y)| \leq C2^{2k}|y|(\|u\|_{L_\infty}|x - x'| + \omega_u(|x - x'|)). \tag{3.27}$$

When $|y| < 2^{-k-3}$, we have $\chi_{B_1}(y) = 1$. Thus similar to the first case,

$$\begin{aligned} &|\xi_k(x, y) - \xi_k(x', y)| \\ &\leq |y| \int_0^1 |(u(x + y) - u(x))D\eta_k(x + ty) - (u(x' + y) - u(x'))D\eta_k(x' + ty)| dt \\ &\quad + |y|^2 \int_0^1 \int_0^1 |u(x)D^2\eta_k(x + tsy) - u(x')D^2\eta_k(x' + tsy)| dt ds := \text{III} + \text{IV}. \end{aligned}$$

Clearly,

$$IV \leq C2^{3k}|y|^2(\omega_u(|x - x'|) + \|u\|_{L_\infty}|x - x'|).$$

When $x, x' \in (B^{k+2})^c$, we have $III \equiv 0$. When $x, x' \in B^{k+2}$, by the triangle inequality,

$$\begin{aligned} III &\leq |y| \int_0^1 |u(x+y) - u(x) - (u(x'+y) - u(x'))| |D\eta_k(x+ty)| dt \\ &\quad + |y| \int_0^1 |u(x'+y) - u(x')| |D\eta_k(x+ty) - D\eta_k(x'+ty)| dt \\ &\leq C2^k|y|^{1+\gamma}|x - x'|^\zeta [u]_{\zeta+\gamma; B^{k+3}} + C2^{2k}|y|\omega_u(|y|)|x - x'|, \end{aligned}$$

where C depends on d , and $\zeta + \gamma < 1$. Here we used the inequality

$$|u(x+y) - u(x) - (u(x'+y) - u(x'))| \leq 2[u]_{\gamma+\zeta}|x - x'|^\zeta|y|^\gamma.$$

Set $\gamma = \zeta = 1/4$. When $x \in B^{k+2}$ and $x' \in (B^{k+2})^c$,

$$\begin{aligned} III &= |y| \int_0^1 |(u(x+y) - u(x))D\eta_k(x+ty)| dt \\ &= |y| \int_0^1 |(u(x+y) - u(x))(D\eta_k(x+ty) - D\eta_k(x'+ty))| dt \\ &\leq C2^{2k}|y|\omega_u(|y|)|x - x'|. \end{aligned}$$

The case when $x' \in B^{k+2}$ and $x \in (B^{k+2})^c$ is similar. Then with the estimates of III and IV above, we obtain that when $|y| < 2^{-k-3}$,

$$\begin{aligned} |\xi_k(x, y) - \xi_k(x', y)| &\leq C2^{3k}|y|^2(\omega_u(|x - x'|) + \|u\|_{L_\infty}|x - x'|) \\ &\quad + C2^k|y|^{5/4}|x - x'|^{1/4}[u]_{1/2; B^{k+3}} + C2^{2k}|y|\omega_u(|y|)|x - x'|, \end{aligned}$$

which, combining with (3.27) for the case when $|y| \geq 2^{-k-3}$, further implies that

$$\begin{aligned} I &\leq C2^{2k}(\omega_u(|x - x'|) + \|u\|_{L_\infty}|x - x'| \\ &\quad + [u]_{1/2; B^{k+3}}|x - x'|^{1/4} + |x - x'| \int_0^1 \frac{\omega_u(r)}{r} dr), \end{aligned}$$

where C depends on d and Λ . Hence, we obtain the estimate of the modulus of continuity of $h_{k\beta}(x)$:

$$\omega_h(r) = C2^{2k} \left(\omega_u(r) + [u]_{1/2; B^{k+3}}r^{1/4} + (\|u\|_{L_\infty} + \int_0^1 \frac{\omega_u(r)}{r} dr)(r + \omega_a(r)) \right).$$

The rest of the proof is the same as the previous case. \square

Conflict of interest statement

The authors state that there is no conflict of interest.

Acknowledgement

The authors would like to thank the referee for his/her careful review as well as many valuable comments and suggestions.

References

- [1] Jiguang Bao, Fully nonlinear elliptic equations on general domains, *Can. J. Math.* 54 (6) (2002) 1121–1141.
- [2] G. Barles, E. Chasseigne, C. Imbert, On the Dirichlet problem for second-order elliptic integro-differential equations, *Indiana Univ. Math. J.* 57 (1) (2008) 213–246.
- [3] Sergio Campanato, Equazioni paraboliche del secondo ordine e spazi $L^{2,\theta}(\Omega, \delta)$, *Ann. Mat. Pura Appl.* (4) 73 (1966) 55–102.
- [4] Charles C. Burch, The Dini condition and regularity of weak solutions of elliptic equations, *J. Differ. Equ.* 30 (3) (1978) 308–323.
- [5] Luis Caffarelli, Luis Silvestre, Regularity theory for fully nonlinear integro-differential equations, *Commun. Pure Appl. Math.* 62 (5) (2009) 597–638.
- [6] Luis Caffarelli, Luis Silvestre, The Evans–Krylov theorem for nonlocal fully nonlinear equations, *Ann. Math.* (2) 174 (2) (2011) 1163–1187.
- [7] Héctor A. Chang-Lara, Gonzalo Dávila, Regularity for solutions of non local parabolic equations, *Calc. Var. Partial Differ. Equ.* 49 (1–2) (2014) 139–172.
- [8] Héctor A. Chang-Lara, Dennis Kriventsov, Further time regularity for nonlocal, fully non-linear parabolic equations, *Commun. Pure Appl. Math.* 70 (5) (2017) 950–977.
- [9] Hongjie Dong, Hong Zhang, On Schauder estimates for a class of nonlocal fully nonlinear parabolic equations, *Calc. Var. Partial Differ. Equ.* (2017), in press, arXiv:1604.00101.
- [10] Hongjie Dong, Doyoon Kim, On L_p -estimates for a class of non-local elliptic equations, *J. Funct. Anal.* 262 (3) (2012) 1166–1199.
- [11] Hongjie Dong, Doyoon Kim, Schauder estimates for a class of non-local elliptic equations, *Discrete Contin. Dyn. Syst.* 33 (6) (2013) 2319–2347.
- [12] Frank Duzaar, Andreas Gastel, Nonlinear elliptic systems with Dini continuous coefficients, *Arch. Math. (Basel)* 78 (1) (2002) 58–73.
- [13] Yong-Cheol Kim, Ki-Ahm Lee, Regularity results for fully nonlinear parabolic integro-differential operators, *Math. Ann.* 357 (4) (2013) 1541–1576.
- [14] Takashi Komatsu, Pseudodifferential operators and Markov processes, *J. Math. Soc. Jpn.* 36 (3) (1984) 387–418.
- [15] Jay Kovats, Fully nonlinear elliptic equations and the Dini condition, *Commun. Partial Differ. Equ.* 22 (11–12) (1997) 1911–1927.
- [16] Tianling Jin, Jingang Xiong, Schauder estimates for solutions of linear parabolic integro-differential equations, *Discrete Contin. Dyn. Syst.* 35 (12) (2015) 5977–5998.
- [17] Tianling Jin, Jingang Xiong, Schauder estimates for nonlocal fully nonlinear equations, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 33 (5) (2016) 1375–1407.
- [18] Yanyan Li, On the C^1 regularity of solutions to divergence form elliptic systems with Dini-continuous coefficients, *Chin. Ann. Math., Ser. B* 38 (2) (2017) 489–496.
- [19] Gary M. Lieberman, Hölder continuity of the gradient of solutions of uniformly parabolic equations with conormal boundary conditions, *Ann. Mat. Pura Appl.* (4) 148 (1987) 77–99.
- [20] Vladimir Maz’ya, Robert McOwen, Differentiability of solutions to second-order elliptic equations via dynamical systems, *J. Differ. Equ.* 250 (2) (2011) 1137–1168.
- [21] R. Mikulevičius, H. Pragarauskas, On the Cauchy problem for certain integro-differential operators in Sobolev and Hölder spaces, *Liet. Mat. Rink.* 32 (2) (1992) 299–331.
- [22] R. Mikulevičius, H. Pragarauskas, On the Cauchy problem for integro-differential operators in Hölder classes and the uniqueness of the martingale problem, *Potential Anal.* 40 (4) (2014) 539–563.
- [23] Chenchen Mou, Interior regularity for nonlocal fully nonlinear equations with Dini continuous terms, *J. Differ. Equ.* 260 (11) (2016) 7892–7922.
- [24] L. Nirenberg, On a generalization of quasi-conformal mappings and its application to elliptic partial differential equations, in: *Contributions to the Theory of Partial Differential Equations*, in: *Annals of Mathematics Studies*, vol. 33, Princeton University Press, Princeton, N. J., 1954, pp. 95–100.
- [25] M.V. Safonov, Classical solution of second-order nonlinear elliptic equations, *Izv. Akad. Nauk SSSR, Ser. Mat.* 52 (6) (1988) 1272–1287, 1328.
- [26] Russel Schwab, Luis Silvestre, Regularity for parabolic integro-differential equations with very irregular kernels, *Anal. PDE* 9 (3) (2016) 727–772.
- [27] Joaquim Serra, $C^{\sigma+\alpha}$ regularity for concave nonlocal fully nonlinear elliptic equations with rough kernels, *Calc. Var. Partial Differ. Equ.* 54 (4) (2015) 3571–3601.
- [28] Emanuel Sperner, Schauder’s existence theorem for α -Dini continuous data, *Ark. Mat.* 19 (2) (1981) 193–216.
- [29] Xu-Jia Wang, Schauder estimates for elliptic and parabolic equations, *Chin. Ann. Math., Ser. B* 27 (6) (2006) 637–642.