



Initial-boundary value problems in a half-strip for two-dimensional Zakharov–Kuznetsov equation [☆]

Andrei V. Faminskii

Peoples' Friendship University of Russia (RUDN University), 6 Miklukho–Maklaya Street, Moscow, 117198, Russian Federation

Received 16 March 2017; received in revised form 5 October 2017; accepted 8 November 2017

Available online 15 November 2017

Abstract

Initial-boundary value problems in a half-strip with different types of boundary conditions for two-dimensional Zakharov–Kuznetsov equation are considered. Results on global existence, uniqueness and long-time decay of weak and regular solutions are established.

© 2017 Elsevier Masson SAS. All rights reserved.

MSC: primary 35Q53; secondary 35B40

Keywords: Zakharov–Kuznetsov equation; Initial-boundary value problem; Global solution; Decay

1. Introduction. Description of main results

The two dimensional Zakharov–Kuznetsov equation (ZK)

$$u_t + bu_x + u_{xxx} + u_{xyy} + uu_x = f(t, x, y) \tag{1.1}$$

(b is a real constant) is a reduction of the three-dimensional one which was derived in [35] for description of ion-acoustic waves in magnetized plasma. Now this equation is considered as a model of two-dimensional nonlinear waves in dispersive media propagating in one preassigned (x) direction with deformations in the transverse (y) direction. A rigorous derivation of the ZK model can be found, for example, in [19,21]. It is one of the variants of multi-dimensional generalizations for Korteweg–de Vries equation (KdV) $u_t + bu_x + u_{xxx} + uu_x = f(t, x)$.

The theory of solubility and well-posedness for ZK equation and its generalizations is most developed for the pure initial-value problem. For the considered two-dimensional case the corresponding results in different functional spaces can be found in [32,6,7,2,26,27,31,15,3,18,30,16,17]. For initial-boundary value problems the theory is most developed for domains of a type $I \times \mathbb{R}$, where I is an interval (bounded or unbounded) on the variable x , that is, the variable y varies in the whole line ([8,9,11,10,33,12,5]).

[☆] The publication was prepared with the support of the “RUDN University Program 5-100” and the Ministry of Education and Science of the Russian Federation (Project 1.962.2017/PCh).

E-mail address: afaminskii@sci.pfu.edu.ru.

On the other hand, from the physical point of view boundary-value problems for this equation in domains, where the variable y varies in a bounded interval, seem at least the same important. Unfortunately certain technique developed for the case $y \in \mathbb{R}$ (especially related to profound investigation of the corresponding linear equation) up to this moment is extended to the case of bounded y only partially. An initial-boundary value problem in a strip $\mathbb{R} \times (0, L)$ with periodic boundary conditions was considered in [28] for ZK equation and local well-posedness result was established in the spaces H^s for $s > 3/2$. This result was improved in [30] where $s \geq 1$, in addition, in the space H^1 appropriate conservation laws provided global well-posedness.

Another way of the study is based on the use of certain weighted spaces. Initial-boundary value problems in a strip $\mathbb{R} \times (0, L)$ with homogeneous boundary conditions of different types – Dirichlet, Neumann or periodic – were considered in [1] for ZK equation with more general nonlinearity and results on global well-posedness in classes of weak solutions with power weights when $x \rightarrow +\infty$ were established. Similar results in the case of exponential weights for ZK equation itself under homogeneous Dirichlet boundary conditions can be found in [14]. Global well-posedness results for ZK equation with certain parabolic regularization also for the initial-boundary value problem in a strip $\mathbb{R} \times (0, L)$ with homogeneous Dirichlet boundary conditions were obtained in [13,14,23,24]. Global well-posedness results for a bounded rectangle can be found in [5,34].

An initial-boundary value problem in a half-strip $\mathbb{R}_+ \times (0, L)$ with homogeneous Dirichlet boundary conditions was studied in [25,22] and global well-posedness in Sobolev spaces with exponential weights when $x \rightarrow +\infty$ was proved.

In the present paper we consider initial-boundary value problems in a domain $\Pi_T^+ = (0, T) \times \Sigma_+$, where $\Sigma_+ = \mathbb{R}_+ \times (0, L) = \{(x, y) : x > 0, 0 < y < L\}$ is a half-strip of a given width L and $T > 0$ is arbitrary, for equation (1.1) with an initial condition

$$u(0, x, y) = u_0(x, y), \quad (x, y) \in \Sigma_+, \tag{1.2}$$

a boundary condition

$$u(t, 0, y) = \mu(t, y), \quad (t, y) \in B_T = (0, T) \times (0, L), \tag{1.3}$$

and boundary conditions for $(t, x) \in \Omega_{T,+} = (0, T) \times \mathbb{R}_+$ of one of the following four types:

$$\begin{aligned} \text{whether} \quad & a) \ u(t, x, 0) = u(t, x, L) = 0, \\ & \text{or} \quad b) \ u_y(t, x, 0) = u_y(t, x, L) = 0, \\ & \text{or} \quad c) \ u(t, x, 0) = u_y(t, x, L) = 0, \\ & \text{or} \quad d) \ u \text{ is an } L\text{-periodic function with respect to } y. \end{aligned} \tag{1.4}$$

We use the notation “problem (1.1)–(1.4)” for each of these four cases.

The main results consist of theorems on global solubility and well-posedness in classes of weak and regular solutions in certain weighted (when $x \rightarrow +\infty$) Sobolev spaces. Both power and exponential weights are allowed. We do not use parabolic regularization of the equation, which is essential for the study in the whole strip in [1,14] and is used, for example, in [25,33,34]. Our approach is based on the investigation of the corresponding linearized problem (in particular, with the use of Fourier technique) and consequent application of the contraction principle and global a priori estimates. The main differences of the obtained results from the already known ones are listed below in Remarks 1.4, 1.6, 1.8.

Besides that, results on large-time decay of small solutions similar to the ones from [25,22], when $\mu \equiv 0, f \equiv 0$, are established in the cases a) and c). In our opinion, the presented proof is more transparent, than the one from [25, 22].

All global existence results are based on conservation laws, which in the case $\mu \equiv 0, f \equiv 0$ for smooth solutions are written as follows:

$$\frac{d}{dt} \iint_{\Sigma_+} u^2 dx dy + \int_0^L u_x^2|_{x=0} dy = 0, \tag{1.5}$$

$$\frac{d}{dt} \iint_{\Sigma_+} (u_x^2 + u_y^2 - \frac{1}{3}u^3) dx dy + \int_0^L (u_{xx}^2 + bu_x^2)|_{x=0} dy = 0. \tag{1.6}$$

In what follows (unless stated otherwise) j, k, l, m, n mean non-negative integers, $p \in [1, +\infty], s \in \mathbb{R}$. For any multi-index $\alpha = (\alpha_1, \alpha_2)$ let $\partial^\alpha = \partial_x^{\alpha_1} \partial_y^{\alpha_2}$,

$$|D^k \varphi| = \left(\sum_{|\alpha| \leq k} (\partial^\alpha \varphi)^2 \right)^{1/2}, \quad |D\varphi| = |D^1 \varphi|.$$

Let $L_{p,+} = L_p(\Sigma_+)$, $W_{p,+}^k = W_p^k(\Sigma_+)$, $H_+^s = H^s(\Sigma_+)$.

Introduce special function spaces taking into account boundary conditions (1.4). Let $\Sigma = \mathbb{R} \times (0, L)$, $\tilde{\mathcal{S}}(\bar{\Sigma})$ be a space of infinitely smooth on $\bar{\Sigma}$ functions $\varphi(x, y)$ such that $(1 + |x|)^n |\partial^\alpha \varphi(x, y)| \leq c(n, \alpha)$ for any n , multi-index α , $(x, y) \in \bar{\Sigma}$ and $\partial_y^{2m} \varphi|_{y=0} = \partial_y^{2m} \varphi|_{y=L} = 0$ in the case a), $\partial_y^{2m+1} \varphi|_{y=0} = \partial_y^{2m+1} \varphi|_{y=L} = 0$ in the case b), $\partial_y^{2m} \varphi|_{y=0} = \partial_y^{2m+1} \varphi|_{y=L} = 0$ in the case c), $\partial_y^m \varphi|_{y=0} = \partial_y^m \varphi|_{y=L}$ in the case d) for any m .

Let \tilde{H}^s be the closure of $\tilde{\mathcal{S}}(\bar{\Sigma})$ in the norm $H^s(\Sigma)$ and \tilde{H}_+^s be the restriction of \tilde{H}^s on Σ_+ .

It is easy to see, that $\tilde{H}_+^0 = L_{2,+}$; for $j \geq 1$ in the case a) $\tilde{H}_+^j = \{\varphi \in H_+^j : \partial_y^{2m} \varphi|_{y=0} = \partial_y^{2m} \varphi|_{y=L} = 0, 2m < j\}$, in the case b) $\tilde{H}_+^j = \{\varphi \in H_+^j : \partial_y^{2m+1} \varphi|_{y=0} = \partial_y^{2m+1} \varphi|_{y=L} = 0, 2m + 1 < j\}$, in the case d) $\tilde{H}_+^j = \{\varphi \in H_+^j : \partial_y^m \varphi|_{y=0} = \partial_y^m \varphi|_{y=L}, m < j\}$.

We also use an anisotropic Sobolev space $\tilde{H}_+^{(0,k)}$ which is defined as the restriction on Σ_+ of a space $\tilde{H}^{(0,k)}$, where the last space is the closure of $\tilde{\mathcal{S}}(\bar{\Sigma})$ in the norm $\sum_{m=0}^k \|\partial_y^m \varphi\|_{L_2(\Sigma)}$.

We say that $\rho(x)$ is an admissible weight function if ρ is an infinitely smooth positive function on $\bar{\mathbb{R}}_+$ such that $|\rho^{(j)}(x)| \leq c(j)\rho(x)$ for each natural j and all $x \geq 0$. Note that such a function satisfies an inequality $\rho(x) \leq ce^{c_0x}$ for certain positive constants c_0, c and all $x \geq 0$. It was shown in [12] that $\rho^s(x)$ for any $s \in \mathbb{R}$ is also an admissible weight function. Any exponent $e^{2\alpha x}$ as well as $(1+x)^{2\alpha}$ are admissible weight functions.

As an another important example of admissible functions, we define $\rho_0(x) \equiv 1 + \frac{2}{\pi} \arctan x$. Note that both ρ_0 and ρ'_0 are admissible weight functions.

For an admissible weight function $\rho(x)$ let $\tilde{H}_+^{k,\rho(x)}$ be a space of functions $\varphi(x, y)$ such that $\varphi \rho^{1/2}(x) \in \tilde{H}_+^k$ (similar definitions for $\tilde{H}_+^{(0,k),\rho(x)}, H_+^{k,\rho(x)}$). Let $L_{2,+}^{\rho(x)} = \tilde{H}_+^{0,\rho(x)} = \{\varphi(x, y) : \varphi \rho^{1/2}(x) \in L_{2,+}\}$. Obviously, $L_{2,+}^{\rho_0(x)} = L_{2,+}$.

We construct solutions to the considered problems in spaces $X_w^{k,\rho(x)}(\Pi_T^+)$ and $X^{k,\rho(x)}(\Pi_T^+)$ for admissible weight functions $\rho(x)$, such that $\rho'(x)$ are also admissible weight functions, consisting of functions $u(t, x, y)$ such that in the case $X_w^{k,\rho(x)}(\Pi_T^+)$

$$\partial_t^j u \in C_w([0, T]; \tilde{H}_+^{k-3j,\rho(x)}) \cap L_2(0, T; \tilde{H}_+^{k-3j+1,\rho'(x)}) \tag{1.7}$$

(the symbol C_w denotes the space of weakly continuous mappings) for $k \geq 3j \geq 0$ (let $X_w^{\rho(x)}(\Pi_T^+) = X_w^{0,\rho(x)}(\Pi_T^+)$), while in the case $X^{k,\rho(x)}(\Pi_T^+)$ the weak continuity with respect to t in (1.7) is substituted by the strong one. The gain of regularity is the reflection of the so-called local smoothing effect (see, for example, [6,7]).

Define also

$$\lambda^+(u; T) = \sup_{x_0 \geq 0} \int_0^T \int_{x_0}^{x_0+1} \int_0^L u^2 dy dx dt. \tag{1.8}$$

Let $\psi_l(y)$, $l = 1, 2, \dots$, be the orthonormal in $L_2(0, L)$ system of the eigenfunctions for the operator $(-\psi'')$ on the segment $[0, L]$ with corresponding boundary conditions $\psi(0) = \psi(L) = 0$ in the case a), $\psi'(0) = \psi'(L) = 0$ in the case b), $\psi(0) = \psi'(L) = 0$ in the case c), $\psi(0) = \psi(L), \psi'(0) = \psi'(L)$ in the case d), λ_l be the corresponding eigenvalues. Such systems are well-known and can be written in trigonometric functions.

For description of properties of the boundary data μ introduce anisotropic functional spaces. Let $B = \mathbb{R}^t \times (0, L)$. Define the functional space $\tilde{\mathcal{S}}(\bar{B})$ similarly to $\tilde{\mathcal{S}}(\bar{\Sigma})$, where the variable x is substituted by t . Let $\tilde{H}^{s/3,s}(B)$ be the closure of $\tilde{\mathcal{S}}(\bar{B})$ in the norm $H^{s/3,s}(B)$.

More exactly, for any $\mu \in \tilde{\mathcal{S}}(\bar{B}), \theta \in \mathbb{R}$ and l let

$$\hat{\mu}(\theta, l) \equiv \iint_B e^{-i\theta t} \psi_l(y) \mu(t, y) dt dy. \tag{1.9}$$

Then the norm in $\tilde{H}^{s/3,s}(B)$ is defined as $\left(\sum_{l=1}^{+\infty} \|(|\theta|^{2/3} + l^2)^{s/2} \widehat{\mu}(\theta, l)\|_{L_2(\mathbb{R}^\theta)}^2\right)^{1/2}$ and the norm in $\tilde{H}^{s/3,s}(I \times (0, L))$ for any interval $I \subset \mathbb{R}$ as the restriction norm.

The use of these norm is justified by the following fact. Let $v(t, x, y)$ be the appropriate solution to the initial value problem

$$v_t + v_{xxx} + v_{yyy} = 0, \quad v|_{t=0} = v_0.$$

Then according to [10] uniformly with respect to $x \in \mathbb{R}$

$$\|D_t^{1/3} v\|_{H_t^{s/3,s}(\mathbb{R}^2)}^2 + \|\partial_x v\|_{H_t^{s/3,s}(\mathbb{R}^2)}^2 + \|\partial_y v\|_{H_t^{s/3,s}(\mathbb{R}^2)}^2 \sim \|v_0\|_{H^s(\mathbb{R}^2)}^2 \tag{1.10}$$

(here D^α denotes the Riesz potential of the order $-\alpha$).

Introduce the notion of weak solutions to the considered problems.

Definition 1.1. Let $u_0 \in L_{2,+}$, $\mu \in L_2(B_T)$, $f \in L_1(0, T; L_{2,+})$. A function $u \in L_\infty(0, T; L_{2,+})$ is called a weak solution to problem (1.1)–(1.4) if for any function $\phi \in L_2(0, T; \tilde{H}_+^2)$, such that $\phi_t, \phi_{xxx}, \phi_{yyy} \in L_2(\Pi_T^+)$, $\phi|_{t=T} = 0$, $\phi|_{x=0} = \phi_x|_{x=0} = 0$, the following equality holds:

$$\begin{aligned} & \iiint_{\Pi_T^+} \left[u(\phi_t + b\phi_x + \phi_{xxx} + \phi_{yyy}) + \frac{1}{2}u^2\phi_x + f\phi \right] dx dy dt \\ & + \iint_{\Sigma_+} u_0\phi|_{t=0} dx dy + \iint_{B_T} \mu\phi_{xx}|_{x=0} dy dt = 0. \end{aligned} \tag{1.11}$$

Remark 1.2. Note that the integrals in (1.11) are well defined (in particular, since $\phi_x \in L_2(0, T; H_+^2) \subset L_2(0, T; L_{\infty,+})$).

Now we can formulate the main results of the paper concerning existence and uniqueness.

Theorem 1.3. Let $u_0 \in L_{2,+}^{\rho(x)}$, $f \in L_1(0, T; L_{2,+}^{\rho(x)})$ for certain $T > 0$ and an admissible weight function $\rho(x)$, such that $\rho'(x)$ is also an admissible weight function. Let $\mu \in \tilde{H}^{s/3,s}(B_T)$ for certain $s > 3/2$. Then there exists a weak solution to problem (1.1)–(1.4) $u \in X_w^{\rho(x)}(\Pi_T^+)$, moreover, $\lambda^+(|Du|; T) < +\infty$. If, in addition, $\rho^{1/2}(x) \leq c\rho'(x) \forall x \geq 0$, then this solution is unique in $X_w^{\rho(x)}(\Pi_T^+)$.

Remark 1.4. The exponential weight $\rho(x) \equiv e^{2\alpha x}$, $\alpha > 0$, satisfies both existence and uniqueness assumptions. The power weight $\rho(x) \equiv (1+x)^{2\alpha}$, $\alpha > 0$, satisfies existence assumptions and for $\alpha \geq 1$ – uniqueness assumptions. If $u_0 \in L_{2,+}$, $f \in L_1(0, T; L_{2,+})$ there exists a weak solution $u \in C_w([0, T]; L_{2,+})$, $\lambda^+(|Du|; T) < +\infty$. Note that weak solutions of the type, constructed in Theorem 1.3, are not considered in [25,22].

Theorem 1.5. Let $u_0 \in \tilde{H}_+^{1,\rho(x)}$, $f \in L_2(0, T; \tilde{H}_+^{1,\rho(x)})$ for certain $T > 0$ and an admissible weight function $\rho(x)$, such that $\rho'(x)$ is also an admissible weight function. Let $\mu \in \tilde{H}^{2/3,2}(B_T)$, $\mu(0, y) \equiv u_0(0, y)$. Then there exists a solution to problem (1.1)–(1.4) $u \in X_w^{1,\rho(x)}(\Pi_T^+)$, moreover, $\lambda^+(|D^2u|; T) < +\infty$. If, in addition, $\rho'(x) \geq 1 \forall x \geq 0$, then this solution is unique in $X_w^{1,\rho(x)}(\Pi_T^+)$.

Remark 1.6. According to (1.10) the assumptions on the boundary data μ are natural. The exponential weight $\rho(x) \equiv e^{2\alpha x}$, $\alpha > 0$, satisfies both existence and uniqueness assumptions. The power weight $\rho(x) \equiv (1+x)^{2\alpha}$, $\alpha > 0$, satisfies existence assumptions and for $\alpha \geq 1/2$ – uniqueness assumptions. If $u_0 \in \tilde{H}_+^1$, $f \in L_1(0, T; \tilde{H}_+^1)$ there exists a weak solution $u \in C_w([0, T]; \tilde{H}_+^1)$, $\lambda^+(|D^2u|; T) < +\infty$. Solutions, similar to the ones from Theorem 1.5, are constructed in [22] in the case of homogeneous Dirichlet boundary conditions and only for exponential weights (which are convenient, but, of course, restrictive). Moreover, for uniqueness results it is also assumed there, that weak solutions are limits of regular ones.

Theorem 1.7. Let $u_0 \in \tilde{H}_+^{3,\rho(x)}$, $f \in L_1(0, T; \tilde{H}_+^{(0,3),\rho(x)}) \cap L_2(0, T; \tilde{H}_+^{1,\rho(x)})$, $f_t \in L_1(0, T; L_{2,+}^{\rho(x)})$ for certain $T > 0$ and an admissible weight function $\rho(x)$, such that $\rho'(x)$ is also an admissible weight function and $\rho'(x) \geq 1 \forall x \geq 0$. Let $\mu \in \tilde{H}^{4/3,4}(B_T)$, $\mu(0, y) \equiv u_0(0, y)$. Then there exists a unique solution to problem (1.1)–(1.4) $u \in X^{3,\rho(x)}(\Pi_T^+)$.

Remark 1.8. According to (1.10) the assumptions on the boundary data μ are natural. Both the exponential weight $\rho(x) \equiv e^{2\alpha x}$, $\alpha > 0$ and the power weight $\rho(x) \equiv (1+x)^{2\alpha}$, $\alpha \geq 1/2$, satisfy the hypothesis of the theorem. In [25] for construction of regular solutions only exponential weights are used and only homogeneous Dirichlet boundary conditions are considered. Moreover, what seems the most important, for the constructed regular solutions existence of $u_{,yy}$, lying in weighted L_2 -spaces uniformly with respect to t , is not obtained there in comparison with Theorem 1.7.

As an application of the obtained results we establish large-time decay of solutions. Here we use only exponential weights.

Theorem 1.9. Let $L_0 = +\infty$ if $b \leq 0$, and if $b > 0$ there exists $L_0 > 0$, such that in both cases for any $L \in (0, L_0)$ there exist $\alpha_0 > 0$, $\epsilon_0 > 0$ and $\beta > 0$, such that if $u_0 \in L_{2,+}^{e^{2\alpha x}}$ for $\alpha \in (0, \alpha_0]$, $\|u_0\|_{L_{2,+}} \leq \epsilon_0$, $f \equiv 0$, $\mu \equiv 0$, in the cases a) and c) in (1.4) the corresponding unique weak solution $u(t, x, y)$ to problem (1.1)–(1.4) from the space $X_w^{e^{2\alpha x}}(\Pi_T^+)$ $\forall T > 0$ satisfies an inequality

$$\|e^{\alpha x} u(t, \cdot, \cdot)\|_{L_{2,+}}^2 \leq e^{-\alpha\beta t} \|e^{\alpha x} u_0\|_{L_{2,+}}^2 \quad \forall t \geq 0. \tag{1.12}$$

If, in addition, $u_0 \in \tilde{H}_+^{1,e^{2\alpha x}}$, $u_0(0, y) \equiv 0$, then for certain constant c , depending on b, α, β , $\|u_0\|_{\tilde{H}_+^{1,e^{2\alpha x}}}$,

$$\|e^{\alpha x} u(t, \cdot, \cdot)\|_{H_+^1}^2 \leq ce^{-\alpha\beta t} \quad \forall t \geq 0. \tag{1.13}$$

Further, let $\eta(x)$ denotes a cut-off function, namely, η is an infinitely smooth non-decreasing function on \mathbb{R} such that $\eta(x) = 0$ when $x \leq 0$, $\eta(x) = 1$ when $x \geq 1$, $\eta(x) + \eta(1-x) \equiv 1$.

Let $\tilde{\mathfrak{S}}(\bar{B}_+)$ be the restriction of $\tilde{\mathfrak{S}}(\bar{B})$ on $\bar{B}_+ = \mathbb{R}_+^t \times [0, L]$.

We drop limits of integration in integrals over the whole half-strip Σ_+ .

The following interpolating inequality generalizing the one from [20] for weighted Sobolev spaces is crucial for the study.

Lemma 1.10. Let $\rho_1(x), \rho_2(x)$ be two admissible weight functions, such that $\rho_1(x) \leq c_0\rho_2(x) \forall x \geq 0$ for some constant $c_0 > 0$. Then for any $q \in [2, +\infty)$ there exists a constant $c > 0$ such that for every function $\varphi(x, y)$, satisfying $|D\varphi|\rho_1^{1/2}(x) \in L_{2,+}$, $\varphi\rho_2^{1/2}(x) \in L_{2,+}$, the following inequality holds:

$$\|\varphi\rho_1^s(x)\rho_2^{1/2-s}(x)\|_{L_{q,+}} \leq c\| |D\varphi|\rho_1^{1/2}(x) \|_{L_{2,+}}^{2s} \|\varphi\rho_2^{1/2}(x)\|_{L_{2,+}}^{1-2s} + c\|\varphi\rho_2^{1/2}(x)\|_{L_{2,+}}, \tag{1.14}$$

where $s = \frac{1}{2} - \frac{1}{q}$. If $\varphi|_{y=0} = 0$ or $\varphi|_{y=L} = 0$, then the constant c in (1.14) is uniform with respect to L .

Proof. For the whole strip $\Sigma = \mathbb{R} \times (0, L)$ this inequality was proved in [14]. For Σ_+ the proof is the same. \square

Lemma 1.11. For an admissible weight function $\rho(x)$ introduce a functional space $H_+^{(-1,0),\rho(x)} = \{\varphi = \varphi_0 + \varphi_{1x} : \varphi_0, \varphi_1 \in L_{2,+}^{\rho(x)}\}$ endowed with the natural norm. Then for $j = 1$ and $j = 2$

$$\|\partial_x^j \varphi\|_{L_{2,+}^{\rho(x)}} \leq c(\rho) (\|\varphi_{xx}\|_{H_+^{(-1,0),\rho(x)}} + \|\varphi\|_{H_+^{j-1,\rho(x)}}). \tag{1.15}$$

Proof. For functions, defined on the real line, the corresponding analogue of such an inequality follows from the Fourier transform approach and an inequality

$$|\xi|^j \leq \frac{|\xi|^3}{1+|\xi|} + (1+|\xi|)^{j-1}.$$

Then the extension for the functions, defined on the half-line, and for weighted spaces is performed via simple but tedious calculations. The presence of the additional variable is unessential. \square

We also use the following obvious interpolating inequality:

$$\int_0^L \varphi^2|_{x=0} dy \leq c \left(\iint \varphi_x^2 \rho' dx dy \right)^{1/2} \left(\iint \varphi^2 \rho dx dy \right)^{1/2} + c \iint \varphi^2 \rho dx dy \tag{1.16}$$

(the constant c depends on the properties of an admissible weight function ρ).

For the decay results, we need Steklov’s inequalities in the following form:

$$\int_0^L \psi^2(y) dy \leq \frac{\sigma L^2}{\pi^2} \int_0^L (\psi'(y))^2 dy, \tag{1.17}$$

where $\sigma = 1$ if $\psi \in H_0^1(0, L)$, $\sigma = 4$ if $\psi \in H^1(0, L)$, $\psi|_{y=0} = 0$.

The paper is organized as follows. Auxiliary linear problems are considered in Section 2. Section 3 is devoted to the existence results for the original problems. Results on uniqueness and continuous dependence are proved in Section 4. Decay of solutions is studied in Section 5.

2. Auxiliary linear problems

Consider an initial-boundary value in Π_T^+ for a linear equation

$$u_t + bu_x + u_{xxx} + u_{xyy} = f(t, x, y) \tag{2.1}$$

with initial and boundary conditions (1.2)–(1.4). Weak solutions to this problem are understood similarly to Definition 1.1, moreover, due to the absence of nonlinearity one can take solutions from more wide space $L_2(\Pi_T^+)$.

Lemma 2.1. *A generalized solution to problem (2.1), (1.2)–(1.4) is unique in the space $L_2(\Pi_T^+)$.*

Proof. The proof of uniqueness is based on existence of smooth solutions to the corresponding adjoint problem. Consider a problem in Π_T^+ for an equation

$$u_t - bu_x - u_{xxx} - u_{xyy} = f(t, x, y) \in C_0^\infty(\Pi_T^+) \tag{2.2}$$

with zero initial data (1.2), boundary data (1.4) and boundary data on B_T

$$u|_{x=0} = u_x|_{x=0} = 0. \tag{2.3}$$

Let $\{\varphi_j(x) : j = 1, 2, \dots\}$ be a set of linearly independent functions complete in the space $\{\varphi \in H^3(\mathbb{R}_+) : \varphi(0) = 0\}$.

We use the Galerkin method and seek an approximate solution in the form $u_k(t, x, y) = \sum_{j,l=1}^k c_{kjl}(t)\varphi_j(x)\psi_l(y)$ via conditions for $i, m = 1, \dots, k, t \in [0, T]$

$$\iint (u_{kt}\varphi_i(x)\psi_m(y) + u_k(b\varphi_i'\psi_m + \varphi_i'''\psi_m + \varphi_i'\psi_m'')) dx dy - \iint f\varphi_i\psi_m dx dy = 0, \tag{2.4}$$

$c_{kjl}(0) = 0$. Multiplying (2.4) by $2c_{kim}(t)$ and summing with respect to i, m , we find that

$$\|u_k\|_{L_\infty(0,T;L_{2,+})} \leq \|f\|_{L_1(0,T;L_{2,+})}. \tag{2.5}$$

Next, putting in (2.4) $t = 0$, multiplying by $c'_{kim}(0)$ and summing with respect to i, m , we derive that $u_{kt}|_{t=0} = 0$. Then differentiating (2.4) with respect to t , multiplying by $2c'_{kim}(t)$ and summing with respect to i, m , we find similarly to (2.5) that

$$\|u_{kt}\|_{L_\infty(0,T;L_{2,+})} \leq \|f_t\|_{L_1(0,T;L_{2,+})}. \tag{2.6}$$

Finally, since $\psi_m^{(2n)}(y) = (-\lambda_m)^n \psi_m(y)$ it follows from (2.4) that similarly to (2.5) for any n

$$\|\partial_y^n u_k\|_{L_\infty(0,T;L_{2,+})} \leq \|\partial_y^n f\|_{L_1(0,T;L_{2,+})}. \tag{2.7}$$

Estimates (2.5), (2.6), (2.7) provide existence of a weak solution $u(t, x, y)$ to the considered problem such that $u, u_t, \partial_y^n u \in L_\infty(0, T; L_{2,+}) \forall n$ in the following sense: for any function $\phi \in L_2(0, T; \tilde{H}_+^2)$, such that $\phi_t, \phi_{xxx}, \phi_{xyy} \in L_2(\Pi_T^+)$, $\phi|_{t=T} = 0, \phi|_{x=0} = 0$, the following equality holds:

$$\iiint_{\Pi_T^+} [u(\phi_t - b\phi_x - \phi_{xxx} - \phi_{xyy}) + f\phi] dx dy dt = 0. \tag{2.8}$$

Note, that the traces of the function u satisfy conditions (1.2) for $u_0 \equiv 0$ and (1.4). Moreover, it follows from (2.8) that $\partial_y^n u_{xxx} \in L_\infty(0, T; H_+^{(-1,0),1}) \forall n$, therefore, inequality (1.15) for $j = 1$ yields that $\partial_y^n u_x \in L_\infty(0, T; L_{2,+}) \forall n$ and one more application of (2.8) yields that $\partial_y^n u_{xxx} \in L_\infty(0, T; L_{2,+}) \forall n$, the function u satisfies equation (2.2) a.e. in Π_T^+ and its traces satisfy (2.3).

The end of the proof of the lemma is standard. \square

With the use of Galerkin method we prove one result on solubility of the considered problem in an infinitely smooth case.

Lemma 2.2. *Let $u_0 \equiv 0, \mu \in C_0^\infty(B_+), f \equiv 0$. Then there exists a solution $u(t, x, y)$ to problem (2.1), (1.2)–(1.4), such that $\partial_t^j \partial^\alpha u \in C_b(\overline{\mathbb{R}}_+^l; L_{2,+})$ for any j and multi-index α (here and further index ‘b’ means a bounded map).*

Proof. Let $v(t, x, y) \equiv u(t, x, y) - \mu(t, y)\eta(1 - x)$, then the original problem is equivalent to the problem for the function v of (2.1), (1.2)–(1.4) type with homogeneous initial-boundary conditions and $f \equiv -\mu_t\eta(1 - x) - b\mu\eta'(1 - x) - \mu\eta'''(1 - x) - \mu_{yy}\eta'(1 - x)$.

Seek an approximate solution in the form $v_k(t, x, y) = \sum_{j,l=1}^k c_{kjl}(t)\varphi_j(x)\psi_l(y)$ (the functions φ_j are the same as in the proof of Lemma 2.1) via conditions for $i, m = 1, \dots, k, t \in [0, T]$

$$\iint (v_{kt} + bv_{kx} + v_{kxxx} + v_{kxyy})\varphi_i(x)\psi_m(y) dx dy - \iint f\varphi_i\psi_m dx dy = 0, \quad c_{kjl}(0) = 0. \tag{2.9}$$

Multiplying (2.9) by $2c_{kim}(t)$ and summing with respect to i, m , we find that

$$\frac{d}{dt} \iint v_k^2 dx dy + \int_0^L v_{kx}^2|_{x=0} dy = 2 \iint f v_k dx dy. \tag{2.10}$$

Note that $\|v_k(t, \cdot, \cdot)\|_{L_{2,+}}$ doesn't increase if $t \geq T$ for certain T . Then the consequent argument from the proof of Lemma 2.1 can be applied here ((2.8) must be substituted by the corresponding analogue of (1.11)). Thus, first existence of a solution v such that $\partial_t^j \partial_y^n v \in C_b(\overline{\mathbb{R}}_+^l; L_{2,+})$ for all j and n is obtained; then with the use of induction with respect to m one can find that $\partial_t^j \partial_y^n \partial_x^{3m} v \in C_b(\overline{\mathbb{R}}_+^l; L_{2,+})$. \square

Before the continuation of the study of the problems in the half-strip consider the corresponding problems in the whole strip.

For $u_0 \in \tilde{\mathcal{S}}(\overline{\Sigma})$ define similarly to (1.9) for $\xi \in \mathbb{R}$ and l

$$\widehat{u}_0(\xi, l) \equiv \iint_{\Sigma} e^{-i\xi x} \psi_l(y) u_0(x, y) dx dy, \tag{2.11}$$

$$S(t, x, y; u_0) \equiv \sum_{l=1}^{+\infty} \frac{1}{2\pi} \int_{\mathbb{R}} e^{it(\xi^3 - b\xi + \lambda_l \xi)} e^{i\xi x} \widehat{u}_0(\xi, l) d\xi \psi_l(y). \tag{2.12}$$

It is easy to see that for all $s \in \mathbb{R}$ the function $S(t, x, y; u_0) \in C_b(\mathbb{R}^t; \tilde{H}^s)$ and for any $t \in \mathbb{R}$

$$\|S(t, \cdot, \cdot; u_0)\|_{\tilde{H}^s} = \|u_0\|_{\tilde{H}^s}. \tag{2.13}$$

This property gives an opportunity to extend the notion of the function $S(t, x, y; u_0)$ to any function $u_0 \in \tilde{H}^s$ for any $s \in \mathbb{R}$ via closure in the space $C_b(\mathbb{R}^t; \tilde{H}^s)$, then, of course, equality (2.13) holds.

Let $\varphi_l(\xi) \equiv \xi^3 - b\xi + \lambda_l \xi$. This function increases monotonically if $\lambda_l \geq b$ on the whole real line and for $\xi < -\sqrt{(b - \lambda_l)/3}$ and $\xi > \sqrt{(b - \lambda_l)/3}$ if $\lambda_l < b$. Let $\kappa_l(\theta) \equiv \varphi_l^{-1}(\theta)$, which is defined for all θ if $\lambda_l \geq b$ and for $|\theta| \geq 2((b - \lambda_l)/3)^{3/2}$ if $\lambda_l < b$ (then $|\kappa_l(\theta)| \geq 2\sqrt{(b - \lambda_l)/3}$).

Lemma 2.3. *If $u_0 \in \tilde{H}^s$ for certain $s \in \mathbb{R}$, then $S(t, x, y; u_0) \in C_b(\mathbb{R}^x; \tilde{H}^{(s+1)/3, s+1}((-T, T) \times (0, L)))$ and for any $x \in \mathbb{R}$*

$$\|S(\cdot, x, \cdot; u_0)\|_{\tilde{H}^{(s+1)/3, s+1}((-T, T) \times (0, L))} \leq c(T)\|u_0\|_{\tilde{H}^s}. \tag{2.14}$$

Proof. Without loss of generality assume that $u_0 \in \tilde{\mathcal{S}}(\bar{\Sigma})$. There exists l_0 such that for $l > l_0$ and all ξ and there exists $\xi_0 \geq 1$ such that for $|\xi| \geq \xi_0$ and all l

$$\varphi'_l(\xi) = 3\xi^2 - b + \lambda_l \geq c(\xi^2 + l^2). \tag{2.15}$$

Divide u_0 into two parts:

$$u_{00}(x, y) \equiv \sum_{l=1}^{l_0} \mathcal{F}_x^{-1}[\widehat{u}_0(\xi, l)\eta(\xi_0 + 1 - |\xi|)](x)\psi_l(y), \quad u_{01}(x, y) \equiv u_0(x, y) - u_{00}(x, y). \tag{2.16}$$

After the change of variables in the corresponding analog of the integral in (2.12) $\theta = \varphi_l(\xi)$ (without loss of generality one can assume also that $\varphi_l^{-1}(\theta) = \kappa_l(\theta)$) we derive that for the obviously defined function $\chi(\xi, l)$ (in particular, $\chi(\xi, l) = 0$ for $l \leq l_0, |\xi| \leq \xi_0$)

$$S(t, x, y; u_{01}) = \sum_{l=1}^{+\infty} \mathcal{F}_t^{-1}\left[e^{i\kappa_l(\theta)x}\widehat{u}_0(\kappa_l(\theta), l)\kappa'_l(\theta)\chi(\kappa_l(\theta), l)\right](t)\psi_l(y) \tag{2.17}$$

and uniformly with respect to x

$$\begin{aligned} &\|S(\cdot, x, \cdot; u_{01})\|_{H^{(s+1)/3, s+1}(B)} \\ &= \left(\sum_{l=1}^{+\infty} \left(\|\varphi_l(\xi)\|^{2/3} + l^2\right)^{(s+1)/2} \widehat{u}_0(\xi, l) (\varphi'_l(\xi))^{-1/2} \chi(\xi, l) \right)_{L_2(\mathbb{R}^\xi)}^2 \Big)^{1/2} \leq c\|u_0\|_{\tilde{H}^s}^2. \end{aligned} \tag{2.18}$$

Finally note that

$$S(t, x, y; u_{00}) = \sum_{l=1}^{l_0} \mathcal{F}_x^{-1}\left[e^{it\varphi_l(\xi)}\widehat{u}_0(\xi, l)\eta(\xi_0 + 1 - |\xi|)\right](x)\psi_l(y)$$

and one can easily show that for any j and n uniformly with respect to $t \in \mathbb{R}$

$$\|\partial_t^j \partial_y^n S(t, \cdot, \cdot; u_{00})\|_{L_2} \leq c(s, j, n)\|u_0\|_{\tilde{H}^s}. \quad \square$$

Next, we introduce the notation

$$K(t, x, y; f) \equiv \int_0^t S(t - \tau, x, y; f(\tau, \cdot, \cdot)) d\tau. \tag{2.19}$$

Obviously, if $f \in L_1(0, T; \tilde{H}^s)$ for certain $s \in \mathbb{R}$, then $K(t, x, y; f) \in C([0, T]; \tilde{H}^s)$ and

$$\|K(\cdot, \cdot, \cdot; f)\|_{C([0, T]; \tilde{H}^s)} \leq \|f\|_{L_1(0, T; \tilde{H}^s)}. \tag{2.20}$$

Lemma 2.4. *If $s \in [-1, 2]$, $f \in L_2(0, T; \tilde{H}^s)$, then the function $K(t, x, y; f) \in C_b(\mathbb{R}^x; \tilde{H}^{(s+1)/3, s+1}(B_T))$ and for any $x \in \mathbb{R}$, $t_0 \in (0, T]$*

$$\|K(\cdot, x, \cdot; f)\|_{\tilde{H}^{(s+1)/3, s+1}(B_{t_0})} \leq c(T)t_0^{1/3-s/6} \|f\|_{L_2(0, t_0; \tilde{H}^s)}. \tag{2.21}$$

Proof. For $s = -1$ it follows from (2.14) that

$$\|K(\cdot, x, \cdot; f)\|_{L_2(B_{t_0})} \leq c(T)\|f\|_{L_1(0, t_0; \tilde{H}^{-1})} \leq c(T)t_0^{1/2} \|f\|_{L_2(0, t_0; \tilde{H}^{-1})}.$$

Next,

$$\begin{aligned} \partial_y^3 K(t, x, y; f) &= K(t, x, y; \partial_y^3 f), \\ \partial_t K(t, x, y; f) &= f(t, x, y) + K(t, x, y; (b\partial_x + \partial_x^3 + \partial_x \partial_y^2)f), \end{aligned}$$

and again applying (2.14) for $s = -1$ we derive that

$$\|K(\cdot, x, \cdot; f)\|_{\tilde{H}^{1,3}(B_{t_0})} \leq c(T)\|f\|_{L_2(0, t_0; \tilde{H}^2)}.$$

For intermediate values of s the result follows by interpolation. \square

If $u_0 \in L_2(\Sigma)$, $f \in L_1(0, T; L_2(\Sigma))$, then a function

$$u(t, x, y) \equiv S(t, x, y; u_0) + K(t, x, y; f) \tag{2.22}$$

is a weak solution to an initial-boundary value problem in a strip Σ to problem (2.1), (1.2) for $(x, y) \in \Sigma$, (1.4) for $(t, x) \in (0, T) \times \mathbb{R}$ (see, for example, [1]).

In what follows, we need some properties of solutions to an algebraic equation

$$z^3 - (\lambda_l - b)z + p = 0, \quad p = \varepsilon + i\theta \in \mathbb{C}. \tag{2.23}$$

For $\varepsilon > 0$ we denote by $z_0(p, l)$ the unique root of this equation, such that $\Re z_0 < 0$.

Lemma 2.5. *There exists*

$$\lim_{\varepsilon \rightarrow +0} z_0(\varepsilon + i\theta, l) = r_0(\theta, l) = p(\theta, l) + iq(\theta, l), \tag{2.24}$$

where $r_0(\cdot, l) \in C(\mathbb{R})$, $r_0(-\theta, l) = \overline{r_0(\theta, l)}$, $p(\theta, l), q(\theta, l) \in \mathbb{R}$ and

$$|r_0(\theta, l)| \leq c(|\theta|^{1/3} + \lambda_l^{1/2} + |b|^{1/2}), \quad c = \text{const} > 0. \tag{2.25}$$

If $\lambda_l \geq b$, then

$$p(\theta, l) \leq -c_0(|\theta|^{1/3} + (\lambda_l - b)^{1/2}), \quad c_0 = \text{const} > 0. \tag{2.26}$$

If $\lambda_l < b$, then for $|\theta| \geq 2((b - \lambda_l)/3)^{3/2}$

$$p(\theta, l) \leq -c_0(|\kappa_l(\theta)| - 2\sqrt{(b - \lambda_l)/3}), \quad c_0 = \text{const} > 0, \tag{2.27}$$

while for $|\theta| < 2((b - \lambda_l)/3)^{3/2}$

$$p(\theta, l) = 0, \quad |q(\theta, l)| \leq \sqrt{(b - \lambda_l)/3}, \quad q(\theta, l) = \varphi_l^{-1}(\theta). \tag{2.28}$$

Proof. This lemma evidently follows from the Cardano formula. In particular, if $\lambda_l < b$ then for $|\theta| > 2((b - \lambda_l)/3)^{3/2}$

$$p(\theta, l) = -\frac{\sqrt{3}}{2} \left[\sqrt[3]{\frac{\theta}{2} + \sqrt{\frac{\theta^2}{4} - \frac{(b - \lambda_l)^3}{27}}} - \sqrt[3]{\frac{\theta}{2} - \sqrt{\frac{\theta^2}{4} - \frac{(b - \lambda_l)^3}{27}}} \right],$$

therefore, it is easily verified that

$$p'(\theta, l) \operatorname{sign} \theta \leq -\frac{1}{2^{4/3}\sqrt{3}}|\theta|^{-2/3}.$$

Since obviously $\varphi_l'(\xi) \geq \varphi_l^{2/3}(\xi)$ for $|\xi| \geq 2\sqrt{(b - \lambda_l)/3}$ and so $\theta^{-2/3} \geq \kappa_l'(\theta)$ inequality (2.27) follows. \square

Now we introduce a special solution of equation (2.1) for $f \equiv 0$ of “boundary potential” type.

Definition 2.6. Let $\mu \in \tilde{\mathcal{S}}(\bar{B})$. Define for $x \geq 0$

$$J(t, x, y; \mu) \equiv \sum_{l=1}^{+\infty} \mathcal{F}_t^{-1} \left[e^{r_0(\theta, l)x} \widehat{\mu}(\theta, l) \right] (t) \psi_l(y), \tag{2.29}$$

where $\widehat{\mu}(\theta, l)$ is given by formula (1.9).

Remark 2.7. Since $\widehat{J}(\theta, x, l; \mu) = e^{r_0(\theta, l)x} \widehat{\mu}(\theta, l)$ and $\Re r_0(\theta, l) \leq 0$, then $J(t, x, l; \mu) \in C_b(\bar{\mathbb{R}}_+^x; \tilde{H}^{s/3, s}(B))$ for any $s \in \mathbb{R}$ and

$$\|J(\cdot, \cdot, \cdot; \mu)\|_{C_b(\bar{\mathbb{R}}_+^x; \tilde{H}^{s/3, s}(B))} \leq \|\mu\|_{\tilde{H}^{s/3, s}}. \tag{2.30}$$

Therefore, the notion of the function $J(t, x, y; \mu)$ can be extended in the space $C_b(\bar{\mathbb{R}}_+^x; \tilde{H}^{s/3, s}(B))$ for any function $\mu \in \tilde{H}^{s/3, s}(B)$ for certain $s \in \mathbb{R}$ with conservation of inequality (2.30). It is obvious, that $J(t, 0, y; \mu) \equiv \mu(t, y)$.

Note, that in the most important for us case $s \geq 0$ the function J can be equivalently defined simply by formula (2.29).

Lemma 2.8. If $\mu \in \tilde{H}^{s/3, s}(B)$ for certain $s \geq 0$, then for any $n \leq s$ there exists $\partial_x^n J(t, x, y; \mu) \in C_b(\bar{\mathbb{R}}_+^x; \tilde{H}^{(s-n)/3, s-n}(B))$ and uniformly with respect to $x \geq 0$

$$\|\partial_x^n J(\cdot, x, \cdot; \mu)\|_{\tilde{H}^{(s-n)/3, s-n}(B)} \leq c(s) \|\mu\|_{\tilde{H}^{s/3, s}(B)}. \tag{2.31}$$

Proof. The proof is similar to the proof of inequality (2.30) with the use of (2.25). \square

Lemma 2.9. If $\mu \in \tilde{H}^{(s+1)/3, s+1}(B)$ for certain $s \geq 0$, then for any $j \leq s/3$ there exists $\partial_t^j J(t, x, y; \mu) \in C_b(\mathbb{R}^t; \tilde{H}_+^{s-3j})$ and uniformly with respect to $t \in \mathbb{R}$

$$\|\partial_t^j J(t, \cdot, \cdot; \mu)\|_{\tilde{H}_+^{s-3j}} \leq c(s) \|\mu\|_{\tilde{H}^{(s+1)/3, s+1}(B)}. \tag{2.32}$$

Proof. Without loss of generality one can assume that $\mu \in \tilde{\mathcal{S}}(\bar{B})$. Let s be integer. Then for $3j + n + m = s$

$$\partial_t^j \partial_x^n \partial_y^m J(t, x, y; \mu) = \sum_{l=1}^{+\infty} \frac{1}{2\pi} \int_{\mathbb{R}} (i\theta)^j r_0^n(\theta, l) e^{it\theta} e^{r_0(\theta, l)x} \widehat{\mu}(\theta, l) d\theta \psi_l^{(m)}(y). \tag{2.33}$$

Divide the expression in the right side of (2.33) into two parts. Let l_0 be such that $\lambda_l < b$ for $l \leq l_0$ and let I_1 be the part of the right side of (2.33) with the sum from $l = 1$ to $l = l_0$ and the integral over $|\theta| < 2((b - \lambda_l)/3)^{3/2}$ (it is absent if $\lambda_l \geq b \forall l$) and let I_2 be the rest part.

First consider I_1 . According to (2.28) $r_0(\theta, l) = iq(\theta, l)$ and changing variables $\xi = q(\theta, l)$ we derive that

$$I_1 = \sum_{l=1}^{l_0} \frac{1}{2\pi} \int_{|\xi| < \sqrt{(b-\lambda_l)/3}} e^{it\varphi_l(\xi)} e^{i\xi x} (i\varphi_l(\xi))^j (i\xi)^n \widehat{\mu}(\varphi_l(\xi), l) \varphi_l'(\xi) d\xi \psi_l^{(m)}(y).$$

Thus, similarly to (2.13) uniformly with respect to $t \in \mathbb{R}$

$$\|I_1\|_{L_2(\Sigma)} \leq c \|\mu\|_{L_2(B)}.$$

For I_2 we use the following fundamental inequality from [4]: if certain continuous function $\gamma(\kappa)$ satisfies an inequality $\Re \gamma(\kappa) \leq -\varepsilon|\kappa|$ for some $\varepsilon > 0$ and all $\kappa \in \mathbb{R}$, then

$$\left\| \int_{\mathbb{R}} e^{\gamma(\kappa)x} f(\kappa) d\kappa \right\|_{L_2(\mathbb{R}_+^x)} \leq c(\varepsilon) \|f\|_{L_2(\mathbb{R})}.$$

Changing variables $\theta = \varphi_l(\kappa)$ we derive with the use of (2.25)–(2.27) that uniformly with respect to $t \in \mathbb{R}$ for $\gamma(\kappa) = -c_0(|\kappa| - 2\sqrt{(b - \lambda_l)/3})$, $\chi(\theta, l) = 1$ for $|\theta| > 2((b - \lambda_l)/3)^{3/2}$ (then $|\kappa| > 2\sqrt{(b - \lambda_l)/3}$) and $\chi(\theta, l) = 0$ for other values of θ if $l \leq l_0$, $\gamma(\kappa) = -c_0|\kappa|$, $\chi(\theta, l) \equiv 1$ if $l > l_0$

$$\begin{aligned} & \|I_2\|_{L_{2,+}} \\ &= \frac{1}{2\pi} \left(\sum_{l=1}^{+\infty} \left\| \int_{\mathbb{R}} \theta^j r_0^n(\theta, l) e^{it\theta} e^{r_0(\theta, l)x} \widehat{\mu}(\theta, l) \chi(\theta, l) d\theta \right\|_{L_2(\mathbb{R}_+^x)}^2 \left\| \psi_l^{(m)} \right\|_{L_2(0, L)}^2 \right)^{1/2} \\ &\leq c \left(\sum_{l=1}^{+\infty} \left\| \int_{\mathbb{R}} (\kappa^2 + l^2)^{(3j+n)/2} e^{\gamma(\kappa)x} |\widehat{\mu}(\varphi_l(\kappa), l)| \chi(\varphi_l(\kappa), l) \varphi_l'(\kappa) d\kappa \right\|_{L_2(\mathbb{R}_+^x)}^2 l^{2m} \right)^{1/2} \\ &\leq c_1 \left(\sum_{l=1}^{+\infty} \left\| (|\theta|^{2/3} + l^2)^{(3j+n+m+1)/2} \widehat{\mu}(\theta, l) \right\|_{L_2(\mathbb{R}^\theta)}^2 \right)^{1/2} = c_1 \|\mu\|_{\widetilde{H}^{(s+1)/3, s+1}(B)}. \end{aligned}$$

Finally, use interpolation for non-integer values of s . \square

Lemma 2.10. Let $\mu \in \widetilde{H}^{s/3, s}(B)$ for certain $s \geq -1/2$. Then for any $T > 0$ and $j \leq s/3 + 1/6$

$$\|\partial_t^j J(\cdot, \cdot, \cdot; \mu)\|_{L_2(0, T; \widetilde{H}_+^{s-3j+1/2})} \leq c(T, s) \|\mu\|_{\widetilde{H}^{s/3, s}(B)}. \tag{2.34}$$

Proof. Without loss of generality one can assume that $\mu \in \widetilde{S}(\overline{B})$. By virtue of (2.26), (2.27) there exists l_0 such that for $l > l_0$ and all θ and there exists $\theta_0 \geq 1$ such that for $|\theta| \geq \theta_0$ and all l

$$p(\theta, l) \leq -c_0(|\theta|^{1/3} + l) \tag{2.35}$$

Similarly to (2.16) divide μ into two parts:

$$\mu_0(t, y) \equiv \sum_{l=1}^{l_0} \mathcal{F}_t^{-1} [\widehat{\mu}(\theta, l) \eta(\theta_0 + 1 - |\theta|)](t) \psi_l(y), \quad \mu_1(t, y) \equiv \mu(t, y) - \mu_0(t, y). \tag{2.36}$$

Let $s + 1/2$ be an integer, then if $3j + n + m = s + 1/2$ we derive from equality (2.33) and inequalities (2.25), (2.35) that for the obviously defined function $\chi(\theta, l)$ (in particular, $\chi(\theta, l) = 0$ for $l \leq l_0$, $|\theta| \leq \theta_0$)

$$\begin{aligned} & \|\partial_t^j \partial_x^n \partial_y^m J(\cdot, \cdot, \cdot; \mu_1)\|_{L_2(\mathbb{R}^t \times \Sigma_+)} \\ &= \left(\sum_{l=1}^{+\infty} \left\| \theta^j r_0^n(\theta, l) \left(\int_{\mathbb{R}_+} e^{-2p(\theta, l)x} dx \right)^{1/2} \widehat{\mu}(\theta, l) \chi(\theta, l) \right\|_{L_2(\mathbb{R}^\theta)}^2 \left\| \psi_l^{(m)} \right\|_{L_2(0, L)}^2 \right)^{1/2} \\ &\leq c \left(\sum_{l=1}^{+\infty} \left\| (|\theta|^j (|\theta|^{2/3} + l^2)^{(n-1)/2} |\widehat{\mu}(\theta, l)| d\theta \right\|_{L_2(\mathbb{R}_+^x)}^2 l^{2m} \right)^{1/2} \leq c_1 \|\mu\|_{\widetilde{H}^{s/3, s}(B)}. \end{aligned} \tag{2.37}$$

For μ_0 inequality (2.32) yields that for any $s_0 \geq 0$

$$\begin{aligned} \|\partial_t^j J(\cdot, \cdot, \cdot; \mu_0)\|_{L_2(0, T; \widetilde{H}_+^{s_0})} &\leq T^{1/2} \|\partial_t^j J(\cdot, \cdot, \cdot; \mu_0)\|_{C_b(\mathbb{R}^t; \widetilde{H}_+^{s_0})} \\ &\leq cT^{1/2} \|\mu_0\|_{\widetilde{H}^{(s_0+1)/3+j, s_0+1+3j}(B)} \leq c_1 T^{1/2} \|\mu\|_{\widetilde{H}^{s/3, s}(B)}. \end{aligned} \tag{2.38}$$

To finish the proof we again use interpolation. \square

Corollary 2.11. Let $\mu \in \widetilde{H}^{s/3, s}(B)$ for certain $s > 1/2$. Then for any $T > 0$ and j, k , such that $3j + k < s - 1/2$,

$$\|\partial_t^j J(\cdot, \cdot, \cdot; \mu)\|_{L_2(0, T; W_{\infty,+}^k)} \leq c(T, s) \|\mu\|_{\widetilde{H}^{s/3, s}(B)}. \tag{2.39}$$

Proof. Estimate (2.39) obviously follows from (2.34) and the well-known embedding $H_+^{k+1+\varepsilon} \subset W_{\infty,+}^k$. \square

Lemma 2.12. Let $\mu \in \tilde{H}^{s/3,s}(B)$ for certain $s \in \mathbb{R}$. Then the function $J(t, x, y; \mu)$ is infinitely differentiable for $x > 0$, $(t, y) \in \bar{B}$ and satisfies equation (2.1), where $f \equiv 0$. Moreover, for any $T > 0$, $x_0 > 0$ and j, n

$$\sup_{x \geq x_0} \|\partial_x^n J(\cdot, x, \cdot; \mu)\|_{\tilde{H}^{j,3j}(B_T)} \leq c(T, x_0, n, j, s) \|\mu\|_{\tilde{H}^{s/3,s}(B)}. \tag{2.40}$$

Proof. Without loss of generality one can assume that $\mu \in \tilde{S}(\bar{B})$. By virtue of (2.29)

$$\partial_x^n J(t, x, y; \mu) \equiv \sum_{l=1}^{+\infty} \mathcal{F}_t^{-1} \left[r_0^n(\theta, l) e^{r_0(\theta, l)x} \widehat{\mu}(\theta, l) \right] (t) \psi_l(y).$$

Again divide μ into two parts as in (2.36). Then by virtue of (2.25) and (2.35)

$$\begin{aligned} & \|\partial_x^n J(\cdot, x, \cdot; \mu_1)\|_{\tilde{H}^{j,3j}(B)}^2 \\ & \leq c \sum_{l=1}^{+\infty} \left(|\theta|^{2/3} + l^2 \right)^{(n+3j)/2} e^{-c_0 x_0 (|\theta|^{1/3} + l)} \widehat{\mu}(\theta, l) \Big|_{L_2(\mathbb{R}^\theta)}^2 \leq c(x_0) \|\mu\|_{\tilde{H}^{s/3,s}(B)}^2. \end{aligned}$$

For μ_0 apply estimate (2.32) similarly to (2.38).

Equality (2.1) for $u \equiv J$, $f \equiv 0$ follows from (2.23), (2.24). \square

Lemma 2.13. Let $\mu \in L_2(B)$ and $\mu(t, y) = 0$ for $t < 0$, then the function $J(t, x, y; \mu)$ for any $T > 0$ is a weak solution (from $L_2(\Pi_T^+)$) to problem (2.1) (for $f \equiv 0$), (1.2) (for $u_0 \equiv 0$), (1.3), (1.4).

Proof. First let $\mu \in C_0^\infty(B_+)$. Consider the smooth solution $u(t, x, y)$ to the considered problem constructed in Lemma 2.2. For any $p = \varepsilon + i\theta$, where $\varepsilon > 0$, define the Laplace–Fourier transform-coefficients

$$\tilde{u}(p, x, l) \equiv \int_{\mathbb{R}_+} \int_0^L e^{-pt} \psi_l(y) u(t, x, y) dy dt$$

(similar definition for $\tilde{\mu}(p, l)$). The function $\tilde{u}(p, x, l)$ solves a problem

$$p\tilde{u}(p, x, l) + b\tilde{u}_x(p, x, l) + \tilde{u}_{xx}(p, x, l) - \lambda_l \tilde{u}(p, x, l) = 0, \tilde{u}(p, 0, l) = \tilde{\mu}(p, l),$$

whence, since $\tilde{u}(p, x, l) \rightarrow 0$ as $x \rightarrow +\infty$, it follows, that

$$\tilde{u}(p, x, l) = \tilde{\mu}(p, l) e^{z_0(p, l)x},$$

where $z_0(p, l)$ is defined in (2.23). Using the formula of inversion of the Laplace transform we find, that

$$u(t, x, y) = \sum_{l=1}^{+\infty} e^{\varepsilon t} \mathcal{F}_t^{-1} \left[\tilde{\mu}(\varepsilon + i\theta, l) e^{z_0(\varepsilon + i\theta, l)x} \right] (t) \psi_l(y).$$

Passing to the limit as $\varepsilon \rightarrow +0$, we derive that $u(t, x, y) \equiv J(t, x, y; \mu)$.

In the general case approximate the function μ by smooth ones, pass to the limit on the basis of estimate (2.34) for $s = 0$ (note, that this estimate is superfluous, the corresponding more weak estimate in $L_2(\Pi_T^+)$ is sufficient) and use the uniqueness result. \square

Corollary 2.14. Let $u_0 \in L_{2,+}$, $\mu \in \tilde{H}^{1/3,1}(B_T)$, $f \in L_2(0, T; L_{2,+})$ for certain $T > 0$. Then there exists a unique solution to problem (2.1), (1.2)–(1.4), such that $u \in C([0, T]; L_{2,+})$, $u, u_x \in C_b(\bar{\mathbb{R}}_+; L_2(B_T))$, given by a formula

$$u(t, x, y) = S(t, x, y; u_0) + K(t, x, y; f) + J(t, x, y; \mu - S(0, \cdot, \cdot; u_0) - K(0, \cdot, \cdot; f)), \tag{2.41}$$

where for the construction of the functions S and K the functions u_0 and f are extended somehow in the same classes for $x < 0$ and for the construction of the function J the function $\mu - S(0, \cdot, \cdot; u_0) - K(0, \cdot, \cdot; f)$ is extended by zero for $t < 0$ and somehow in the same class as μ for $t > T$.

Proof. This assertion directly succeeds from (2.13), (2.14), (2.20), (2.21), (2.22), (2.31), (2.32) and Lemma 2.13. \square

We introduce certain additional function space. Let $\tilde{\mathcal{S}}_{exp}(\overline{\Sigma}_+)$ denotes a space of infinitely smooth functions $\varphi(x, y)$ in $\overline{\Sigma}_+$, such that $e^{nx}|\partial^\alpha \varphi(x, y)| \leq c(n, \alpha)$ for any n , multi-index α , $(x, y) \in \overline{\Sigma}_+$ and $\partial_y^{2m} \varphi|_{y=0} = \partial_y^{2m} \varphi|_{y=L} = 0$ in the case a), $\partial_y^{2m+1} \varphi|_{y=0} = \partial_y^{2m+1} \varphi|_{y=L} = 0$ in the case b), $\partial_y^{2m} \varphi|_{y=0} = \partial_y^{2m+1} \varphi|_{y=L} = 0$ in the case c), $\partial_y^m \varphi|_{y=0} = \partial_y^m \varphi|_{y=L} = 0$ in the case d) for any m .

Let $\tilde{\Phi}_0(x, y) \equiv u_0(x, y)$ and for $j \geq 1$

$$\tilde{\Phi}_j(x, y) \equiv \partial_t^{j-1} f(0, x, y) - (b\partial_x + \partial_x^3 + \partial_x \partial_y^2) \tilde{\Phi}_{j-1}(x, y).$$

Lemma 2.15. Let $u_0 \in \tilde{\mathcal{S}}(\overline{\Sigma}) \cap \tilde{\mathcal{S}}_{exp}(\overline{\Sigma}_+)$, $f \in C^\infty([0, 2T]; \tilde{\mathcal{S}}(\overline{\Sigma}) \cap \tilde{\mathcal{S}}_{exp}(\overline{\Sigma}_+))$, $\mu \in \tilde{\mathcal{S}}(\overline{B}_+)$ and $\partial_t^j \mu(0, y) \equiv \tilde{\Phi}_j(0, y)$ for any j . Then there exists a unique solution to problem (2.1), (1.2)–(1.4) $u \in C^\infty([0, T]; \tilde{\mathcal{S}}_{exp}(\overline{\Sigma}_+))$.

Proof. Let $w(t, x, y) = S(t, x, y; u_0) + K(t, x, y; f)$ be the solution to initial-boundary value problem (2.1), (1.2), (1.4) from the space $u \in C^\infty([0, 2T]; \tilde{\mathcal{S}}(\overline{\Sigma}) \cap \tilde{\mathcal{S}}_{exp}(\overline{\Sigma}_+))$ (see, for example, [1]).

Let $\tilde{\mu}(t, y) \equiv (\mu(t, y) - w(t, 0, y))\eta(2 - t/T)$. Extend this function to the whole strip B by zero for $t < 0$. Such an extension can be performed by virtue of the compatibility conditions on the line $t = 0, x = 0$. Then $\tilde{\mu} \in \tilde{\mathcal{S}}(\overline{B})$.

Then formula (2.41) provides the solution to the considered problem such that $\partial_t^j u \in C([0, T], \tilde{H}_+^k)$ for all j and k (see Lemma 2.9).

Finally, let $v(t, x, y) \equiv u(t, x, y)\eta(x - 1)$. The function v solves an initial value problem in a strip Σ of (2.1), (1.2) type, where f, u_0 are substituted by corresponding functions F, v_0 from the same classes and [1] provides that $v \in C^\infty([0, T]; \tilde{\mathcal{S}}_{exp}(\overline{\Sigma}_+))$. \square

Remark 2.16. In further lemmas of this section we first consider smooth solutions constructed in Lemma 2.15 and then pass to the limit on the basis of obtained estimates.

Lemma 2.17. Let $\mu \equiv 0$, $\rho(x)$ be an admissible weight function, such that $\rho'(x)$ is also an admissible weight function, $u_0 \in L_{2,+}^{\rho(x)}$, $f \equiv f_0 + f_{1x} + f_2$, where $f_0 \in L_1(0, T; L_{2,+}^{\rho(x)})$, $f_1 \in L_2(0, T; L_{2,+}^{\rho^2(x)/\rho'(x)})$, $f_2 \rho^{3/4}(x)/(\rho'(x))^{1/4} \in L_{4/3}(\Pi_T^+)$. Then there exist a (unique) weak solution to problem (2.1), (1.2)–(1.4) from the space $X^{\rho(x)}(\Pi_T^+)$ and a function $v \in L_2(B_T)$, such that for any function $\phi \in L_2(0, T; \tilde{H}_+^2)$, $\phi_t, \phi_{xxx}, \phi_{xyy} \in L_2(\Pi_T^+)$, $\phi|_{t=T} = 0$, $\phi|_{x=0} = 0$, the following equality holds:

$$\begin{aligned} & \iint_{\Pi_T^+} [u(\phi_t + b\phi_x + \phi_{xxx} + \phi_{xyy}) + (f_0 + f_2)\phi - f_1\phi_x] dx dy dt \\ & + \iint u_0 \phi|_{t=0} dx dy - \iint_{B_T} v \phi_x|_{x=0} dy dt = 0. \end{aligned} \tag{2.42}$$

Moreover, for $t \in (0, T]$

$$\begin{aligned} \|u\|_{X^{\rho(x)}(\Pi_T^+)} + \|v\|_{L_2(B_t)} & \leq c(T) \left(\|u_0\|_{L_{2,+}^{\rho(x)}} + \|f_0\|_{L_1(0,t;L_{2,+}^{\rho(x)})} \right. \\ & \left. + \|f_1\|_{L_2(0,t;L_{2,+}^{\rho^2(x)/\rho'(x)})} + \|f_2 \rho^{3/4}(x)/(\rho'(x))^{1/4}\|_{L_{4/3}(\Pi_t^+)} \right), \end{aligned} \tag{2.43}$$

$$\begin{aligned} & \iint u^2(t, x, y) \rho(x) dx dy + \int_0^t \iint (3u_x^2 + u_y^2 - bu^2) \rho'(x) dx dy d\tau + \rho(0) \iint_{B_t} v^2 dy d\tau \\ & = \iint u_0^2 \rho(x) dx dy + \int_0^t \iint u^2 \rho'''(x) dx dy d\tau + 2 \int_0^t \iint (f_0 + f_2) u \rho(x) dx dy d\tau \end{aligned}$$

$$-2 \int_0^t \iint f_1(u\rho(x))_x dx dy d\tau. \tag{2.44}$$

If $f_1 = f_2 \equiv 0$, then in equality (2.44) one can put $\rho \equiv 1$.

Proof. Multiplying (2.1) by $2u(t, x, y)\rho(x)$ and integrating over Σ_+ we find that

$$\begin{aligned} & \frac{d}{dt} \iint u^2 \rho dx dy + \rho(0) \int_0^L u_x^2|_{x=0} dy + \iint (3u_x + u_y^2 - bu^2)\rho' dx dy \\ &= \iint u^2 \rho''' dx dy + 2 \iint (f_0 + f_2)u\rho dx dy - 2 \iint f_1(u\rho)_x dx dy. \end{aligned} \tag{2.45}$$

Note that

$$\begin{aligned} \left| \iint f_2 u \rho dx dy \right| &\leq \|u(\rho'\rho)^{1/4}\|_{L_{4,+}} \|f_2 \rho^{3/4}(\rho')^{-1/4}\|_{L_{4/3,+}} \\ &\leq c \left[\|Du(\rho')^{1/2}\|_{L_{2,+}}^{1/2} \|u\rho^{1/2}\|_{L_{2,+}}^{1/2} + \|u\rho^{1/2}\|_{L_{2,+}} \right] \|f_2 \rho^{3/4}(\rho')^{-1/4}\|_{L_{4/3,+}} \\ &\leq \varepsilon \iint |Du|^2 \rho' dx dy + c(\varepsilon) \|f_2 \rho^{3/4}(\rho')^{-1/4}\|_{L_{4/3,+}}^{4/3} \left(\iint u^2 \rho dx dy \right)^{1/3} \\ &\quad + c \|f_2 \rho^{3/4}(\rho')^{-1/4}\|_{L_{4/3,+}} \left(\iint u^2 \rho dx dy \right)^{1/2}, \end{aligned} \tag{2.46}$$

$$\begin{aligned} \left| \iint f_1(u\rho)_x dx dy \right| &\leq c \|f_1 \rho(\rho')^{-1/2}\|_{L_{2,+}} \|(|u_x| + |u|)(\rho')^{1/2}\|_{L_{2,+}} \\ &\leq \varepsilon \iint (u_x^2 + u^2)\rho' dx dy + c(\varepsilon) \|f_1\|_{L_{2,+}^{\rho^2(x)/\rho'(x)}}^2, \end{aligned} \tag{2.47}$$

where $\varepsilon > 0$ can be chosen arbitrarily small. Equality (2.45) and inequalities (2.46), (2.47) imply that that for smooth solutions

$$\|u\|_{X^{\rho(x)}(\Pi_T^+)} + \|u_x|_{x=0}\|_{L_2(B_T)} \leq c.$$

The end of the proof is standard. \square

Remark 2.18. The method of construction of weak solution in Lemma 2.17 via closure ensures that $u|_{x=0} = 0$ in the trace sense (this fact can be also easily derived from equality (2.42), since $u_x \in L_2((0, T) \times (0, x_0) \times (0, L))$ for certain $x_0 > 0$). Moreover, if it is known, in addition, that $u_x \in C_w([0, x_0]; L_2(B_T))$ for certain $x_0 > 0$, then equality (2.42) yields that $u_x|_{x=0} = v$ (for example, one can put $\varphi \equiv x\eta(1 - x/h)\omega(t, y)$ for $h > 0$ and any $\omega \in C_0^\infty(B_T)$ and then tend h to zero).

Lemma 2.19. Let $\mu \equiv 0$, $\rho(x)$ be an admissible weight function, such that $\rho'(x)$ is also an admissible weight function, $u_0 \in \tilde{H}_+^{1,\rho(x)}$, $u_0|_{x=0} \equiv 0$, $f \equiv f_0 + f_1$, where $f_0 \in L_2(0, T; \tilde{H}_+^{1,\rho(x)})$, $f_1 \in L_2(0, T; L_{2,+}^{\rho^2(x)/\rho'(x)})$. Then there exist a (unique) weak solution to problem (2.1), (1.2)–(1.4) from the space $X^{1,\rho(x)}(\Pi_T^+)$ and a function $v_1 \in L_2(B_T)$ such that for any $t \in (0, T]$

$$\begin{aligned} & \|u\|_{X^{1,\rho(x)}(\Pi_T^+)} + \|v_1\|_{L_2(B_T)} \\ & \leq c(T) \left(\|u_0\|_{\tilde{H}_+^{1,\rho(x)}} + \|f_0\|_{L_2(0,t;\tilde{H}_+^{1,\rho(x)})} + \|f_1\|_{L_2(0,t;L_{2,+}^{\rho^2(x)/\rho'(x)})} \right), \end{aligned} \tag{2.48}$$

$$\iint (u_x^2 + u_y^2)\rho dx dy + \int_0^t \iint (3u_{xx}^2 + 4u_{xy}^2 + u_{yy}^2 - bu_x^2 - bu_y^2)\rho' dx dy d\tau$$

$$\begin{aligned}
 & + \iint_{B_t} (v_1^2 \rho + 2v_1 u_x \rho' + bu_x^2 \rho - u_x^2 \rho'')|_{x=0} dyd\tau = \iint (u_{0x}^2 + u_{0y}^2) \rho dx dy \\
 & + \int_0^t \iint (u_x^2 + u_y^2) \rho''' dx dy d\tau + 2 \int_0^t \iint (f_{0x} u_x + f_{0y} u_y) \rho dx dy d\tau \\
 & + 2 \iint_{B_t} (f_0 u_x \rho)|_{x=0} dyd\tau - 2 \int_0^t \iint f_1 [(u_x \rho)_x + u_{yy} \rho] dx dy d\tau.
 \end{aligned} \tag{2.49}$$

Proof. In the smooth case multiplying (2.1) by $-2((u_x(t, x, y)\rho(x))_x + u_{yy}(t, x, y)\rho(x))$ and integrating over Σ_+ , one obtains an equality:

$$\begin{aligned}
 & \frac{d}{dt} \iint (u_x^2 + u_y^2) \rho dx dy + \iint (3u_{xx}^2 + 4u_{xy}^2 + u_{yy}^2 - bu_x^2 - bu_y^2) \rho' dx dy \\
 & - \iint (u_x^2 + u_y^2) \rho''' dx dy + \int_0^L (u_{xx}^2 \rho + 2u_{xx} u_x \rho' - u_x^2 \rho'' + bu_x^2 \rho)|_{x=0} dy \\
 & = 2 \iint (f_{0x} u_x + f_{0y} u_y) \rho dx dy + 2 \int_0^L (f_0 u_x \rho)|_{x=0} dy - 2 \iint f_1 [(u_x \rho)_x + u_{yy} \rho] dx dy.
 \end{aligned} \tag{2.50}$$

Since the trace of u_x on the plane $x = 0$ is already estimated in (2.43) (here $v = u_x|_{x=0}$, see Remark 2.18) equality (2.50) provides that

$$\|u\|_{X^{1,\rho(x)}(\Pi_T^+)} + \|u_{xx}|_{x=0}\|_{L_2(B_T)} \leq c. \quad \square$$

Lemma 2.20. Let the hypothesis of Lemma 2.19 be satisfied in the case $\rho(x) \equiv e^{2\alpha x}$ for certain $\alpha > 0$. Consider the weak solution $u \in X^{1,\rho(x)}(\Pi_T^+)$ to problem (2.1), (1.2)–(1.4). Then for any $t \in (0, T]$ the following equality holds:

$$\begin{aligned}
 & -\frac{1}{3} \iint u^3(t, x, y) \rho(x) dx dy + \frac{b}{3} \int_0^t \iint u^3 \rho' dx dy d\tau \\
 & + 2 \int_0^t \iint (u_{xx} + u_{yy}) u u_x \rho dx dy d\tau + \int_0^t \iint (u_{xx} + u_{yy}) u^2 \rho' dx dy d\tau \\
 & = -\frac{1}{3} \iint u_0^3 \rho dx dy - \int_0^t \iint f u^2 \rho dx dy d\tau.
 \end{aligned} \tag{2.51}$$

Proof. In the smooth case multiplying (2.1) by $-u^2(t, x, y)\rho(x)$ and integrating one instantly obtains equality (2.51).

In the general case this equality is established via closure. Note that by virtue of (1.14) (for $q = 4$, $\rho_1 = \rho_2 \equiv \rho$) if $u \in X^{1,\rho(x)}(\Pi_T^+)$ then

$$u \in C([0, T]; L_{4,+}^{\rho(x)}), \quad |Du| \in L_2(0, T; L_{4,+}^{\rho(x)})$$

and this passage to the limit is easily justified. \square

Lemma 2.21. Let $\mu \equiv 0$, $\rho(x)$ be an admissible weight function, such that $\rho'(x)$ is also an admissible weight function, $u_0 \in \tilde{H}_+^{2,\rho(x)}$, $u_0|_{x=0} \equiv 0$ and $u_{0xx}, u_{0yy} \in L_{2,+}^{\rho(x)}$, $f \in C([0, T]; L_{2,+}^{\rho(x)})$, moreover, $f \equiv f_0 + f_{1x}$, where $f_0, f_{0t} \in L_1(0, T; L_{2,+}^{\rho(x)})$, $f_1, f_{1t} \in L_2(0, T; L_{2,+}^{\rho^2(x)/\rho'(x)})$. Then for the (unique) weak solution to problem (2.1), (1.2)–(1.4)

from the space $X^{\rho(x)}(\Pi_T^+)$ there exists $u_t \in X^{\rho(x)}(\Pi_T^+)$, which is the weak solution to problem of (2.1), (1.2)–(1.4) type, where f is substituted by $f_t, u_0 - by (f|_{t=0} - bu_{0x} - u_{0x_{xx}} - u_{0x_{yy}}), \mu \equiv 0$.

Proof. The proof for the function $v \equiv u_t$ is similar to Lemma 2.17. \square

Lemma 2.22. Let the hypotheses of Lemma 2.19 and Lemma 2.21 be satisfied and, in addition, $f \in L_1(0, T; \tilde{H}_+^{(0,2),\rho(x)})$. Then there exists a (unique) solution to problem (2.1), (1.2)–(1.4) from the space $X^{2,\rho(x)}(\Pi_T^+)$ and for any $t \in [0, T]$

$$\begin{aligned} \|u\|_{X^{2,\rho(x)}(\Pi_T^+)}^2 &\leq c(T) \left(\|u_{0yy}\|_{L_{2,+}^{\rho(x)}}^2 + \|f\|_{C([0,t];L_{2,+}^{\rho(x)})}^2 + \|u_t\|_{C([0,t];L_{2,+}^{\rho(x)})}^2 \right. \\ &\quad + \|u\|_{C([0,t];\tilde{H}_+^{1,\rho(x)})}^2 + \sup_{\tau \in (0,t]} \left| \int_0^\tau \iint f_{yy} u_{yy} \rho \, dx dy ds \right| + \int_0^t \iint (u_{xx}^2 + u_{yy}^2) \rho \, dx dy d\tau \\ &\quad \left. + \int_0^t \iint (f^2 + u_t^2 + b^2 u_x^2 + bu_{yy}^2) \rho' \, dx dy d\tau \right). \end{aligned} \tag{2.52}$$

Proof. For smooth solutions differentiating equality (2.1) twice with respect to y , multiplying the obtained equality by $2u_{yy}(t, x, y)\rho(x)$ and integrating over Σ_+ we derive similarly to (2.45) that

$$\begin{aligned} \frac{d}{dt} \iint u_{yy}^2 \rho \, dx dy + \rho(0) \int_0^L u_{xyy}^2|_{x=0} \, dy + \iint (3u_{xyy}^2 + u_{yyy}^2 - bu_{yy}^2) \rho' \, dx dy \\ = \iint u_{yy}^2 \rho''' \, dx dy + 2 \iint f_{yy} u_{yy} \rho \, dx dy, \end{aligned} \tag{2.53}$$

whence obviously follows that

$$\|u_{yy}\|_{X^{\rho(x)}(\Pi_T^+)} \leq c. \tag{2.54}$$

Hence, for the weak solution also $u_{yy} \in X^{\rho(x)}(\Pi_T^+)$. Lemmas 2.19 and 2.21 provide that $u \in X^{1,\rho(x)}(\Pi_T^+), u_t \in X^{\rho(x)}(\Pi_T^+)$. Write equality (2.1) in the form

$$u_{xxx} = f - u_t - bu_x - u_{xyy}. \tag{2.55}$$

Then, inequality (1.15) for $j = 2$ and (2.55) yield that

$$\begin{aligned} \|u_{xx}\|_{L_{2,+}^{\rho(x)}} &\leq c(\rho) (\|u_{xxx}\|_{H^{(-1,0),\rho(x)}} + \|u\|_{\tilde{H}_+^{1,\rho(x)}}) \\ &\leq c(\rho, b) (\|f\|_{L_{2,+}^{\rho(x)}} + \|u_t\|_{L_{2,+}^{\rho(x)}} + \|u_{yy}\|_{L_{2,+}^{\rho(x)}} + \|u\|_{\tilde{H}_+^{1,\rho(x)}}). \end{aligned} \tag{2.56}$$

Since

$$\iint u_{xy}^2 \rho \, dx dy = \iint u_{xx} u_{yy} \rho \, dx dy + \iint u_{yy} u_x \rho' \, dx dy,$$

estimates (2.54) and (2.56) yield that $u \in C([0, T]; \tilde{H}_+^{2,\rho(x)})$ and

$$\|u(t, \cdot, \cdot)\|_{\tilde{H}_+^{2,\rho(x)}} \leq c(\|f\|_{L_{2,+}^{\rho(x)}} + \|u_t\|_{L_{2,+}^{\rho(x)}} + \|u_{yy}\|_{L_{2,+}^{\rho(x)}} + \|u\|_{\tilde{H}_+^{1,\rho(x)}}). \tag{2.57}$$

Next,

$$\begin{aligned} \iint u_{xxy}^2 \rho' \, dx dy &= \iint u_{xxx} u_{xyy} \rho' \, dx dy + \iint u_{xyy} u_{xx} \rho'' \, dx dy \\ &\quad + \int_0^L (u_{xyy} u_{xx} \rho')|_{x=0} \, dy \end{aligned}$$

and inequality (1.16) provides that

$$\iint u_{xx}^2 \rho' dx dy \leq \iint (u_{xxx}^2 + u_{xyy}^2) \rho' dx dy + \int_0^L u_{xyy}^2|_{x=0} dy + c \iint u_{xx}^2 \rho dx dy. \tag{2.58}$$

From equality (2.55) we derive that

$$\iint u_{xxx}^2 \rho' dx dy \leq c \iint (f^2 + u_t^2 + b^2 u_x^2 + u_{xyy}^2) \rho' dx dy \tag{2.59}$$

and combining (2.53), (2.57)–(2.59) finish the proof. \square

Lemma 2.23. *Let $\mu \equiv 0$, $\rho(x)$ be an admissible weight function, such that $\rho'(x)$ is also an admissible weight function, $u_0 \in \tilde{H}_+^{3,\rho(x)}$, $u_0(0, y) \equiv 0$, $f \in C([0, T]; L_{2,+}^{\rho(x)})$ and $f \equiv f_0 + f_{1x}$, where $f_0 \in L_2(0, T; \tilde{H}_+^{1,\rho(x)}) \cap L_1(0, T; \tilde{H}_+^{(0,3),\rho(x)})$, $f_{0t} \in L_1(0, T; L_{2,+}^{\rho(x)})$, $f_1 \in L_2(0, T; L_{2,+}^{\rho^2(x)/\rho'(x)})$, $f_{1x} \in L_2(0, T; \tilde{H}_+^{(0,2),\rho^2(x)/\rho'(x)})$, $f_{1t} \in L_2(0, T; L_{2,+}^{\rho^2(x)/\rho'(x)})$. Then there exists a (unique) solution to problem (2.1), (1.2)–(1.4) from the space $X^{3,\rho(x)}(\Pi_T^+)$ and for any $t \in (0, T]$*

$$\begin{aligned} \|u\|_{X^{3,\rho(x)}(\Pi_T^+)} \leq c(T) & (\|u_0\|_{\tilde{H}_+^{3,\rho(x)}} + \|f\|_{C([0,t];L_{2,+}^{\rho(x)})} + \|f_0\|_{L_2(0,t;\tilde{H}_+^{1,\rho(x)})} \\ & + \|f_{0yy}\|_{L_1(0,t;L_{2,+}^{\rho(x)})} + \|f_{0t}\|_{L_1(0,t;L_{2,+}^{\rho(x)})} + \|f_1\|_{L_2(0,t;L_{2,+}^{\rho^2(x)/\rho'(x)})} \\ & + \|f_{1x}\|_{L_2(0,t;L_{2,+}^{\rho^2(x)/\rho'(x)})} + \|f_{1xyy}\|_{L_2(0,t;L_{2,+}^{\rho^2(x)/\rho'(x)})} + \|f_{1t}\|_{L_2(0,t;L_{2,+}^{\rho^2(x)/\rho'(x)})}). \end{aligned} \tag{2.60}$$

Proof. First of all note that hypotheses of Lemmas 2.17 (for $f_2 \equiv 0$), Lemmas 2.19, 2.21 and 2.22 are satisfied. Therefore, taking into account also Remark 2.18 we derive for smooth solutions that

$$\|u\|_{X^{2,\rho(x)}(\Pi_T^+)} + \|u_x|_{x=0}\|_{L_2(B_T)} + \|u_t\|_{X^{\rho(x)}(\Pi_T^+)} + \|u_{tx}|_{x=0}\|_{L_2(B_T)} \leq c. \tag{2.61}$$

Next, differentiating equality (2.1) twice with respect to y , multiplying the obtained equality by $-2u_{yyyy}(t, x, y)\rho(x)$ and integrating over Σ_+ we derive similarly to (2.53) that

$$\begin{aligned} & \frac{d}{dt} \iint u_{yyy}^2 \rho dx dy + \rho(0) \int_0^L u_{xyyy}^2|_{x=0} dy + \iint (3u_{xyyy} + u_{yyyy} - bu_{yyy}^2) \rho' dx dy \\ & = \iint u_{yyy}^2 \rho''' dx dy + 2 \iint f_{0yyy} u_{yyy} \rho dx dy - 2 \iint f_{1yy} u_{yyyy} \rho dx dy. \end{aligned} \tag{2.62}$$

Here

$$\left| 2 \iint f_{1yy} u_{yyyy} \rho dx dy \right| \leq \varepsilon \iint u_{yyyy}^2 \rho' dx dy + \frac{1}{\varepsilon} \iint f_{1yy}^2 \frac{\rho^2}{\rho'} dx dy,$$

where $\varepsilon > 0$ can be chosen arbitrarily small, and equality (2.62) yields that

$$\|u_{yyy}\|_{X^{\rho(x)}(\Pi_T^+)} + \|u_{xyyy}|_{x=0}\|_{L_2(B_T)} \leq c. \tag{2.63}$$

Again apply equality (2.55). Then it follows from (2.63) that we have the suitable estimate on u_y in the space $L_2(0, T; \tilde{H}_+^{3,\rho'(x)})$ and, in particular, on u_{xxyy} in $L_2(0, T; L_{2,+}^{\rho'(x)})$ (for similar argument see (2.58)). One more application of (2.55) yields the estimate on u_{xxxx} in $L_2(0, T; L_{2,+}^{\rho'(x)})$. As a result

$$\|u\|_{L_2(0,T;\tilde{H}_+^{4,\rho'(x)})} \leq c. \tag{2.64}$$

Consider the extensions of the functions u and f for $y \in (L, 2L]$ and $y \in [-L, 0)$ in the case a) by the even reflections through $y = L$ and $y = 0$, in the case b) – by the odd ones, in the case c) – by the corresponding combination of these methods, in the case d) – by the periodic extension. Then the functions u and f remain smooth in the more

wide domain $[0, T] \times \overline{\mathbb{R}}_+ \times [-L, 2L]$, and equality (2.1) also remains valid. Let $\eta_L(y) \equiv \eta(1 + y/L)\eta(2 - y/L)$, $\tilde{u}(t, x, y) \equiv u(t, x, y)\eta_L(y)$, $\tilde{f}(t, x, y) \equiv f(t, x, y)\eta_L(y)$. Now we apply the inequality (see, e.g. [29]) for the domain $\mathbb{R}_+^2 = \{(x, y); x > 0\}$

$$\|g\|_{H^2(\mathbb{R}_+^2)} \leq c(\|\Delta g\|_{L_2(\mathbb{R}_+^2)} + \|g|_{x=0}\|_{H^{3/2}(\mathbb{R})} + \|g\|_{H^1(\mathbb{R}_+^2)})$$

for the function $g \equiv \tilde{u}_x \rho^{1/2}(x)$. Note that

$$\Delta_{x,y}(\tilde{u}_x \rho^{1/2}) = (\tilde{f} - \tilde{u}_t - b\tilde{u}_x + 2u_{xy}\eta'_L + u_x\eta''_L)\rho^{1/2} + 2\tilde{u}_{xx}(\rho^{1/2})' + \tilde{u}_x(\rho^{1/2})''.$$

It follows from (2.61) that

$$\|\Delta_{x,y}(\tilde{u}_x \rho^{1/2})\|_{C([0,T];L_2(\mathbb{R}_+^2))} \leq c.$$

Moreover, by virtue of (2.61), (2.63) and embedding $H^2(\mathbb{R}_+^2) \subset H^{3/2}(\{x=0\} \times \mathbb{R})$ (see [29])

$$\begin{aligned} \|u_x|_{x=0}\|_{C([0,T];H^{3/2}(\mathbb{R}))} &\leq \|u_{0x}|_{x=0}\|_{H^{3/2}(\mathbb{R})} \\ &\quad + 2\|u_{tx}|_{x=0}\|_{L_2((0,T)\times\mathbb{R})}^{1/2} \|u_x|_{x=0}\|_{L_2(0,T;H^3(\mathbb{R}))}^{1/2} \leq c. \end{aligned}$$

Therefore,

$$\|u_x\|_{C([0,T];H^{2,\rho(x)})} \leq c. \tag{2.65}$$

Estimates (2.61), (2.63)–(2.65) provide the desired result. \square

3. Existence of solutions

Consider an auxiliary equation

$$u_t + bu_x + u_{xxx} + u_{xyy} + (g(u))_x + (\psi(t, x, y)u)_x = f(t, x, y). \tag{3.1}$$

The notion of a weak solution to problem (3.1), (1.2)–(1.4) is similar to Definition 1.1.

Lemma 3.1. *Let $g \in C^1(\mathbb{R})$, $g(0) = 0$, $|g'(u)| \leq c \forall u \in \mathbb{R}$, $\psi \in L_2(0, T; L_{\infty,+})$, $u_0 \in L_{2,+}$, $f \in L_1(0, T; L_{2,+})$ and $u_0(x, y) = f(t, x, y) = 0$ if $x > R$ for certain $R > 0$, $\mu \equiv 0$. Then problem (3.1), (1.2)–(1.4) has a unique weak solution $u(t, x, y)$, such that $u \in X^{\rho(x)}(\Pi_T^+)$ for any $\alpha > 0$ and $\rho(x) \equiv e^{2\alpha x}$.*

Proof. We apply the contraction principle. Fix $\alpha > 0$ and $\rho(x) \equiv e^{2\alpha x}$. For $t_0 \in (0, T]$ define a mapping Λ on $X^{\rho(x)}(\Pi_{t_0}^+)$ as follows: $u = \Lambda v \in X^{\rho(x)}(\Pi_{t_0}^+)$ is a weak solution to a linear problem

$$u_t + bu_x + u_{xxx} + u_{xyy} = f - g(v)_x - (\psi v)_x \tag{3.2}$$

in $\Pi_{t_0}^+$ with initial and boundary conditions (1.2)–(1.4).

Note that $\rho^2(\rho')^{-1/2} \sim \rho$, $|g(v)| \leq c|v|$ and, therefore, Lemma 2.17 provides that the mapping Λ exists. Moreover, for functions $v, \tilde{v} \in X^{\rho(x)}(\Pi_{t_0}^+)$ according to inequality (2.43) (where $f_2 \equiv 0$)

$$\|\Lambda v - \Lambda \tilde{v}\|_{X^{\rho(x)}(\Pi_{t_0}^+)} \leq c(T)\omega(t_0)\|v - \tilde{v}\|_{X^{\rho(x)}(\Pi_{t_0}^+)},$$

where $\omega(t_0) \rightarrow 0$ as $t_0 \rightarrow +0$ and ω depends on the properties of continuity of the primitives of the function $\|\psi(t, \cdot, \cdot)\|_{L_{\infty,+}}^2$ on $[0, T]$. Since the constant in the right side of this inequality is uniform with respect to u_0 and f , one can construct the solution on the whole time segment $[0, T]$ by the standard argument. \square

Now we pass to the results of existence in Theorem 1.3.

Proof of Existence Part of Theorem 1.3. First of all we make zero boundary data for $x = 0$. Let

$$\psi(t, x, y) \equiv J(t, x, y; \mu)\eta(2 - x), \tag{3.3}$$

where for the construction of the boundary potential J the function μ is extended to the whole strip B in the same class. Then the results of Section 2 provide that

$$\begin{cases} \tilde{\psi} \equiv \psi_t + b\psi_x + \psi_{xxx} + \psi_{xyy} \in C^\infty(\overline{\Pi_T^+}), & \psi = 0 \quad \text{for } x \geq 2, \\ \psi \in C([0, T]; L_{2,+}) \cap L_2(0, T; \tilde{H}_+^1) \cap L_2(0, T; W_{\infty,+}^1), \\ \psi \in C([0, 2]; \tilde{H}^{1/3,1}(B_T)), \quad \psi|_{x=0} = \mu, \quad \psi_x \in C([0, 2]; L_2(B_T)). \end{cases} \tag{3.4}$$

Consider a function

$$U(t, x, y) \equiv u(t, x, y) - \psi(t, x, y). \tag{3.5}$$

Then $u \in X_w^{\rho(x)}(\Pi_T^+)$ is a weak solution to problem (1.1)–(1.4) iff $U \in X_w^{\rho(x)}(\Pi_T^+)$ is a weak solution to an initial-boundary value problem in Π_T^+ for an equation

$$U_t + bU_x + U_{xxx} + U_{xyy} + UU_x + (\psi U)_x = F \equiv f - \tilde{\psi} - \psi\psi_x, \tag{3.6}$$

with initial and boundary conditions

$$U|_{t=0} = U_0 \equiv u_0 - \psi|_{t=0}, \quad U|_{x=0} = 0 \tag{3.7}$$

and the same boundary conditions on $\Omega_{T,+}$ as (1.4). Note also that the functions U_0, F satisfy the same assumptions as the corresponding functions u_0, f in the hypothesis of the theorem.

For $h \in (0, 1]$ consider a set of initial-boundary value problems in Π_T^+

$$U_t + bU_x + U_{xxx} + U_{xyy} + (gh(U))_x + (\psi U)_x = F_h \tag{3.8}$$

with boundary conditions (1.4) and

$$U|_{t=0} = U_{0h}, \quad U|_{x=0} = 0, \tag{3.9}$$

where

$$F_h(t, x, y) \equiv F(t, x, y)\eta(1/h - x), \quad U_{0h}(x, y) \equiv U_0\eta(1/h - x) \tag{3.10}$$

and

$$g_h(u) \equiv \int_0^u \left[\theta\eta(2 - h|\theta|) + \frac{2 \operatorname{sign} \theta}{h} \eta(h|\theta| - 1) \right] d\theta.$$

Note that $g_h(u) = u^2/2$ if $|u| \leq 1/h$, $|g'_h(u)| \leq 2/h \forall u \in \mathbb{R}$ and $|g'_h(u)| \leq 2|u|$ uniformly with respect to h .

According to Lemma 3.1, there exists a unique solution to this problem $U_h \in X^{e^{2\alpha x}}(\Pi_T^+)$ for any $\alpha > 0$.

Next, establish appropriate estimates for functions U_h uniform with respect to h (we drop the index h in intermediate steps for simplicity). First, note that $g'(U)U_x, \psi U_x, \psi_x U, F \in L_1(0, T; L_{2,+}^{\rho(x)})$ and so the hypothesis of Lemma 2.17 is satisfied (for $f_1 = f_2 \equiv 0$). Then the equality (2.44) provides that

$$\begin{aligned} & \iint U^2 \rho \, dx dy + \int_0^t \iint (3U_x^2 + U_y^2) \rho' \, dx dy d\tau \\ & \leq \iint U_0^2 \rho \, dx dy + c \int_0^t \iint U^2 \rho \, dx dy d\tau + 2 \int_0^t \iint F U \rho \, dx dy d\tau \\ & \quad + \int_0^t \iint (\psi \rho' - \psi_x \rho) U^2 \, dx dy d\tau + 2 \int_0^t \iint \left(\int_0^U g'(\theta) \theta \, d\theta \right) \rho' \, dx dy d\tau. \end{aligned} \tag{3.11}$$

Choosing $\rho \equiv 1$, we obtain, that uniformly with respect to h (and also uniformly with respect to L)

$$\|u_h\|_{C([0,T];L_{2,+})} \leq c. \tag{3.12}$$

Note that

$$\left| \int_0^U g'(\theta)\theta \, d\theta \right| \leq c|U|^3. \tag{3.13}$$

Applying interpolating inequality (1.14) for $\rho_1 = \rho_2 \equiv \rho'$, we obtain that

$$\begin{aligned} \iint |U|^3 \rho' \, dx dy &\leq \left(\iint U^2 \, dx dy \iint U^4 (\rho')^2 \, dx dy \right)^{1/2} \leq c \left(\iint U^2 \, dx dy \right)^{1/2} \\ &\times \left[\left(\iint |DU|^2 \rho' \, dx dy \right)^{1/2} \left(\iint U^2 \rho' \, dx dy \right)^{1/2} + \iint U^2 \rho' \, dx dy \right] \end{aligned} \tag{3.14}$$

(note that here the constant c is also uniform with respect to L in the cases a) and c)). Since the norm of the functions u_h in the space L_2 is already estimated in (3.12), it follows from (3.11)–(3.14) that uniformly with respect to h

$$\|u_h\|_{X^{\rho(x)}(\Pi_T^+)} \leq c. \tag{3.15}$$

Finally, write down the analogue of (3.11), where $\rho(x)$ is substituted by $\rho_0(x - x_0)$ for any $x_0 \geq 0$. Then it easily follows that (see (1.8))

$$\lambda^+(|Du_h|; T) \leq c. \tag{3.16}$$

From equation (3.8) itself, estimate (3.12) and the well-known embedding $L_{1,+} \subset H_+^{-2}$, it follows that uniformly with respect to h

$$\|u_{ht}\|_{L_1(0,T;H_+^{-3})} \leq c. \tag{3.17}$$

Estimates (3.15)–(3.17) by the standard argument provide existence of a weak solution to problem (1.1)–(1.4) $u \in X^{\rho(x)}(\Pi_T)$, $\lambda^+(|Du|; T) < \infty$ (see, for example, [12]) as a limit of functions u_h when $h \rightarrow +0$. \square

We now proceed to solutions in spaces $\tilde{H}^{1,\rho(x)}$ and first estimate a lemma analogous to Lemma 3.1.

Lemma 3.2. *Let $g(u) \equiv u^2/2$, $\psi \in L_2(0, T; W_{\infty,+}^1) \cap L_2(0, T; \tilde{H}_+^2)$, $u_0 \in \tilde{H}_+^1$, $u_0|_{x=0} \equiv 0$, $f \in L_2(0, T; \tilde{H}_+^1)$ and $u_0(x, y) = f(t, x, y) = \psi(t, x, y) = 0$ if $x > R$ for certain $R > 0$, $\mu \equiv 0$. Then problem (3.1), (1.2)–(1.4) has a unique weak solution $u(t, x, y)$, such that $u \in X^{1,\rho(x)}(\Pi_T^+)$ for any $\alpha > 0$ and $\rho(x) \equiv e^{2\alpha x}$.*

Proof. Fix $\alpha > 0$ and $\rho(x) \equiv e^{2\alpha x}$. For $t_0 \in (0, T]$ define a mapping Λ on $X^{1,\rho(x)}(\Pi_{t_0}^+)$ as follows: $u = \Lambda v \in X^{1,\rho(x)}(\Pi_{t_0}^+)$ is a weak solution to a linear problem

$$u_t + bu_x + u_{xxx} + u_{xyy} = f - vv_x - (\psi v)_x \tag{3.18}$$

in Π_{t_0} with initial and boundary conditions (1.2)–(1.4).

Note that by virtue of (1.14) for $\rho_1 = \rho_2 \equiv \rho \geq 1$

$$\|vv_x\|_{L_2(0,t_0;L_{2,+}^{\rho(x)})} \leq \left[\int_0^{t_0} \|v_x \rho^{1/2}\|_{L_{4,+}}^2 \|v \rho^{1/2}\|_{L_{4,+}}^2 \, dt \right]^{1/2} \leq ct_0^{1/4} \|v\|_{X^{1,\rho(x)}(\Pi_{t_0}^+)}^2. \tag{3.19}$$

In particular, the hypothesis of Lemma 2.19 is satisfied (since $\rho^2(\rho')^{-1/2} \sim \rho$) and, therefore, the mapping Λ exists. Moreover, by virtue of (2.48) it is easy to see, that for small t_0 , depending on $\|u_0\|_{\tilde{H}_+^{1,\rho(x)}}$, it is a contraction on a certain ball, which verifies existence of the unique weak solution to the considered problem in the space $X^{1,\rho(x)}(\Pi_{t_0}^+)$.

In order to finish the proof, we establish the following a priori estimate: if $u \in X^{1,\rho(x)}(\Pi_{T'}^+)$ is a solution to the considered problem for some $T' \in (0, T]$ and $\psi(t, x, y) = 0$ for $x > R$, then

$$\|u\|_{X^{1,\rho(x)}(\Pi_{T'}^+)} \leq c \left(\|u_0\|_{\tilde{H}_+^{1,\rho(x)}}, \|f\|_{L_2(0,T;\tilde{H}_+^{1,\rho(x)})}, \|\psi\|_{L_2(0,T;W_{\infty,+}^1 \cap \tilde{H}_+^2)} \right). \tag{3.20}$$

First of all note that similarly to (3.12), (3.15) one can derive from (2.44) that

$$\|u\|_{X^{\rho(x)}(\Pi_T^+)} \leq c. \tag{3.21}$$

Next, since the hypotheses of Lemma 2.19 and, consequently, Lemma 2.20 are satisfied, write down the corresponding analogues of equalities (2.49), (2.51) and sum them, then

$$\begin{aligned} & \iint (u_x^2 + u_y^2 - \frac{1}{3}u^3)\rho \, dx dy + \int_0^t \iint (3u_{xx}^2 + 4u_{xy}^2 + u_{yy}^2)\rho' \, dx dy d\tau \\ & \leq \iint (u_{0x}^2 + u_{0y}^2 - \frac{u_0^3}{3})\rho \, dx dy + c \int_0^t \iint (u_x^2 + u_y^2)\rho \, dx dy d\tau + 2 \int_0^t \iint uu_x^2\rho' \, dx dy d\tau \\ & \quad - \int_0^t \iint (u_{xx} + u_{yy})u^2\rho' \, dx dy d\tau + \int_0^t \iint (2f_xu_x + 2f_yu_y - fu^2)\rho \, dx dy d\tau \\ & \quad + \int_{B_t} f^2|_{x=0} dy d\tau + c \iint_{B_t} u_x^2|_{x=0} dy d\tau - \int_0^t \iint (\frac{b}{3}u^3 + \frac{1}{4}u^4)\rho' \, dx dy d\tau \\ & \quad - \iint_{B_t} (\psi u_x^2\rho)|_{x=0} dy d\tau - \int_0^t \iint \psi_x(3u_x^2 + u_y^2)\rho \, dx dy d\tau + \int_0^t \iint \psi(u_x^2 + u_y^2)\rho' \, dx dy d\tau \\ & \quad - 2 \int_0^t \iint \psi_y u_x u_y \rho \, dx dy d\tau - 2 \int_0^t \iint (\psi_{xx}uu_x + \psi_{xy}uu_y)\rho \, dx dy d\tau. \end{aligned} \tag{3.22}$$

Interpolating inequality (1.16) provides that

$$\begin{aligned} \int_0^L ((1 + |\psi|)u_x^2)|_{x=0} dy & \leq \varepsilon \iint u_{xx}^2\rho' \, dx dy \\ & \quad + c(\varepsilon)(1 + \sup_{(x,y) \in \Sigma_+} \psi^2) \iint u_x^2\rho \, dx dy, \end{aligned} \tag{3.23}$$

where $\varepsilon > 0$ can be chosen arbitrarily small. Next, since $\rho(x) \leq c(R)\rho'(x)$ for $x \in [0, R]$

$$\begin{aligned} \left| \iint \psi_{xx}uu_x\rho \, dx dy \right| & \leq c \left(\iint \psi_{xx}^2 \, dx dy \right)^{1/2} \left(\iint u^4\rho^2 \, dx dy \iint u_x^4\rho' \, dx dy \right)^{1/4} \\ & \leq \varepsilon \iint |Du_x|^2\rho' \, dx dy + c(\varepsilon) \left[\iint \psi_{xx}^2 \, dx dy + 1 \right] \left(\iint |Du|^2\rho \, dx dy + 1 \right). \end{aligned} \tag{3.24}$$

Other integrals from (3.22) can be estimated, for example, similarly to the ones from the proof of Theorem 1.2 from [14], where for the sake of the use in the sequel can be assumed only that ρ and ρ' are admissible weight functions. Then (3.20) follows. \square

Proof of Existence Part of Theorem 1.5. Introduce the function ψ by formula (3.3). Then in addition to properties (3.4) it follows from the results of Section 2 that

$$\begin{cases} \psi \in C([0, T]; \tilde{H}_+^1) \cap L_2(0, T; \tilde{H}_+^2) \cap C([0, 2]; \tilde{H}^{2/3,2}(B_T)), \\ \psi_x \in C([0, 2]; \tilde{H}^{1/3,2}(B_T)), \quad \psi_{xx} \in C([0, 2]; L_2(B_T)). \end{cases} \tag{3.25}$$

Again introduce the function $U(t, x, y)$ by formula (3.5) and consider problem (3.6), (3.7), (1.4) instead of (1.1)–(1.4). Note that here (3.4), (3.25) provide that the properties of the functions U_0, F , are the same as the corresponding ones for the functions u_0, f in the hypothesis of the theorem.

For $h \in (0, 1]$, consider a set of initial-boundary value problems in Π_T^+

$$U_t + bU_x + U_{xxx} + U_{xyy} + UU_x + (\psi U)_x = F_h \tag{3.26}$$

with boundary conditions (3.7), (1.4), where F_h and U_{0h} are given by (3.10).

Repeating the argument in (3.11)–(3.14) for $g(u) \equiv u^2/2$ and (3.22)–(3.24) we derive that uniformly with respect to h

$$\|u_h\|_{X^{1,\rho(x)}(\Pi_T^+)} \leq c. \tag{3.27}$$

Similarly to (3.16) one can obtain that

$$\lambda^+(|D^2 u_h|; T) \leq c. \tag{3.28}$$

Estimates (3.27), (3.28) and (3.17) provide existence of a weak solution to the considered problem $u \in X^{1,\rho(x)}(\Pi_T^+)$. \square

Finally, we consider regular solutions.

Lemma 3.3. *Let $g(u) \equiv u^2/2$, $\mu \equiv 0$, the functions u_0 and f satisfy the hypothesis of Theorem 1.7, $\psi \in X^{3,\rho(x)}(\Pi_T^+) \cap L_2(0, T; W_{\infty,+}^3)$, $\psi_t \in L_2(0, T; L_{\infty,+})$ and $\psi(t, x, y) = 0$ if $x > R$ for certain $R > 0$. Then problem (3.1), (1.2)–(1.4) has a unique solution $u \in X^{3,\rho(x)}(\Pi_T^+)$.*

Proof. For $t_0 \in (0, T]$, $v \in X^{3,\rho(x)}(\Pi_{t_0}^+)$ let $u = \Lambda v \in X^{3,\rho(x)}(\Pi_{t_0}^+)$ be a solution to linear problem (3.18), (1.2)–(1.4).

Apply Lemma 2.23, where f stands for $f_0, -(v^2/2 + \psi v) -$ for f_1 . We have:

$$\begin{aligned} \|v v_x + \psi v_x + \psi_x v\|_{C([0,t_0]; L_{2,+}^{\rho(x)})} &\leq \|u_0 u_{0x} + (\psi|_{t=0} u_0)_x\|_{L_{2,+}^{\rho(x)}} \\ &\quad + \|(v v_x)_t + (\psi v)_{tx}\|_{L_1(0,t_0; L_{2,+}^{\rho(x)})}, \end{aligned} \tag{3.29}$$

where with the use of the embedding $H_+^2 \subset L_{\infty,+}$

$$\|u_0 u_{0x}\|_{L_{2,+}^{\rho(x)}} \leq c \|u_0\|_{L_{\infty,+}} \|u_{0x}\|_{L_{2,+}^{\rho(x)}} \leq c_1 \|u_0\|_{\tilde{H}_+^{3,\rho(x)}}^2, \tag{3.30}$$

$$\|(\psi|_{t=0} u_0)_x\|_{L_{2,+}^{\rho(x)}} \leq c (\|\psi_x|_{t=0}\|_{L_{2,+}} + \|\psi|_{t=0}\|_{L_{\infty,+}}) \|u_0\|_{\tilde{H}_+^{3,\rho(x)}}, \tag{3.31}$$

then since $\rho'(x) \geq 1$

$$\begin{aligned} \|(v v_x)_t\|_{L_1(0,t_0; L_{2,+}^{\rho(x)})} &\leq \int_0^{t_0} \left[\|v \rho^{1/2}\|_{L_{\infty,+}} \|v_{tx}\|_{L_{2,+}^{\rho(x)}} \right. \\ &\quad \left. + \|v_x \rho^{1/2}\|_{L_{4,+}} \|v_t (\rho' \rho)^{1/4}\|_{L_{4,+}} \right] dt \leq c t_0^{1/2} \|v\|_{X^{2,\rho(x)}(\Pi_{t_0}^+)} \|v\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)}. \end{aligned} \tag{3.32}$$

Next,

$$\begin{aligned} \|v v_t\|_{L_2(0,t_0; L_{2,+}^{\rho^2(x)/\rho'(x)})} &\leq \left(\int_0^{t_0} \|v \rho^{1/2}\|_{L_{\infty,+}}^2 \|v_t\|_{L_{2,+}^{\rho(x)}}^2 dt \right)^{1/2} \\ &\leq c t_0^{1/2} \|v\|_{X^{2,\rho(x)}(\Pi_{t_0}^+)} \|v\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)}, \end{aligned} \tag{3.33}$$

$(v v_x)_{yy} = v v_{xyy} + 2v_y v_{xy} + v_x v_{yy}$, where similarly to (3.33)

$$\|v v_{xyy}\|_{L_2(0,t_0; L_{2,+}^{\rho^2(x)/\rho'(x)})} \leq c t_0^{1/2} \|v\|_{X^{2,\rho(x)}(\Pi_{t_0}^+)} \|v\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)}, \tag{3.34}$$

$$\begin{aligned} \|v_y v_{xy}\|_{L_2(0,t_0; L_{2,+}^{\rho^2(x)/\rho'(x)})} &\leq \left(\int_0^{t_0} \|v_y \rho^{1/2}\|_{L_{4,+}}^2 \|v_{xy} \rho^{1/2}\|_{L_{4,+}}^2 dt \right)^{1/2} \\ &\leq c t_0^{1/2} \|v\|_{X^{2,\rho(x)}(\Pi_{t_0}^+)} \|v\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)}. \end{aligned} \tag{3.35}$$

and similar estimate holds for v_x, v_{yy} . Moreover, the assumptions on the function ψ ensure that the corresponding boundary conditions on the function f_1 are satisfied for $y = 0$ and $y = L$. Therefore, the mapping Λ exists and one can use estimate (2.60) to derive inequalities

$$\|\Lambda v\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)} \leq \tilde{c} + c(T)t_0^{1/2}(\|\psi\|_{X^{3,\rho(x)}(\Pi_T^+)}\|v\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)} + \|v\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)}^2), \tag{3.36}$$

$$\begin{aligned} \|\Lambda v - \Lambda \tilde{v}\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)} &\leq c(T)t_0^{1/2}\left(\|\psi\|_{X^{3,\rho(x)}(\Pi_T^+)}\|v - \tilde{v}\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)} \right. \\ &\quad \left. + (\|v\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)} + \|\tilde{v}\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)})\|v - \tilde{v}\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)}\right), \end{aligned} \tag{3.37}$$

where the constant \tilde{c} depends on the properties of functions u_0, f, ψ . Hence, existence of the unique solution to the considered problem in the space $X^{3,\rho(x)}(\Pi_{t_0}^+)$ on the time interval $[0, t_0]$, depending on $\|u_0\|_{\tilde{H}_+^{3,\rho(x)}}$, follows by the standard argument.

Now establish the following a priori estimate: if $u \in X^{3,\rho(x)}(\Pi_{T'}^+)$ is a solution to the considered problem for some $T' \in (0, T]$, then

$$\|u\|_{X^{3,\rho(x)}(\Pi_{T'}^+)} \leq c, \tag{3.38}$$

where the constant c depends on T and the properties of the functions u_0, f, ψ from the hypothesis of the present lemma.

According to (3.20)

$$\|u\|_{X^{1,\rho(x)}(\Pi_{T'}^+)} \leq c. \tag{3.39}$$

Next, since the hypothesis of Lemma 2.21 is fulfilled write down the corresponding analogue of equality (2.44) for the function u_t :

$$\begin{aligned} &\iint u_t^2 \rho \, dx dy + \int_0^t \iint (3u_{tx}^2 + u_{ty}^2) \rho' \, dx dy d\tau \\ &\leq \iint (f - bu_x - u_{xxx} - u_{xyy} - uu_x - (\psi u)_x)^2|_{t=0} \rho \, dx dy + c \int_0^t \iint u_t^2 \rho \, dx dy d\tau \\ &\quad + 2 \int_0^t \iint (f - (\psi u)_x)_t u_t \rho \, dx dy d\tau + 2 \int_0^t \iint uu_t (u_t \rho)_x \, dx dy d\tau. \end{aligned} \tag{3.40}$$

Here since $\rho' \geq 1$ and estimate (3.39) holds

$$\begin{aligned} 2 \iint uu_t (u_t \rho)_x \, dx dy &= \iint (u\rho' - u_x \rho) u_t^2 \, dx dy \\ &\leq c \left(\iint \left(u_x^2 \frac{\rho}{\rho'} + u^2 \right) \, dx dy \iint u_t^4 \rho' \rho \, dx dy \right)^{1/2} \\ &\leq c_1 \left[\left(\iint |Du_t|^2 \rho' \, dx dy \iint u_t^2 \rho \, dx dy \right)^{1/2} + \iint u_t^2 \rho \, dx dy \right], \\ &\left| \iint \psi_{tx} uu_t \rho \, dx dy \right| \\ &\leq c(R) \left(\iint \psi_{tx}^2 \, dx dy \right)^{1/2} \left(\iint u^4 \rho^2 \, dx dy \iint u_t^4 \rho' \rho \, dx dy \right)^{1/4} \\ &\leq \varepsilon \iint |Du_t|^2 \rho' \, dx dy + c(\varepsilon, R) \left[\iint \psi_{tx}^2 \, dx dy + 1 \right] \left(\iint u_t^2 \rho \, dx dy + 1 \right), \end{aligned}$$

where $\varepsilon > 0$ can be chosen arbitrarily small. Other terms in (3.40) are estimated in a obvious way and, consequently,

$$\|u_t\|_{X^{\rho(x)}(\Pi_{T'}^+)} \leq c. \tag{3.41}$$

Now apply Lemma 2.22, then inequality (2.52) and estimates (3.39), (3.41) yield that for any $t \leq T'$

$$\begin{aligned} \|u\|_{X^{2,\rho(x)}(\Pi_t^+)}^2 &\leq c + c\|uu_x\|_{C([0,t];L_{2,+}^{\rho(x)})}^2 \\ &\quad + c \sup_{\tau \in (0,t]} \left| \int_0^\tau \iint (f - uu_x - (\psi u)_x)_{yy} u_{yy} \rho \, dx dy ds \right| + c \int_0^t \iint (u_{xx}^2 + u_{yy}^2) \rho \, dx dy \end{aligned} \quad (3.42)$$

We have

$$\|uu_x\|_{L_{2,+}^{\rho(x)}}^2 \leq c\|u_x \rho^{1/2}\|_{L_{4,+}}^2 \leq \varepsilon \| |Du_x| \|_{L_{2,+}^{\rho(x)}}^2 + c(\varepsilon),$$

where $\varepsilon > 0$ can be chosen arbitrarily small;

$$\iint (uu_x)_{yy} u_{yy} \rho \, dx dy = \frac{1}{2} \iint (u_x \rho - u \rho') u_{yy}^2 \, dx dy + 2 \iint u_y u_{xy} u_{yy} \rho \, dx dy,$$

where again since $\rho' \geq 1$

$$\begin{aligned} \left| \iint u_y u_{xy} u_{yy} \rho \, dx dy \right| &\leq \left(\iint u_y^2 \rho \, dx dy \iint (u_{xy}^4 + u_{yy}^4) \rho' \rho \, dx dy \right)^{1/2} \\ &\leq c_1 \left(\iint |D^3 u|^2 \rho' \, dx dy \iint |D^2 u|^2 \rho \, dx dy \right)^{1/2}. \end{aligned}$$

Integral of $(u_x \rho - u \rho') u_{yy}^2$ is estimated in a similar way and it follows from (3.42) that

$$\|u\|_{X^{2,\rho(x)}(\Pi_{T'}^+)} \leq c. \quad (3.43)$$

Finally, apply Lemma 2.23 on the basis of the already obtained estimates (3.41), (3.43), then inequality (2.60) and estimates (3.29)–(3.35) applied to $v \equiv u$ provide similarly to (3.36) that for any $t_0 \in (0, T']$

$$\|u\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)} \leq \tilde{c} + c(T) t_0^{1/2} (\|\psi\|_{X^{3,\rho(x)}(\Pi_T^+)} + \|u\|_{X^{2,\rho(x)}(\Pi_T^+)}) \|u\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)},$$

whence (3.38) follows. \square

Proof of Theorem 1.7. Introduce the functions ψ, U by formulas (3.3), (3.5) and consider problem (3.6), (3.7), (1.4). Then the functions $\psi, F \sim f$ and $U_0 \sim u_0$ satisfy the hypothesis of Lemma 3.3 and the result is immediate. \square

4. Uniqueness and continuous dependence

Theorem 4.1. Let $\rho(x)$ be an admissible weight function, such that $\rho'(x)$ is also an admissible weight function and $\rho^{1/2}(x) \leq c_0 \rho'(x) \forall x \geq 0$ for certain positive constant c_0 . Then for any $T > 0$ and $M > 0$ there exists a constant $c = c(T, M)$, such that for any two weak solutions $u(t, x, y)$ and $\tilde{u}(t, x, y)$ to problem (1.1)–(1.4), satisfying $\|u\|_{X_w^{\rho(x)}(\Pi_T^+)}, \|\tilde{u}\|_{X_w^{\rho(x)}(\Pi_T^+)} \leq M$, with corresponding data $u_0, \tilde{u}_0 \in L_{2,+}^{\rho(x)}, \mu, \tilde{\mu} \in \tilde{H}^{1/3,1}(B_T), f, \tilde{f} \in L_1(0, T; L_{2,+}^{\rho(x)})$ the following inequality holds:

$$\|u - \tilde{u}\|_{X_w^{\rho(x)}(\Pi_T^+)} \leq c (\|u_0 - \tilde{u}_0\|_{L_{2,+}^{\rho(x)}} + \|\mu - \tilde{\mu}\|_{H^{1/3,1}(B_T)} + \|f - \tilde{f}\|_{L_1(0,T;L_{2,+}^{\rho(x)})}). \quad (4.1)$$

Proof. Let the function ψ is defined by formula (3.3), the function $\tilde{\psi}$ in a similar way for $\tilde{\mu}$ and $\Psi \equiv \psi - \tilde{\psi}$. Then, in particular,

$$\|\Psi\|_{X^{\rho(x)}(\Pi_T^+)} \leq c(T) \|\mu - \tilde{\mu}\|_{H^{1/3,1}(B_T)}. \quad (4.2)$$

Let $U_0 \equiv u_0 - \tilde{u}_0 - \Psi|_{t=0}, F \equiv f - \tilde{f} - (\Psi_t + b\Psi_x + \Psi_{xx} + \Psi_{xy})$, then

$$\|U_0\|_{L_{2,+}^{\rho(x)}} \leq \|u_0 - \tilde{u}_0\|_{L_{2,+}^{\rho(x)}} + c(T) \|\mu - \tilde{\mu}\|_{H^{1/3,1}(B_T)}, \quad (4.3)$$

$$\|F\|_{L_1(0,T;L_{2,+}^{\rho(x)})} \leq \|f - \tilde{f}\|_{L_1(0,T;L_{2,+}^{\rho(x)})} + c(T) \|\mu - \tilde{\mu}\|_{H^{1/3,1}(B_T)}. \quad (4.4)$$

The function $U(t, x, y) \equiv u(t, x, y) - \tilde{u}(t, x, y) - \Psi(t, x, y)$ is a weak solution to an initial-boundary value problem in Π_T^+ for an equation

$$U_t + bU_x + U_{xxx} + U_{xyy} = F - (uu_x - \tilde{u}\tilde{u}_x) \tag{4.5}$$

with initial and boundary conditions (1.4),

$$U|_{t=0} = U_0, \quad U|_{x=0} = 0. \tag{4.6}$$

Apply Lemma 2.17 where $f_2 = -(uu_x - \tilde{u}\tilde{u}_x)$. Note that assumptions on the function ρ provide that $\rho(\rho')^{-1/3} \leq c\rho^{1/3}\rho'$ and by virtue of (1.14)

$$\begin{aligned} \|uu_x\rho^{3/4}(\rho')^{-1/4}\|_{L_{4/3}(\Pi_T^+)}^{4/3} &\leq c \int_0^T \left(\iint u^4\rho'\rho \, dx dy \right)^{1/3} \left(\iint u_x^2\rho' \, dx dy \right)^{2/3} dt \\ &\leq c_1 \|u\|_{L_\infty(0,T;L_{2,+}^{\rho(x)})}^{1/3} \|u\|_{L_2(0,T;H_+^{1,\rho'(x)})} < +\infty. \end{aligned}$$

Therefore, we derive from (2.44) that for $t \in (0, T]$

$$\begin{aligned} &\iint U^2\rho \, dx dy + \int_0^t \iint (3U_x^2 + U_y^2)\rho' \, dx dy d\tau \\ &\leq \iint U_0^2\rho \, dx dy + c \int_0^t \iint U^2\rho \, dx dy d\tau + 2 \int_0^t \iint (F - (uu_x - \tilde{u}\tilde{u}_x))U\rho \, dx dy d\tau. \end{aligned} \tag{4.7}$$

Here

$$-2 \iint (uu_x - \tilde{u}\tilde{u}_x)U\rho \, dx dy = \iint (u + \tilde{u})(U + \Psi)(U\rho)_x \, dx dy. \tag{4.8}$$

Then by virtue of (1.14) and the assumptions on the function ρ (which yield that $(\rho/\rho')^3 \leq c\rho'\rho$)

$$\begin{aligned} &\iint |u(U + \Psi)U_x|\rho \, dx dy \\ &\leq c \left(\iint u^4(\rho/\rho')^3 \, dx dy \iint (U^4 + \Psi^4)\rho'\rho \, dx dy \right)^{1/4} \left(\iint U_x^2\rho' \, dx dy \right)^{1/2} \\ &\leq c_1 \left(\|u\|_{H_+^{1,\rho'(x)}}^{1/2} \|u\|_{L_{2,+}^{\rho(x)}}^{1/2} + \|u\|_{L_{2,+}^{\rho(x)}} \right) \left[\iint (|DU|^2 + |D\Psi|^2)\rho' \, dx dy \right]^{3/4} \\ &\quad \times \left[\iint (U^2 + \Psi^2)\rho \, dx dy \right]^{1/4} + \iint (U^2 + \Psi^2)\rho \, dx dy \end{aligned} \tag{4.9}$$

and, therefore,

$$\begin{aligned} \int_0^t \iint |u(U + \Psi)U_x|\rho \, dx dy d\tau &\leq \varepsilon \int_0^t \iint (|DU|^2 + |D\Psi|^2)\rho' \, dx dy d\tau \\ &\quad + c(\varepsilon) \int_0^t \gamma(\tau) \iint (U^2 + \Psi^2)\rho \, dx dy d\tau, \end{aligned} \tag{4.10}$$

where $\varepsilon > 0$ can be chosen arbitrarily small and $\gamma \equiv 1 + \|u\|_{H_+^{1,\rho'(x)}}^2 \in L_1(0, T)$. Then estimates (4.2)–(4.4), (4.10) and inequality (4.7) provide the desired result. \square

Remark 4.2. Theorems 1.3 and 4.1 show that under the hypothesis of Theorem 1.3 problem (1.1)–(1.4) is globally well-posed in the space $X_w^{\rho(x)}(\Pi_T^+)$.

Theorem 4.3. Let $\rho(x)$ be an admissible weight function, such that $\rho'(x)$ is also an admissible weight function and $\rho'(x) \geq c_0 \forall x \geq 0$ for certain positive constant c_0 . Then for any $T > 0$ and $M > 0$ there exists a constant $c = c(T, M)$, such that for any two weak solutions $u(t, x, y)$ and $\tilde{u}(t, x, y)$ to problem (1.1)–(1.4), satisfying $\|u\|_{X_w^{1,\rho(x)}(\Pi_T^+)} , \|\tilde{u}\|_{X_w^{1,\rho(x)}(\Pi_T^+)} \leq M$, with corresponding data $u_0, \tilde{u}_0 \in L_{2,+}^{\rho(x)}$, $\mu, \tilde{\mu} \in \tilde{H}^{1/3,1}(B_T)$, $f, \tilde{f} \in L_1(0, T; L_{2,+}^{\rho(x)})$ inequality (4.1) holds.

Proof. The proof mostly repeats the proof of Theorem 4.1. The difference is related only to the nonlinear term. Here we apply Lemma 2.17 where $f_1 = -(u^2 - \tilde{u}^2)/2$. Note that for any $t \in [0, T]$

$$\|u^2\|_{L_{2,+}^{\rho^2(x)/\rho'(x)}} \leq \|u\|_{H_+^{1,\rho(x)}}^2,$$

in particular, $u^2 \in L_\infty(0, T; L_{2,+}^{\rho^2(x)/\rho'(x)})$. Write down inequality (4.7). In comparison with (4.8) we transform the integral of the nonlinear term in the following way:

$$\begin{aligned} 2 \iint (uu_x - \tilde{u}\tilde{u}_x)U\rho dx dy &= \frac{1}{2} \iint (u + \tilde{u})_x U^2 \rho dx dy \\ &\quad - \frac{1}{2} \iint (u + \tilde{u})U^2 \rho' dx dy + \iint ((u + \tilde{u})\Psi)_x U\rho dx dy. \end{aligned}$$

Here

$$\begin{aligned} \iint |u_x|(U^2 + \Psi^2)\rho dx dy &\leq \left(\iint u_x^2 \frac{\rho}{\rho'} dx dy \iint (U^4 + \Psi^4)\rho' \rho dx dy \right)^{1/2} \\ &\leq c \left[\iint (|DU|^2 + |D\Psi|^2)\rho' dx dy \iint (U^2 + \Psi^2)\rho dx dy \right]^{1/2} \\ &\quad + \iint (U^2 + \Psi^2)\rho dx dy, \\ \iint |u\Psi_x U|\rho dx dy &\leq c \left(\iint u^4 \rho^2 dx dy \iint U^4 \rho' \rho dx dy \right)^{1/4} \left(\iint \Psi_x^2 \rho dx dy \right)^{1/2} \\ &\leq \varepsilon \iint |DU|^2 \rho' dx dy + c(\varepsilon) \iint U^2 \rho' dx dy + \iint \Psi_x^2 \rho dx dy. \end{aligned}$$

Note that similarly to (4.2)

$$\|\Psi_x\|_{L_2(0,T;L_{2,+}^{\rho(x)})} \leq c(T)\|\mu - \tilde{\mu}\|_{H^{1/3,1}(B_T)} \tag{4.11}$$

since $\Psi = 0$ for $x \geq 2$. Then the desired result succeeds from inequality (4.7). \square

Theorem 4.4. Let $\rho(x)$ be an admissible weight function, such that $\rho'(x)$ is also an admissible weight function and $\rho^{1/3}(x) \leq c_0\rho'(x) \forall x \geq 0$ for certain positive constant c_0 . Then for any $T > 0$ and $M > 0$ there exists a constant $c = c(T, M)$, such that for any two weak solutions $u(t, x, y)$ and $\tilde{u}(t, x, y)$ to problem (1.1)–(1.4), satisfying $\|u\|_{X_w^{1,\rho(x)}(\Pi_T^+)} , \|\tilde{u}\|_{X_w^{1,\rho(x)}(\Pi_T^+)} \leq M$, with corresponding data $u_0, \tilde{u}_0 \in \tilde{H}_+^{1,\rho(x)}$, $\mu, \tilde{\mu} \in \tilde{H}^{2/3,2}(B_T)$, $f, \tilde{f} \in L_2(0, T; \tilde{H}_+^{1,\rho(x)})$, $u_0(0, y) \equiv \mu(0, y)$, $\tilde{u}_0(0, y) \equiv \tilde{\mu}(0, y)$, the following inequality holds:

$$\|u - \tilde{u}\|_{X_w^{1,\rho(x)}(\Pi_T^+)} \leq c(\|u_0 - \tilde{u}_0\|_{H_+^{1,\rho(x)}} + \|\mu - \tilde{\mu}\|_{H^{2/3,2}(B_T)} + \|f - \tilde{f}\|_{L_2(0,T;H_+^{1,\rho(x)})}). \tag{4.12}$$

Proof. First of all note that the hypothesis of Theorem 4.3 holds and, consequently, inequality (4.1) is satisfied. Introduce the same functions Ψ, U_0, F, U as in the proof of Theorem 4.1. Note that

$$\begin{aligned} \|\Psi\|_{X^{1,\rho(x)}(\Pi_T^+)} &\leq c(T)\|\mu - \tilde{\mu}\|_{H^{2/3,2}(B_T)}, \\ \|U_0\|_{H_+^{1,\rho(x)}} &\leq \|u_0 - \tilde{u}_0\|_{H_+^{1,\rho(x)}} + c(T)\|\mu - \tilde{\mu}\|_{H^{2/3,2}(B_T)}, \\ \|F\|_{L_2(0,T;H_+^{1,\rho(x)})} &\leq \|f - \tilde{f}\|_{L_2(0,T;H_+^{1,\rho(x)})} + c(T)\|\mu - \tilde{\mu}\|_{H^{2/3,2}(B_T)}. \end{aligned}$$

Apply Lemma 2.19. Note that since $\rho^2(\rho')^{-1} \leq c\rho^{3/2}(\rho')^{1/2}$

$$\begin{aligned} \iint u^2 u_x^2 \rho^2(\rho')^{-1} dx dy &\leq c \left(\iint u^4 \rho^2 dx dy \iint u_x^4 \rho' dx dx \right)^{1/2} \\ &\leq c_1 \iint (|DU_x|^2 \rho' + u_x^2 \rho) dx dy. \end{aligned}$$

In particular, $uu_x \in L_2(0, T; L_{2,+}^{\rho^2(x)/\rho'(x)})$. Then inequality (2.49) for $\gamma \equiv 1$ (together with (1.16)) yields that for $t \in (0, T]$

$$\begin{aligned} &\iint (U_x^2 + U_y^2) \rho dx dy + \int_0^t \iint |D^2 U|^2 \rho' dx dy d\tau \\ &\leq \iint (U_{0x}^2 + U_{0y}^2) \rho dx dy + c \int_0^t \iint (U_x^2 + U_y^2) \rho dx dy d\tau + \int_0^t \iint (F_x^2 + F_y^2 + F^2) \rho dx dy d\tau \\ &\quad + 2 \int_0^t \iint (uu_x - \tilde{u}\tilde{u}_x) [(U_x \rho)_x + U_{yy} \rho] dx dy d\tau. \end{aligned} \tag{4.13}$$

The last integral in the right side of (4.13) is not greater than

$$\begin{aligned} &\varepsilon \int_0^t \iint (U_{xx}^2 + U_{yy}^2 + U_x^2) \rho' dx dy d\tau \\ &\quad + c(\varepsilon) \int_0^t \iint [(u_x^2 + \tilde{u}_x^2)(U^2 + \Psi^2) + (u^2 + \tilde{u}^2)(U_x^2 + \Psi_x^2)] \frac{\rho^2}{\rho'} dx dy d\tau, \end{aligned}$$

where $\varepsilon > 0$ can be chosen arbitrarily small. Here again since $\rho^2(\rho')^{-1} \leq c\rho^{3/2}(\rho')^{1/2}$

$$\begin{aligned} &\iint u_x^2 (U^2 + \Psi^2) \frac{\rho^2}{\rho'} dx dy \\ &\leq c \left(\iint u_x^4 \rho' dx dy \iint (U^4 + \Psi^4) \rho^2 dx dy \right)^{1/2} \\ &\leq c_1 \iint (|DU|^2 + |D\Psi|^2) \rho dx dy + c_1 \left[\iint |DU_x|^2 \rho' dx dy + 1 \right] \iint (U^2 + \Psi^2) \rho dx dy, \end{aligned}$$

where the first multiplier in the last term belongs to the space $L_1(0, T)$ and the second one is estimated uniformly with respect to t according to (4.1) and (4.2). Finally,

$$\begin{aligned} &\iint u^2 (U_x^2 + \Psi_x^2) \frac{\rho^2}{\rho'} dx dy \\ &\leq c \left(\iint u^4 \rho^2 dx dy \iint (U_x^4 + \Psi_x^4) \rho' \rho dx dy \right)^{1/2} \\ &\leq c_1 \left(\iint (|DU_x|^2 + |D\Psi_x|^2) \rho' dx dy \iint (U_x^2 + \Psi_x^2) \rho dx dy \right)^{1/2} + \iint (U_x^2 + \Psi_x^2) \rho dx dy. \end{aligned}$$

As a result, the statement of the theorem follows from inequality (4.13). \square

Remark 4.5. Theorems 1.5, 4.3 and 4.4 show that under the hypothesis of Theorem 1.5 and additional assumption $\rho^{1/3}(x) \leq c_0 \rho'(x) \forall x \geq 0$ problem (1.1)–(1.4) is globally well-posed in the space $X_w^{1,\rho(x)}(\Pi_T^+)$. This additional assumption holds for any exponential weight $e^{2\alpha x}$, $\alpha > 0$, and for the power weight $(1+x)^{2\alpha}$ if $\alpha \geq 3/4$.

For regular solutions we prefer to present well-posedness in another form.

Theorem 4.6. *Let $T > 0$ and $\rho(x)$ be an admissible weight function, such that $\rho'(x)$ is also an admissible weight function and $\rho'(x) \geq 1 \forall x \geq 0$. Denote by \mathcal{F} the space of functions $f(t, x, y)$, defined on Π_T^+ and satisfying the hypothesis of Theorem 1.7, endowed with the natural norm. Then the mapping $(u_0, \mu, f) \mapsto u$, where u is the corresponding solution of problem (1.1)–(1.4) and $u_0(0, y) \equiv \mu(0, y)$, is Lipschitz continuous on any ball in the norm of the mapping $\tilde{H}_+^{3,\rho(x)} \times \tilde{H}^{4/3,4}(B_T) \times \mathcal{F} \rightarrow X^{3,\rho(x)}(\Pi_T^+)$.*

Proof. Let $M > 0$, let the functions u_0, μ, f satisfy the hypothesis of Theorem 1.7 and

$$\|(u_0, \mu, f)\|_{\tilde{H}_+^{3,\rho(x)} \times \tilde{H}^{4/3,4}(B_T) \times \mathcal{F}} \leq M,$$

then it follows from (3.38) that $\|u\|_{X^{3,\rho(x)}(\Pi_T^+)} \leq c_0(M)$. Define the functions ψ and U by formulas (3.3) and (3.5). Let the triplet $(\tilde{u}_0, \tilde{\mu}, \tilde{f})$ be another one satisfying the same assumptions, define similarly the functions $\tilde{\psi}$ and \tilde{U} . Then similarly to (3.29)–(3.37) for $t_0 \in (0, T]$

$$\begin{aligned} \|U - \tilde{U}\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)} &\leq c(M) (\|u_0 - \tilde{u}_0\|_{\tilde{H}_+^{3,\rho(x)}} + \|\mu - \tilde{\mu}\|_{\tilde{H}^{4/3,4}(B_T)} + \|f - \tilde{f}\|_{\mathcal{F}} \\ &\quad + t_0^{1/2} \|U - \tilde{U}\|_{X^{3,\rho(x)}(\Pi_{t_0}^+)}). \end{aligned}$$

Taking into account also that $\|\psi - \tilde{\psi}\|_{X^{3,\rho(x)}(\Pi_T^+)} \leq c(T) \|\mu - \tilde{\mu}\|_{\tilde{H}^{4/3,4}(B_T)}$ we finish the proof by the standard argument. \square

5. Large-time decay of small solutions

Proof of Theorem 1.9. Let $\alpha > 0$, $\rho(x) \equiv e^{2\alpha x}$, $u_0 \in \tilde{H}_+^{1,\rho(x)}$, $u_0(0, y) \equiv 0$, $\mu \equiv 0$, $f \equiv 0$. Consider the solution to problem (1.1)–(1.4) (in the cases a) and c)) $u \in X_w^{1,\rho(x)}(\Pi_T^+) \forall T$. Note that $uu_x \in L_2(0, T; L_{2,+}^{\rho(x)})$ (see, for example, (3.19)).

Apply Lemma 2.17, then equality (2.44) for $\rho \equiv 1$ provides, in fact, the conservation law (1.5), in particular,

$$\|u(t, \cdot, \cdot)\|_{L_{2,+}} \leq \|u_0\|_{L_{2,+}} \quad \forall t \geq 0. \tag{5.1}$$

Next, write down equality (2.44) for $\rho \equiv e^{2\alpha x}$:

$$\begin{aligned} \iint_{B_t} u^2 \rho \, dx dy + \iint_{B_t} u_x^2|_{x=0} \, dy d\tau + 2\alpha \int_0^t \iint (3u_x^2 + u_y^2) \rho \, dx dy d\tau \\ - 2\alpha(b + 4\alpha^2) \int_0^t \iint u^2 \rho \, dx dy d\tau = \iint u_0^2 \rho \, dx dy + \frac{2\alpha}{3} \int_0^t \iint u^3 \rho \, dx dy d\tau. \end{aligned} \tag{5.2}$$

Since $u^3 \rho \in L_\infty(0, T; L_{1,+})$ equality (5.2) provides the following equality in a differential form: for a.e. $t > 0$

$$\begin{aligned} \frac{d}{dt} \iint u^2 \rho \, dx dy + \int_0^L u_x^2|_{x=0} \, dy + 2\alpha \iint (3u_x^2 + u_y^2) \rho \, dx dy \\ - 2\alpha(b + 4\alpha^2) \iint u^2 \rho \, dx dy = \frac{2\alpha}{3} \iint u^3 \rho \, dx dy. \end{aligned} \tag{5.3}$$

Continuing inequality (3.14), we find with the use of (5.1) that uniformly with respect to L

$$\frac{2}{3} \iint u^3 \rho \, dx dy \leq \frac{1}{2} \iint |Du|^2 \rho \, dx dy + c(\|u_0\|_{L_{2,+}} + \|u_0\|_{L_{2,+}}^2) \iint u^2 \rho \, dx dy. \tag{5.4}$$

Inequality (1.17) yields that for certain constant c_0

$$\frac{1}{2} \iint u_y^2 \rho \, dx dy \geq \frac{c_0}{L^2} \iint u^2 \rho \, dx dy. \tag{5.5}$$

Combining (5.3)–(5.5) we find that uniformly with respect to α and L

$$\begin{aligned} & \frac{d}{dt} \iint u^2 \rho \, dx dy + \int_0^L u_x^2|_{x=0} \, dy + \alpha \iint |Du|^2 \rho \, dx dy \\ & + \alpha \left(\frac{c_0}{L^2} - 2b - 8\alpha^2 - c(\|u_0\|_{L_{2,+}} + \|u_0\|_{L_{2,+}}^2) \right) \iint u^2 \rho \, dx dy \leq 0. \end{aligned} \tag{5.6}$$

Choose $L_0 = \frac{1}{2} \sqrt{\frac{c_0}{b}}$ if $b > 0$, $\alpha_0 = \frac{\sqrt{c_0}}{8L}$, $\epsilon > 0$ satisfying an inequality $\epsilon_0 + \epsilon_0^2 \leq \frac{c_0}{8cL^2}$, $\beta = \frac{c_0}{4L^2}$. Then it follows from (5.6) that

$$\frac{d}{dt} \iint u^2 \rho \, dx dy + \int_0^L u_x^2|_{x=0} \, dy + \alpha \iint |Du|^2 \rho \, dx dy + \alpha \beta \iint u^2 \rho \, dx dy \leq 0. \tag{5.7}$$

In particular, inequality (5.7) provides estimate (1.12) if $u_0 \in \tilde{H}_+^{1,\rho(x)}$, $u_0(0, y) \equiv 0$. In the general case $u_0 \in L_{2,+}^{\rho(x)}$ this estimate is obtained via closure with the use of Theorem 4.1.

Moreover, since inequality (5.7) can be written in a form

$$\frac{d}{dt} \left[e^{\alpha\beta t} \iint u^2 \rho \, dx dy \right] + e^{\alpha\beta t} \left[\int_0^L u_x^2|_{x=0} \, dy + \alpha \iint |Du|^2 \rho \, dx dy \right] \leq 0,$$

we find (again if $u_0 \in \tilde{H}_+^{1,\rho(x)}$, $u_0(0, y) \equiv 0$) that

$$\int_0^t e^{\alpha\beta\tau} \left[\int_0^L u_x^2|_{x=0} \, dy + \alpha \iint |Du|^2 \rho \, dx dy \right] d\tau \leq \|u_0\|_{L_{2,+}^{\rho(x)}}^2. \tag{5.8}$$

Summing equality (2.49) (for $f_1 \equiv -uu_x$) and equality (2.51) (for $f \equiv -uu_x$), differentiating, multiplying by $e^{\alpha\beta t}$ and again integrating with respect to t , we derive the following inequality:

$$\begin{aligned} & e^{\alpha\beta t} \iint (u_x^2 + u_y^2 - \frac{1}{3}u^3) \rho \, dx dy + 2\alpha \int_0^t e^{\alpha\beta\tau} \iint (3u_{xx}^2 + 4u_{xy}^2 + u_{yy}^2) \rho \, dx dy d\tau \\ & \leq \alpha(2b + \beta + 8\alpha^2) \int_0^t e^{\alpha\beta\tau} \iint (u_x^2 + u_y^2) \rho \, dx dy d\tau + (8\alpha^2 - b) \int_0^t e^{\alpha\beta\tau} \int_0^L u_x^2|_{x=0} \, dy d\tau \\ & + 4\alpha \int_0^t e^{\alpha\beta\tau} \iint uu_x^2 \rho \, dx dy d\tau - 2\alpha \int_0^t e^{\alpha\beta\tau} \iint u^2(u_{xx} + u_{yy}) \rho \, dx dy d\tau \\ & - \frac{\alpha}{3}(\beta + 2b) \int_0^t e^{\alpha\beta\tau} \iint u^3 \rho \, dx dy d\tau. \end{aligned} \tag{5.9}$$

Estimating the integrals in the right side of (5.9) with the help of (1.12) and (5.8) yields:

$$e^{\alpha\beta t} \iint (u_x^2 + u_y^2 - \frac{1}{3}u^3) \rho \, dx dy \leq c,$$

where

$$\frac{1}{3} \iint u^3 \rho \, dx dy \leq c \left[\left(\iint |Du|^2 \rho \, dx dy \right)^{1/2} \iint u^2 \rho \, dx dy + \left(\iint u^2 \rho \, dx dy \right)^{3/2} \right],$$

and (1.13) follows. \square

Conflict of interest statement

No conflict of interest.

References

- [1] E.S. Baykova, A.V. Faminskii, On initial-boundary-value problems in a strip for the generalized two-dimensional Zakharov–Kuznetsov equation, *Adv. Differ. Equ.* 18 (2013) 663–686.
- [2] H.A. Biagioni, F. Linares, Well-posedness for the modified Zakharov–Kuznetsov equation, *Prog. Nonlinear Differ. Equ. Appl.* 54 (2003) 181–189.
- [3] E. Bustamante, J. Jimenez, J. Mejia, The Zakharov–Kuznetsov equation in weighted Sobolev spaces, *J. Math. Anal. Appl.* 433 (2016) 149–175.
- [4] J.L. Bona, S. Sun, B.-Y. Zhang, The initial-boundary-value problem for the KdV equation on a quarter plane, *Trans. Am. Math. Soc.* 354 (2001) 427–490.
- [5] G.G. Doronin, N.A. Larkin, Stabilization of regular solutions for the Zakharov–Kuznetsov equation posed on bounded rectangles and on a strip, *Proc. Edinb. Math. Soc.* 58 (2015) 661–682.
- [6] A.V. Faminskii, The Cauchy problem for quasilinear equations of odd order, *Mat. Sb.* 180 (1989) 1183–1210; English transl. in *Math. USSR Sb.* 68 (1991) 31–59.
- [7] A.V. Faminski, The Cauchy problem for the Zakharov–Kuznetsov equation, *Differ. Uravn.* 31 (1995) 1070–1081; English transl. in *Differ. Equ.* 31 (1995) 1002–1012.
- [8] A.V. Faminskii, On the mixed problem for quasilinear equations of the third order, *J. Math. Sci.* 110 (2002) 2476–2507.
- [9] A.V. Faminskii, On the nonlocal well-posedness of a mixed problem for the Zakharov–Kuznetsov equation, *J. Math. Sci.* 147 (2007) 6524–6537.
- [10] A.V. Faminskii, Well-posed initial-boundary value problems for the Zakharov–Kuznetsov equation, *Electron. J. Differ. Equ.* 127 (2008) 1–23.
- [11] A.V. Faminskii, I.Yu. Bashlykova, Weak solutions to one initial-boundary value problem with three boundary conditions for quasilinear equations of the third order, *Ukr. Math. Bull.* 5 (2008) 83–98.
- [12] A.V. Faminskii, Weak solutions to initial-boundary-value problems for quasilinear evolution equations of an odd order, *Adv. Differ. Equ.* 17 (2012) 421–470.
- [13] A.V. Faminskii, An initial-boundary value problem in a strip for two-dimensional Zakharov–Kuznetsov–Burgers equation, *Nonlinear Anal.* 116 (2015) 132–144.
- [14] A.V. Faminskii, An initial-boundary value problem in a strip for two-dimensional equations of Zakharov–Kuznetsov type, *Contemp. Math.* 653 (2015) 137–162.
- [15] L.G. Farah, F. Linares, A. Pastor, A note on the 2D generalized Zakharov–Kuznetsov equation: local, global and scattering results, *J. Differ. Equ.* 253 (2012) 2558–2571.
- [16] G. Fonseca, M. Panchón, Well-posedness for the two dimensional generalized Zakharov–Kuznetsov equation in anisotropic weighted Sobolev spaces, *J. Math. Anal. Appl.* 443 (2016) 566–584.
- [17] A. Grünrock, On the generalized Zakharov–Kuznetsov equation at critical regularity, arXiv:1509.09146v1 [math.AP], 30 Sep. 2015.
- [18] A. Grünrock, S. Herr, The Fourier restriction norm method for the Zakharov–Kuznetsov equation, *Discrete Contin. Dyn. Syst., Ser. A* 34 (2014) 2061–2068.
- [19] D. Han-Kwan, From Vlasov–Poisson to Korteweg–de Vries and Zakharov–Kuznetsov, *Commun. Math. Phys.* 324 (2013) 961–993.
- [20] O.A. Ladyzhenskaya, V.A. Solonnikov, N.N. Uraltseva, *Linear and Quasilinear Equations of Parabolic Type*, *Trans. of Math. Monogr.*, vol. 23, American Math. Soc., Providence, RI, 1968.
- [21] D. Lannes, F. Linares, J.-C. Saut, The Cauchy Problem for the Euler–Poisson System and Derivation of the Zakharov–Kuznetsov Equation, *Progress Nonlinear Differential Equ. Appl.*, vol. 84, 2013, pp. 183–215.
- [22] N.A. Larkin, Exponential decay of the H^1 -norm for the 2D Zakharov–Kuznetsov equation, *J. Math. Anal. Appl.* 405 (2013) 326–335.
- [23] N.A. Larkin, The 2D Zakharov–Kuznetsov–Burgers equation with variable dissipation on a strip, *Electron. J. Differ. Equ.* (60) (2015) 1–20.
- [24] N.A. Larkin, The 2D Zakharov–Kuznetsov–Burgers equation on a strip, *Bol. Soc. Parana. Mat.* (3) 34 (2016) 151–172.
- [25] N.A. Larkin, E. Tronco, Regular solutions of the 2D Zakharov–Kuznetsov equation on a half-strip, *J. Differ. Equ.* 254 (2013) 81–101.
- [26] F. Linares, A. Pastor, Well-posedness for the two-dimensional modified Zakharov–Kuznetsov equation, *SIAM J. Math. Anal.* 41 (2009) 1323–1339.
- [27] F. Linares, A. Pastor, Well-posedness for the 2D modified Zakharov–Kuznetsov equation, *J. Funct. Anal.* 260 (2011) 1060–1085.
- [28] F. Linares, A. Pastor, J.-C. Saut, Well-posedness for the Zakharov–Kuznetsov equation in a cylinder and on the background of a KdV soliton, *Commun. Partial Differ. Equ.* 35 (2010) 1674–1689.
- [29] J.-L. Lions, E. Magenes, *Problèmes aux limites non homogènes et applications*, Dunod, Paris, 1968.
- [30] L. Molinet, D. Pilod, Bilinear Strichartz estimates for the Zakharov–Kuznetsov equation and applications, *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* 32 (2015) 347–371.
- [31] F. Ribaud, S. Vento, A note on the Cauchy problem for the 2D generalized Zakharov–Kuznetsov equation, *C. R. Acad. Sci. Paris* 350 (2012) 499–503.
- [32] J.-C. Saut, Sur quelques generalizations de l’équation de Korteweg–de Vries, *J. Math. Pures Appl.* 58 (1979) 21–61.

- [33] J.-C. Saut, R. Temam, An initial boundary value problem for the Zakharov–Kuznetsov equation, *Adv. Differ. Equ.* 15 (2010) 1001–1031.
- [34] J.-C. Saut, R. Temam, C. Wang, An initial and boundary-value problem for the Zakharov–Kuznetsov equation in a bounded domain, *J. Math. Phys.* 53 (2012) 115612.
- [35] V.E. Zakharov, E.A. Kuznetsov, On three-dimensional solutions, *Zh. Èksp. Teor. Fiz.* 66 (1974) 594–597; English transl. in *Sov. Phys. JETP* 39 (1974) 285–288.