

MONODROMY OF A FAMILY OF HYPERSURFACES CONTAINING A GIVEN SUBVARIETY

BY ANIA OTWINOWSKA AND MORIHIKO SAITO

ABSTRACT. – For a subvariety of a smooth projective variety, consider the family of smooth hypersurfaces of sufficiently large degree containing it, and take the quotient of the middle cohomology of the hypersurfaces by the cohomology of the ambient variety and also by the cycle classes of the irreducible components of the subvariety. Using Deligne’s semisimplicity theorem together with Steenbrink’s theory for semistable degenerations, we give a simpler proof of the first author’s theorem (with a better bound of the degree of hypersurfaces) that this monodromy representation is irreducible.

© 2005 Published by Elsevier SAS

RÉSUMÉ. – Soit Z une sous-variété d’une variété projective lisse Y . Nous considérons la famille des hypersurfaces lisses X de Y contenant Z et de degré suffisamment grand. Nous étudions le quotient de la cohomologie de X par l’image de la cohomologie de Y et par les classes des composantes irréductibles de Z . Nous donnons alors une démonstration plus simple (et avec une meilleure borne sur le degré des hypersurfaces) d’un théorème du premier auteur qui affirme que la représentation de monodromie sur le quotient est irréductible. Nous utilisons le théorème de semi-simplicité de Deligne et la théorie de Steenbrink des dégénérescences semi-stables.

© 2005 Published by Elsevier SAS

Introduction

It is well known after Noether and Lefschetz that, for a *general* smooth hypersurface X in $\mathbb{P}_{\mathbb{C}}^3$, the subspace of Hodge cycles in the middle cohomology $H^2(X, \mathbb{Q})$ is generated by that of $\mathbb{P}_{\mathbb{C}}^3$ (i.e. it is 1-dimensional) if the degree of the hypersurface is at least 4. This follows from the irreducibility of the monodromy representation on the primitive cohomology of the hypersurfaces. Sometimes we want to consider a family of hypersurfaces containing a given closed subvariety Z , and ask if an analogue of the above assertion holds.

More generally, let Y be an irreducible smooth complex projective variety embedded in a projective space, and Z be a (possibly reducible) closed subvariety of Y . Let $Z_{\{j\}} = \{x \in Z : \dim T_x Z = j\}$, where $T_x Z$ denotes the Zariski tangent space. In this paper we assume

$$(0.1) \quad \dim Z_{\{j\}} + j \leq \dim Y - 1 \quad \text{for any } j \leq \dim Y.$$

In particular, $2 \dim Z \leq \dim Y - 1$. Let \mathcal{I}_Z be the ideal sheaf of Z , and δ be a positive integer such that $\mathcal{I}_Z(\delta) := \mathcal{I}_Z \otimes \mathcal{O}_Y(\delta)$ is generated by global sections, where $\mathcal{O}_Y(i)$ denotes the restriction of $\mathcal{O}(i)$ on the ambient projective space. S. Kleiman and A. Altman [13] then proved that condition (0.1) implies the existence of a smooth hypersurface section X of degree d of Y

containing Z for any $d > \delta$ (where their condition for δ is slightly different from ours). Actually, condition (0.1) is also a necessary condition, see [19] (or 1.4 below).

Let $m = \dim X$, and $H^m(X, \mathbb{Q})^{\text{van}}$ denote the orthogonal complement of the injective image of $H^m(Y, \mathbb{Q})$ in $H^m(X, \mathbb{Q})$, which is called the vanishing part or the vanishing cohomology, because it is generated by the vanishing cycles of a Lefschetz pencil using the Picard–Lefschetz formula. Let $H^m(X, \mathbb{Q})_Z^{\text{van}}$ denote the subspace of $H^m(X, \mathbb{Q})^{\text{van}}$ generated by the cycle classes of the maximal dimensional irreducible components of Z modulo the image of $H^m(Y, \mathbb{Q})$ (using the orthogonal decomposition) if $m = 2 \dim Z$, and $H^m(X, \mathbb{Q})_Z^{\text{van}} = 0$ otherwise. Let $H^m(X, \mathbb{Q})_{\perp Z}^{\text{van}}$ be its orthogonal complement in $H^m(X, \mathbb{Q})^{\text{van}}$. The first author [19] made the following.

0.2. CONJECTURE. – Assume $\deg X \geq \delta + 1$. Then the monodromy representation on $H^m(X, \mathbb{Q})_{\perp Z}^{\text{van}}$ for the family of hypersurface sections X containing Z as above is irreducible.

In the case Z is smooth, an easy proof was given by C. Voisin, see 1.6. In general, using a degeneration argument inspired by [10,16], the first author proved the following.

0.3. THEOREM [19]. – *There exists a positive real number C such that Conjecture 0.2 holds if $\deg X \geq C(\delta + 1)$.*

This was used in an essential way for the proof of the main theorem in [20], which implies the Hodge conjecture for a hypersurface section of sufficiently large degree belonging to some open subset of an irreducible component of the Noether–Lefschetz locus of low codimension, and whose argument generalizes that of [18]. In the proof of Theorem 0.3, however, an asymptotic argument is used, and C can be quite large. In this paper, we give a simpler proof of the theorem below by using the theory of nearby cycles [5,24] together with Hodge theory [2,4,22,25,29] and some cohomological properties of Lefschetz pencils [6,12,14,15,28].

0.4. THEOREM. – *Let δ be as above, and d be a positive integer. Assume either $d \geq 2 + \delta$ or a general hypersurface section of degree $d - \delta$ of Y has a nontrivial differential form of the highest degree. Then Conjecture 0.2 holds with the assumption replaced by $\deg X = d$.*

The first assumption $d \geq \delta + 2$ may be replaced with a weaker condition that the 2-jets at each point are generated by the global sections of $\mathcal{O}_Y(d - \delta)$. The second assumption on a differential form is stable by hypersurface sections, and it is satisfied for any $d \geq \delta + 1$ if Y has a nontrivial differential form of the highest degree. The proof of Theorem 0.4 uses Deligne’s semisimplicity theorem [4] which implies that irreducibility is equivalent to indecomposability. We take a semistable degeneration as in [10], and calculate the graded pieces of the weight filtration on the nearby cycles using [24]. Then we can proceed by induction on m showing the nontriviality of certain extension classes by calculating the cohomology of a Lefschetz pencil or a non-Lefschetz fibration. In the non-Lefschetz case, we use a special kind of degeneration generalizing a construction in the surface case in [19] (see 2.7 below) if the condition $d \geq \delta + 2$ is satisfied, and use Hodge theory (see 3.2, 3.3 below) if the second assumption on a differential form is satisfied. In the Lefschetz case, the argument is rather easy, see 2.3, 2.5 below. (We can generalize Theorem 0.4 by replacing $\mathcal{O}_Y(\delta)$, $\mathcal{O}_Y(d - \delta)$ with two ample line bundles \mathcal{L}_1 , \mathcal{L}_2 satisfying appropriate conditions even if they are linearly independent in the Picard group of Y tensored with \mathbb{Q} , see 4.8 below.)

By a standard argument, Theorem 0.4 implies

0.5. COROLLARY. – *Under the assumption of Theorem 0.4, assume further that $m = 2 \dim Z$, and the vanishing cohomology of a general hypersurface section of degree d does not have a Hodge structure of type $(m/2, m/2)$. Then the Hodge cycles in the middle cohomology of*

a general hypersurface section X of degree d of Y containing Z are generated by the image of the Hodge cycles on Y together with the cycle classes of the irreducible components of Z . In particular, the Hodge conjecture for X is reduced to that for Y .

In Section 1, we review the theory of hypersurface sections containing a subvariety (this is mainly a reproduction from [19]). In Section 2, we prove the nonvanishing of some extension classes using a topological method, and in Section 3, we do it using a Hodge-theoretic method. In Section 4, we study the graded pieces of the weight filtration on the nearby cycles, and prove Theorem 0.4.

In this paper, a variety means a (not necessarily reduced nor irreducible) separated scheme of finite type over \mathbb{C} , and a point of a variety means a closed point. We say that a member of a family parametrized by the points of a variety is general if the point corresponding to the member belongs to some dense open subvariety of the variety.

1. Hypersurfaces containing a subvariety

In this section, we review some basic facts in the theory of hypersurface sections containing a subvariety, which will be needed in the proof of Theorem 0.4. We mainly reproduce the arguments in [19, Section 1] see also [13]. Our argument works only in the case of characteristic 0.

1.1. Hypersurface sections

Let Y be an $(m + 1)$ -dimensional irreducible smooth projective variety embedded in a projective space \mathcal{P} . For a positive integer i , let $\mathcal{O}_Y(i)$ be the restriction to Y of $\mathcal{O}_{\mathcal{P}}(i)$ on the projective space \mathcal{P} , and define

$$\mathcal{A}^i = \Gamma(Y, \mathcal{O}_Y(i)).$$

Note that the restriction morphism

$$\Gamma(\mathcal{P}, \mathcal{O}_{\mathcal{P}}(i)) \rightarrow \Gamma(Y, \mathcal{O}_Y(i))$$

is not surjective in general (e.g. if Y is a hypersurface section of degree i of an Abelian variety contained in \mathcal{P}). For $P \in \mathcal{A}^i$, we denote by X_P the associated hypersurface section of degree i (in a generalized sense unless P belongs to the image of the above morphism). Let

$$\mathcal{V}^i = \{P \in \mathcal{A}^i \setminus \{0\} : X_P \text{ is smooth}\}.$$

It is identified with a smooth variety, and $\mathcal{V}^i/\mathbb{C}^*$ parametrizes the smooth hypersurface sections of Y of degree i .

Let Z be a closed subvariety of Y , and put

$$\mathcal{A}_Z^i = \{P \in \mathcal{A}^i : Z \subset X_P\}, \quad \mathcal{V}_Z^i = \mathcal{A}_Z^i \cap \mathcal{V}^i.$$

In this section we do not necessarily assume condition (0.1) in Introduction. Let δ be a positive integer such that $\mathcal{I}_Z(\delta)$ is generated by \mathcal{A}_Z^δ . It is shown by Kleiman and Altman [13] that condition (0.1) in Introduction implies the existence of a smooth hypersurface of degree j containing Z for any $j > \delta'$, where the definition of δ' in loc. cit. uses the ideal of Z in the

projective space \mathcal{P} instead of \mathcal{I}_Z (in particular $\delta \leq \delta'$, and we may have a strict inequality if the degrees of the defining equations of Y in \mathcal{P} are bigger than those for Z in Y). More precisely, they showed the following theorem for $d_i \geq \delta' + 1$ and $V_i = \mathcal{A}_Z^{d_i}$, except possibly for the estimate of $\dim \text{Sing } X_{\mathbf{P}}$, which we need later. Concerning the connectedness, we will need it for the proof of Theorem 0.4 only in the cases where $a \geq 0$ and either $r = 1$ or 2 . Note that the argument is simpler in these cases because X_{P_1} is smooth.

In Theorem 1.2 below, we will take vector subspaces V_i of $\mathcal{A}_Z^{d_i}$, and consider the following condition:

- (C_i) $d_i > \delta$ and there exist a vector subspace V'_i of \mathcal{A}_Z^δ generating $\mathcal{O}_Y(\delta)$ outside Z and a vector subspace V''_i of $\mathcal{A}^{d_i-\delta}$ giving an embedding of Y into a projective space such that V_i is the image of $V'_i \otimes V''_i$.

1.2. THEOREM [13,19]. – *Let $a = \dim Y - 1 - \max\{\dim Z_{\{j\}} + j\}$ so that condition (0.1) is satisfied if and only if $a \geq 0$. Let $d_1 \leq \dots \leq d_r$ be integers with $r \leq c_Z := \text{codim}_Y Z$. Assume $d_i \geq \delta$, and let V_i be a vector subspace of $\mathcal{A}_Z^{d_i}$ generating $\mathcal{I}_Z \otimes \mathcal{O}_Y(d_i)$. Take a general $\mathbf{P} = (P_1, \dots, P_r)$ in $V_1 \times \dots \times V_r$, and put $X_{\mathbf{P}} = \bigcap_{1 \leq i \leq r} X_{P_i}$. Then $X_{\mathbf{P}}$ is a complete intersection, $\dim \text{Sing } X_{\mathbf{P}} \leq r - a - 2$, and $X_{\mathbf{P}} \setminus Z$ is smooth.*

Assume furthermore that $\dim X_{\mathbf{P}} \geq 1$, the above condition (C_i) is satisfied for any integer i in $[c_Z, r]$ in the case where $\dim Y \leq 2 \dim Z + a + 2$ or Z is reduced on the complement of a closed subvariety of dimension $< \dim Z$, and (C_i) is satisfied for any integer i in $[c_Z - 1, r]$ otherwise. Then $X_{\mathbf{P}} \setminus Z$ is connected. Here $[c_Z, r] = \emptyset$ if $r < c_Z$.

Proof. – Let $\Pi_r = V_1 \times \dots \times V_r$, and

$$\Sigma_r = \{(\mathbf{P}, x) \in \Pi_r \times Z : \text{rank}(dP_1, \dots, dP_r)_x < r\},$$

where the dP_i are defined by taking a local trivialization of $\mathcal{O}(1)$, and the rank is independent of the trivialization because the X_{P_i} contain Z . We have

$$T_x^* Z = m_{Y,x} / (\mathcal{I}_{Z,x} + m_{Y,x}^2),$$

(where $m_{Y,x}$ denotes the maximal ideal), and the $P \in V_i$ generate $(\mathcal{I}_{Z,x} + m_{Y,x}^2) / m_{Y,x}^2$ for any i by the definition of V_i (taking a local trivialization of $\mathcal{O}(1)$). Let

$$\Sigma_{r,x} = \Sigma_r \cap (\Pi_r \times \{x\}).$$

By considering the fiber of the projection $\Pi_r \rightarrow \Pi_{r-1}$ and by induction on r we get for any $x \in Z_{\{\dim Y - j\}}$,

$$\text{codim}_{\Pi_r} \Sigma_{r,x} = \max(j - r + 1, 0) \geq j - r + 1.$$

By definition of a we have $\text{codim}_Y Z_{\{\dim Y - j\}} \geq \dim Y - j + a + 1$, hence

$$\text{codim}_{\Pi_r \times Y} \Sigma_r \geq \dim Y - r + a + 2.$$

For a general $\mathbf{P} \in \Pi_r$, this implies

$$\text{codim}_Y \Sigma_r \cap (\{\mathbf{P}\} \times Z) \geq \dim Y - r + a + 2.$$

So it remains to show the assertion on the smoothness and the connectedness.

For the smoothness, we see that a subvariety of $(Y \setminus Z) \times \Pi_r$ defined by the relation $x \in X_{\mathbf{P}}$ for $(x, \mathbf{P}) \in (Y \setminus Z) \times \Pi_r$ is smooth. (Indeed, in the case $r = 1$, the variety is defined by $\sum_i t_i P_i$

where (P_i) is a basis of V_1 as a vector space, and (t_i) is the corresponding coordinate system of V_1 . Furthermore, for any $x \in Y \setminus Z$, some P_i does not vanish on a neighborhood of x , where the above equation can be divided by this P_i . The argument is similar for $r > 1$.) Then the smoothness follows from the Bertini theorem in characteristic 0.

For the connectedness, we proceed by induction on r . Let $Y' = X_{\mathbf{P}'}$ for a general $\mathbf{P}' \in \Pi_{r-1}$, where $Y' = Y$ if $r = 1$. Set $b = 0$ if $\dim Y \leq 2 \dim Z + a + 2$ or Z is reduced on the complement of a closed subvariety of dimension $< \dim Z$, and $b = 1$ otherwise. Assume first $r < \text{codim}_Y Z - b$. Since $Y' \setminus X_{\mathbf{P}'}$ is a smooth affine variety of dimension ≥ 2 , its first cohomology with compact supports vanishes by the weak Lefschetz theorem together with Poincaré duality. So $X_{\mathbf{P}'}$ is connected. By Hartshorne's connectedness theorem (see 1.3 below) the connectedness of $X_{\mathbf{P}'} \setminus Z$ is then reduced to that $\text{codim}_{X_{\mathbf{P}'}} \text{Sing } X_{\mathbf{P}'} > 1$.

Since $\text{Sing } X_{\mathbf{P}'} \subset Z$, the last condition is trivially satisfied in the case $r < \text{codim}_Y Z - 1$. If $r = \text{codim}_Y Z - 1$ and Z is generically reduced as above so that $b = 0$, then $X_{\mathbf{P}'}$ is smooth at a general point of Z because each V_i generates $\mathcal{I}_Z \otimes \mathcal{O}_Y(d_i)$. Thus the assertion is proved. Similarly, if $r = \text{codim}_Y Z - 1$ and $\dim Y \leq 2 \dim Z + a + 2$, then $\dim \text{Sing } X_{\mathbf{P}'} \leq r - a - 2 < \dim Z$ and the assertion follows.

Assume now $r \geq \text{codim}_Y Z - b$. Consider a rational map of Y' to a projective space defined by the restriction of the linear system V_r to Y' . It induces an embedding of $Y' \setminus Z$ because $V_r'|_{Y'}$ generates $\mathcal{O}_{Y'}(\delta)$ outside Z and $V_r''|_{Y'}$ induces an embedding. So $X_{\mathbf{P}'} \setminus Z$ is isomorphic to a general hyperplane section of $Y' \setminus Z$ for this embedding. Since $Y' \setminus Z$ is a smooth connected variety of dimension at least 2, the Bertini theorem in characteristic 0 implies that $X_{\mathbf{P}'} \setminus Z$ is connected. This completes the proof of Theorem 1.2. \square

1.3. Complement to the proof of Theorem 1.2

A local complete intersection is irreducible if it is connected and the singular locus has codimension > 1 . This follows from Hartshorne's connectedness theorem (see, e.g., [9, Theorem 18.12]), because a local complete intersection is Cohen–Macaulay. It also follows from the theory of perverse sheaves [1], considering the long exact sequence of perverse cohomology sheaves associated to the distinguished triangle $\mathbb{Q}_{X_0} \rightarrow \mathbb{Q}_{X_1} \oplus \mathbb{Q}_{X_2} \rightarrow \mathbb{Q}_{X_3} \rightarrow$ where $X_0 = X_1 \cup X_2$, $X_3 = X_1 \cap X_2$. Indeed, if X_0 is a local complete intersection of dimension n and X_3 has dimension $\leq n - 2$, then $\mathbb{Q}_{X_0}[n]$ is a perverse sheaf (i.e. ${}^p\mathcal{H}^n \mathbb{Q}_{X_0} = \mathbb{Q}_{X_0}[n]$, see, e.g., [7, Theorem 5.1.19]) and ${}^p\mathcal{H}^j \mathbb{Q}_{X_3} = 0$ for $j \geq n - 1$, see [1]. So $\mathbb{Q}_{X_0}[n]$ is the direct sum of ${}^p\mathcal{H}^n \mathbb{Q}_{X_i}$ for $i = 1, 2$, and it is a contradiction if X_0 is connected.

The following gives a converse of [13] (and we reproduce the arguments here for the convenience of the reader).

1.4. THEOREM [19]. – *Assume $d \geq \delta + 1$. If there exists a smooth hypersurface section of degree d containing Z , then condition (0.1) in Introduction is satisfied.*

Proof. – Let U be a nonempty smooth open subvariety of $Z_{\{\dim Y - j\}}$ such that $\dim U = \dim Z_{\{\dim Y - j\}}$ and $\mathcal{O}(1)$ is trivialized over U . Let

$$\mathcal{E} = \text{Ker} \left(T^*Y|_U \rightarrow \bigcup_{x \in U} T_x^*Z \right).$$

It is a vector bundle of rank j over U , and the associated coherent sheaf $\mathcal{O}_U(\mathcal{E})$ is generated by dP for $P \in \mathcal{A}_Z^\delta$ (where we fix a trivialization of $\mathcal{O}(1)$ over U). So we may assume that \mathcal{E} is trivialized by $P_1, \dots, P_j \in \mathcal{A}_Z^\delta$.

Since $d \geq \delta + 1$, we see that the dP for $P \in \mathcal{A}_Z^d$ generate the 1-jets of $\mathcal{O}_U(\mathcal{E})$, i.e. they generate $\mathcal{O}_U(\mathcal{E})_x \otimes \mathcal{O}_{U,x}/m_{U,x}^2$ over \mathbb{C} for any $x \in U$ (because there are smooth hypersurface sections of degree $d - \delta$ which give local coordinates of U at x). \square

If the assumption of 1.4 is satisfied, then dP for a general $P \in \mathcal{V}_Z^d$ gives a nowhere vanishing section of \mathcal{E} over U because X_P is smooth. Since condition (0.1) is equivalent to $\dim Z_{\{\dim Y - j\}} < j$ for any j , the assertion is reduced to the following:

1.5. LEMMA. – *Let X be a smooth variety of dimension $\geq j$, and \mathcal{E} be a trivial vector bundle of rank j over X . Let V be a finite dimensional vector subspace of $\Gamma(X, \mathcal{E})$ which generates the 1-jets at every point of X . Then for a general $\sigma \in V$, the zero locus $\sigma^{-1}(0) \subset X$ is nonempty.*

Proof. – Since \mathcal{E} is trivial, we may identify $\sigma \in \Gamma(X, \mathcal{E})$ with a morphism of X to a vector space E . We have a natural morphism $\theta: V \times X \rightarrow E$ which sends (σ, x) to $\sigma(x)$. Let $Y = \theta^{-1}(0)$. Then it is enough to show that Y is dominant over V .

Since V generates the 1-jets and $\dim X \geq j$, there exists $y = (\sigma, x) \in Y$ such that the differential $d_x \sigma: T_x X \rightarrow T_{\sigma(x)} E = E$ of $\sigma: X \rightarrow E$ (by the above identification) is surjective. Consider the commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & T_x X & \longrightarrow & T_y(V \times X) & \longrightarrow & T_\sigma V \longrightarrow 0 \\
 & & \downarrow d_x \sigma & & \downarrow d_y \theta & & \downarrow \\
 0 & \longrightarrow & E & \longrightarrow & E & \longrightarrow & 0 \longrightarrow 0
 \end{array}$$

where the top row is induced by the inclusion $\{\sigma\} \times X \rightarrow V \times X$ and the projection $V \times X \rightarrow V$. Then the surjectivity of $d_x \sigma$ implies that of $\text{Ker } d_y \theta \rightarrow T_\sigma V$ by the snake lemma. So the assertion follows because $\text{Ker } d_y \theta = T_y Y$. This completes the proof of Theorem 1.4. \square

1.6. Voisin’s proof of Conjecture 0.2 in the smooth case

Consider a rational morphism of Y to a projective space \mathcal{P} defined by the linear system $\mathcal{A}_Z^{\delta+1}$. It induces an embedding of $Y \setminus Z$, in particular, it is birational to the image. Let \tilde{Y} be the closure of the image. If Z is smooth, we see that \tilde{Y} is the blow-up of Y along Z . For a smooth hyperplane section \tilde{Y}_s of \mathcal{P} and the corresponding smooth hypersurface section Y_s of Y , there is a morphism $H^m(\tilde{Y}_s)^{\text{van}} \rightarrow H^m(Y_s)^{\text{van}}$ by the functoriality of the Gysin morphisms, and it is injective by the irreducibility of $\{H^m(\tilde{Y}_s)^{\text{van}}\}$. Calculating the cohomology of the blow-up, we see that the dimension of its cokernel coincides with the number of the $(m/2)$ -dimensional irreducible components of Z . So Conjecture 0.2 is proved in this case.

2. Topological method

In this section, we prove the nonvanishing of certain extension classes when the condition $d \geq \delta + 2$ in Theorem 0.4 is satisfied.

2.1. Exact sequences

Let Y be a connected smooth complex algebraic variety, and X be a divisor on Y with the inclusion $i: X \rightarrow Y$. Put $U = Y \setminus X$ with the inclusion $j: U \rightarrow Y$. Let $f: Y \rightarrow S$ be a proper morphism to a smooth variety S , and let $g = f \circ i: X \rightarrow S, h = f \circ j: U \rightarrow S$ be the restrictions of f . Put $m = \dim Y - \dim S$. We have a long exact sequence of constructible sheaves

$$(2.1.1) \quad \begin{aligned} &\rightarrow R^{m-1}f_*\mathbb{Q}_Y \xrightarrow{\gamma_{m-1}} R^{m-1}g_*\mathbb{Q}_X \\ &\rightarrow R^m h_! \mathbb{Q}_U \rightarrow R^m f_* \mathbb{Q}_Y \xrightarrow{\gamma_m} R^m g_* \mathbb{Q}_X \rightarrow, \end{aligned}$$

using the distinguished triangle $j_! \mathbb{Q}_U \rightarrow \mathbb{Q}_Y \rightarrow i_* \mathbb{Q}_X \xrightarrow{+1}$ together with the functor $\mathbf{R}f_* = \mathbf{R}f_!$. Since f, g are proper, the base change holds so that the stalk of the direct image is isomorphic to the (relative) cohomology of the fiber. Let

$$\mathcal{F} = R^m h_! \mathbb{Q}_U, \quad \mathcal{F}' = \text{Coker } \gamma_{m-1}, \quad \mathcal{F}'' = \text{Ker } \gamma_m,$$

so that we have a short exact sequence of constructible sheaves

$$(2.1.2) \quad 0 \rightarrow \mathcal{F}' \rightarrow \mathcal{F} \rightarrow \mathcal{F}'' \rightarrow 0.$$

Let $X_s = g^{-1}(s)$, etc. If f, g are smooth projective and X_s is a hypersurface section of Y_s for $s \in S$, then we have

$$(2.1.3) \quad \mathcal{F}'_s = H^{m-1}(X_s, \mathbb{Q})^{\text{van}}, \quad \mathcal{F}_s = H^m(Y_s, X_s, \mathbb{Q}), \quad \mathcal{F}''_s = H^m(Y_s, \mathbb{Q})^{\text{prim}}.$$

We will assume that they are nonzero (because otherwise the extension class vanishes).

If f, g are smooth morphisms, then (2.1.1) and (2.1.2) are exact sequences of local systems, which underlie naturally variation of mixed Hodge structures, see [4,25] (and also [22]). In the application we will also consider the dual of (2.1.2)

$$0 \rightarrow \mathcal{F}''^* \rightarrow \mathcal{F}^* \rightarrow \mathcal{F}'^* \rightarrow 0,$$

where $*$ denotes the dual variation of mixed Hodge structure. Note that

$$\mathcal{F}'^* = \mathcal{F}'(m-1), \quad \mathcal{F}''^* = \mathcal{F}''(m), \quad \mathcal{F}^* = H^m(Y_s \setminus X_s, \mathbb{Q})(m),$$

where (m) denotes the Tate twist, see [4].

2.2. Lefschetz pencils

With the above notation, assume $S = \mathbb{P}^1$ and $g: X \rightarrow S$ is a Lefschetz pencil of a smooth projective variety Y_0 . Let $H^{m-1}(X_s, \mathbb{Q})^{\text{van}}$ be the subgroup of $H^{m-1}(X_s, \mathbb{Q})$ generated by the vanishing cycles, and assume it nonzero. Let s_0 be any point in the discriminant of the Lefschetz pencil. Since the discriminant in the parameter space of hypersurfaces is irreducible, the last assumption is equivalent to the surjectivity of the restriction morphism

$$(2.2.1) \quad H^{m-1}(X_s, \mathbb{Q}) \rightarrow H^{m-1}(B_s, \mathbb{Q})$$

for s sufficiently near s_0 , where B is a small ball in X around the unique singular point of X_{s_0} , and $B_s := B \cap X_s$ is called a Milnor fiber, see [17]. This implies that the cospecialization morphism

$$(2.2.2) \quad H^m(X_{s_0}, \mathbb{Q}) \rightarrow H^m(X_s, \mathbb{Q})$$

is an isomorphism, using a long exact sequence. So we get

$$(2.2.3) \quad R^j g_* \mathbb{Q}_X \text{ is a constant sheaf on } S \text{ for any } j \neq m-1,$$

see [12], where the case $j = m$ follows from the above argument, and the other cases are easy. As a corollary, we get in this case

$$(2.2.4) \quad \mathcal{F}'' \text{ in (2.1.2) is a constant sheaf.}$$

The above argument also implies for $j = m - 1$ that $R^{m-1}g_*\mathbb{Q}_X$ is a (shifted) intersection complex, i.e.

$$(2.2.5) \quad R^{m-1}g_*\mathbb{Q}_X = j_*j^*R^{m-1}g_*\mathbb{Q}_X,$$

where $j: S' \rightarrow S$ is the inclusion of a dense open subvariety over which g is smooth. (This also follows from the local invariant cycle theorem [3] or the decomposition theorem [1].)

The following proposition was proved in [19] using a generalization of the Picard–Lefschetz formula together with an assertion concerning the vanishing cycles of a Lefschetz pencil and related to the classical work of Lefschetz and Poincaré (see [14,15,28]). We give here a simple proof of the proposition using the above cohomological property of the Lefschetz pencil.

2.3. PROPOSITION (Lefschetz pencil case) [19]. – *With the notation and the assumptions of 2.1, assume $S = \mathbb{P}^1$, $Y = Y_0 \times S$, $f = \text{pr}_2$, and $g: X \rightarrow S$ is a Lefschetz pencil of Y_0 . Let S' be any nonempty open subvariety of S over which g is smooth. Then for any nonzero local subsystem \mathcal{G} of $\mathcal{F}''|_{S'}$, the composition of the inclusion $\mathcal{G} \rightarrow \mathcal{F}''|_{S'}$ with the extension class defined by the restriction of the short exact sequence (2.1.2) to S' is nontrivial as an extension of local systems.*

Proof. – Since the local system $\mathcal{F}''|_{S'}$ is constant, we may assume that \mathcal{G} has rank 1, and is generated by $u \in H^0(S', \mathcal{F}''|_{S'}) = H^m(Y_0, \mathbb{Q})^{\text{prim}}$. Assume u is the image of $v \in H^0(S', \mathcal{F}|_{S'})$. Then it gives a section of (2.1.2) on \mathcal{G} . So it is enough to show that $u = 0$ in this case.

We see that \mathcal{F} is a (shifted) intersection complex by (2.1.2), because \mathcal{F}' and \mathcal{F}'' are (shifted) intersection complexes with support S . So $H^0(S, \mathcal{F}) = H^0(S', \mathcal{F}|_{S'})$, and we may replace S' with S or any nonempty open subvariety of S . Thus we may assume that $S' = S \setminus \{s_0\}$ and X_{s_0} is smooth.

Let $h': U' \rightarrow S'$ be the restriction of h over S' , where $U' = Y' \setminus X'$. Consider the Leray spectral sequence

$$(2.3.1) \quad E_2^{p,q} = H^p(S', R^q h'_! \mathbb{Q}_{U'}) \implies H^{p+q}(Y', X'; \mathbb{Q}).$$

This degenerates at E_2 , because $E_2^{p,q} = 0$ unless $p = 0$ or 1 . Thus we get $w \in H^m(Y', X'; \mathbb{Q})$ whose image in $H^0(S', \mathcal{F}|_{S'})$ is v . Its image in $H^0(S', \mathcal{F}''|_{S'}) = H^m(Y', \mathbb{Q}) = H^m(Y_0, \mathbb{Q})$ is u , where the last isomorphism follows from $Y' = Y_0 \times \mathbb{A}^1$. Then the image of u in $H^m(X', \mathbb{Q})$ vanishes. But this is induced by the restriction morphism under the birational morphism $X' \rightarrow Y_0$. So we can verify that u belongs to the image of the Gysin morphism under the inclusion $X_{s_0} \rightarrow Y_0$, and we get $u = 0$ because u is primitive. This completes the proof of Proposition 2.3. \square

We also give an outline of the original proof of Proposition 2.3. We start with the explanation of a generalized Picard–Lefschetz formula.

2.4. Generalized Picard–Lefschetz formula

Let \mathcal{F} be a constructible sheaf on a curve S with a local coordinate t . Let $\psi_t \mathcal{F}, \varphi_t \mathcal{F}$ denote the nearby and vanishing cycles, see [5]. Then we have natural morphisms

$$\text{can}: \psi_t \mathcal{F} \rightarrow \varphi_t \mathcal{F}, \quad \text{var}: \varphi_t \mathcal{F} \rightarrow \psi_t \mathcal{F},$$

such that

$$(2.4.1) \quad T - \text{id} = \text{var} \circ \text{can} : \psi_t \mathcal{F} \rightarrow \psi_t \mathcal{F},$$

where T is the monodromy. It is well known that the functors ψ_t, φ_t commute with the functor assigning the dual, and duality exchanges can and var up to a sign, see e.g. [11,21]. In the case of a Lefschetz pencil, we can identify the morphism can with the restriction to the Milnor fiber (2.2.1).

Assume that $\varphi_t \mathcal{F} \simeq \mathbb{Q}$, and can, var are nonzero. Let γ_0 be a generator of $\varphi_t \mathcal{F}$, and γ be its image in $\psi_t \mathcal{F}$ by var . Let \mathcal{F}^* be the (shifted) dual of \mathcal{F} which is defined by $\mathbf{R}\mathcal{H}om(\mathcal{F}, \mathbb{Q}_S)$. Let γ_0^* be the generator of $\varphi_t \mathcal{F}^*$ such that $\langle \gamma_0^*, \gamma_0 \rangle = 1$. Let γ^* be its image in $\psi_t \mathcal{F}^*$ by var . Then we have a generalized Picard–Lefschetz formula

$$(2.4.1) \quad T(u) - u = \pm \langle \gamma^*, u \rangle \gamma \quad \text{for } u \in \psi_t \mathcal{F},$$

because $\langle \gamma_0^*, \text{can}(u) \rangle = \pm \langle \text{var}(\gamma_0^*), u \rangle$, see also [6]. This was proved in [19] for the cohomology of the complement of a hypersurface section.

2.5. Outline of the original proof of Proposition 2.3 (see [19])

It is enough to show that $\mathcal{F}|_{S'}$ has no global sections. In this case, the stalk of $\mathcal{F}^*|_{S'}$ is $H_m(Y_s, X_s)$, and γ^* can be constructed explicitly using the ball and the Milnor fiber around the critical point (and this coincides with the construction in [14,15,28]), because we can identify the morphism can with the restriction to the complement of the Milnor fiber in the ball, see [19]. Furthermore, considering γ^* at any points of the discriminant of the Lefschetz pencil, they generate $H_m(Y_s, X_s)$. (This is closely related to the classical work of Lefschetz and Poincaré, and seems to have been known to some people, see [14] and also [15,28].) So the local system $\mathcal{F}|_{S'}$ has no global section, and the assertion follows.

2.6. Non-Lefschetz fibration case

Let Y be a connected smooth projective variety embedded in a projective space \mathcal{P} , and X be a hypersurface section of Y with at most isolated singularities. We assume $m := \dim X = \dim Y - 1 \geq 1$. Let Z be an irreducible component of X (hence $Z = X$ if $m > 1$). Let d be an integer ≥ 2 , and S be the parameter space of hypersurfaces of degree d of \mathcal{P} whose intersections Z_s, Y_s with Z, Y are smooth divisors on $Z \setminus \text{Sing } X$ and Y respectively (in particular, the hypersurfaces parametrized by S do not meet $\text{Sing } X$).

Let $\{H^{m-1}(Z_s)_{\mathbb{Z}}^{\text{van}}\}_{s \in S}$ be the local subsystem of $\{H^{m-1}(Z_s)\}_{s \in S}$ generated by the vanishing cycles at general points of the discriminant of the morphism $\bigsqcup_{s \in \bar{S}} Z_s \rightarrow \bar{S}$, where \bar{S} is the parameter space of all the hypersurfaces of degree d of \mathcal{P} , and $\bigsqcup_{s \in \bar{S}} Z_s$ denotes the total space of the associated family of hypersurfaces. If $m > 1$ and $X = Z$, let $\{H^{m-1}(Z_s)_{\mathbb{Z}}^{\text{van}}\}_{s \in S}$ be the local subsystem generated by the vanishing cycles for the inclusion $Z_s \rightarrow Y_s$. By the Picard–Lefschetz formula, the latter is the orthogonal complement of the injective image of the cohomology of Y_s (or Y using the weak Lefschetz theorem), and hence contains the former. If $X = Z$ is smooth, they coincide because they are the orthogonal complement of the injective image of $H^{m-1}(Y)$. If $m = 1$, let $H^{m-1}(Z_s)_{\mathbb{Z}}^{\text{van}} = H^{m-1}(Z_s)_{\mathbb{Z}}^{\text{van}} (= \tilde{H}^{m-1}(Z_s))$.

Let S' be a dense open subvariety of S , and \tilde{L} be any local system on S' such that

$$\{H^{m-1}(Z_s)_{\mathbb{Z}}^{\text{van}}\}_{s \in S'} \subset \tilde{L} \subset \{H^{m-1}(Z_s)_{\mathbb{Z}}^{\text{van}}\}_{s \in S'},$$

and the restriction of the intersection pairing to \tilde{L} is nondegenerate. Let \tilde{L}^\perp be the orthogonal complement of \tilde{L} in $\{H^{m-1}(Z_s)\}$. Note that the restrictions of the intersection pairing to the injective image of $H^{m-1}(Y)$ and to $H^{m-1}(Z_s)^{\text{van}}$ are nondegenerate using Hodge theory (or [1] because it is essentially equivalent to the hard Lefschetz theorem).

Consider the kernel of the composition

$$H^m(Y_s, Z_s) \rightarrow H^m(Y_s) \rightarrow H^{m+2}(Y)(1),$$

where the last morphism is the Gysin morphism. Let $H^m(Y_s, Z_s)_{\tilde{L}}^{\text{van}}$ be the quotient of the kernel by the image of \tilde{L}^\perp .

The following is a generalization of a construction in the surface case in [19], and gives a topological proof of variants of Propositions 3.2 in the non-Lefschetz case and 3.3 in the surface case.

2.7. PROPOSITION (Non-Lefschetz fibration case). – *With the above notation and assumptions, we have a short exact sequence of local systems on S'*

$$(2.7.1) \quad 0 \rightarrow \tilde{L} \rightarrow \{H^m(Y_s, Z_s)_{\tilde{L}}^{\text{van}}\} \rightarrow \{H^m(Y_s)^{\text{van}}\} \rightarrow 0,$$

and it does not split if the first and last terms are nonzero.

Proof. – The exactness of (2.7.1) is clear by definition. To show the nonsplitting of (2.7.1), we may assume $S' = S$ using the direct image by $S' \rightarrow S$. We take a smooth point O of X contained in Z , and consider a hypersurface H_0 in the ambient projective space which intersects X, Y transversely at smooth points outside O and whose intersections with Z, Y have an ordinary double point at O . (Such a hypersurface exists because the degree of the hypersurface is at least 2.) Then we consider a generic deformation $\{H_\lambda\}$ of H_0 , parametrized by $\Lambda := \Delta^{m+2}$ where Δ is a sufficiently small open disk. Using the embedding by $\mathcal{O}_Y(d)$ (which changes hypersurfaces into hyperplanes) and an appropriate projection between projective spaces, we may assume locally

$$\begin{aligned} Y &= \left\{ x_{m+2} = \sum_{1 \leq i \leq m+1} x_i^2 + \Psi(x_1, \dots, x_{m+1}) \right\}, \\ Z &= \{x_{m+1} = 0\} \subset Y, \\ H_\lambda &= \left\{ x_{m+2} = \sum_{1 \leq i \leq m+1} 2a_i x_i + a_{m+2} \right\} \quad \text{for } \lambda = (a_i) \in \Delta^{m+2}, \end{aligned}$$

in a $(m + 2)$ -dimensional projective space, where (x_i) is an affine coordinate system and $\Psi(x_1, \dots, x_{m+1}) \in (x_1, \dots, x_{m+1})^3$.

Let $Z_\lambda = Z \cap H_\lambda, Y_\lambda = Y \cap H_\lambda$. The discriminant of $\{Z_\lambda\}_{\lambda \in \Lambda}$ and $\{Y_\lambda\}_{\lambda \in \Lambda}$ in Λ are defined respectively by

$$\begin{aligned} D_Z &= \left\{ a_{m+2} + \sum_{1 \leq i \leq m} a_i^2 + \Phi_Z(a_1, \dots, a_m) = 0 \right\}, \\ D_Y &= \left\{ a_{m+2} + \sum_{1 \leq i \leq m+1} a_i^2 + \Phi_Y(a_1, \dots, a_{m+1}) = 0 \right\}, \end{aligned}$$

where $\Phi_Z(a_1, \dots, a_m) \in (a_1, \dots, a_m)^3, \Phi_Y(a_1, \dots, a_{m+1}) \in (a_1, \dots, a_{m+1})^3$.

Take $c \in \Delta \setminus \{0\}$, and put $0' = (0) \in \Delta^m$. Define

$$\Lambda_c = \{0'\} \times \{c\} \times \Delta, \quad \{\alpha\} = \Lambda_c \cap D_Z, \quad \{\beta\} = \Lambda_c \cap D_Y,$$

$$\Lambda'_c = \Lambda_c \setminus \{\alpha, \beta\}.$$

Let B be a sufficiently small open ball around O in the ambient space, and assume Δ is much smaller. Let

$$Z_B = Z \cap B, \quad Z_{B,\lambda} = Z_\lambda \cap B \quad (\text{similarly for } Y).$$

For $\lambda \in \Lambda_c \setminus \{\beta\}$, it is well known that

$$(2.7.2) \quad \begin{aligned} \tilde{H}^j(Y_{B,\lambda}) &= \mathbb{Q} \quad \text{for } j = m, \text{ and } 0 \text{ otherwise,} \\ \tilde{H}^j(Y_{B,\beta}) &= 0 \quad \text{for any } j \end{aligned}$$

and similarly for $Z_{B,\lambda}$ with β replaced by α , and m by $m - 1$.

The local monodromy of $H^{m-1}(Z_{B,\lambda})$ (respectively $H^m(Y_{B,\lambda})$) around α (respectively β) is $(-1)^m$ (respectively $(-1)^{m+1}$). By (2.7.2) we have the short exact sequences

$$(2.7.3) \quad 0 \rightarrow H^{m-1}(Z_{B,\lambda}) \rightarrow H^m(Y_{B,\lambda}, Z_{B,\lambda}) \rightarrow H^m(Y_{B,\lambda}) \rightarrow 0,$$

$$(2.7.4) \quad 0 \rightarrow H^{m-1}(Z_{B,\lambda}) \rightarrow H^m_c(Y_{B,\lambda} \setminus Z_{B,\lambda}) \rightarrow H^m_c(Y_{B,\lambda}) \rightarrow 0.$$

By definition, we have an injective morphism of (2.7.4) to (2.7.1). Taking the dual, we get a surjective morphism of (2.7.1) to (2.7.3) because the intersection form is nondegenerate on \tilde{L} . Let e, e' denote the extension classes defined by (2.7.3) and (2.7.4) respectively. Then the assertion is reduced to Lemma 2.8 below. Indeed, if m is odd and the composition of e with the surjection

$$\{H^m(Y_\lambda)^{\text{van}}\}_{\lambda \in \Lambda'_c} \rightarrow \{H^m(Y_{B,\lambda})\}_{\lambda \in \Lambda'_c}$$

vanishes, then $\{H^m(Y_{B,\lambda}, Z_{B,\lambda})\}$ is isomorphic to a subquotient of $\{H^m(Y_\lambda)^{\text{van}}\}$ (by the nontriviality of e) and there is a surjection from the kernel of the above morphism to $\{H^{m-1}(Z_{B,\lambda})\}$, but this contradicts the nontriviality of the monodromy of $\{H^{m-1}(Z_{B,\lambda})\}$ around α . The argument is similar for m even. \square

2.8. LEMMA. – *The extension classes e, e' defined by (2.7.3), (2.7.4) do not vanish if m is odd or even respectively.*

Proof. – We first show the nonsplitting of (2.7.3) for m odd. In this case, the monodromy of $\{H^m(Y_{B,\lambda}, Z_{B,\lambda})\}$ around β is unipotent, and we have to show that it is not the identity. Let $D \subset \Delta_c$ be a sufficiently small open disk with center β , and restrict the local systems to $D^* := D \setminus \{\beta\}$. The short exact sequence of local systems (2.7.3) on D^* is naturally extended to D as constructible sheaves, by using the higher direct images as in 2.1 instead of cohomology.

More precisely, let $Z_{B,D}, Y_{B,D}$ be the restriction of Z_B, Y_B over D . Let $U = Y_{B,D} \setminus Z_{B,D}$ with the inclusion $j_U : U \rightarrow Y_{B,D}$. Then the short exact sequence of local systems on D^* defined by (2.7.3) is naturally extended to a short exact sequence of constructible sheaves on D

$$(2.8.1) \quad 0 \rightarrow \mathcal{G}' \rightarrow \mathcal{G} \rightarrow \mathcal{G}'' \rightarrow 0.$$

This is also defined by the higher direct images of $(j_U)_! \mathbb{Q}_U$, etc. under $Y_{B,D} \rightarrow D$. Here the base change holds, because (Y_λ, Z_λ) is transversal to the boundary of B . In particular, the stalk at β is given by (2.7.3).

Using (2.7.2), we see that $\mathcal{G}' = \mathbb{Q}_D, \mathcal{G}'' = (j_D)_! \mathbb{Q}_{D^*}$, where $j_D : D^* \rightarrow D$ is the inclusion morphism. Let t be a local coordinate around β . Then $\varphi_t \mathcal{G}' = 0$ and $\psi_t \mathcal{G}' = \psi_t \mathcal{G}'' = \varphi_t \mathcal{G}'' = \mathbb{Q}$

using the bijectivity of can (see 2.4) for \mathcal{G}'' . The last assertion follows from the distinguished triangle of functors

$$(2.8.2) \quad i_0^* \rightarrow \psi_t \xrightarrow{\text{can}} \varphi_t \rightarrow,$$

where $i_0: \{\beta\} \rightarrow D$ denotes an inclusion, see [5]. Since ψ_t, φ_t are exact functors, we get $\text{rank } \psi_t \mathcal{G} = 2, \text{rank } \varphi_t \mathcal{G} = 1$ together with the surjectivity of $\text{can}: \psi_t \mathcal{G} \rightarrow \varphi_t \mathcal{G}$.

By (2.4.1) the assertion is thus reduced to the injectivity of $\text{var}: \varphi_t \mathcal{G} \rightarrow \psi_t \mathcal{G}$. By duality (see 2.4) it is equivalent to the surjectivity of $\text{can}: \psi_t \mathcal{G}^* \rightarrow \varphi_t \mathcal{G}^*$, where \mathcal{G}^* is the (shifted) dual of \mathcal{G} which is defined by $\mathbf{R}\mathcal{H}om(\mathcal{G}, \mathbb{Q}_D)$.

By duality [27] \mathcal{G}^* is isomorphic to the derived direct image with compact supports of $\mathbf{R}(j_U)_* \mathbb{Q}_U$. The stalk at β of the higher direct image sheaves is

$$(2.8.3) \quad H_c^j(Y_{B,\beta}, \mathbf{R}(j_\beta)_* \mathbb{Q}_{Y_\beta \setminus Z_\beta}),$$

(where $j_\beta: Y_\beta \setminus Z_\beta \rightarrow Y_\beta$ denotes a natural inclusion), because Z_β and Y_β are smooth on a neighborhood of Z_β . So the assertion is reduced to the vanishing of (2.8.3) for $j = m + 1$ by the distinguished triangle (2.8.2) because it gives the cokernel of can .

Using a one-parameter deformation $t\Psi(t \in \mathbb{C})$, we can reduce the assertion to the case $\Psi = 0$. So we may assume that $Y_{B,\beta}$ is the intersection of B with an affine cone Y_β of a nonsingular conic, and $Z_{B,\beta}$ is its intersection with a linear space passing near the origin. Then we may omit the subscript B by replacing B with arbitrary large open balls and taking the limit. So the assertion follows from Artin’s generalization of the weak Lefschetz theorem [1]. Indeed, the closure \overline{Z}_β of Z_β in the projective space intersects transversely the divisor at infinity $\overline{Y}_\beta \setminus Y_\beta$ of Y_β so that we can first take the direct image with compact supports by $Y_\beta \setminus Z_\beta \rightarrow \overline{Y}_\beta \setminus \overline{Z}_\beta$, and then the usual direct image by $\overline{j}_\beta: \overline{Y}_\beta \setminus \overline{Z}_\beta \rightarrow \overline{Y}_\beta$. Thus the assertion is proved for m odd.

In the case m is even, let D be a sufficiently small neighborhood of α in Δ_c . For a variety V of pure dimension r , let $\tilde{H}_c^j(V) = H_c^j(V)$ for $j \neq 2r$, and define $\tilde{H}_c^{2r}(V)$ to be the kernel of $\text{Tr}: H_c^{2r}(V) \rightarrow \mathbb{Q}(-r)$ which is induced by the canonical morphism of $\mathbb{Q}_V(r)[2r]$ to the dualizing complex \mathbb{D}_V . Then $\tilde{H}_c^j(Z_{B,\alpha}) = \mathbb{Q}$ for $j = m - 1, m$ by calculating the cohomology of $\partial Z_{B,\alpha}$ and using the long exact sequence $\rightarrow H^{j-1}(\partial Z_{B,\alpha}) \rightarrow H_c^j(Z_{B,\alpha}) \rightarrow H^j(Z_{B,\alpha}) \rightarrow$. In particular, (2.7.4) for $\lambda = \alpha$ is not exact, and is extended to a long exact sequence. So we have to consider a distinguished triangle in the derived category of sheaves on D , which is defined by using the direct images with compact supports under the morphisms of $Z_{B,D}, Y_{B,D} \setminus Z_{B,D}, Y_{B,D}$ to D (in particular, the base change holds). They contain some shifted constant sheaves which are annihilated by taking the reduced cohomology in (2.7.2). So deleting them, we get a short exact sequence of shifted perverse sheaves as in (2.8.1) such that the stalk of the 0th cohomology sheaf is given by (2.7.4). By the above calculation, we have $\mathcal{G}' = \mathbf{R}(j_D)_* \mathbb{Q}_{D^*}$ and $\mathcal{G}'' = \mathbb{Q}_D$ in this case.

Using an argument similar to the case m odd, we see that $\text{var}: \varphi_t \mathcal{G} \rightarrow \psi_t \mathcal{G}$ is injective (reducing to the assertion for \mathcal{G}' in this case) where t is a coordinate around α . So it is enough to show the nontriviality of $\text{can}: \psi_t \mathcal{G} \rightarrow \varphi_t \mathcal{G}$. By the triangle (2.8.2) this is equivalent to $H_c^m(Y_{B,\alpha} \setminus Z_{B,\alpha}) = \mathbb{Q}$, because it gives the kernel of can . By duality it is further equivalent to

$$(2.8.4) \quad H^m(Y_{B,\alpha} \setminus Z_{B,\alpha}) = \mathbb{Q}.$$

By the same argument as above, we may omit the subscript B by assuming that Y_α is the restriction of a nonsingular conic \overline{Y}_α in \mathbb{P}^{m+1} to the affine space \mathbb{A}^{m+1} , Z_α is its intersection with a hyperplane H , which is an affine cone of a nonsingular conic in $\mathbb{P}^{m-1} = H \cap \mathbb{P}^m$, and the divisor at infinity $\partial Y_\alpha = \overline{Y}_\alpha \setminus Y_\alpha$ of Y_α is smooth. Then, using a projection from the vertex

of the affine cone Z_α , we see that $Y_\alpha \setminus Z_\alpha$ is isomorphic to the complement of the union of ∂Y_α and $H \cap \mathbb{P}^m$ in $\mathbb{P}^m = \mathbb{P}^{m+1} \setminus \mathbb{A}^{m+1}$. So (2.8.4) follows considering the long exact sequence containing the Gysin morphism by $\partial Y_\alpha \setminus H \rightarrow \mathbb{A}^m = \mathbb{P}^m \setminus H$. This completes the proofs of Lemma 2.8 and Proposition 2.7. \square

3. Hodge-theoretic method

In this section, we prove the nonvanishing of certain extension classes when the assumption on a differential form in Theorem 0.4 is satisfied.

3.1. Extension groups

With the notation and the assumptions of 2.1, we assume in this section that f, g are smooth (by restricting S), and $\mathcal{F}', \mathcal{F}''$ are nonzero. We will consider whether (2.1.2) splits in the category of local systems. Let

$$\mathcal{H} = \mathcal{H}om(\mathcal{F}'', \mathcal{F}').$$

Then there is a canonical isomorphism

$$(3.1.1) \quad \text{Ext}^1(\mathbb{Q}_S, \mathcal{H}) = \text{Ext}^1(\mathcal{F}'', \mathcal{F}'),$$

where the extension group is taken in the category of admissible variations of mixed Hodge structures [25] (or equivalently, in that of mixed Hodge modules [22]). These groups are identified (using [2]) with the scalar extension of the group of admissible normal functions, which are sections of the family of Jacobians $\{J(\mathcal{H}_s)\}_{s \in S}$ satisfying some good conditions [23]. (Here \mathcal{H} also denotes a variation of mixed \mathbb{Z} -variation of Hodge structure whose scalar extension is \mathcal{H} .) We have furthermore a short exact sequence

$$(3.1.2) \quad 0 \rightarrow \text{Ext}^1(\mathbb{Q}, H^0(S, \mathcal{H})) \rightarrow \text{Ext}^1(\mathbb{Q}_S, \mathcal{H}) \xrightarrow{r} \text{Hom}(\mathbb{Q}, H^1(S, \mathcal{H})) \rightarrow 0,$$

see [22,29], where Ext and Hom are taken in the category of mixed Hodge structures or that of admissible variations of mixed Hodge structures. (Using [22], this follows from the adjoint relation between the direct image and the pull-back of mixed Hodge modules by $S \rightarrow pt$. Using [29], we get the above short exact sequence with the cohomology in the last term replaced by the intersection cohomology, which is a subgroup of the cohomology in this case. But this is enough for our purpose, although we can show that these two give the same by taking Hom , see also [8].)

Let $\mathcal{F}_\mathbb{Q}, \mathcal{H}_\mathbb{Q}$, etc. denote the underlying local systems. Then we have

$$(3.1.3) \quad \text{Ext}^1(\mathcal{F}_\mathbb{Q}'', \mathcal{F}_\mathbb{Q}') = H^1(S, \mathcal{H}_\mathbb{Q}).$$

This is compatible with the last morphism r in (3.1.2). Let e be the extension class defined by the short exact sequence (2.1.2). Then (2.1.2) splits in the category of local systems if and only if $r(e) = 0$.

3.2. PROPOSITION (Non-Lefschetz pencil case). – *With the notation and the assumptions of 2.1, assume $Y = Y_0 \times S$, $f = \text{pr}_2$, S is an affine rational curve, and $\text{pr}_1 \circ i : X \rightarrow Y_0$ is birational, where pr_i is the i th projection. Assume furthermore that Y_0 has a nontrivial differential form of*

the highest order. Then (2.1.2) does not split in the category of local systems. Furthermore, if there is a direct sum decomposition of variations of Hodge structures $\mathcal{F}^l = \mathcal{F}'_1 \oplus \mathcal{F}'_2$ such that \mathcal{F}'_1 is a constant variation, then the extension class between \mathcal{F}'' and \mathcal{F}'_2 does not vanish in the category of local systems.

Proof. – Consider the short exact sequence

$$(3.2.1) \quad \mathrm{Gr}_m^W H^m(Y, X; \mathbb{Q}) \xrightarrow{\alpha} \mathrm{Gr}_m^W H^m(Y, \mathbb{Q}) \xrightarrow{\beta} \mathrm{Gr}_m^W H^m(X, \mathbb{Q}),$$

where W is the weight filtration of mixed Hodge structure [4]. Since S is a smooth affine rational curve, $H^1(S, \mathbb{Q})$ has weights > 1 , and we have by the Künneth decomposition

$$\mathrm{Gr}_m^W H^m(Y, \mathbb{Q}) = H^m(Y_0, \mathbb{Q}).$$

Then the last morphism β in (3.2.1) is identified with the restriction morphism

$$H^m(Y_0, \mathbb{Q}) \rightarrow \mathrm{Gr}_m^W H^m(X, \mathbb{Q})$$

by $X \rightarrow Y_0$, and its kernel is annihilated by the restriction morphism to a nonempty open subvariety of Y_0 , because $X \rightarrow Y_0$ is birational.

Thus the kernel of β has level $< m$ (where the level of a Hodge structure is the difference between the maximal and minimal numbers p such that the p th graded piece of the Hodge filtration does not vanish, see [4]). This implies that $\mathrm{Im} \beta \neq 0$, because $H^m(Y_0, \mathbb{Q})$ has level m by the hypothesis on the highest form. So we have a nonzero element

$$u \in \mathrm{Gr}_m^W H^m(Y, \mathbb{Q}) = H^0(S, R^m f_* \mathbb{Q}_Y)$$

such that $\beta(u) \neq 0$. Using the semisimplicity of polarizable Hodge structures [4], we may assume that $\gamma_m(u) = 0$ in $H^0(S, R^m g_* \mathbb{Q}_X)$ (or equivalently, in $H^m(X_s, \mathbb{Q})$ for $s \in S$), because $H^m(X_s, \mathbb{Q})$ has level $< m$. Thus u belongs to $H^0(S, \mathcal{F}'')$.

For the first morphism α in (3.2.1), consider the Leray spectral sequence in the category of mixed Hodge structures

$$(3.2.2) \quad E_2^{p,q} = H^p(S, R^q h_! \mathbb{Q}_U) \implies H^{p+q}(Y, X; \mathbb{Q}).$$

This degenerates at E_2 , because $E_2^{p,q} = 0$ unless $p = 0$ or 1 . So we get the surjection

$$(3.2.3) \quad \mathrm{Gr}_m^W H^m(Y, X; \mathbb{Q}) \rightarrow \mathrm{Gr}_m^W H^0(S, \mathcal{F}) \quad (\text{see 2.1 for } \mathcal{F}).$$

Now assume that (2.1.2) splits in the category of local systems. Then there exists $v \in H^0(S, \mathcal{F})$ whose image in $H^0(S, \mathcal{F}'')$ is u . Here we may assume $v \in \mathrm{Gr}_m^W H^0(S, \mathcal{F})$, because $H^0(S, \mathcal{F}'')$ is pure of weight m , and the image commutes with Gr_m^W . But this contradicts the nonvanishing of $\beta(u)$ using the surjectivity of (3.2.3). So the first assertion follows.

For the last assertion, it is enough to show the vanishing of the extension class between \mathcal{F}'' and \mathcal{F}'_1 as local systems. But this follows from (3.1.2) and (3.1.3) because the first cohomology of a constant variation of Hodge structures of weight -1 on a rational curve has weight > 0 (if it is nonzero). This completes the proof of Proposition 3.2. \square

3.3. PROPOSITION (Surface case). – *With the notation and the assumptions of 2.1, assume $m = 1$ (i.e. Y is a surface and X is a curve), S is an affine rational curve, and the fiber Y_s of f is not a rational curve. Assume further that $f: Y \rightarrow S$ can be extended to a proper smooth morphism $f': Y' \rightarrow S'$ so that Y, S are open subvarieties of Y', S' respectively, and the closure X' of X in Y' is nonsingular, but it is not smooth over S' . Then (2.1.2) does not split in the category of local systems.*

Proof. – In this case we have

$$\mathcal{F}'_s = \tilde{H}^0(X_s, \mathbb{Q}), \quad \mathcal{F}''_s = H^1(Y_s, \mathbb{Q}).$$

Then in the notation of (3.1.3) we have

$$H^0(S, \mathcal{H}) = \text{Hom}(\mathcal{F}''_{\mathbb{Q}}, \mathcal{F}'_{\mathbb{Q}}) = 0,$$

considering the monodromy of \mathcal{F}'_s around $S' \setminus S$. So by (3.1.2) it is enough to show that the corresponding normal function in (3.1.1) is nontorsion, and the assertion is local on S in the classical topology.

Let $s_0 \in S' \setminus S$, and Δ be an open disk around s_0 in S' such that $\Delta \cap S = \Delta^* (:= \Delta \setminus \{s_0\})$. Then the assumption on S' implies that there exist continuously $\Lambda_s = \{x_s, y_s\} \subset X_s$ for $s \in \Delta^*$ such that the action of the monodromy around s_0 on Λ_s is nontrivial (replacing s_0 if necessary). Locally on Δ^* , the difference $[x_s] - [y_s]$ defines an element $u_s \in \tilde{H}^0(X_s)$ and also a point ξ_s of the Jacobian $J(Y_s)$ of Y_s . Note that ξ_s corresponds to the pull-back of the dual of the short exact sequence of mixed Hodge structures $(2.1.2)_s$ by u_s (using [2]), where $(2.1.2)_s$ is the stalk at s of the exact sequence (2.1.2).

We have $\xi_s \neq 0$ in $J(Y_s)$, because Y_s is nonrational. If a nonzero multiple of ξ_s vanishes locally on Δ^* , it defines a locally constant section of the division points of the family of Jacobians over Δ (because f is proper smooth over Δ), and we get a contradiction by considering the monodromy around s_0 . So the assertion follows. \square

3.4. Complement on the assumption of 3.3

The last assumption of Proposition 3.3 is satisfied in the case of a generic Lefschetz pencil as follows. Let Y be a smooth surface embedded in a projective space $\mathcal{P} := \mathbb{P}^r_{\mathbb{C}}$ ($r > 2$). Let X be a (locally closed) smooth curve on Y . Then there is a hyperplane of \mathcal{P} which intersects Y transversely, but is tangent to X .

Indeed, let \mathcal{P}^* denote the dual projective space of \mathcal{P} consisting of hyperplanes H of \mathcal{P} , and D_Y, D_X be the set of hyperplanes tangent to Y, X respectively. By definition, D_Y is the image of a \mathbb{P}^{r-3} -bundles P_Y over Y (where $P_{Y,y}$ consists of the hyperplanes tangent to Y at y), and similarly for X with \mathbb{P}^{r-3} replaced by \mathbb{P}^{r-2} . Let $x \in X$, and assume $P_{X,x} \subset D_Y$. Then there exist a pencil of hyperplanes $\{H_t\}_{t \in \mathbb{P}^1}$ contained in $P_{X,x}$ and a smooth analytic curve C locally defined on Y together with a nonconstant holomorphic map $\rho: C \rightarrow \mathbb{P}^1$ such that $H_{\rho(c)}$ is tangent to Y (and hence to C) at any $c \in C$. But this implies the constancy of ρ , which is a contradiction.

4. Degeneration and nearby cycles

In this section, we calculate the weight filtration of Steenbrink, and prove Theorem 0.4 using the results in Sections 2 and 3.

4.1. Family of hypersurfaces

With the notation of 1.1, assume condition 0.1 in Introduction. Let V_1, V_2, V_3 be vector subspaces of $\mathcal{A}_Z^\delta, \mathcal{A}^{d-\delta}, \mathcal{A}_Z^d$ such that V_1, V_3 generate $\mathcal{I}_Z \otimes \mathcal{O}_Y(\delta), \mathcal{I}_Z \otimes \mathcal{O}_Y(d)$ respectively and V_2 gives an embedding of Y into a projective space. We assume V_3 is the image of $V_3' \otimes V_3''$ where V_3' is a vector subspace of \mathcal{A}_Z^δ generating $\mathcal{O}_Y(\delta)$ outside Z and V_3'' is a vector subspace of $\mathcal{A}^{d-\delta}$ giving an embedding of Y into a projective space. In the proof of Theorem 0.4, V_1, V_3' and V_2, V_3'' will be respectively the restrictions of $\mathcal{A}_{Z'}^\delta$ and $\mathcal{A}^{d-\delta}$ defined for some smooth projective variety Y' containing Y where $Z' \subset Y'$ is a subvariety whose intersection with Y is Z . (Actually, it is possible that Z is empty, but Z' is not.) These are necessary to carry out an inductive argument in 4.7.

We will identify V_1, V_2, V_3 with the corresponding affine spaces. Let $\tilde{V}_1, \tilde{V}_2, \tilde{V}_3$ be their intersections with $\mathcal{V}_Z^\delta, \mathcal{V}^{d-\delta}, \mathcal{V}_Z^d$. Put $X_{P,Q} = X_P \cap X_Q, X_{P,Q,R} = X_P \cap X_Q \cap X_R$ for $P \in \mathcal{A}^i$, etc. Let

$$S_0 = \{(P, Q, R) \in \tilde{V}_1 \times \tilde{V}_2 \times \tilde{V}_3: X_{P,Q}, X_{Q,R}, X_{P,Q,R} \text{ are SCI}\},$$

where SCI means smooth complete intersection. By [19] (or 1.2), S_0 is nonempty, and $X_{P,R}$ has at most isolated singularities, see also [13]. Here we can replace S_0 with a nonempty subvariety because of 4.2 below. Let

$$S' = \{(P, Q, R, t) \in S_0 \times \mathbb{C}^*: X_{PQ+tR} \text{ is smooth}\},$$

and $S'' = \mathcal{V}_Z^d / \mathbb{C}^*$. We have a local system L on S'' whose stalks are given by $H^m(X, \mathbb{Q})_{\perp Z}^{\text{van}}$ in Conjecture 0.2. Consider a morphism $\rho: S' \rightarrow S''$ which associates $PQ + tR$ to (P, Q, R, t) . We apply the reduction argument in 4.2 below to this so that the proof of Theorem 0.4 is reduced to the pull-back of the local system to S' . Here it is enough to show the *indecomposability* of L , because L is semisimple by Deligne [4].

4.2. Reduction argument

Let L be a local system on a connected analytic space S'' . Then it is simple (respectively indecomposable) if there exists a morphism $\rho: S' \rightarrow S''$ such that ρ^*L is simple (respectively indecomposable). This follows from the fact that the functor ρ^* is exact and faithful (or using the corresponding representation of the fundamental group). A similar assertion also holds for the nearby cycle functor ψ .

4.3. Semistable degeneration

With the notation of 4.1, there is a smooth projective morphism $f': \mathcal{X}' \rightarrow S'$ whose fiber $\mathcal{X}_s := f'^{-1}(s)$ is X_{PQ+tR} for $s = (P, Q, R, t) \in S'$. Let L' be a subsheaf of $R^m f'_* \mathbb{Q}_{\mathcal{X}'}$ whose stalk is the orthogonal complement of the subspace generated by $H^m(Y, \mathbb{Q})$ together with the cycle classes of the irreducible components of the closed subvariety Z . Then $L' = \rho^*L$. Let

$$S = \{(P, Q, R, t) \in S_0 \times \mathbb{C}: X_{PQ+tR} \text{ is smooth with } t \neq 0 \text{ or } t = 0\},$$

i.e. S is the disjoint union of S' and S_0 . Then f' is naturally extended to $f: \mathcal{X} \rightarrow S$. However, \mathcal{X} has certain singularities.

Let \mathcal{C} be the closed subvariety of \mathcal{X} whose fiber over $(P, Q, R, t) \in S$ is $X_{Q,R}$. Note that its restriction over S' is a locally principal divisor. Let $\pi: \tilde{\mathcal{X}} \rightarrow \mathcal{X}$ be the blow-up along \mathcal{C} . Let $\tilde{f}: \tilde{\mathcal{X}} \rightarrow S$ denote the composition with f . We also denote by t the function defined by

the last component of (P, Q, R, t) . Let $\tilde{\mathcal{X}}_0, \mathcal{X}_0, S_0$ be the subvarieties of $\tilde{\mathcal{X}}, \mathcal{X}, S$ defined by $t = 0$ (this is compatible with the previous definition of S_0). We have the induced morphism $\tilde{f}_0 : \tilde{\mathcal{X}}_0 \rightarrow S_0$.

Let \mathcal{X}_P be the closed subvariety of \mathcal{X} whose fiber over $(P, Q, R) \in S_0$ is X_P (using the projection $\mathcal{X} \rightarrow S_0$), and similarly for $\mathcal{X}_{P,Q}, \mathcal{X}_{P,Q,R}$, etc. (Note that $\mathcal{X}_P, \mathcal{X}_Q$ are contained in \mathcal{X}_0 .) Let $\mathcal{X}_{\tilde{P}}$ be the blow-up of \mathcal{X}_P along $\mathcal{X}_{P,Q,R}$. Then $\tilde{\mathcal{X}}_0$ is a divisor with normal crossings on a smooth variety $\tilde{\mathcal{X}}$, and its irreducible components are $\mathcal{X}_{\tilde{P}}$ and \mathcal{X}_Q , see [10].

Let ψ denotes the nearby cycle functor, see [5]. Then

$$(4.3.1) \quad R^m \tilde{f}_* \psi_t \mathbb{Q}_{\tilde{\mathcal{X}}} = \psi_t R^m \tilde{f}_* \mathbb{Q}_{\tilde{\mathcal{X}}} (= \psi_t R^m f'_* \mathbb{Q}_{\mathcal{X}'}),$$

because the nearby cycle functor commutes with the direct image under a proper morphism.

Let W be the weight filtration on $\psi_t \mathbb{Q}_{\tilde{\mathcal{X}}}$. By Steenbrink [24], we have

$$(4.3.2) \quad \begin{aligned} \mathrm{Gr}_{m-1}^W \psi_t \mathbb{Q}_{\tilde{\mathcal{X}}} &= \mathbb{Q}_{\mathcal{X}_{P,Q}}[-1], \\ \mathrm{Gr}_m^W \psi_t \mathbb{Q}_{\tilde{\mathcal{X}}} &= \mathbb{Q}_{\mathcal{X}_Q} \oplus \mathbb{Q}_{\mathcal{X}_{\tilde{P}}}, \\ \mathrm{Gr}_{m+1}^W \psi_t \mathbb{Q}_{\tilde{\mathcal{X}}} &= \mathbb{Q}_{\mathcal{X}_{P,Q}}(-1)[-1], \end{aligned}$$

and $\mathrm{Gr}_k^W \psi_t \mathbb{Q}_{\tilde{\mathcal{X}}} = 0$ for $|k - m| > 1$.

4.4. Weight spectral sequence

Consider the weight spectral sequence in the category of local systems

$$(4.4.1) \quad E_1^{-k, j+k} = R^j \tilde{f}_* \mathrm{Gr}_k^W \psi_t \mathbb{Q}_{\tilde{\mathcal{X}}} \implies \psi_t R^j f'_* \mathbb{Q}_{\mathcal{X}'}$$

This degenerates at E_2 by [24]. (In this case, it also follows from an easy calculation.) The local system $\{H^j(X_P)\}$ on S_0 can be identified with $R^j \tilde{f}_* \mathbb{Q}_{\mathcal{X}_P}$, and similarly for $\{H^j(X_{\tilde{P}})\}$, etc., where the cohomology groups are with rational coefficients unless otherwise stated explicitly. We have

$$H^j(X_{\tilde{P}}) = H^j(X_P) \oplus H^{j-2}(X_{P,Q,R})(-1),$$

because

$$(4.4.2) \quad \mathbf{R}\pi_* \mathbb{Q}_{\mathcal{X}_{\tilde{P}}} = \mathbb{Q}_{\mathcal{X}_P} \oplus \mathbb{Q}_{\mathcal{X}_{P,Q,R}}(-1)[-2],$$

by the decomposition theorem [1]. Let $\mathcal{Z}_{\tilde{P}} = \pi^{-1}(X_{P,Q,R})$. Then a canonical morphism $\mathbb{Q}_{\mathcal{X}_{P,Q,R}}(-1)[-2] \rightarrow \mathbf{R}\pi_* \mathbb{Q}_{\mathcal{X}_{\tilde{P}}}$ and its right inverse (up to a sign) are given by the compositions

$$(4.4.3) \quad \begin{aligned} \mathbb{Q}_{\mathcal{X}_{P,Q,R}}(-1)[-2] &\rightarrow \mathbf{R}\pi_* \mathbb{Q}_{\mathcal{Z}_{\tilde{P}}}(-1)[-2] \rightarrow \mathbf{R}\pi_* \mathbb{Q}_{\mathcal{X}_{\tilde{P}}}, \\ \mathbf{R}\pi_* \mathbb{Q}_{\mathcal{X}_{\tilde{P}}} &\rightarrow \mathbf{R}\pi_* \mathbb{Q}_{\mathcal{Z}_{\tilde{P}}} \rightarrow \mathbb{Q}_{\mathcal{X}_{P,Q,R}}(-1)[-2]. \end{aligned}$$

The differential d_1 of the spectral sequence is induced by the extension classes between the Gr_k^W which are given by the restriction and Gysin morphisms. Indeed, the extension class between Gr_{m-1}^W and Gr_m^W corresponds to a morphism $\mathrm{Gr}_m^W \rightarrow \mathrm{Gr}_{m-1}^W$ [1] in the derived category of \mathbb{Q} -modules on $\tilde{\mathcal{X}}$ (see [26]) and it is given by the restriction morphisms

$$\begin{aligned} \mathbb{Q}_{\mathcal{X}_Q} &\rightarrow \mathbb{Q}_{\mathcal{X}_{P,Q}}, \\ \mathbb{Q}_{\mathcal{X}_P^\sim} &\rightarrow \mathbb{Q}_{\mathcal{X}_{P,Q}}. \end{aligned}$$

The direct image by π of the last morphism is given by the restriction and Gysin morphisms

$$\begin{aligned} \mathbb{Q}_{\mathcal{X}_P} &\rightarrow \mathbb{Q}_{\mathcal{X}_{P,Q}}, \\ \mathbb{Q}_{\mathcal{X}_{P,Q,R}}(-1)[-2] &\rightarrow \mathbb{Q}_{\mathcal{X}_{P,Q}} \end{aligned}$$

(up to signs), using (4.4.2). We have a similar assertion between Gr_m^W and Gr_{m+1}^W .

Let $H^{m-1}(X_{P,Q})^{\text{van}}$ denote the orthogonal complement of the image of the (injective) restriction morphism $H^{m-1}(Y) \rightarrow H^{m-1}(X_{P,Q})$ where $H^{m-1}(Y)$ can be replaced with $H^{m-1}(X_P)$ by the weak Lefschetz theorem, and similarly for $H^{m-2}(X_{P,Q,R})^{\text{van}}$ and also for $H^m(X_P)^{\text{van}}$. Then we have the direct sum decompositions

$$(4.4.4) \quad \begin{aligned} H^m(X_P) &= H^m(X_P)^{\text{van}} \oplus H^m(Y)^{\text{prim}} \oplus H^{m-2}(Y)(-1), \\ H^m(X_Q) &= H^m(X_Q)^{\text{van}} \oplus H^m(Y)^{\text{prim}} \oplus H^{m-2}(Y)(-1), \\ H^{m-2}(X_{P,Q,R}) &= H^{m-2}(X_{P,Q,R})^{\text{van}} \oplus H^{m-2}(Y), \end{aligned}$$

together with the isomorphisms

$$H^{m-2}(X_{P,Q})(-1) = H^m(X_{P,Q}) = H^{m-2}(Y)(-1).$$

Here $H^m(Y)^{\text{prim}}$ denotes the primitive cohomology, and $H^{m-2}(Y)(-1)$ in the first two decompositions of (4.4.4) is actually the image of $H^{m-2}(Y)(-1)$ by the action of the cohomology class of the hyperplane section.

We see that the stalk of $\text{Gr}_m^W \psi_t R^m f'_* \mathbb{Q}_{\mathcal{X}'}$ is the cohomology of the complex

$$(4.4.5) \quad H^{m-2}(X_{P,Q})(-1) \rightarrow H^m(X_P) \oplus H^m(X_Q) \oplus H^{m-2}(X_{P,Q,R})(-1) \rightarrow H^m(X_{P,Q})$$

where the morphisms are induced by the restriction and Gysin morphisms up to signs. Using (4.4.4), the first morphism of (4.4.5) is given by the identity on $H^{m-2}(Y)(-1)$ up to signs, and the last morphism is given by the multiplications on $H^{m-2}(Y)(-1)$ by $d - \delta, \delta, d$ up to signs. Indeed, the composition of the Gysin and restriction morphisms

$$H^{m-2}(X_{P,Q})(-1) \rightarrow H^m(X_P) \rightarrow H^m(X_{P,Q})$$

coincides with the action of the restriction of the cycle class of $X_{P,Q}$ in X_P .

Similarly we can verify

$$\begin{aligned} \text{Gr}_{m-1}^W \psi_t R^m f'_* \mathbb{Q}_{\mathcal{X}'} &= \{H^{m-1}(X_{P,Q})^{\text{van}}\}, \\ \text{Gr}_{m+1}^W \psi_t R^m f'_* \mathbb{Q}_{\mathcal{X}'} &= \{H^{m-1}(X_{P,Q})^{\text{van}}(-1)\}. \end{aligned}$$

Here the first assertion is easy, and the second follows from it using duality.

Note that the extension classes between the Gr_k^W are induced by the restriction and Gysin morphisms in the derived category of sheaves on \mathcal{X} as above.

4.5. Limit of the invariant part

Taking the nearby cycle functor (i.e. passing to the limit by $t \rightarrow 0$), the restriction morphism $H^m(Y) \rightarrow H^m(X_{PQ+tR})$ induces a morphism of $H^m(Y)$ to the middle term of the complex (4.4.5), which is defined (up to signs) by using the isomorphism

$$(4.5.1) \quad H^m(Y) = H^m(Y)^{\text{prim}} \oplus H^{m-2}(Y)(-1)$$

together with (4.4.4). So Gr_m^W of the limit of $H^m(X_{PQ+tR})/H^m(Y)$ is given by

$$(4.5.2) \quad (H^m(X_P)^{\text{prim}} \oplus H^m(X_Q)^{\text{prim}} \oplus H^{m-2}(X_{P,Q,R})^{\text{van}}(-1))/H^m(Y)^{\text{prim}}.$$

Note that

$$(4.5.3) \quad H^m(X_P)^{\text{prim}} = H^{m-2}(X_P)^{\text{van}} \oplus H^m(Y)^{\text{prim}}$$

(similarly for $H^m(X_Q)^{\text{prim}}$), and the division by $H^m(Y)^{\text{prim}}$ is defined by using the diagonal morphism. In particular, the quotient (4.5.2) is isomorphic to

$$(4.5.4) \quad H^m(X_P)^{\text{prim}} \oplus H^m(X_Q)^{\text{van}} \oplus H^{m-2}(X_{P,Q,R})^{\text{van}}(-1).$$

Here $\{H^m(X_Q)^{\text{van}}\}$ is an irreducible local system on S_0 . Assuming an appropriate inductive hypothesis (see 4.7 below), this also holds for the quotient of $H^{m-2}(X_{P,Q,R})^{\text{van}}$ by the subspace generated by the cycle classes of the intersection with X_Q of the $(m/2)$ -dimensional irreducible components of Z if $m > 2$.

4.6. Cycle classes of the irreducible components

The cycle classes of the irreducible components of Z in the limit of $H^m(X_{PQ+tR})$ are given by using

$$(4.6.1) \quad H_{\text{BM}}^{m-2}(X_{P,R})(-1) \rightarrow H^m(X_P) \oplus H^{m-2}(X_{P,Q,R})(-1),$$

which is induced by the Gysin and restriction morphisms up to signs. (This can be verified by using (4.4.3).) Here $H_{\text{BM}}^j(V)$ denotes Borel–Moore cohomology for an equidimensional variety V in general, and is defined by $H^j(V, \mathbb{D}_V(-n)[-2n])$, where \mathbb{D}_V is the dualizing complex and $n = \dim V$. If V is compact, it is isomorphic to $H_{2n-j}(V)(-n)$. The restriction of the dualizing complex to the smooth part V_{reg} is isomorphic to $\mathbb{Q}_{V_{\text{reg}}}(n)[2n]$, and we get the restriction morphism $H_{\text{BM}}^j(V) \rightarrow H^j(V')$ for any subvariety V' of V_{reg} .

The morphism to the second factor of (4.6.1) is injective, i.e. the Gysin morphism $H^{m-2}(X_{P,Q,R})(-1) \rightarrow H^m(X_{P,R})$ is surjective. This follows from Artin’s theorem (see [1]) which asserts the vanishing of $H^m(X_{P,R} \setminus X_{P,Q,R})$ because $X_{P,R} \setminus X_{P,Q,R}$ is affine and the constant sheaf on it is semi-perverse up to a shift of complex by $m - 1$.

We can verify that the limit of the cycle class of Z is then defined by using the cycle class of the cycle in X_P and the cycle class in $X_{P,Q,R}$ of the intersection of the cycle with X_Q .

4.7. Proof of Theorem 0.4

Assume first $m = 2$ (and Y' in (4.1) is Y). In this case, $X_{P,Q,R}$ has dimension 0 and is not connected. We first fix P, R and consider $\{X_{P,Q}\}_Q$ and $\{X_{P,Q,R}\}_Q$. Let $Z_i (i > 0)$ be the 1-

dimensional irreducible components of Z . Then $X_{P,R}$ is the union of $\bigcup_{i>0} Z_i$ and an irreducible curve Z_0 , because $X_{P,R} \setminus Z$ is smooth and connected by 1.2 (where $b = 0$ if $\dim Z = 1$), see also [13,19].

We apply the dual of Proposition 2.7 or 3.3 to $\{X_{P,Q}\}_Q$ and $\{Z_{i,Q}\}_Q$ for $i \geq 0$, where $Z_{i,Q} = Z_i \cap X_Q$. Then we get the nontriviality of the extension class between $\text{Gr}_{m-1}^W = \{H^{m-1}(X_{P,Q})^{\text{van}}\}$ and $\{\tilde{H}^{m-2}(Z_{i,Q})(-1)\}$ for $i \geq 0$, using 4.2. Similarly we apply Proposition 2.7 or 3.2 to show the nontriviality of the extension class between Gr_{m-1}^W and $\{H^m(X_Q)^{\text{van}}\}$, where we fix Q, R or P, R to apply the propositions. We use Proposition 2.3 for the extension between Gr_{m-1}^W and any simple factor of $\{H^m(X_P)^{\text{prim}}\}$. (Note that $\{H^m(X_P)^{\text{prim}}\}$ is semisimple by [4].)

Thus we get the nontriviality of the extension class between $\text{Gr}_{m-1}^W \psi_t L'$ and each simple factor of $\text{Gr}_m^W \psi_t L'$ in the notation of 4.3, because $\text{Gr}_m^W \psi_t L'$ is isomorphic to the quotient of (4.5.4) by the image of the cycle classes of the $Z_i (i > 0)$ (where the last term vanishes unless $\dim Z = 1$), see 4.6. We have the dual argument for the extension between Gr_m^W and Gr_{m+1}^W . Note that the $\{\tilde{H}^{m-2}(Z_{i,Q})\}$ are not isomorphic to each other (considering the monodromy around Q such that X_Q is tangents to Z_j at a smooth point for one j , but intersects Z_i transversely at smooth points for $i \neq j$). We also see that any simple factor of $\{H^m(X_P)^{\text{prim}}\}$ is not isomorphic to $\{\tilde{H}^{m-2}(Z_{i,Q})\}$, nor to $\{H^m(X_Q)^{\text{van}}\}$ (fixing P), and similarly between the last two.

Assume that there is a decomposition $\psi_t L' = L_1 \oplus L_2$ in the notation of 4.3. We may assume that the L_i are stable by the action of the monodromy T , because the decomposition is induced by that of L' . In our case the weight filtration W is defined by $W^{m-1} = \text{Im } N$ and $W^m = \text{Ker } N$. Here $N = T - \text{id}$ because $N^2 = 0$. If, for example, L_1 is not contained in W^m , then $NL_1 = W^{m-1}$, because $NL_1 \subset W^{m-1}$ is nonzero and $W^{m-1} = \text{Gr}_{m-1}^W$ is simple. In this case L_2 is contained in W^m , because otherwise it also contains W^{m-1} . So the decomposition induces that of Gr_m^W , and the nontriviality of the above extension classes implies the triviality of the decomposition. Thus, using Deligne's semisimplicity theorem, Theorem 0.4 is proved for $m = 2$.

If $m = 1$, then X_P is smooth and connected by Theorem 1.2 with $r = 1$, and $X_{P,Q,R}$ is empty. So the argument is essentially same as above.

Assume now $m > 2$. We apply an inductive hypothesis to $X_{P,Q}$ and $Z \cap X_Q$ to show the irreducibility of $\{H^{m-2}(X_{P,Q,R})^{\text{van}}\}_R$ divided by the image of the cycle classes of the irreducible components of Z , see 4.6. To carry out this induction we take general P_i, Q_i, R_i such that $R_i = P_{i+1}Q_{i+1} + R_{i+1}$ and Z is contained in X_{P_i}, X_{R_i} . Note that the singular locus of $\bigcap_{1 \leq j \leq i} X_{P_j}$ has dimension $\leq i - 2$, see [19] (or 1.2). We apply the above argument to the restriction of (P_i, Q_i, R_i) to $Y'' := \bigcap_{1 \leq j < i} X_{P_j, Q_j}$ for each i , and proceed by decreasing induction on i . This is allowed by the definition of S_0 in 4.1 (where Y', Y in 4.1 is Y, Y'' here). Here we use Proposition 2.7 or 3.2 (instead of 3.3) to show the nontriviality of the extension between Gr_W^{m-1} and $\{H^{m-2}(X_{P,Q,R})^{\text{van}}(-1)\}$ divided by the cycle classes as above. Note that even if $Y'' \cap Z$ is empty, we still have some restriction to (P_i, Q_i, R_i) coming from Z . At the first step of the induction, we have $m = 2$ or 1 . This completes the proof of Theorem 0.4.

4.8. Generalization of Theorem 0.4

The assertion also holds for smooth zero loci of sections of $\mathcal{L}_1 \otimes \mathcal{L}_2$ containing Z where $\mathcal{L}_1, \mathcal{L}_2$ are line bundles such that $\mathcal{L}_1 \otimes \mathcal{I}_Z$ is generated by its global sections, and \mathcal{L}_2 is very ample and satisfies one of the following two conditions: either a general smooth hyperplane section of \mathcal{L}_2

has a nontrivial differential form of the highest degree, or the 2-jets at each point is generated by the global sections of \mathcal{L}_2 (e.g. it is the $(d - \delta)$ -ple tensor of a very ample line bundle with $d \geq \delta + 2$).

REFERENCES

- [1] BEILINSON A., BERNSTEIN J., DELIGNE P., Faisceaux pervers, *Astérisque*, vol. **100**, Soc. Math. France, Paris, 1982.
- [2] CARLSON J., Extensions of mixed Hodge structures, in: *Journées de géométrie algébrique d'Angers 1979, Sijthoff-Noordhoff Alphen a/d Rijn*, 1980, pp. 107–128.
- [3] CLEMENS H., Degeneration of Kähler manifolds, *Duke Math. J.* **44** (1977) 215–290.
- [4] DELIGNE P., Théorie de Hodge I, *Actes Congrès Intern. Math.* **1** (1970) 425–430;
DELIGNE P., Théorie de Hodge II, *Publ. Math. IHES* **40** (1971) 5–57;
DELIGNE P., Théorie de Hodge III, *Publ. Math. IHES* **44** (1974) 5–77.
- [5] DELIGNE P., Le formalisme des cycles évanescents, in: *SGA7 XIII and XIV*, in: *Lecture Notes in Math.*, vol. **340**, Springer, Berlin, 1973, pp. 82–115 and 116–164.
- [6] DELIGNE P., La formule de Picard–Lefschetz, in: *SGA7 XV*, in: *Lecture Notes in Math.*, vol. **340**, Springer, Berlin, 1973, pp. 165–197.
- [7] DIMCA A., *Sheaves in Topology*, Universitext, Springer, Berlin, 2004.
- [8] DIMCA A., SAITO M., Monodromy at infinity and the weights of cohomology, *Compositio Math.* **138** (2003) 55–71.
- [9] EISENBUD D., *Commutative Algebra with a View Toward Algebraic Geometry*, Springer, New York, 1995.
- [10] GRIFFITHS P., HARRIS J., On the Noether–Lefschetz theorem and some remarks on codimension two cycles, *Math. Ann.* **271** (1985) 31–51.
- [11] ILLUSIE L., Autour du théorème de monodromie locale, *Astérisque* **223** (1994) 9–57.
- [12] KATZ N., Étude cohomologique des pinceaux de Lefschetz, in: *Lecture Notes in Math.*, vol. **340**, Springer, Berlin, 1973, pp. 254–327.
- [13] KLEIMAN S., ALTMAN A., Bertini theorems for hypersurface sections containing a subscheme, *Comm. Algebra* **7** (1979) 775–790.
- [14] LEFSCHETZ S., *L'analysis situs et la géométrie algébrique*, Gauthier-Villars, Paris, 1924.
- [15] LEWIS J.D., *A Survey of the Hodge Conjecture*, second ed., *Monograph Series*, vol. **10**, American Mathematical Society, Providence RI, 1999.
- [16] LOPEZ A.F., Noether–Lefschetz Theory and the Picard Group of Projective Surfaces, *Mem. Amer. Math. Soc.*, vol. **89**, American Mathematical Society, Providence, RI, 1991.
- [17] MILNOR J., *Singular Points of Complex Hypersurfaces*, *Ann. of Math. Stud.*, vol. **61**, Princeton University Press, Princeton, NJ, 1968.
- [18] OTWINOWSKA A., Composantes de petite codimension du lieu de Noether–Lefschetz; un argument en faveur de la conjecture de Hodge pour les hypersurfaces, *J. Algebraic Geom.* **12** (2003) 307–320.
- [19] OTWINOWSKA A., *Monodromie d'une famille d'hypersurfaces*, Preprint.
- [20] OTWINOWSKA A., *Sur les variétés de Hodge des hypersurfaces*, math.AG/0401092.
- [21] SAITO M., Modules de Hodge polarisables, *Publ. RIMS, Kyoto Univ.* **24** (1988) 849–995.
- [22] SAITO M., Mixed Hodge modules, *Publ. RIMS, Kyoto Univ.* **26** (1990) 221–333.
- [23] SAITO M., Admissible normal functions, *J. Algebra Geom.* **5** (1996) 235–276.
- [24] STEENBRINK J.H.M., Limits of Hodge structures, *Invent. Math.* **31** (1975/76) 229–257.
- [25] STEENBRINK J.H.M., ZUCKER S., Variation of mixed Hodge structure I, *Invent. Math.* **80** (1985) 489–542.
- [26] VERDIER J.-L., Catégories dérivées, in: *SGA 4 1/2*, in: *Lecture Notes in Math.*, vol. **569**, Springer, Berlin, 1977, pp. 262–311.

- [27] VERDIER J.-L., Dualité dans la cohomologie des espaces localement compacts, *Sém. Bourbaki (1965/66), Exp. n° 300 Collection hors série de la SMF* **9** (1995) 337–349.
- [28] VOISIN C., *Hodge Theory and Complex Algebraic Geometry, II*, Cambridge University Press, Cambridge, 2003.
- [29] ZUCKER S., Hodge theory with degenerating coefficients, L_2 -cohomology in the Poincaré metric, *Ann. of Math.* **109** (1979) 415–476.

(Manuscrit reçu le 23 avril 2004 ;
accepté, après révision, le 31 mars 2005.)

Ania OTWINOWSKA
Laboratoire de Mathématiques,
Université Paris-Sud,
Bât. 425,
91405 Orsay cedex, France
E-mail: ania.otwinowska@math.u-psud.fr

Morihiko SAITO
RIMS,
Kyoto University,
Kyoto 606-8502, Japan
E-mail: msaito@kurims.kyoto-u.ac.jp