



Mathematical analysis

Distances between classes of sphere-valued Sobolev maps

*Distances entre classes d'applications de Sobolev à valeurs dans une sphère*Haïm Brézis^{b,c}, Petru Mironescu^a, Itai Shafrir^c^a Université de Lyon, Université Lyon-1, CNRS UMR 5208, Institut Camille-Jordan, 43, bd. du-11-Novembre-1918, 69622 Villeurbanne cedex, France^b Department of Mathematics, Rutgers University, USA^c Department of Mathematics, Technion – I.I.T., 32 000 Haifa, Israel

ARTICLE INFO

Article history:

Received 23 April 2016

Accepted 24 April 2016

Available online 18 May 2016

Presented by Haïm Brézis

ABSTRACT

We introduce an equivalence relation on $W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$ involving the topological degree, and we evaluate the distances (in the usual sense and in the Hausdorff sense) between the equivalence classes. In some special cases, we even obtain exact formulas. Next we discuss related issues for $W^{1,p}(\Omega; \mathbb{S}^1)$.

© 2016 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

R É S U M É

On introduit une relation d'équivalence sur $W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$ liée au degré topologique, et on présente des estimées pour les distances (au sens usuel et au sens de Hausdorff) entre les classes d'équivalence. Dans certains cas particuliers, il s'agit même de formules exactes. On considère ensuite des questions semblables pour $W^{1,p}(\Omega; \mathbb{S}^1)$.

© 2016 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

We report on two recent works [5] and [4] concerning distances between classes of Sobolev maps taking their values in \mathbb{S}^N , in two different settings. In the first part, we deal with distances in $W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$ [5]. The classes in this case correspond to the equivalence relation $f \sim g$ if and only if $\deg f = \deg g$. As we will recall below, the topological degree makes sense not only for continuous maps, but also for VMO-maps, and in particular for maps in $W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$ with $sp \geq N$. In the second part, we consider distances between classes of $W^{1,p}(\Omega; \mathbb{S}^1)$, where Ω is a smooth bounded simply connected domain in \mathbb{R}^N , $N \geq 2$ and $p \in [1, 2)$. In contrast with the first setting, the decomposition into classes is not due to the nontrivial topology of the domain (\mathbb{S}^N in the first setting), but instead it is related to the location and topological degree of the singularities of the maps in each class. More precisely, the classes are defined according to an equivalence relation:

E-mail addresses: brezis@math.rutgers.edu (H. Brézis), mironescu@math.univ-lyon1.fr (P. Mironescu), shafrir@math.technion.ac.il (I. Shafrir).

<http://dx.doi.org/10.1016/j.crma.2016.05.001>

1631-073X/© 2016 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

$u \sim v$ if and only if there exists $\varphi \in W^{1,1}(\Omega; \mathbb{R})$ such that $u = e^{i\varphi} v$. This definition is analogous to the one used in the first part when $N = 1$, see [Remark 1](#) below. This is an indication of a deep connection between the two parts. Actually, we are going to see (in [Remark 5](#)) that the results concerning $W^{1,1}(\mathbb{S}^1; \mathbb{S}^1)$ in [Section 2](#) can be viewed as special cases of analogous results about $W^{1,1}(\Omega; \mathbb{S}^1)$ in [Section 3](#).

2. Distances between homotopy classes of $W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$

The first author and L. Nirenberg [\[6\]](#) developed a concept of topological degree for map in $VMO(\mathbb{S}^N; \mathbb{S}^N)$, $N \geq 1$, which applies in particular to the (integer or fractional) Sobolev spaces $W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$ with

$$s > 0, 1 \leq p < \infty \text{ and } sp \geq N. \quad (1)$$

We will make assumption [\(1\)](#) throughout this section. We define an equivalence relation on $W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$ by $f \sim g$ if and only if $\deg f = \deg g$. It is known that the homotopy classes of $W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$ are precisely the equivalence classes for the relation $f \sim g$ and are given by

$$\mathcal{E}_d := \{f \in W^{s,p}(\mathbb{S}^N; \mathbb{S}^N); \deg f = d\} \text{ where } d \in \mathbb{Z}; \quad (2)$$

these classes depend not only on d , but also on s and p , but, in order to keep notations simple, we do not mention the dependence on s and p .

Remark 1. When $N = 1$ there is an alternative description of the equivalence relation $f \sim g$. Given $f, g \in W^{s,p}(\mathbb{S}^1; \mathbb{S}^1)$, we have (see [\[2\]](#)):

$$f \sim g \text{ if and only if } f = e^{i\varphi} g \text{ for some } \varphi \in W^{s,p}(\mathbb{S}^1; \mathbb{R}). \quad (3)$$

Therefore, it makes sense to denote also $\mathcal{E}_d = \mathcal{E}(f)$ when $\deg f = d$. We shall use this notation in [Remark 2](#) below.

Our purpose is to investigate the usual distance and the Hausdorff distance (in $W^{s,p}$) between the classes \mathcal{E}_d . For that matter we introduce the $W^{s,p}$ -distance between two maps $f, g \in W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$ by

$$d_{W^{s,p}}(f, g) := |f - g|_{W^{s,p}}, \quad (4)$$

where for $h \in W^{s,p}(\mathbb{S}^N; \mathbb{R}^{N+1})$ we let

$$|h|_{W^{s,p}} := \left\| h - \int_{\mathbb{S}^N} h \right\|_{W^{s,p}},$$

and $\| \cdot \|_{W^{s,p}}$ is any one of the standard norms on $W^{s,p}$. Let $d_1 \neq d_2$ and define the following two quantities:

$$\text{dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) := \inf_{f \in \mathcal{E}_{d_1}} \inf_{g \in \mathcal{E}_{d_2}} d_{W^{s,p}}(f, g), \quad (5)$$

and

$$\text{Dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) := \sup_{f \in \mathcal{E}_{d_1}} \inf_{g \in \mathcal{E}_{d_2}} d_{W^{s,p}}(f, g). \quad (6)$$

It is conceivable that

$$\text{Dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = \text{Dist}_{W^{s,p}}(\mathcal{E}_{d_2}, \mathcal{E}_{d_1}), \forall d_1, d_2 \in \mathbb{Z}, \quad (7)$$

but we have not been able to prove this equality (see [Open Problem 1](#) below). Therefore we consider also the symmetric version of [\(6\)](#), which is nothing but the Hausdorff distance between the two classes:

$$H - \text{dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = \max \{ \text{Dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}), \text{Dist}_{W^{s,p}}(\mathcal{E}_{d_2}, \mathcal{E}_{d_1}) \}. \quad (8)$$

The usual distance $\text{dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2})$ in certain (non-fractional) Sobolev spaces was investigated in works by J. Rubinstein and I. Shafrir [\[8\]](#), when $s = 1$, $p \geq N = 1$, and S. Levi and I. Shafrir [\[7\]](#), when $s = 1$, $p \geq N \geq 2$. In particular, they obtained exact formulas for the distance (see [\[8, Remark 2.1\]](#), [\[7, Theorem 3.4\]](#)) and tackled the question whether this distance is achieved (see [\[8, Theorem 1\]](#), [\[7, Theorem 3.4\]](#)).

We pay special attention to the case where $N = 1$ and $s = 1$. In this case, we have several sharp results when we take

$$d_{W^{1,p}}(f, g) = |f - g|_{W^{1,p}} := \left(\int_{\mathbb{S}^1} |\dot{f} - \dot{g}|^p \right)^{1/p}. \quad (9)$$

The following result was obtained in [8] (see also [5]).

$$\begin{aligned} \text{dist}_{W^{1,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) &= \inf_{f \in \mathcal{E}_{d_1}} \inf_{g \in \mathcal{E}_{d_2}} d_{W^{1,p}}(f, g) \\ &= \left(\frac{2}{\pi}\right) \min_{h \in \mathcal{E}_{d_1-d_2}} \left(\int_{\mathbb{S}^1} |\dot{h}|^p\right)^{1/p} \\ &= 2^{(1/p)+1} \pi^{(1/p)-1} |d_1 - d_2|. \end{aligned} \tag{10}$$

In particular,

$$\text{dist}_{W^{1,1}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = 4 |d_1 - d_2|. \tag{11}$$

It is natural to ask whether, given $d_1 \neq d_2$, the infimum in (10) is achieved. The answer is given by the following result, proved in [8] when $p = 2$.

Theorem 1. *Let $N = 1$. Let $d_1, d_2 \in \mathbb{Z}, d_1 \neq d_2$.*

1. *When $p = 1$, the infimum in (10) is always achieved.*
2. *When $1 < p < 2$, the infimum in (10) is achieved if and only if $d_2 = -d_1$.*
3. *When $p \geq 2$, the infimum in (10) is not achieved.*

For $s = 1, N \geq 2, p \geq N$, and for the semi-norm $|f - g|_{W^{1,p}} = \|\nabla f - \nabla g\|_{L^p}$, the exact value of the $W^{1,p}$ distance $\text{dist}_{W^{1,p}}$ between the classes \mathcal{E}_{d_1} and $\mathcal{E}_{d_2}, d_1 \neq d_2$, has been computed by S. Levi and I. Shafrir [7]. By contrast with (10), this distance does not depend on d_1 and d_2 , but only on p (and N).

We now turn to the case $s \neq 1$ and $N \geq 1$. Here, we will only obtain the order of magnitude of the distances $\text{dist}_{W^{s,p}}$, and thus our results are not sensitive to the choice of a specific distance among various equivalent ones. When $0 < s < 1$, a standard distance is associated with the Gagliardo $W^{s,p}$ semi-norm

$$d_{W^{s,p}}(f, g) := \left(\int_{\mathbb{S}^N} \int_{\mathbb{S}^N} \frac{|[f(x) - g(x)] - [f(y) - g(y)]|^p}{|x - y|^{N+sp}} dx dy \right)^{1/p}. \tag{12}$$

We start with $\text{dist}_{W^{s,p}}$.

Theorem 2. *We have*

1. *If $N \geq 1$ and $1 < p < \infty$, then*

$$\text{dist}_{W^{N/p,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = 0, \quad \forall d_1, d_2 \in \mathbb{Z}. \tag{13}$$

2. *If $N = 1, s > 0, 1 \leq p < \infty$ and $sp > 1$, then*

$$C'_{s,p} |d_1 - d_2|^s \leq \text{dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) \leq C_{s,p} |d_1 - d_2|^s. \tag{14}$$

3. *If $N \geq 2, [1 < p < \infty$ and $s > N/p]$ or $[p = 1$ and $s \geq N]$, then*

$$C'_{s,p,N} \leq \text{dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) \leq C_{s,p,N}, \quad \forall d_1, d_2 \in \mathbb{Z} \text{ such that } d_1 \neq d_2. \tag{15}$$

In the above, C, C' are positive constants independent of d_1, d_2 .

We now turn to $\text{Dist}_{W^{s,p}}$.

Theorem 3. *We have*

1. *If $N = 1, s = 1$ and $p = 1$, then*

$$\text{Dist}_{W^{1,1}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = 2\pi |d_1 - d_2|, \quad \forall d_1, d_2 \in \mathbb{Z}. \tag{16}$$

2. *If $N \geq 1$ and $1 \leq p < \infty$, then*

$$\text{Dist}_{W^{N/p,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) \leq C_{p,N} |d_1 - d_2|^{1/p}, \quad \forall d_1, d_2 \in \mathbb{Z}. \tag{17}$$

3. If $N \geq 1$ and $sp > N$, then

$$\text{Dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = \infty, \forall d_1, d_2 \in \mathbb{Z} \text{ such that } d_1 \neq d_2. \tag{18}$$

The detailed proofs appeared in [5]. We call the attention of the reader to a new idea which yields inequality “ \geq ” in (16) in a “more uniform” way; this will become clear in Remark 5 below.

Remark 2. For later use it is convenient to reformulate (11) and (16) as follows. Assume $N = 1, s = 1$ and $p = 1$. Then $\forall f, g \in W^{1,1}(\mathbb{S}^1; \mathbb{S}^1)$ we have

$$\text{dist}_{W^{1,1}}(\mathcal{E}(f), \mathcal{E}(g)) = 4|\text{deg}(f\bar{g})| \tag{19}$$

and

$$\text{Dist}_{W^{1,1}}(\mathcal{E}(f), \mathcal{E}(g)) = 2\pi|\text{deg}(f\bar{g})|. \tag{20}$$

In particular, $\forall f \in W^{1,1}(\mathbb{S}^1; \mathbb{S}^1)$ we have

$$4|\text{deg } f| \leq \text{dist}_{W^{1,1}}(f, \mathcal{E}_0) \leq 2\pi|\text{deg } f|. \tag{21}$$

Moreover the constants 4 and 2π in (21) are optimal.

Here are two natural questions that we could not solve.

Open Problem 1. Is it true that for every $d_1, d_2 \in \mathbb{Z}, N \geq 1, s > 0, 1 \leq p < \infty$,

$$\text{Dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) = \text{Dist}_{W^{s,p}}(\mathcal{E}_{d_2}, \mathcal{E}_{d_1})? \tag{22}$$

Or even better:

$$\text{Does } \text{Dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) \text{ depend only on } |d_1 - d_2| \text{ (and } s, p, N)? \tag{23}$$

There are several cases where we have an explicit formula for $\text{Dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2})$ and in all such cases (23) holds. We may also ask questions similar to (23) for $\text{dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2})$ and for $H - \text{dist}_{W^{s,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2})$ (assuming that the answer to (23) is negative). A striking special case still open when $N = 1$ is: does $\text{dist}_{W^{2,1}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2})$ depend only on $|d_1 - d_2|$?

Open Problem 2. Is it true that for every $N \geq 1$ and every $1 \leq p < \infty$, there exists some $C'_{p,N} > 0$ such that

$$H - \text{dist}_{W^{N/p,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) \geq C'_{p,N}|d_1 - d_2|^{1/p}, \forall d_1, d_2 \in \mathbb{Z} \tag{24}$$

Even better, do we have

$$\text{Dist}_{W^{N/p,p}}(\mathcal{E}_{d_1}, \mathcal{E}_{d_2}) \geq C'_{p,N}|d_1 - d_2|^{1/p}, \forall d_1, d_2 \in \mathbb{Z} \tag{25}$$

Some partial answers to these open problems are presented in [5].

3. Distances between classes in $W^{1,p}(\Omega; \mathbb{S}^1)$

Let Ω be a smooth bounded domain in $\mathbb{R}^N, N \geq 2$. In order to simplify the presentation, we assume throughout that Ω is simply connected; however, many of the results remain valid without this assumption (see, e.g., Remark 5 below). In this section, we decompose $W^{1,p}(\Omega; \mathbb{S}^1)$ into equivalence classes and study their distances. We start with the case $p = 1$ and recall two basic “negative” facts originally discovered by F. Bethuel and X. Zheng [1] (see also [2] for an updated and more detailed presentation).

Fact 1. Maps u of the form $u = e^{i\varphi}$ with $\varphi \in W^{1,1}(\Omega; \mathbb{R})$ belong to $W^{1,1}(\Omega; \mathbb{S}^1)$. However they do not exhaust $W^{1,1}(\Omega; \mathbb{S}^1)$: there exist maps in $W^{1,1}(\Omega; \mathbb{S}^1)$ which cannot be written as $u = e^{i\varphi}$ for some $\varphi \in W^{1,1}(\Omega; \mathbb{R})$. We set

$$X = \{u \in W^{1,1}(\Omega; \mathbb{S}^1); u = e^{i\varphi} \text{ for some } \varphi \in W^{1,1}(\Omega; \mathbb{R})\}. \tag{26}$$

Fact 2. Maps in $C^\infty(\overline{\Omega}; \mathbb{S}^1)$ are not dense in $W^{1,1}(\Omega; \mathbb{S}^1)$. In fact (see, e.g., [2]) we have

$$X = \overline{C^\infty(\overline{\Omega}; \mathbb{S}^1)}^{W^{1,1}}. \tag{27}$$

We now introduce an equivalence relation in $W^{1,1}(\Omega; \mathbb{S}^1)$:

$$u \sim v \text{ if and only if } u = e^{i\varphi} v \text{ for some } \varphi \in W^{1,1}(\Omega; \mathbb{R}). \tag{28}$$

We denote by $\mathcal{E}(u)$ the equivalence class of an element $u \in W^{1,1}(\Omega; \mathbb{S}^1)$. In particular, if $u = 1$, then $\mathcal{E}(u) = X$.

A useful device for constructing maps in the same equivalence class is the following (see [4]). Let $T \in \text{Lip}(\mathbb{S}^1; \mathbb{S}^1)$ be a map of degree one. Then

$$T \circ u \sim u \quad \forall u \in W^{1,1}(\Omega; \mathbb{S}^1). \tag{29}$$

With each $u \in W^{1,1}(\Omega; \mathbb{S}^1)$, we associate a number $\Sigma(u) \geq 0$ defined by

$$\Sigma(u) = \inf_{v \in \mathcal{E}(u)} \int_{\Omega} |\nabla v|. \tag{30}$$

Note that

$$\Sigma(u) = \inf_{\psi \in W^{1,1}(\Omega; \mathbb{R})} \int_{\Omega} |u \wedge \nabla u - \nabla \psi|. \tag{31}$$

This follows from the identities

$$(uv) \wedge \nabla(uv) = u \wedge \nabla v + v \wedge \nabla u \quad \forall u, v \in W^{1,1}(\Omega; \mathbb{S}^1), \tag{32}$$

$$e^{i\varphi} \wedge \nabla(e^{i\varphi}) = \nabla \varphi \quad \forall \varphi \in W^{1,1}(\Omega; \mathbb{R}), \tag{33}$$

$$\bar{u} \wedge \nabla \bar{u} = -u \wedge \nabla u \quad \forall u \in W^{1,1}(\Omega; \mathbb{S}^1). \tag{34}$$

The quantity $\Sigma(u)$ was originally introduced in [3] when $N = 2$. It plays an extremely important role in many questions involving $W^{1,1}(\Omega; \mathbb{S}^1)$ (see [2]). In some sense, it measures how much a given $u \in W^{1,1}(\Omega; \mathbb{S}^1)$ “deviates” from X . By (30) we have, $u \sim v \implies \Sigma(u) = \Sigma(v)$. Moreover we have (see [2]):

$$u \sim 1 \iff \Sigma(u) = 0, \tag{35}$$

and

$$\Sigma(u\bar{v}) \geq |\Sigma(u) - \Sigma(v)| \quad \forall u, v \in W^{1,1}(\Omega; \mathbb{S}^1). \tag{36}$$

Given $u_0, v_0 \in W^{1,1}(\Omega; \mathbb{S}^1)$ such that u_0 is not equivalent to v_0 , it is of interest to consider the distance of u_0 to $\mathcal{E}(v_0)$ defined by

$$d_{W^{1,1}}(u_0, \mathcal{E}(v_0)) = \inf_{v \sim v_0} \int_{\Omega} |\nabla u_0 - \nabla v|,$$

and define, analogously to (5)–(6),

$$\begin{aligned} \text{dist}_{W^{1,1}}(\mathcal{E}(u_0), \mathcal{E}(v_0)) &:= \inf_{u \sim u_0} d_{W^{1,1}}(u, \mathcal{E}(v_0)) \\ &= \inf_{u \sim u_0} \inf_{v \sim v_0} \int_{\Omega} |\nabla(u - v)|, \end{aligned} \tag{37}$$

$$\begin{aligned} \text{Dist}_{W^{1,1}}(\mathcal{E}(u_0), \mathcal{E}(v_0)) &:= \sup_{u \sim u_0} d_{W^{1,1}}(u, \mathcal{E}(v_0)) \\ &= \sup_{u \sim u_0} \inf_{v \sim v_0} \int_{\Omega} |\nabla(u - v)|. \end{aligned} \tag{38}$$

The next theorem provides explicit formulas for these two quantities.

Theorem 4. For every $u_0, v_0 \in W^{1,1}(\Omega; \mathbb{S}^1)$, we have

$$\text{dist}_{W^{1,1}}(\mathcal{E}(u_0), \mathcal{E}(v_0)) = \frac{2}{\pi} \Sigma(u_0 \bar{v}_0) \tag{39}$$

and

$$\text{Dist}_{W^{1,1}}(\mathcal{E}(u_0), \mathcal{E}(v_0)) = \Sigma(u_0 \bar{v}_0). \tag{40}$$

The two assertions in Theorem 4 look very simple but the proofs are quite tricky (see [4]). Note in particular that it follows from (40) that $\text{Dist}_{W^{1,1}}$ is symmetric, which is not clear from its definition (compare with Open Problem 1).

Remark 3. There is an alternative point of view on the equivalence relation $u \sim v$ using the Jacobian of u . For every $u \in W^{1,1}(\Omega; \mathbb{S}^1)$ consider the antisymmetric matrix with coefficients in \mathcal{D}' defined by

$$Ju := \frac{1}{2} \left[\frac{\partial}{\partial x_i} \left(u \wedge \frac{\partial u}{\partial x_j} \right) - \frac{\partial}{\partial x_j} \left(u \wedge \frac{\partial u}{\partial x_i} \right) \right].$$

One can show (see [2]) that $\forall u, v \in W^{1,1}(\Omega; \mathbb{S}^1)$,

$$u \sim v \text{ if and only if } J(u\bar{v}) = Ju - Jv = 0. \tag{41}$$

Remark 4. In order to have a feeling for the equivalence relation $u \sim v$, it is instructive to understand what it means, when $N = 2$ and Ω is simply connected, for $u, v \in \mathcal{R}$ where

$$\mathcal{R} = \{u \in W^{1,1}(\Omega; \mathbb{S}^1); u \text{ is smooth except at a finite number of points}\}.$$

The class \mathcal{R} plays an important role since it is dense in $W^{1,1}(\Omega; \mathbb{S}^1)$ (see [1,2]). If $u \in \mathcal{R}$ then $Ju = \pi \sum k_j \delta_{a_j}$, where a_j are the singular points of u and $k_j = \text{deg}(u, a_j)$. In particular, when $u, v \in \mathcal{R}$, then $u \sim v$ if and only if u and v have the same singularities and the same degree for each singularity.

A special case of interest is the distance of a given $u \in W^{1,1}(\Omega; \mathbb{S}^1)$ to the class $\mathcal{E}(1) = X = \overline{C^\infty(\overline{\Omega}; \mathbb{S}^1)}^{W^{1,1}}$ (see (26)–(27)) that we denote for convenience

$$d(u, X) = d_{W^{1,1}}(u, X) = \inf \left\{ \int_{\Omega} |\nabla u - \nabla(e^{i\varphi})|; \varphi \in W^{1,1}(\Omega; \mathbb{R}) \right\}. \tag{42}$$

An immediate consequence of Theorem 4 is that for every $u \in W^{1,1}(\Omega; \mathbb{S}^1)$, we have

$$\frac{2}{\pi} \Sigma(u) \leq d(u, X) \leq \Sigma(u), \tag{43}$$

and the bounds are optimal in the sense that

$$\sup_{\Sigma(u) > 0} \frac{d(u, X)}{\Sigma(u)} = 1, \tag{44}$$

and

$$\inf_{\Sigma(u) > 0} \frac{d(u, X)}{\Sigma(u)} = \frac{2}{\pi}. \tag{45}$$

The proof of (40) actually provides an explicit recipe for constructing “maximizing sequences” for $\text{Dist}_{W^{1,1}}$. In order to describe it, we first introduce, for each $n \geq 3$, a map $T_n \in \text{Lip}(\mathbb{S}^1; \mathbb{S}^1)$ with $\text{deg } T_n = 1$ by $T_n(e^{i\theta}) = e^{i\tau_n(\theta)}$, with τ_n defined on $[0, 2\pi]$ by setting $\tau_n(0) = 0$ and

$$\tau'_n(\theta) = \begin{cases} n, & \theta \in [2j\pi/n^2, (2j+1)\pi/n^2] \\ -(n-2), & \theta \in ((2j+1)\pi/n^2, (2j+2)\pi/n^2] \end{cases}, \quad j = 0, 1, \dots, n^2 - 1. \tag{46}$$

A basic ingredient in the proof of (40) in Theorem 4 is the following Theorem.

Theorem 5. For every $u_0, v_0 \in W^{1,1}(\Omega; \mathbb{S}^1)$ such that $u_0 \not\sim v_0$, we have

$$\lim_{n \rightarrow \infty} \frac{d_{W^{1,1}}(T_n \circ u_0, \mathcal{E}(v_0))}{\Sigma(u_0 \bar{v}_0)} = 1 \tag{47}$$

and the limit is uniform over all such u_0 and v_0 .

From (29), it is clear that Theorem 5 implies inequality “ \geq ” in (40). The inequality “ \leq ” in (40) is an immediate consequence of the following result established in [4]:

$$d_{W^{1,1}}(u, \mathcal{E}(v_0)) \leq \Sigma(u_0 \bar{v}_0), \quad \forall u_0, v_0 \in W^{1,1}(\Omega; \mathbb{S}^1), \quad \forall u \in \mathcal{E}(u_0). \tag{48}$$

We now discuss briefly the proof of [Theorem 5](#). Inequality “ \leq ” in [\(47\)](#) is a consequence of [\(48\)](#). The heart of the proof of inequality “ \geq ” in [\(47\)](#) is the next lemma.

Lemma 6. For each $\delta > 0$ there exists $n_1 = n_1(\delta)$ such that for every $u, v \in W^{1,1}(\Omega; \mathbb{S}^1)$ and $n \geq n_1$ there holds

$$\int_{\Omega} |\nabla(T_n \circ u) - \nabla v| \geq (1 - \delta)\Sigma(u\bar{v}).$$

Remark 5. Note the similarity between the definitions of $f \sim g$ in $W^{s,p}(\mathbb{S}^1; \mathbb{S}^1)$ (see [\(3\)](#)) and $u \sim v$ in $W^{1,1}(\Omega; \mathbb{S}^1)$ (see [\(28\)](#)) and also the analogy between [\(19\)–\(20\)](#) and [Theorem 4](#) where $|\deg f|$ plays a role similar to $\Sigma(u)$. In fact, the analogy goes beyond the formal resemblance of the formulas. In the above, we could replace Ω by a manifold \mathcal{M} (with or without boundary, simply connected or not). [Theorem 4](#) holds as is and this is the case also for [Theorem 5](#) and [Lemma 6](#). Choosing $\mathcal{M} = \mathbb{S}^1$ one sees easily that $\Sigma(u) = 2\pi|\deg u|$, for all $u \in W^{1,1}(\mathbb{S}^1; \mathbb{S}^1)$. Indeed, denoting $k = \deg u$, we have, on the one hand, for $w(z) = z^k$, $\int_{\mathbb{S}^1} |\dot{w}| = \int_{\mathbb{S}^1} |w \wedge \dot{w}| = 2\pi|k|$, and, on the other hand, for all $v \in \mathcal{E}(u)$,

$$\int_{\mathbb{S}^1} |\dot{v}| = \int_{\mathbb{S}^1} |v \wedge \dot{v}| \geq \left| \int_{\mathbb{S}^1} v \wedge \dot{v} \right| = 2\pi|k|.$$

Hence some results from [Section 2](#) about $W^{1,1}(\mathbb{S}^1; \mathbb{S}^1)$ become special cases of [Theorem 4](#). It is interesting to write explicitly the statement of [Theorem 5](#) for the special case $\Omega = \mathbb{S}^1$. It improves upon [\[5, Lemma 3.1\]](#) by providing a “more uniform” estimate:

For all $f, g \in W^{1,1}(\mathbb{S}^1; \mathbb{S}^1)$ with $\deg f \neq \deg g$, we have

$$\lim_{n \rightarrow \infty} \frac{d_{W^{1,1}}(T_n \circ f, \mathcal{E}(g))}{2\pi|\deg f - \deg g|} = 1$$

and the limit is *uniform* over all such f and g .

There are many challenging open problems concerning the question whether the supremum and the infimum in various formulas above are achieved. Here are some brief comments, restricted to the case $N = 2$; we refer to [\[2,4\]](#) for further discussions.

(i) The question whether the infimum in [\(30\)](#) is achieved is extensively studied in [\[2\]](#). The answer is delicate and depends heavily on Ω and u .

(ii) Concerning the infimum in [\(42\)](#), the answer is positive when Ω is the unit disc and $u(x) = \frac{x}{|x|}$, and in some other cases satisfying $d(u, X) = (2/\pi)\Sigma(u)$ (see [\[4\]](#)). In general, the question is widely open.

(iii) Concerning the infimum in [\(45\)](#), the answer seems to depend on the shape of Ω . We know that when Ω is the unit disc, the infimum in [\(45\)](#) is achieved by $u(x) = \frac{x}{|x|}$. On the other hand, it seems plausible that if Ω is the interior of a non-circular ellipse, then the infimum in [\(45\)](#) is not achieved.

(iv) The question whether the supremum in [\(44\)](#) is achieved is widely open. We suspect that the supremum in [\(44\)](#) is achieved in *every* domain, but we do not know *any* domain in which the supremum is achieved.

Finally, we turn to the classes in $W^{1,p}(\Omega; \mathbb{S}^1)$, $1 < p < 2$, defined in the same way as in the $W^{1,1}$ -case. The distances between the classes are defined analogously to [\(37\)–\(38\)](#) by

$$\text{dist}_{W^{1,p}}(\mathcal{E}(u_0), \mathcal{E}(v_0)) := \inf_{u \sim u_0} \inf_{v \sim v_0} \|\nabla(u - v)\|_{L^p(\Omega)} \tag{49}$$

and

$$\text{Dist}_{W^{1,p}}(\mathcal{E}(u_0), \mathcal{E}(v_0)) := \sup_{u \sim u_0} \inf_{v \sim v_0} \|\nabla(u - v)\|_{L^p(\Omega)}. \tag{50}$$

We first establish a lower bound for $\text{dist}_{W^{1,p}}$:

Proposition 1. For every $u_0, v_0 \in W^{1,p}(\Omega; \mathbb{S}^1)$, we have

$$\text{dist}_{W^{1,p}}(\mathcal{E}(u_0), \mathcal{E}(v_0)) \geq \left(\frac{2}{\pi}\right) \inf_{w \sim u_0 \bar{v}_0} \|\nabla w\|_{L^p(\Omega)}. \tag{51}$$

Remark 6. For $p > 1$ the infimum on the R.H.S. of (51) is actually a minimum, see [2].

We do not know whether the lower bound in (51) is optimal:

Open Problem 3. Is there equality in (51) for every $u_0, v_0 \in W^{1,p}(\Omega; \mathbb{S}^1)$?

We suspect that the answer might be negative in general. We are able to prove that the answer is positive in the case of the distance to smooth maps:

Theorem 7. For every $u_0 \in W^{1,p}(\Omega; \mathbb{S}^1)$, $p \in (1, 2)$, we have

$$\text{dist}_{W^{1,p}}(\mathcal{E}(u_0), \mathcal{E}(1)) = \left(\frac{2}{\pi}\right) \inf_{w \sim u_0 \bar{v}_0} \|\nabla w\|_{L^p(\Omega)}. \quad (52)$$

By analogy with item 3 in Theorem 3, we have the following result:

Theorem 8. For every $u_0, v_0 \in W^{1,p}(\Omega; \mathbb{S}^1)$ such that $u_0 \not\sim v_0$, we have

$$\text{Dist}_{W^{1,p}}(\mathcal{E}(u_0), \mathcal{E}(v_0)) = \infty. \quad (53)$$

The detailed proofs of the results on $W^{1,p}(\Omega; \mathbb{S}^1)$, $p \in [1, 2)$, will appear in [4].

Acknowledgements

The first author (HB) was partially supported by NSF grant DMS-1207793. The second author (PM) was partially supported by the LABEX MILYON (ANR-10-LABX-0070) of Université de Lyon, within the program “Investissements d’Avenir” (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR). The third author (IS) was supported by the Israel Science Foundation (Grant No. 999/13).

References

- [1] F. Bethuel, X.M. Zheng, Density of smooth functions between two manifolds in Sobolev spaces, *J. Funct. Anal.* 80 (1988) 60–75.
- [2] H. Brézis, P. Mironescu, Sobolev maps with values into the circle, Birkhäuser, in preparation.
- [3] H. Brézis, P. Mironescu, A. Ponce, $W^{1,1}$ -maps with values into \mathbb{S}^1 , in: *Geometric Analysis of PDE and Several Complex Variables*, in: *Contemp. Math.*, vol. 368, American Mathematical Society, Providence, RI, USA, 2005, pp. 69–100.
- [4] H. Brézis, P. Mironescu, I. Shafrir, Distances between classes in $W^{1,1}(\Omega; \mathbb{S}^1)$, in preparation.
- [5] H. Brézis, P. Mironescu, I. Shafrir, Distances between homotopy classes of $W^{s,p}(\mathbb{S}^N; \mathbb{S}^N)$, *ESAIM Control Optim. Calc. Var.* (2016), in press, hal-01257581.
- [6] H. Brézis, L. Nirenberg, Degree theory and BMO. I. Compact manifolds without boundaries, *Sel. Math. New Ser.* 1 (1995) 197–263.
- [7] S. Levi, I. Shafrir, On the distance between homotopy classes of maps between spheres, *J. Fixed Point Theory Appl.* 15 (2014) 501–518.
- [8] J. Rubinstein, I. Shafrir, The distance between homotopy classes of \mathbb{S}^1 -valued maps in multiply connected domains, *Isr. J. Math.* 160 (2007) 41–59.