



ELSEVIER

Contents lists available at ScienceDirect

C. R. Acad. Sci. Paris, Ser. I

www.sciencedirect.com



Functional analysis

## Exhaustive families of representations of $C^*$ -algebras associated with $N$ -body Hamiltonians with asymptotically homogeneous interactions

*Familles exhaustives de représentations des  $C^*$ -algèbres associées aux hamiltoniens du problème à  $N$  corps à potentiel asymptotiquement homogène*

Jérémy Mougel, Nicolas Prudhon

Université de Lorraine, UFR MIM, 3, rue Augustin-Fresnel, 57045 Metz, France

## ARTICLE INFO

## Article history:

Received 26 November 2018

Accepted after revision 4 February 2019

Available online 18 February 2019

Presented by the Editorial Board

## ABSTRACT

We continue the analysis of algebras introduced by Georgescu, Nistor, and their coauthors, in order to study  $N$ -body type Hamiltonians with interactions. More precisely, let  $Y \subset X$  be a linear subspace of a finite-dimensional Euclidean space  $X$ , and  $v_Y$  be a continuous function on  $X/Y$  that has uniform homogeneous radial limits at infinity. We consider, in this paper, Hamiltonians of the form  $H = -\Delta + \sum_{Y \in \mathcal{S}} v_Y$ , where the subspaces  $Y \subset X$  belong to some given family  $\mathcal{S}$  of subspaces. Georgescu and Nistor have considered the case when  $\mathcal{S}$  consists of *all* subspaces  $Y \subset X$ , and Nistor and coauthors considered the case when  $\mathcal{S}$  is a finite semilattice and Georgescu generalized these results to any family. In this paper, we develop new techniques to prove their results on the spectral theory of the Hamiltonian to the case where  $\mathcal{S}$  is any family of subspaces also, and extend those results to other operators affiliated to a larger algebra of pseudodifferential operators associated with the action of  $X$  introduced by Connes. In addition, we exhibit Fredholm conditions for such elliptic operators. We also note that the algebras we consider answer a question of Melrose and Singer.

© 2019 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## R É S U M É

Nous poursuivons l'étude des algèbres introduites par Georgescu, Nistor et leurs collaborateurs pour étudier les hamiltoniens du problème à  $N$  corps avec interactions. Plus précisément, soit  $Y \subset X$  un sous-espace d'un espace euclidien  $X$  de dimension finie, et soit  $v_Y$  une fonction continue sur  $X/Y$  possédant des limites radiales à l'infini. Nous considérons ici des hamiltoniens de la forme  $H = -\Delta + \sum_{Y \in \mathcal{S}} v_Y$ , où  $\mathcal{S}$  est une famille donnée de sous-espaces de  $X$ . Georgescu et Nistor ont étudié en détail le cas où la famille  $\mathcal{S}$  contient tous les sous-espaces de  $X$ ; Nistor et ses co-auteurs ont étudié le cas d'une famille finie stable par intersections, et Georgescu a généralisé certains de ces résultats à des familles quelconques. Dans cette note, nous développons de nouvelles techniques pour étendre les

E-mail addresses: [jeremy.mougel@univ-lorraine.fr](mailto:jeremy.mougel@univ-lorraine.fr) (J. Mougel), [nicolas.prudhon@univ-lorraine.fr](mailto:nicolas.prudhon@univ-lorraine.fr) (N. Prudhon).

<https://doi.org/10.1016/j.crma.2019.01.010>

1631-073X/© 2019 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

résultats spectraux précédents à des familles quelconques également et pour des opérateurs pseudo-différentiels (ou affiliés) associés à l'action de  $X$  par translation, introduits par Connes. En outre, nous obtenons, pour les opérateurs elliptiques, une caractérisation comme étant des opérateurs de Fredholm. Nous faisons remarquer également que ces algèbres répondent à une question de Melrose et Singer.

© 2019 Académie des sciences. Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

A new approach to the study of Hamiltonians of  $N$ -body type with interactions that are asymptotically homogeneous at infinity on a finite dimensional Euclidean space  $X$  was initiated by Georgescu and Nistor [3,7,4].

For any finite real vector space  $Z$ , we let  $\bar{Z}$  denote its spherical compactification. A function in  $C(\bar{Z})$  is thus a continuous function on  $Z$  that has uniform radial limits at infinity. Let  $\mathbb{S}_Z$  be the set of half-lines in  $Z$ , that is  $\mathbb{S}_Z := \{\hat{a}, a \in Z, a \neq 0\}$  where  $\hat{a} := \{ra, r > 0\}$ . We identify  $\mathbb{S}_Z = \bar{Z} \setminus Z$ .

For any subspace  $Y \subset X$ ,  $\pi_Y : X \rightarrow X/Y$  denotes the canonical projection. Let

$$H = -\Delta + \sum_{Y \in \mathcal{S}} v_Y, \tag{1}$$

where  $v_Y \in C(\overline{X/Y})$  is seen as a bounded continuous function on  $X$  via the projection  $\pi_Y : X \rightarrow X/Y$ . The sum is over all subspaces  $Y \subset X$ ,  $Y \in \mathcal{S}$  and is assumed to be uniformly convergent. One of the main results of [7,10] describes the essential spectrum of  $H$  extending the celebrated HVZ theorem [14]. The goal of this paper is to explain how these results can be extended to any family of subspaces that contains  $\{0\}$  and to more general operators using  $C^*$ -algebras techniques.

Let  $\mathcal{S}$  be a family of subspaces of  $X$  with  $0 \in \mathcal{S}$ . We define the commutative sub- $C^*$ -algebra  $\mathcal{E}_{\mathcal{S}}(X)$  of the commutative  $C^*$ -algebra  $C_b^u(X)$  of bounded uniformly continuous functions on  $X$  by

$$\mathcal{E}_{\mathcal{S}}(X) = \langle C(\overline{X/Y}), Y \in \mathcal{S} \rangle \subset C_b^u(X). \tag{2}$$

The algebras  $\mathcal{E}_{\mathcal{S}}(X)$  give an answer to a question of Melrose and Singer [9].

**Theorem 1.** *Let  $n$  be an integer. Let  $\mathcal{S}^n$  be the semilattice of subspaces of  $X^n$  generated by  $\mathcal{S}_i^n \cup \mathcal{S}_{ij}^n$  where*

$$\begin{aligned} \mathcal{S}_i^n &= \{(x_1, \dots, x_n) \in X^n; x_i = 0\} \\ \mathcal{S}_{ij}^n &= \{(x_1, \dots, x_n) \in X^n; x_i = x_j\}. \end{aligned}$$

*Then the spectrum  $\Omega_{\mathcal{S}^n}$  of  $\mathcal{E}_{\mathcal{S}^n}(X^n)$  is a compactification of  $X^n$  satisfying the following properties:*

- (1)  $\Omega_{\mathcal{S}^n}$  is the spherical compactification  $\overline{X^n}$ .
- (2) The action of the symmetric group  $\mathfrak{S}_n$  on  $X^n$  extends continuously to  $\Omega_{\mathcal{S}^n}$ .
- (3) The projections  $p_i^{n,k} : X^n \rightarrow X^k$ ,  $p_i^{n,k}(x_1, \dots, x_n) = (x_{i_1}, \dots, x_{i_k})$  extend continuously to  $p_i^{n,k} : \Omega_{\mathcal{S}^n} \rightarrow \Omega_{\mathcal{S}^k}$ .
- (4) The difference maps  $\delta_{ij}(x_1, \dots, x_n) = x_i - x_j$  from  $X^n$  to  $X$  extend continuously to the compactifications.

Actually, the spectrum  $\Omega_{\mathcal{S}^n}$  have very strong connection with the space built by Vasy in [16] and generalized by Kottke in the last section of [8].

The additive group  $X$  acts by translation on  $C_b^u(X)$  and the subalgebra  $\mathcal{E}_{\mathcal{S}}(X)$  is invariant. So a crossed product  $C^*$ -algebra is obtained

$$\mathcal{E}_{\mathcal{S}}(X) \rtimes X, \tag{3}$$

which can be regarded as an algebra of operators on  $L^2(X)$ . Thanks to the assumption  $0 \in \mathcal{S}$ , the algebra  $C_0(X)$  belongs  $\mathcal{E}_{\mathcal{S}}(X)$ . Hence  $C_0(X) \rtimes X$  is contained in  $\mathcal{E}_{\mathcal{S}}(X) \rtimes X$ . It follows from the definition of crossed products algebras that the  $C^*$ -algebra  $\mathcal{E}_{\mathcal{S}}(X) \rtimes X$  is generated by two kinds of operators: multiplication operators  $m_f$  associated with functions  $f \in \mathcal{E}_{\mathcal{S}}(X)$ , and convolution operators

$$C_\phi u(x) := \int_X \phi(y) u(x - y) dy$$

with  $\phi \in C_c(X)$ , a continuous compactly supported function. An immediate computation shows that  $m_f c_\phi$  (resp.  $c_\phi m_f$ ) is a kernel operator with kernel

$$K(x, y) = f(x) \phi(y - x), \quad (\text{resp. } K(x, y) = f(y) \phi(y - x)). \tag{4}$$

**Proposition 2.** (i) *The subalgebra  $C_0(X) \rtimes X$  is the algebra  $\mathcal{K}(X)$  of compact operators on  $L^2(X)$ .  
(ii) *For  $f \in C(\overline{X})$  and  $\phi \in C_c(X)$  the commutator  $[m_f, c_\phi]$  is compact.**

The point (i) is a consequence of equation (4) because the kernel  $K$  has compact support when  $f$  does and the result follows by density. Again, thanks to Eq. (4), one sees that the commutator is a kernel operator with kernel

$$K(x, y) = \phi(y - x)(f(x) - f(y)).$$

Hence, in view of  $\phi \in C_c(X)$ , the support of  $K$  is contained in a band around the diagonal. The distance between the border of the band and the diagonal is bounded. Moreover,  $K$  goes to 0 at infinity because  $f$  has radial limits. So the commutator is a limit of Hilbert–Schmidt operators, and hence is compact.

Recall that a self-adjoint operator  $P$  on  $L^2(X)$  is said to be affiliated to a  $C^*$ -algebra  $A$  of bounded operators on  $L^2(X)$  if for some (and hence any) function  $h \in C_0(\mathbb{R})$ , then  $h(P)$  belongs to  $A$ . For example, it follows from the identity

$$(H + i)^{-1} = (-\Delta + i)^{-1} (1 + V(-\Delta + i)^{-1})^{-1},$$

that  $H$  is affiliated to  $\mathcal{E}_S(X) \rtimes X$ . More generally, for any  $C^*$ -algebra  $A$ , a morphism  $h : C_0(\mathbb{R}) \rightarrow A$  is called an operator affiliated to  $A$ . Following Connes [2] and Baaj [1], we introduce the algebra  $\Psi^\infty(\mathcal{E}_S(X); X)$  of pseudodifferential operators associated with the action of  $X$  on  $\mathcal{E}_S(X)$ . We shall need the  $C^*$ -algebra of  $\Psi\text{DO}(\mathcal{E}_S(X), X)$  given by the norm closure of  $\Psi^0(\mathcal{E}_S(X); X)$  and the exact sequence

$$0 \rightarrow \mathcal{E}_S(X) \rtimes X \rightarrow \Psi\text{DO}(\mathcal{E}_S(X), X) \xrightarrow{\sigma_0} C(\mathbb{S}_X \times \widehat{\mathcal{E}_S(X)}) \rightarrow 0, \tag{5}$$

where  $\sigma_0$  is the principal symbol map. Positive-order pseudodifferential operators are examples of operators affiliated to the algebra of non-positive-order pseudodifferential operators  $\Psi\text{DO}(\mathcal{E}_S(X), X)$ .

Let  $\alpha \in \mathbb{S}_X$ . For each  $x \in X$ , we let  $(T_x f)(y) = f(y - x)$  denote the translation on  $L^2(X)$ . For any operator  $P$  on  $L^2(X)$ , we let

$$\tau_\alpha(P) = \lim_{r \rightarrow +\infty} T_{ra}^* P T_{ra}, \quad \text{if } \alpha = \hat{a} \in \mathbb{S}_X, \tag{6}$$

whenever the strong limit exists.

**Lemma 3.** *For  $f \in C(\overline{X/Y})$ , one has*

$$\tau_\alpha(f)(x) = \begin{cases} f(x) & \text{if } Y \supset \alpha, \\ f(\pi_Y(\alpha)) & \text{else.} \end{cases}$$

We define  $\mathcal{S}_\alpha = \{Y \in \mathcal{S}; \alpha \subset Y\}$ . It follows from the previous lemma that, on  $\mathcal{E}_S(X)$ ,  $\tau_\alpha$  is the projection on the subalgebra  $\mathcal{E}_{\mathcal{S}_\alpha}(X)$ ,

$$\tau_\alpha : \mathcal{E}_S(X) \rightarrow \mathcal{E}_{\mathcal{S}_\alpha}(X).$$

**Theorem 4.**

- (1) *Let  $P$  be a self-adjoint operator affiliated to  $\Psi\text{DO}(\mathcal{E}_S(X), X)$  and  $\alpha = \hat{a} \in \mathbb{S}_X$ . Then the limit  $\tau_\alpha(P) := \lim_{r \rightarrow +\infty} T_{ra}^* P T_{ra}$  exists.*
- (2) *Let  $P \in \Psi\text{DO}(\mathcal{E}_S(X), X)$ . Then  $P$  is a Fredholm operator if and only if  $P$  is elliptic (i.e.  $\sigma_0(P)$  is invertible) and for all  $\alpha \in \mathbb{S}_X$ ,  $\tau_\alpha(P)$  is invertible.*
- (3) *If  $P \in \Psi\text{DO}(\mathcal{E}_S(X), X)$ ,*

$$\text{Spec}_{\text{ess}}(P) = \bigcup_{\alpha \in \mathbb{S}_X} \text{Spec}(\tau_\alpha(P)) \cup \text{Im}(\sigma_0(P)).$$

- (4) *If  $P \in \Psi^m(\mathcal{E}_S(X); X)$ ,  $m > 0$ , is elliptic, then,*

$$\text{Spec}_{\text{ess}}(P) = \bigcup_{\alpha \in \mathbb{S}_X} \text{Spec}(\tau_\alpha(P)).$$

Note that in classical results on the  $N$ -body problem, one usually has the closure of the union in the spectral decomposition. See, however [10,4]. See also [7,10] for related results, where operators affiliated to  $\mathcal{E}_S(X) \rtimes X$  were considered. In [10], only finite semi-lattice  $\mathcal{S}$  are considered. The closure of the union means that the family  $(\tau_\alpha)$  is a faithful family of morphism of  $\mathcal{E}_S(X) \rtimes X$ . The stronger result of [10] is obtained by showing that the family  $(\tau_\alpha \rtimes X)_{\alpha \in \mathbb{S}_X}$  is actually an exhaustive family of representations of  $\mathcal{E}_S(X) \rtimes X$ , when  $\mathcal{S}$  is a finite semi-lattice. In [5], pseudodifferential operators on  $\mathbb{R}$  were considered (see Remark 3.23 of that paper). In the framework of admissible locally compact groups, a decomposition of an essential spectrum involving exhaustive families can be found in [11] [12]. In fact, by [13, Proposition 3.12], exhaustive families are also strictly spectral families in the following sense.

**Definition 5** ([13,15]).

- (1) A family  $(\phi_i)_{i \in I}$  of morphisms of a  $C^*$ -algebra  $A$  is said to be *exhaustive* if any primitive ideal contains at least  $\ker \phi_i$  for some  $i \in I$ .
- (2) A family  $(\phi_i)_{i \in I}$  of morphisms of a unital  $C^*$ -algebra  $A$  is said to be *strictly spectral* if

$$(\forall a \in A) \quad \text{Spec}(a) = \cup_{i \in I} \text{Spec}(\phi_i(a)).$$

**Theorem 6.** *Let  $\mathcal{S}$  be a family of subspaces of  $X$  with  $0 \in \mathcal{S}$ . Then the family  $(\tau_\alpha \rtimes X)_{\alpha \in \mathbb{S}_X}$  is an exhaustive family of  $\mathcal{E}_{\mathcal{S}}(X) \rtimes X/\mathcal{K}(X)$ .*

Let us prove this result. Let  $\pi$  be an irreducible representation of  $\mathcal{E}_{\mathcal{S}}(X) \rtimes X/\mathcal{K}(X)$ . It extends to an irreducible representation of  $\mathcal{E}_{\mathcal{S}}(X) \rtimes X$  as well as to their multipliers algebras  $\mathcal{M}(\mathcal{E}_{\mathcal{S}}(X) \rtimes X/\mathcal{K}(X))$  and  $\mathcal{M}(\mathcal{E}_{\mathcal{S}}(X) \rtimes X)$ . By Proposition 2(i), one obtains the following commutative diagram:

$$\begin{array}{ccccc}
 C(\overline{X}) & \hookrightarrow & \mathcal{E}_{\mathcal{S}}(X) & \longrightarrow & \mathcal{M}(\mathcal{E}_{\mathcal{S}}(X) \rtimes X) \\
 \downarrow p & & & & \downarrow \\
 C(\mathbb{S}_X) & \xrightarrow{\phi} & \mathcal{M}(\mathcal{E}_{\mathcal{S}}(X) \rtimes X/\mathcal{K}(X)) & \xrightarrow{\pi} & \mathcal{B}(\mathcal{H}_\pi)
 \end{array} \tag{7}$$

**Lemma 7.** *The image  $\phi(C(\mathbb{S}_X))$  is central in  $\mathcal{M}(\mathcal{E}_{\mathcal{S}}(X) \rtimes X/\mathcal{K}(X))$ .*

In fact, it is enough to show that any  $f \in C(\overline{X})$  commutes with any element of  $\mathcal{E}_{\mathcal{S}}(X) \rtimes X$  modulo a compact operator. But the result is true on the generators by Proposition 2(ii), so the lemma follows by density.

By the Schur Lemma, we deduce that  $\pi \circ \phi$  is a character of  $C(\mathbb{S}_X)$ . Hence there exists some  $\alpha \in \mathbb{S}_X$  such that  $\pi|_{C(\overline{X})} = \chi_\alpha I$ , where  $\chi_\alpha$  is the character of  $C(\overline{X})$  given by the evaluation at  $\alpha \in \mathbb{S}_X$ .

**Proposition 8.** *One has  $\ker \tau_\alpha = (\ker \chi_\alpha)\mathcal{E}_{\mathcal{S}}(X)$ .*

**Proof.** We need to show that  $\mathcal{E}_{\mathcal{S}}(X)/\ker \tau_\alpha = \mathcal{E}_{\mathcal{S}_\alpha}(X)$  and  $\mathcal{E}_{\mathcal{S}}(X)/(\ker \chi_\alpha)\mathcal{E}_{\mathcal{S}}(X)$  have the same characters. By definition, for any character  $\chi$  of  $\mathcal{E}_{\mathcal{S}_\alpha}(X)$ , there exists a unique character  $\chi'$  of  $\mathcal{E}_{\mathcal{S}}(X)$  such that  $\chi' = \chi \circ \tau_\alpha$ . In view of Lemma 3, this is equivalent to the following:

$$(\forall Y \in \mathcal{S}, \alpha \notin Y, \forall u \in C(\overline{X/Y})) \quad \chi(u) = u(\pi_Y(\alpha)). \tag{8}$$

In particular, for  $Y = 0$ , we see that  $\chi|_{C(\overline{X})} = \chi_\alpha$ . Reciprocally, it follows from [7, Lemma 6.7] that if  $\chi|_{C(\overline{X})} = \chi_\alpha$ , then relation (8) is true. On the other hand, the characters of  $\mathcal{E}_{\mathcal{S}}(X)/(\ker \chi_\alpha)\mathcal{E}_{\mathcal{S}}(X)$  are precisely the characters  $\chi$  of  $\mathcal{E}_{\mathcal{S}}(X)$  such that  $\chi|_{C(\overline{X})} = \chi_\alpha$ . So  $\ker \tau_\alpha = (\ker \chi_\alpha)\mathcal{E}_{\mathcal{S}}(X)$  as claimed.  $\square$

Now if  $\pi|_{C(\overline{X})} = \chi_\alpha$ , one has  $\ker \pi \supset (\ker \chi_\alpha)\mathcal{E}_{\mathcal{S}}(X) = \ker \tau_\alpha$ . Finally,

$$\ker(\tau_\alpha \rtimes X) = (\ker \tau_\alpha) \rtimes X \subset \ker \pi.$$

It follows that  $(\tau_\alpha \rtimes X)_{\alpha \in \mathbb{S}_X}$  is an exhaustive family of morphisms.

**Remark 9.** The results presented here can easily be extended to pseudodifferential operators with matrix coefficients. For example, Dirac operators  $D_V = D + V$ , with potentials  $V$  as in (1), may be considered to satisfy the condition of Theorem 4.

See also [6, Example 6.35] for others physical interesting operators.

**Acknowledgements**

The authors thank Vladimir Georgescu and Victor Nistor for useful discussions.

**References**

- [1] S. Baaj, Calcul pseudo-différentiel et produits croisés de  $C^*$ -algèbres. I, C. R. Acad. Sci. Paris, Ser. I 307 (11) (1988) 581–586.
- [2] A. Connes,  $C^*$  algèbres et géométrie différentielle, C. R. Acad. Sci. Paris, Ser. A–B 290 (13) (1980) A599–A604.
- [3] V. Georgescu, On the structure of the essential spectrum of elliptic operators on metric spaces, J. Funct. Anal. 260 (6) (2011) 1734–1765.
- [4] V. Georgescu, On the essential spectrum of elliptic differential operators, J. Math. Anal. Appl. 468 (2) (2018) 839–864.

- [5] V. Georgescu, A. Iftimovici,  $C^*$ -algebras of quantum Hamiltonians, in: *Operator Algebras and Mathematical Physics*, Constanța, 2001, Theta, Bucharest, 2003, pp. 123–167.
- [6] V. Georgescu, A. Iftimovici, Localizations at infinity and essential spectrum of quantum Hamiltonians. I. General theory, *Rev. Math. Phys.* 18 (4) (2006) 417–483.
- [7] V. Georgescu, V. Nistor, On the essential spectrum of  $N$ -body Hamiltonians with asymptotically homogeneous interactions, *J. Oper. Theory* 77 (2) (2017) 333–376.
- [8] C. Kottke, Functorial compactification of linear spaces, arXiv:1712.03902, 2017.
- [9] R. Melrose, M. Singer, Scattering configuration spaces, arXiv:0808.2022, 2008.
- [10] J. Mougel, V. Nistor, N. Prudhon, A refined HVZ-theorem for asymptotically homogeneous interactions and finitely many collision planes, *Rev. Roum. Math. Pures Appl.* 62 (1) (2017) 287–308.
- [11] M. Măntoiu,  $C^*$ -algebras, dynamical systems at infinity and the essential spectrum of generalized Schrödinger operators, *J. Reine Angew. Math.* 550 (2002) 211–229.
- [12] M. Măntoiu, Essential spectrum and Fredholm properties for operators on locally compact groups, *J. Oper. Theory* 77 (2) (2017) 481–501.
- [13] V. Nistor, N. Prudhon, Exhaustive families of representations and spectra of pseudodifferential operators, *J. Oper. Theory* 78 (2) (2017) 247–279.
- [14] M. Reed, B. Simon, *Methods of Modern Mathematical Physics. IV. Analysis of Operators*, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1978.
- [15] S. Roch, Algebras of approximation sequences: structure of fractal algebras, in: *Singular Integral Operators, Factorization and Applications*, in: *Operator Theory: Advances and Applications*, vol. 142, Birkhäuser, Basel, 2003, pp. 287–310.
- [16] A. Vasy, Propagation of singularities in many-body scattering, *Ann. Sci. Éc. Norm. Supér.* (4) 34 (3) (2001) 313–402.