

CONVEX COMBINATIONS OF LOW EIGENVALUES, FRAENKEL ASYMMETRIES AND ATTAINABLE SETS *

DARIO MAZZOLENI¹ AND DAVIDE ZUCCO^{2,3}

Abstract. We consider the problem of minimizing convex combinations of the first two eigenvalues of the Dirichlet–Laplacian among open sets of \mathbb{R}^N of fixed measure. We show that, by purely elementary arguments, based on the minimality condition, it is possible to obtain informations on the *geometry* of the minimizers of convex combinations: we study, in particular, when these minimizers are no longer convex, and the optimality of balls. As an application of our results we study the boundary of the attainable set for the Dirichlet spectrum. Our techniques involve symmetry results *à la* Serrin, explicit constants in quantitative inequalities, as well as a purely geometrical problem: the minimization of the Fraenkel 2-asymmetry among *convex* sets of fixed measure.

Mathematics Subject Classification. 47A75, 49G05, 49Q10, 49R05.

Received November 27, 2015. Revised April 5, 2016. Accepted March 30, 2016.

1. INTRODUCTION

Spectral optimization problems have received a lot of attention in the last years, with a particular emphasis to extremum problems for eigenvalues of elliptic operators (see the books [11, 23, 25]). A typical problem consists in the minimization of a functional defined in terms of the eigenvalues of the Laplace operator among sets of fixed measure. Here to simplify the exposition we will always consider the measure constraint equal one. The first issue for this kind of problems concerns the *existence* of an optimal shape: a result proved in the 1990s by Buttazzo and Dal Maso [16] is even now a cornerstone of the matter, and, for a large class of functionals, it ensures the existence of a solution in the class of quasi-open sets of fixed measure (*a priori* contained into a given box, which provides the necessary compactness to prove existence). Moreover, the *regularity* of an optimal shape is a highly difficult problem and a general regularity theory is nowadays not available: even a proof which guarantees that an optimal shape is open, and not merely quasi-open, is far from being trivial, see [15]. Another important point consists in proving some *geometric* properties of optimal shapes, such as connectedness, convexity, symmetry with respect to some axis, and so on. By the way, only for few special functionals optimal

Keywords and phrases. Eigenvalues, Dirichlet Laplacian, Fraenkel asymmetry, attainable set.

* *The first author was supported by the ERC advanced grant n. 339958 “COMPAT”. The second author was supported by the INdAM-GNAMPA through the Project 2015 “Critical phenomena in the mechanics of materials: a variational approach”.*

¹ Dipartimento di Matematica G. Peano, Università degli Studi di Torino, via Carlo Alberto, 10–10123 Torino, Italy. dariocesaresevero.mazzoleni@unito.it

² Scuola Internazionale Superiore di Studi Avanzati, via Bonomea, 265–34136 Trieste, Italy. davide.zucco@sissa.it

³ Dipartimento di Scienze Matematiche G.L. Lagrange, Politecnico di Torino, Corso Duca degli Abruzzi, 24–10129 Torino, Italy. davide.zucco@polito.it

shapes are explicitly known: classical examples are the lowest eigenvalues of the Dirichlet–Laplacian. We recall that, for a given integer $N \geq 2$ and an open set $\Omega \subset \mathbb{R}^N$ with finite measure, the *first* and *second eigenvalues* of the Dirichlet–Laplacian can be defined as

$$\lambda_1(\Omega) := \min_{u \in H_0^1(\Omega) \setminus \{0\}} \frac{\int_{\Omega} |\nabla u(x)|^2 dx}{\int_{\Omega} |u(x)|^2 dx}, \quad \lambda_2(\Omega) := \min_{\substack{u \in H_0^1(\Omega) \setminus \{0\} \\ \int_{\Omega} uu_1 = 0}} \frac{\int_{\Omega} |\nabla u(x)|^2 dx}{\int_{\Omega} |u(x)|^2 dx},$$

where these minima are attained, respectively, by the *first* and *second eigenfunctions* u_1 and u_2 (which are unique, up to a multiplicative constant).

The interest in the minimization of the first eigenvalue goes back to a conjecture due to Lord Rayleigh in 1877, then proved by Faber and Krahn in the 1920s. The *Faber–Krahn inequality* asserts that of all open sets of fixed measure, the ball has the minimum first eigenvalue: in formula, for every open set $\Omega \subset \mathbb{R}^N$ with unit measure

$$\lambda_1(\Omega) \geq \lambda_1(B) = \omega_N^{2/N} j_{N/2-1}^2, \tag{1.1}$$

where ω_N denotes the measure of the ball in \mathbb{R}^N with unit radius, j_n the first positive zero of the Bessel function J_n , and B the open ball of unit measure in \mathbb{R}^N . Equality in (1.1) holds if and only if Ω is that ball (up to sets of capacity zero). The same issue for the second eigenvalue is known as the *Krahn–Szegő inequality*, which asserts that two disjoint open balls of half measure each are the unique (up to sets of capacity zero) minimizer, namely for every open set $\Omega \subset \mathbb{R}^N$ with unit measure

$$\lambda_2(\Omega) \geq \lambda_2(B_- \cup B_+) = 2^{2/N} \lambda_1(B) = (2\omega_N)^{2/N} j_{N/2-1}^2, \tag{1.2}$$

where $B_- \cup B_+$ is the union of two equal and disjoint open balls of half measure each, and equality in (1.2) holds if and only if $\Omega = B_- \cup B_+$.

Starting with the important work of Keller and Wolf [27], there was a strong interest for *convex combinations* of the first two eigenvalues of the Dirichlet–Laplacian, namely the functional F_t defined, for every $t \in (0, 1)$, as

$$F_t(\Omega) := t\lambda_1(\Omega) + (1 - t)\lambda_2(\Omega), \tag{1.3}$$

where $\Omega \subset \mathbb{R}^N$ is an open set of finite measure. Then, the corresponding spectral optimization problem writes as

$$\min \{ F_t(\Omega) : \Omega \subset \mathbb{R}^N, \Omega \text{ open}, |\Omega| = 1 \}. \tag{1.4}$$

The existence of a minimizer for this problem is now well understood and is guaranteed by a general theory recently developed in the works [10, 15, 29], all based on the above mentioned result [16], but with the new difficulty of working in the entire space \mathbb{R}^N . Notice that, all these results guarantee the existence of an optimal shape in the larger class of quasi-open sets, and only *a posteriori* one proves that a minimizer of problem (1.4) is in fact *open*, and so problem (1.4) is well-posed. Moreover, in [28] it was proved that, for every $t \in (0, 1)$, minimizers of (1.4) are *connected* (more generally, this topological property was studied for minimizers of convex combinations of the first three eigenvalues). In two dimensions ($N = 2$), some numerical computations on the shape of these minimizers appeared in [26]. We sum up all these results in the following theorem.

Theorem 1.1. *For every $t \in (0, 1)$, there exists a minimizer in (1.4). Moreover, every minimizer Ω_t is a connected set of finite perimeter with uniformly bounded diameter (depending only on the dimension N).*

Our goal here is to show that, by purely elementary arguments essentially based on the minimality condition, it is possible to obtain interesting informations on the geometry of the minimizers for problem (1.4), and to recover some known results on the boundary of the attainable set for the Dirichlet spectrum (see [2, 3, 9, 27]).

Notice that, if $t = 1$ the convex combination (1.3) is minimized by the ball with unit measure (because of the Faber–Krahn inequality (1.1)), while if $t = 0$, by two equal balls of half measure each (because of the

Krahn–Szegő inequality (1.2)). Therefore, as t moves from 1 to 0, one expects the shape of a minimizer Ω_t deforming from a ball of unit measure to two balls of half measure each; in particular, it is natural to conjecture that at some value of t the convexity of all the minimizers in (1.4) is lost (as was numerically observed in [26], in two dimensions, the critical value for t is expected to be 1/2). This question also appeared in [23] as the Open Problem 21. We give a first answer to this question, though non-optimal. All the results of this paper, unless otherwise specified, will hold in every dimension $N \geq 2$.

Theorem 1.2. *There exists a threshold $T > 0$ such that, for all $t \in (0, T)$, every minimizer in (1.4) is no longer convex.*

We provide a *quantitative* proof of this theorem, namely we explicitly construct the threshold T via the eigenvalues of the Dirichlet–Laplacian (3.12) (cf. with [14], Sect. 5.3 where a similar question was analyzed in a different context, and whose strategy of proof could be adapted to prove Theorem 1.2, however without getting an explicit value for T). To be more concrete, in two dimensions, we provide a numerical lower bound on T using a quantitative Krahn–Szegő inequality involving the so-called Fraenkel 2-asymmetry. Therefore, we are naturally led to consider a purely geometrical problem, which is probably the most innovative part of the paper: the minimization of the Fraenkel 2-asymmetry among *convex* sets of given area. We show that the *mobile*, i.e., the intersection of the convex hull of two tangent balls with a strip, see Definition 2.2, is the unique minimizer satisfying an isoperimetric inequality for the Fraenkel 2-asymmetry (2.10). An explicit value for the constant in the quantitative Krahn–Szegő inequality will be also needed. This opens a new area of application for quantitative inequalities.

As second question, we analyze the optimality of a special convex set: the ball, generalizing a result from [27].

Theorem 1.3. *For all $t \in (0, 1)$ the ball B is never a minimizer in (1.4).*

We prove more generally that the second eigenvalue of a minimizer in (1.4) has to be simple and, as a consequence of the multiplicity of the second eigenvalue over balls, we immediately get the result in Theorem 1.3. The proof of the simplicity of the second eigenvalue relies on some ideas developed in [13, 23], with the help of a classical symmetry result due to Serrin [31] (see also [20]).

As an application of our results, we show how to get informations on the shape of the *attainable set*, namely the subset of the plane described by the range of the first two eigenvalues of the Dirichlet–Laplacian

$$\mathcal{E} := \{(x, y) \in \mathbb{R}^2 : x = \lambda_1(\Omega), y = \lambda_2(\Omega), \Omega \subset \mathbb{R}^N, \Omega \text{ open}, |\Omega| = 1\}. \tag{1.5}$$

This set was introduced in [27], and then deeply studied in [12] (see also [2, 3, 9]), where several geometrical properties of \mathcal{E} were discussed (the closedness of \mathcal{E} is important for the existence of optimal shapes for non monotone functionals see [12]).

The link between problem (1.4) and the set \mathcal{E} is the following: for a fixed $t \in (0, 1)$ a minimizer Ω_t in (1.4) corresponds to the first point of \mathcal{E} of coordinates $(\lambda_1(\Omega_t), \lambda_2(\Omega_t))$ that we reach with a line $tx + (1 - t)y = a$ increasing the value a , that is $P_{\Omega_t} := (\lambda_1(\Omega_t), \lambda_2(\Omega_t))$ is one of the intersection points of the tangent line to \mathcal{E} with slope $t/(t - 1)$. In particular, if $t = 1$ the tangent line $x = \lambda_1(B)$ has a *unique* intersection point corresponding to the ball B (because of the Faber–Krahn inequality (1.1)), while if $t = 0$, the tangent line $y = \lambda_2(\Theta)$ has a *unique* intersection point corresponding to the two balls $B_- \cup B_+$ (because of the Krahn–Szegő inequality (1.2)).

Therefore, in Theorem 4.2, we will present a new strategy for studying the asymptotic behavior of the boundary of \mathcal{E} near the points corresponding to B and $B_- \cup B_+$, extending to all dimensions a result proved in [27] only in two dimensions, and recovering the result proved in [9]. Indeed, according to [2, 3, 9, 27], the common strategy to study the asymptotic behavior of $\partial\mathcal{E}$ consists of two steps: the construction, through a parameter $\epsilon > 0$, of an explicit perturbation of the set corresponding to the limit point on the boundary of \mathcal{E} (i.e., a ball or two balls) and then the computation of the corresponding limit as $\epsilon \rightarrow 0$. Here instead, to compute

such a limit, we rely on the minimality condition of the minimizers of convex combinations (3.1) without any explicit construction. Moreover, looking at the boundary of the attainable set through convex combinations is very useful, since it works in all dimensions, and can be applied to other attainable sets with different constraints.

We suspect that to properly understand the boundary behavior of the attainable set, one has to carefully analyze problem (1.4). For this reason we restate the long-standing conjecture about the convexity of the attainable set in the language of the minimizers of convex combinations of the lowest Dirichlet eigenvalues. Notice that the link of problem (1.4) with the attainable set is not new, in fact it was used in [27] (and more recently in [2, 3]) to draw numerically the boundary of the attainable set. However, up to our knowledge, it was never used to study analytic results.

The paper is organized as follows. In Section 2 we derive an isoperimetric inequality for the Fraenkel 2-asymmetry. In Section 3 we prove Theorems 1.2 and 1.3. In Section 4 we show some applications to the attainable set. In Appendix A we compute explicit constants in quantitative inequalities.

Notation. Throughout the paper, we always denote by Ω_t a minimizer of problem (1.4) for $t \in (0, 1)$, by B the open ball of unit measure and by $B_- \cup B_+$ two open balls of half measure each, saving the particular notation Θ when their centers are on the x -axis and they are *tangent* in the origin (i.e., their closures are touching in the origin). We use the symbol \subset to denote the *strict* inclusion between sets, while \subseteq if the inclusion holds possibly with the equality, and we use Δ for the symmetric difference between sets. We write $\text{diam}(\cdot)$ and $\text{hull}(\cdot)$ to denote the diameter and the convex envelope of a set, respectively. We use \approx to denote an approximate value for a constant and we always consider only three decimal digits.

2. AN ISOPERIMETRIC INEQUALITY FOR THE FRAENKEL 2-ASYMMETRY

Quantitative inequalities are refinements of isoperimetric inequalities: they measure how far a set is from the optimal shape in terms of the deviation of the functional to its minimum value (for a brief overview on the topic see Appendix A). It is then important to look at the right quantity that provides such a measure.

The most well-known example is the so called *Fraenkel asymmetry*, which is often used when balls are optimal in an isoperimetric inequality: for an open set $\Omega \subset \mathbb{R}^N$ with unit measure, it is defined as

$$\mathcal{A}(\Omega) := \min \{ |\Omega \Delta B| : B \text{ open ball, } |B| = 1 \}. \tag{2.1}$$

In this paper we will rely on the Krahn–Szegő inequality (1.2) in a quantitative form, and, since in this case two disjoint balls of equal measure are the optimal set, a different asymmetry is needed. According to [7], the *Fraenkel 2-asymmetry* is, for an open set $\Omega \subset \mathbb{R}^N$ with unit measure, defined as

$$\mathcal{A}_2(\Omega) := \min \{ |\Omega \Delta (B_- \cup B_+)| : B_-, B_+ \text{ disjoint open balls, } |B_-| = |B_+| = 1/2 \}. \tag{2.2}$$

We point out that if Ω and E are two measurable sets of \mathbb{R}^N of the same measure $|\Omega| = |E|$, then

$$|\Omega \Delta E| = 2|\Omega \setminus E| = 2|E \setminus \Omega|, \tag{2.3}$$

and this allows to write (2.1) and (2.2) in a slightly different way, choosing $E = B$ or $E = B_- \cup B_+$. The relation (2.3) will be used several times in the sequel.

A quantitative Krahn–Szegő inequality was proved by Brasco and Pratelli in [7]: they showed the existence of a constant $C_{KS} > 0$ (depending on the dimension) such that for all open sets $\Omega \subset \mathbb{R}^N$ of finite measure $|\Omega| = |B_- \cup B_+|$,

$$\frac{\lambda_2(\Omega)}{\lambda_2(B_- \cup B_+)} - 1 \geq C_{KS} \mathcal{A}_2(\Omega)^{2(N+1)}. \tag{2.4}$$

Notice that C_{KS} can be explicitly computed, see Appendix A. In two dimensions a quantitative Faber–Krahn inequality proved by Bhattacharya in [6], allows to improve the exponent of (2.4), still with a constant that

can be explicitly computed: more precisely, if $N = 2$ there exists $\beta_{\text{KS}} > 0$ such that for all open sets $\Omega \subset \mathbb{R}^2$ of finite measure $|\Omega| = |B_- \cup B_+|$,

$$\frac{\lambda_2(\Omega)}{\lambda_2(B_- \cup B_+)} - 1 \geq \beta_{\text{KS}} \mathcal{A}_2(\Omega)^{9/2}, \tag{2.5}$$

where the constant β_{KS} , according to Appendix A, can be chosen as

$$\beta_{\text{KS}} = \frac{1}{24j_0^2} \frac{\pi^{3/2}}{(\pi + 2 + 2\pi^{1/3})^{9/2}} \times 10^{-5} \approx 3.331 \times 10^{-11}. \tag{2.6}$$

In this section we analyze the following purely geometrical problem: minimize the Fraenkel 2-asymmetry among *convex* sets

$$\inf\{\mathcal{A}_2(\Omega) : \Omega \subset \mathbb{R}^N, \Omega \text{ open and convex}, |\Omega| = 1\}. \tag{2.7}$$

The reason for studying problem (2.7) is to quantitatively answer to Theorem 1.2. We focus only on the two dimensional case, since in this case we are able to identify the unique minimizer for this problem, although some of the results that we prove hold in any dimension (see Rem. 2.5).

The condition for the balls B_-, B_+ to be disjoint in (2.2) prevents two balls to overlap, but not their closures to be tangent. This occurs, in particular, whenever the set Ω is convex.

Lemma 2.1. *Let $\Omega \subset \mathbb{R}^2$ be a convex open set with unit measure. The minimum in the definition of the Fraenkel 2-asymmetry (2.2) is attained by two tangent balls (i.e. those whose closures are touching in a point).*

Proof. We prove that tangent balls are always better competitors than non-tangent balls. Let B_- and B_+ be two admissible balls for (2.2) which are non-tangent. We may assume both $\Omega \cap B_-$ and $\Omega \cap B_+$ to be non-empty sets, otherwise the quantity $|\Omega \Delta (B_- \cup B_+)|$ can be decreased by any other couple of non-tangent balls satisfying this property. Therefore, the boundary of the convex envelope hull($(\Omega \cap B_-) \cup (\Omega \cap B_+)$) outside the balls B_- and B_+ is made exactly by two segments. By moving and rotating the coordinate system we may assume these segments to be onto the half lines $y = \pm mx$ with $x \geq 0$, or in the limiting case $y = \pm m$ with $x \geq 0$, for a suitable choice of the constant $m > 0$. Possibly exchanging the role of B_- and B_+ , we can also suppose B_- to be on the left with respect to B_+ . All these assumptions combined with the convexity of Ω guarantee the following fact: to every point $P_1 = (x_1, y) \in \Omega \cap B_-$ there exists a point (with the same ordinate) $P_2 = (x_2, y) \in \Omega \cap B_+$, and moreover, the whole segment $\overline{P_1 P_2} \subseteq \Omega$. Therefore, if τ denotes the translation toward the right, mapping the ball B_- into the (left) tangent ball $\tau(B_-)$ to B_+ , this means that $\tau(\Omega \cap B_-) = \tau(\Omega) \cap \tau(B_-) \subseteq \Omega \cap \tau(B_-)$. In particular $|\Omega \cap B_-| = |\tau(\Omega \cap B_-)| \leq |\Omega \cap \tau(B_-)|$, and so passing to the complementary sets yields that $|(B_- \cup B_+) \setminus \Omega| \geq |(\tau(B_-) \cup B_+) \setminus \Omega|$, which, together with (2.3) for $E = B_- \cup B_+$, implies that the functional to be minimized has been not increased on tangent balls. We remark that the configuration of the two optimal balls is not always unique (one can think, for example to Ω as a long rectangle), but also in this case, two tangent balls are one of the admissible optimal configuration, and we choose them for the next steps. \square

We are ready to analyze in detail problem (2.7). Therefore, we introduce the following definition.

Definition 2.2. We call *mobile* the open convex set M given by the intersection of the stadium hull(Θ) (centered in the origin) with the horizontal strip $\{(x, y) \in \mathbb{R}^2 : -h \leq y \leq h\}$, where $h > 0$ is chosen so that $|M| = 1$ (see Fig. 1).

For the sake of the reader we immediately compute the right value of h deducing the value of the Fraenkel 2-asymmetry for the mobile M .

Lemma 2.3. *The height h in Definition 2.2 is approximately 0.336. Therefore, the Fraenkel 2-asymmetry for the mobile M is*

$$\mathcal{A}_2(M) = \frac{16h}{\sqrt{2\pi}} - 2 \approx 0.147. \tag{2.8}$$




FIGURE 1. The stadium $\text{hull}(\Theta)$ and the mobile M .

Proof. From Lemma 2.1 and Definition 2.2 it is clear that $\mathcal{A}_2(M) = |M \triangle \Theta|$. Denoting by X the region delimited by the points O, Q, S (see Fig. 2), and by Y the set $\{(x, y) \in \Theta : x < 0, y > h\}$ yields that $|M \setminus \Theta| = 4|X|$ and $|\Theta \setminus M| = 4|Y|$, therefore, to compute $\mathcal{A}_2(M)$ it is sufficient to compute the area of X and of Y . To this purpose, we introduce the smallest angle α made by the radius PQ and the segment PR , where the height h of Definition 2.2 is linked to α by

$$h = \frac{1}{\sqrt{2\pi}} \cos \alpha, \tag{2.9}$$

since Θ is made by two balls of radius $1/\sqrt{2\pi}$. Now the area of the rectangle $OPRS$, of the triangle PQR and of the circular sector OPQ are, respectively,

$$|OPRS| = \frac{h}{\sqrt{2\pi}}, \quad |PQR| = \frac{h}{2} \sqrt{\frac{1}{2\pi} - h^2}, \quad |OPQ| = \frac{1}{4\pi} \left(\frac{\pi}{2} - \alpha \right),$$

and then

$$|X| = \frac{h}{\sqrt{2\pi}} - \frac{h}{2} \sqrt{\frac{1}{2\pi} - h^2} - \frac{1}{4\pi} \left(\frac{\pi}{2} - \alpha \right), \quad |Y| = \frac{\alpha}{2\pi} - h \sqrt{\frac{1}{2\pi} - h^2}.$$

The constraint $|M| = |\Theta| = 1$ yields (2.3) with $E = \Theta$, thus imposing $|X| = |Y|$ and recalling (2.9) the height h have to satisfy the following equality:

$$h \sqrt{\frac{1}{2\pi} - h^2} + \frac{2h}{\sqrt{2\pi}} - \frac{\arccos(\sqrt{2\pi}h)}{2\pi} = \frac{1}{4},$$

which is solved by $h \approx 0.336$. Moreover, plugging this equation and (2.9) into the formula for $|Y|$ we obtain $\mathcal{A}_2(M) = 8|X| = 8|Y|$ which provides (2.8). \square

Theorem 2.4 (Isoperimetric inequality for the Fraenkel 2-asymmetry). *Among all convex open planar sets with unit area, the mobile has the minimum Fraenkel 2-asymmetry, that is for every convex open set $\Omega \subset \mathbb{R}^2$ with $|\Omega| = 1$*

$$\mathcal{A}_2(\Omega) \geq \mathcal{A}_2(M), \tag{2.10}$$

and equality holds if and only if $\Omega = M$.

Proof. For every convex and open set Ω , moving and rotating the coordinate system, from Lemma 2.1 we may always assume the minimum in the definition of the Fraenkel 2-asymmetry to be reached by Θ , namely the two balls of equal measure which are tangent in the origin and with centers on the x -axis. Therefore, recalling (2.3) with $E = \Theta$ and noticing that $\mathcal{A}_2(M) = 2|\Theta \setminus M|$, in order to prove (2.10) it is sufficient to prove the auxiliary inequality

$$|\Theta \setminus \Omega| \geq |\Theta \setminus M| \quad \text{for every convex open set } \Omega \subset \mathbb{R}^2 \text{ with } |\Omega| = 1. \tag{2.11}$$




FIGURE 2. The Fraenkel 2-asymmetry of the mobile M .

We show this inequality in several steps, starting with the existence result, and then listing several necessary conditions that have to be satisfied by an optimal set.

(i) *Existence of an optimal set.*

The existence of an optimal set Ω^* for the minimization problem

$$\min\{|\Theta \setminus \Omega| : \Omega \subset \mathbb{R}^2, \Omega \text{ open and convex, } |\Omega| = 1\}, \tag{2.12}$$

is a straightforward consequence of the direct method of the Calculus of Variations. Indeed take a minimizing sequence $\{\Omega_n\}$ made of open and convex sets of unit area. The diameter $\text{diam}(\Omega_n)$ is uniformly bounded, otherwise from the area and the convexity constraints $|\Omega_n \cap \Theta| \downarrow 0$, and so $|\Theta \setminus \Omega_n| \uparrow 1$, which is strictly greater to the value attained by the mobile $|\Theta \setminus M| \approx 0.073$ (see Lem. 2.3).

Therefore, we can apply Proposition 2.4.3 and 2.4.4 of [11] to infer that we are minimizing a continuous functional $\Omega \mapsto |\Theta \setminus \Omega|$ over a compact class of sets with respect to the uniform convergence.

(ii) *An optimal set is included in the stadium hull(Θ).*

Assume, by contradiction, that Ω^* is not contained in the stadium hull(Θ), namely the following strict inclusion holds:

$$\Omega^* \cap \text{hull}(\Theta) \subset \Omega^*. \tag{2.13}$$

Consider the non-decreasing (in the sense of set inclusion) family of dilated sets

$$\Omega(r) := \text{hull}(r\Omega^* \cap \Theta), \quad \text{with } r \in [1, \infty). \tag{2.14}$$

For $r = 1$, the corresponding set $\Omega(1) \subset \Omega^*$ and, in particular, $|\Omega(1)| < |\Omega^*|$. Indeed, the convex set $\Omega^* \cap \text{hull}(\Theta)$ contains $\Omega^* \cap \Theta$, thus, by definition of the convex envelope, $\text{hull}(\Omega^* \cap \Theta) \subseteq \Omega^* \cap \text{hull}(\Theta)$, which combined with the assumption (2.13) provides the strict inclusion $\Omega(1) \subset \Omega^*$. For $r = \text{diam} \Theta$, $\Omega(\text{diam} \Theta) \supset \text{hull}(\Theta)$, and, in particular, $|\Omega(\text{diam} \Theta)| > |\Omega^*|$. Since the function $|\Omega(r)|$ is continuous over $[1, \text{diam}(\Theta)]$, i.e. for a sequence of real numbers $r_n \rightarrow r$, the convex sets $\Omega(r_n)$ converge to $\Omega(r)$ for the L^1 convergence of characteristic functions, there exists $r_1 > 1$ such that the convex set $\Omega(r_1)$ defined by (2.14) with $r = r_1$ satisfies the volume constraint $|\Omega(r_1)| = |\Omega| = |\Theta|$. Therefore, $\Omega(r_1)$ is an admissible competitor in (2.11) which, by definition (2.14), is so that $\Omega^* \cap \Theta \subset \Omega(r_1) \cap \Theta$ (note that r_1 is strictly greater than 1). Passing to the complementary sets, this strict inclusion yields that

$$|\Theta \setminus \Omega^*| > |\Theta \setminus \Omega(r_1)|,$$

contradicting the optimality of Ω^* .

(iii) *The closure of the boundary of an optimal set which is not included in $\partial\Theta$ is made by two connected components, each of which is a segment.*

From the previous step (ii), in particular (2.14), we know that the boundary of an optimal set which is not included in the two balls Θ is made by exactly two connected components, each of which is a segment. Then, we can consider the two supporting lines of Ω^* passing through these two segments, denoting by E the region between these two lines containing Ω^* . The region E is a strip, in the case the supporting lines are parallel, otherwise a cone with a vertex outside $\overline{\text{hull}(\Theta)}$ (if these lines would intersect in a point into $\overline{\text{hull}(\Theta)}$, we would have that $|E \cap \text{hull}(\Theta)| < |\Theta|$ and since $\Omega^* \subseteq E \cap \text{hull}(\Theta)$ this would contradict the area constraint). Now assuming, by contradiction, that point iii) does not hold, we would have the strict inequality

$$|E \cap \text{hull}(\Theta)| > |\Omega^*|. \tag{2.15}$$

Then, we consider the contracted family of convex sets

$$\Omega(r) := rE \cap \text{hull}(\Theta), \quad \text{with } r \in (0, 1), \tag{2.16}$$

where now rE denotes the contraction of the cone E of factor r that keeps fixed its vertex, and we notice that the function $|\Omega(r)|$ is continuous over $(0, 1)$, *i.e.* for a sequence of real numbers $r_n \rightarrow r$ in $(0, 1)$, the convex sets $\Omega(r_n)$ converge to $\Omega(r)$ for the L^1 convergence of characteristic functions. Moreover, as $r \rightarrow 1$, from (2.15), we have that $|\Omega(r)| > |\Omega^*|$, while as $r \rightarrow 0$, the set rE shrinks to a line, thus in particular, $|\Omega(r)| \rightarrow 0$. Therefore, there exists $r_2 < 1$ such that the convex set $\Omega(r_2)$ defined in (2.16) satisfies the area constraint $|\Omega(r_2)| = |\Omega|$ and the strict inclusion $\Omega(r_2) \setminus \Theta \subset \Omega^* \setminus \Theta$ gives $|\Omega(r_2) \setminus \Theta| < |\Omega^* \setminus \Theta|$ (since r_2 is strictly smaller than 1). Recalling (2.3) with $E = \Theta$ yields

$$|\Theta \setminus \Omega^*| > |\Theta \setminus \Omega(r_2)|,$$

contradicting the optimality of Ω^* .

(iv) *The mobile is the optimal set.*

Combining the steps (i) and (ii) with (iii) we know that an optimal set Ω^* has the following form

$$\Omega^* = E \cap \text{hull}(\Theta), \tag{2.17}$$

where E , as before, is a strip or a cone with a vertex outside $\overline{\text{hull}(\Theta)}$ such that $|E \cap \text{hull}(\Theta)| = 1$. Now, the boundary of this set is made by four pieces: two segments and two arc of circles. If for every cone E_{cone} one can find a strip E_{strip} which decreases the functional in (2.12), then optimizing among all sets of the form $\Omega^* = E_{\text{strip}} \cap \text{hull}(\Theta)$ would give that the symmetric strip is the best one, that is, according to Definition 2.2, the mobile. Indeed, assume that the optimal set Ω^* as in (2.17) is generated by a cone $E = E_{\text{cone}}$, and, up to a change of coordinates we may assume that it has on the boundary a segment which belongs to the half-plane $\{y > 0\}$ and that is not parallel to the line $\{y = 0\}$. We show how it is possible to rotate this segment decreasing the functional in (2.12). Let us call l_1 the supporting line generating this segment, P be the point on this line with $x_P = 0$, and assume that the following quantity is positive

$$d := |\{y > 0\} \cap \{x > 0\} \cap (E \cap \text{hull}(\Theta))| - |\{y > 0\} \cap \{x < 0\} \cap (E \cap \text{hull}(\Theta))| > 0.$$

Rotating the line l_1 around the point P , so as to decrease d until $d = 0$, yields the line l_2 parallel to $\{y = 0\}$. Calling E_2 the new set obtained from E_{cone} with l_2 in place of l_1 we can see that the area has been increased, namely $|E_2 \cap \text{hull}(\Theta)| > |E_{\text{cone}} \cap \text{hull}(\Theta)|$, and moreover $|\text{hull}(\Theta) \setminus E_2| < |\text{hull}(\Theta) \setminus E_{\text{cone}}|$. Now, we replace the line l_2 with a parallel line l_3 and define a new set E_3 so that $|E_3 \cap \text{hull}(\Theta)| = |E_{\text{cone}} \cap \text{hull}(\Theta)|$. Therefore we have constructed a set E_3 which satisfies the area constraint and so that the functional has been decreased $|\text{hull}(\Theta) \setminus E_3| < |\text{hull}(\Theta) \setminus E_{\text{cone}}|$. The same strategy can be adapted if the quantity $d < 0$, and for the other segment on the boundary.

At the end, we constructed a better competitor E_{strip} and optimizing among all E_{strip} will provide that the mobile is the unique minimizer, and the theorem is concluded. \square

Remark 2.5. If $N > 2$ it seems more difficult to find the optimal set in (2.7). Nevertheless, some of the above results can immediately generalized to all dimensions, such as Lemma 2.1 as well as the steps (i)–(iii) in the proof of Theorem 2.4. As a by product, for $N > 2$, an optimal set Ω^* in (2.7) is so that $\Omega^* = E \cap \text{hull}(\Theta)$ where E is an N dimensional cone or strip.

3. CONVEX COMBINATIONS OF THE LOWEST DIRICHLET EIGENVALUES

3.1. Basic properties

We start discussing some properties of the functional F_t defined in (1.3) that follow from the definition of minimality: we say that Ω_t is a minimizer in (1.4) if for every admissible competitor Ω

$$t\lambda_1(\Omega_t) + (1 - t)\lambda_2(\Omega_t) \leq t\lambda_1(\Omega) + (1 - t)\lambda_2(\Omega), \tag{3.1}$$

and equivalently, rearranging the terms

$$\lambda_1(\Omega_t) - \lambda_1(\Omega) + \lambda_2(\Omega) - \lambda_2(\Omega_t) \leq \frac{1}{t} \left(\lambda_2(\Omega) - \lambda_2(\Omega_t) \right). \tag{3.2}$$

Lemma 3.1. For $s, t \in (0, 1)$ with $s < t$, let Ω_s and Ω_t be minimizers of the functionals F_s and F_t , respectively. The following properties hold:

- (i) the gap non-decreases $\lambda_2(\Omega_s) - \lambda_1(\Omega_s) \leq \lambda_2(\Omega_t) - \lambda_1(\Omega_t)$;
- (ii) the first eigenvalue non-increases $\lambda_1(\Omega_s) \geq \lambda_1(\Omega_t)$;
- (iii) the second eigenvalue non-decreases $\lambda_2(\Omega_s) \leq \lambda_2(\Omega_t)$;
- (iv) a rescaled convex combination increases $F_s(\Omega_s)/(1 - s) < F_t(\Omega_t)/(1 - t)$;
- (v) the convex combination decreases $F_s(\Omega_s) > F_t(\Omega_t)$.

Proof. The minimality (3.1) of Ω_s for F_s with the competitor $\Omega = \Omega_t$ writes as

$$s\lambda_1(\Omega_s) + (1 - s)\lambda_2(\Omega_s) \leq s\lambda_1(\Omega_t) + (1 - s)\lambda_2(\Omega_t), \tag{3.3}$$

while the minimality (3.1) of Ω_t for F_t with the competitor $\Omega = \Omega_s$, as

$$t\lambda_1(\Omega_t) + (1 - t)\lambda_2(\Omega_t) \leq t\lambda_1(\Omega_s) + (1 - t)\lambda_2(\Omega_s). \tag{3.4}$$

Summing up (3.3) with (3.4) we get to

$$(t - s)(\lambda_1(\Omega_t) - \lambda_1(\Omega_s)) \leq (t - s)(\lambda_2(\Omega_t) - \lambda_2(\Omega_s)), \tag{3.5}$$

and by the assumption $t - s > 0$, we immediately obtain the first point i) of this lemma. If $\lambda_1(\Omega_s) < \lambda_1(\Omega_t)$ the right-hand side of (3.5) would be positive, hence $\lambda_2(\Omega_s) < \lambda_2(\Omega_t)$, which would contradict (3.4). Then, necessarily, the second point ii) of this lemma holds. Moreover, if $\lambda_2(\Omega_s) > \lambda_2(\Omega_t)$, using point (ii) we get a contradiction with (3.3), and so also the third point iii) holds. Now, from (3.3) and the fact that $s/(1 - s) < t/(1 - t)$, we have

$$\frac{s}{(1 - s)}\lambda_1(\Omega_s) + \lambda_2(\Omega_s) \leq \frac{s}{(1 - s)}\lambda_1(\Omega_t) + \lambda_2(\Omega_t) < \frac{t}{(1 - t)}\lambda_1(\Omega_t) + \lambda_2(\Omega_t).$$

Therefore, $F_t(\Omega)/(1 - t) = t/(1 - t)\lambda_1(\Omega) + \lambda_2(\Omega)$ and we get point iv). To prove the last point v) we assume, by contradiction, that $F_s(\Omega_s) \leq F_t(\Omega_t)$. Then recalling (3.4) we have

$$s\lambda_1(\Omega_s) + (1 - s)\lambda_2(\Omega_s) \leq t\lambda_1(\Omega_t) + (1 - t)\lambda_2(\Omega_t) \leq t\lambda_1(\Omega_s) + (1 - t)\lambda_2(\Omega_s),$$

which leads to

$$(t - s)\lambda_2(\Omega_s) \leq (t - s)\lambda_1(\Omega_s).$$

The assumption $t - s > 0$ implies that $\lambda_1(\Omega_s) = \lambda_2(\Omega_s)$, contradicting the connectedness of Ω_s proved in [28]. □

Lemma 3.2. *Let $s, t \in (0, 1)$ with $s < t$. If X is a minimizer of both F_s and F_t , then X is also a minimizer of F_r , for every $r \in (s, t)$.*

Proof. We assume, by contradiction, that X is not a minimizer of F_r , for some fixed $r \in (s, t)$, and call by Y a minimizer of the corresponding functional F_r . As in the proof of the previous lemma, using the minimality of X for F_s and of Y for F_r we arrive to (3.5) with $t = r$, namely since $s < r$,

$$(\lambda_1(Y) - \lambda_1(X)) < (\lambda_2(Y) - \lambda_2(X)),$$

and, similarly, using the minimality of X for F_t and again of Y for F_r

$$(\lambda_1(X) - \lambda_1(Y)) < (\lambda_2(X) - \lambda_2(Y)),$$

where the strict inequalities are a consequence of the assumption $F_r(Y) < F_r(X)$. Summing up these two inequalities, we reach a contradiction. □

3.2. On the non-convexity of the minimizers

Recently, Henrot and Oudet in [24] investigated the problem of minimizing the second eigenvalue of the Dirichlet–Laplacian among sets of fixed measure and with an additional *convexity* constraint. Finding an explicit minimizer in this class seems a very difficult problem: a possible candidate to be the optimum is the stadium (*i.e.*, the convex hull of two tangent balls), but this conjecture was refuted in [24]. Indeed any set which contains on the boundary some pieces of balls can not be a minimizer. Nevertheless, in [24] it was proved the existence of a convex minimizer Ω_{HO} so that, for every open and *convex* set $\Omega \subset \mathbb{R}^N$ with unit area,

$$\lambda_2(\Omega) \geq \lambda_2(\Omega_{\text{HO}}), \tag{3.6}$$

(*cf.* (3.6) with the Krahn–Szegő inequality, where no-convexity constraint is required). Notice that, since Ω_{HO} has no pieces of balls on its boundary, in particular $\Omega_{\text{HO}} \neq B$ and

$$\omega_N^{2/N} j_{N/2}^2 = \lambda_2(B) > \lambda_2(\Omega_{\text{HO}}).$$

In two dimensions, Oudet in [30] and, more recently, Antunes and Henrot in [2], made some numerical computations, showing the shape of the optimal set Ω_{HO} and highlighting that Ω_{HO} is very close to the stadium, both from a geometrical and a numerical point of view; in particular

$$\lambda_2(B_- \cup B_+) = 2\pi j_0^2 \approx 36.336, \quad \lambda_2(\Omega_{\text{HO}}) \approx 37.987, \quad \lambda_2(\Omega_{\text{stadium}}) \approx 38.002, \tag{3.7}$$

where Ω_{stadium} is the stadium with $|\Omega_{\text{stadium}}| = 1$, *i.e.*, a contracted version of the set $\text{hull}(\Theta)$.

We are now in position to prove Theorem 1.2.

Proof of Theorem 1.2. From the Krahn–Szegő inequality (1.2) and the connectedness of Ω_t it follows that

$$\lambda_2(B_- \cup B_+) < \lambda_2(\Omega_t), \tag{3.8}$$

which plugged into (3.1) with $\Omega = B_- \cup B_+$ yields

$$\lambda_1(\Omega_t) < \lambda_1(B_- \cup B_+). \tag{3.9}$$

Taking $\Omega = B_- \cup B_+$ also in (3.2) and dividing therein by the negative quantity $\lambda_2(B_- \cup B_+) - \lambda_2(\Omega_t)$ (recall (3.8)) we get to

$$\frac{\lambda_1(B_- \cup B_+) - \lambda_1(\Omega_t)}{\lambda_2(\Omega_t) - \lambda_2(B_- \cup B_+)} + 1 \geq \frac{1}{t}. \tag{3.10}$$

From (3.8) and (3.9), the ratio on the left-hand side of this inequality turns out to be a positive number; therefore, we can use the Faber–Krahn inequality (1.1) to estimate $\lambda_1(\Omega_t)$ at the numerator of this ratio. Moreover, if Ω_t would be a convex set, we could also use (3.6) to estimate $\lambda_2(\Omega_t)$ at the denominator of this ratio, obtaining the following uniform bound on t :

$$t \geq \frac{1}{\frac{\lambda_1(B_- \cup B_+) - \lambda_1(B)}{\lambda_2(\Omega_{HO}) - \lambda_2(B_- \cup B_+)} + 1}. \tag{3.11}$$

Calling T the quantity on the right-hand side of this inequality, the Krahn–Szegő inequality for convex sets gives $\lambda_2(\Omega_{HO}) - \lambda_2(B_- \cup B_+) > 0$, thus $T > 0$. Therefore, if $t < T$, Ω_t can not be convex. \square

The proof of Theorem 1.2 is constructive and reveals an explicit expression for the threshold T in terms of the eigenvalues of the Dirichlet–Laplacian.

Corollary 3.3. *The threshold T in Theorem 1.2 has the following expression:*

$$T = 1 - \frac{(2^{2/N} - 1)\lambda_1(B)}{\lambda_2(\Omega_{HO}) - \lambda_1(B)}, \tag{3.12}$$

where Ω_{HO} is a minimizer in (3.6). In two dimensions, it turns out that

$$T \geq 1 - \frac{1}{1 + 2\beta_{KS}\mathcal{A}_2(M)^{9/2}} \approx 1.192 \times 10^{-14}, \tag{3.13}$$

where the constants β_{KS} and $\mathcal{A}_2(M)$ are as in (2.6) and (2.10) respectively.

Proof. Define as T the quantity on the right-hand side of (3.11). Noticing that $\lambda_1(B_- \cup B_+) = \lambda_2(B_- \cup B_+) = 2^{2/N}\lambda_1(B)$ and simplifying, we reach (3.12). Moreover, plugging, the quantitative Krahn–Szegő inequality (2.4) into (3.12), yields a lower bound on T , which is independent on the eigenvalues of the Dirichlet–Laplacian

$$T \geq 1 - \frac{(2^{2/N} - 1)}{(2^{2/N} - 1) + 2^{2/N}C_{KS}\mathcal{A}_2(\Omega_{HO})^{2(N+1)}}.$$

In two dimensions, on the other hand, one can use (2.5), (2.6) and Theorem 2.4 together with (2.8) to get the explicit value in (3.13). \square

Remark 3.4. The explicit value for the lower bound to the threshold T is not very accurate, mostly due to the fact that the constant β_{KS} is not the optimal one, but we believe it is important to show that a numerical value can actually be provided. Moreover, if $N = 2$, plugging the numerical computation of $\lambda_2(\Omega_{HO})$ recalled in (3.7) into (3.12) and using $\lambda_1(B) = \pi j_0^2 \approx 18.168$, reveals a numerical approximation for the threshold defined by (3.12):

$$T \approx 0.083.$$

3.3. The ball never minimizes the convex combination

For the proof of Theorem 1.3 we need the following result.

Proposition 3.5. *For a fixed $t \in (0, 1)$, let Ω_t be a minimizer of problem (1.4). If the boundary of Ω_t is of class C^2 and connected, then $\lambda_2(\Omega_t)$ is simple, namely $\lambda_1(\Omega_t) < \lambda_2(\Omega_t) < \lambda_3(\Omega_t)$. Moreover, on the boundary of Ω_t , the following optimality condition holds:*

$$t|\nabla u_1(x)|^2 + (1-t)|\nabla u_2(x)|^2 = \frac{2F_t(\Omega_t)}{N}, \quad x \in \partial\Omega_t. \tag{3.14}$$

Proof. The fact that $\lambda_1(\Omega_t) < \lambda_2(\Omega_t)$ holds since Ω_t is connected, as it was proved in [28]. To prove that $\lambda_2(\Omega_t)$ is simple we proceed as in ([13], Lem. 2.1) (see also [23], Lem. 2.5.9) assuming, by contradiction, that $\lambda_2(\Omega_t)$ is a multiple eigenvalue, in the sense that $\lambda_2(\Omega_t) = \dots = \lambda_k(\Omega_t)$ for some $k \geq 3$ (see [23] for a precise definition of the higher eigenvalues). We use the results of derivability of multiple eigenvalues, see for instance ([23], Thm. 2.5.8). We deform the domain Ω_t with a regular vector field $\phi_\epsilon(x) = x + \epsilon V(x)$ and we denote with $\{u_i\}_{2 \leq i \leq k}$ the orthonormal (for the L^2 scalar product) family of eigenfunctions associated to $\lambda_2(\Omega_t)$. The directional derivative of the map $\epsilon \mapsto \lambda_2(\phi_\epsilon(\Omega_t))$ at $\epsilon = 0$ is one of the eigenvalues of the $(k - 1) \times (k - 1)$ matrix

$$A = \left(- \int_{\partial\Omega_t} \frac{\partial u_i}{\partial \nu} \frac{\partial u_j}{\partial \nu} V \cdot \nu \, d\sigma \right)_{2 \leq i, j \leq k}, \tag{3.15}$$

where $V \cdot \nu$ denotes the normal displacement of the boundary induced by the vector field V . Moreover, as observed before, the first eigenvalue is simple at Ω_t , then it is differentiable and the derivative is a linear form in $V \cdot \nu$ supported on $\partial\Omega_t$ (see e.g. [23], Thm. 2.5.1)

$$d\lambda_1(\Omega_t; V) = - \int_{\partial\Omega_t} \left(\frac{\partial u_1}{\partial \nu} \right)^2 V \cdot \nu \, d\sigma, \tag{3.16}$$

while the derivative of the volume is given by

$$d\text{Vol}(\Omega_t; V) = \int_{\partial\Omega_t} V \cdot \nu \, d\sigma. \tag{3.17}$$

Now, since Ω_t is a minimizer in (1.4), then it is also a minimizer for the Lagrangian

$$\mathcal{L}(\Omega) = t\lambda_1(\Omega) + (1 - t)\lambda_2(\Omega) + \mu|\Omega|, \quad \text{with} \quad \mu := \frac{2F_t(\Omega_t)}{N}. \tag{3.18}$$

Indeed, for such a μ , the function $f(s) = \mathcal{L}(s\Omega)$ of the real variable $s > 0$ achieves its minimum in $s = 1$, and since Ω_t minimizes F_t this implies that $\mathcal{L}(\Omega_t) \leq \mathcal{L}(\Omega)$ for every bounded open set $\Omega \subset \mathbb{R}^N$ (see for instance [18], Rem. 3.6). Then, we can differentiate the lagrangian \mathcal{L} without taking care of the volume constraint, and from (3.15), (3.16) and (3.17) the derivative of (3.18) is the smallest eigenvalue of the matrix

$$A_t = t d\lambda_1(\Omega_t; V) Id + (1 - t)A + \mu d\text{Vol}(\Omega; V) Id,$$

where Id is the $(k - 1) \times (k - 1)$ identity matrix. Therefore, in order to get a contradiction with the optimality of Ω_t , it is enough to prove that there is always a vector field V such that the matrix A_t has a negative eigenvalue. To this purpose let us consider two points P and Q on $\partial\Omega_t$ and two small neighborhoods γ_P and γ_Q of these two points of same length, say 2δ . Let us choose any regular function φ defined on $(-\delta, \delta)$, vanishing on the boundary of the interval, and a deformation field V such that:

$$V \cdot \nu = \varphi \text{ on } \gamma_P, \quad V \cdot \nu = -\varphi \text{ on } \gamma_Q, \quad V \cdot \nu = 0 \text{ elsewhere on } \partial\Omega_t.$$

With this choice of the field V , we have that the matrix A in (3.15) becomes the difference of two matrices

$$A = A(P) - A(Q) := \left(- \int_{\gamma_P} \frac{\partial u_i}{\partial \nu} \frac{\partial u_j}{\partial \nu} \varphi \, d\sigma \right)_{2 \leq i, j \leq k} + \left(\int_{\gamma_Q} \frac{\partial u_i}{\partial \nu} \frac{\partial u_j}{\partial \nu} \varphi \, d\sigma \right)_{2 \leq i, j \leq k}$$

and so the matrix A_t becomes the difference of two matrices $A_t(P) - A_t(Q)$, where

$$A_t(P) = -t \int_{\gamma_P} \left(\frac{\partial u_1}{\partial \nu} \right)^2 \varphi \, d\sigma Id + (1 - t)A(P) + \mu \int_{\gamma_P} \varphi \, d\sigma Id,$$

and $A_t(Q)$ defined analogously. The only case in which we cannot choose two points P, Q and a function φ such that the matrix has a negative eigenvalue is when $A_t(P) = A_t(Q)$, and this implies that, for every $2 \leq i, j \leq k$

$$\int_{\gamma_P} \frac{\partial u_i}{\partial \nu} \frac{\partial u_j}{\partial \nu} \varphi \, d\sigma = \int_{\gamma_Q} \frac{\partial u_i}{\partial \nu} \frac{\partial u_j}{\partial \nu} \varphi \, d\sigma,$$

and

$$\int_{\gamma_P} \left[\left(\frac{\partial u_i}{\partial \nu} \right)^2 - \left(\frac{\partial u_j}{\partial \nu} \right)^2 \right] \varphi \, d\sigma = \int_{\gamma_Q} \left[\left(\frac{\partial u_i}{\partial \nu} \right)^2 - \left(\frac{\partial u_j}{\partial \nu} \right)^2 \right] \varphi \, d\sigma,$$

for every $P, Q \in \partial\Omega_t$ and for every regular function φ . It means, in particular, that the product $(\partial u_2/\partial \nu)(\partial u_3/\partial \nu)$ and the difference $(\partial u_2/\partial \nu)^2 - (\partial u_3/\partial \nu)^2$ should be constant on $\partial\Omega_t$, and then $(\partial u_2/\partial \nu)^2 = c^2$ is constant on $\partial\Omega_t$. If the constant c is equal to zero, we get a contradiction thanks to the Holmgren uniqueness theorem. On the other hand, if $c \neq 0$, we have that $\partial u_2/\partial \nu = c$, because the boundary of Ω_t is connected. Then we can apply the classical Serrin Theorem [31] to u_2 (or possibly to $-u_2$), in the slightly more general version ([20], Thm. 17), which gives that Ω_t must be a ball, that is a contradiction since $\partial u_2/\partial \nu$ cannot be constant on the boundary of a ball (see for example [23], Sect. 1.2.5). Then, also $\lambda_2(\Omega_t)$ has to be simple.

At Ω_t the derivative of the Lagrangian (3.18) has to be zero: then (3.16), (3.17), and the corresponding formula for the derivative of the second eigenvalue λ_2 (3.15) with $k = 2$ (since it is simple) yields that

$$t \int_{\partial\Omega_t} \left(\frac{\partial u_1}{\partial \nu} \right)^2 V \cdot \nu \, d\sigma + (1-t) \int_{\partial\Omega_t} \left(\frac{\partial u_2}{\partial \nu} \right)^2 V \cdot \nu \, d\sigma = \mu \int_{\partial\Omega_t} V \cdot \nu \, d\sigma$$

for every regular vector field V . From this, recalling that in the Dirichlet case $|\nabla u_i|^2 = (\partial u_i/\partial \nu)^2$ with $i = 1, 2$, we obtain (3.14). □

Remark 3.6. The simplicity of $\lambda_2(\Omega_t)$ can also be proved under other regularity assumptions on Ω_t . The weakest assumption that is needed in order to differentiate the domain is that the boundary $\partial\Omega_t$ contains a part

$$\Gamma \text{ which is nonempty, relatively open in } \partial\Omega_t, \text{ connected and of class } C^{1,1}. \tag{3.19}$$

Then, it is enough to repeat the same proof by taking care that the vector field V constructed therein is chosen with support contained in Γ , and one obtains an overdetermined condition only along that part Γ of the boundary. If a Serrin principle is also available for a *partially overdetermined* condition we are done. Indeed Proposition 3.5 holds under different regularity assumptions on Ω_t , for instance:

- if Ω_t is *convex* with Γ as in (3.19) (using [20], Thm. 7);
- if $\partial\Omega_t$ is connected with Γ as in (3.19) and analytically continuable (using [20], Thm. 1).

Proof of Theorem 1.3. The proof is a straightforward consequence of Proposition 3.5: in every dimension, the second eigenvalue $\lambda_2(B)$ is not simple, therefore the ball B can not be a minimizer for any $t \in (0, 1)$. □

Remark 3.7. In two dimensions, the fact that balls are never minimizers was implicitly contained in the work [27]. For an arbitrary $\varepsilon > 0$ small enough, in [27] the authors constructed a nearly spherical competitor B_ε , with $|B_\varepsilon| = 1$, such that

$$\lambda_1(B_\varepsilon) \leq \lambda_1(B) + d_1 \varepsilon^2, \quad \text{while} \quad \lambda_2(B_\varepsilon) \leq \lambda_2(B) - d_2 \varepsilon,$$

for some positive constants d_1, d_2 . Therefore, for every $t \in (0, 1)$, it is possible to find $\varepsilon > 0$ so small so that

$$t\lambda_1(B_\varepsilon) + (1-t)\lambda_2(B_\varepsilon) < t\lambda_1(B) + (1-t)\lambda_2(B).$$




FIGURE 3. The attainable set for the lowest eigenvalues.

4. REMARKS ON THE ATTAINABLE SET OF THE LOWEST DIRICHLET EIGENVALUES

We start listing the most important properties that are known on the attainable set \mathcal{E} defined in (1.5) (see Fig. 3):

- (1) lies above the bisector $y = x$ (since by definition $\lambda_2(\Omega) \geq \lambda_1(\Omega)$ for every $\Omega \subset \mathbb{R}^N$).
- (2) lies on the right of the line $x = \lambda_1(B)$ (for the Faber–Krahn inequality (1.1)).
- (3) lies above the line $y = \lambda_2(B_- \cup B_+)$ (for the Krahn–Szegő inequality (1.2)).
- (4) lies below the line $y = \frac{\lambda_2(B)}{\lambda_1(B)}x$ (for the Ashbaugh–Benguria inequality [4]).
- (5) is conical with respect to the origin.

The numerical picture provided by Keller and Wolff suggests the evidence that the attainable set \mathcal{E} is convex, but this is a long-standing conjecture which is still unsolved.

Conjecture 4.1. *The attainable set \mathcal{E} is convex.*

The most important result in the direction of this conjecture was proposed by Bucur *et al.* in [12]. These authors proved that the attainable set (1.5), constructed through quasi-open set instead of open set, is convex in the vertical and in the horizontal direction and, as a consequence, that it is closed. Nevertheless the vertical and horizontal convexity *do not* imply convexity (think, for example to an L-shaped set).

From the properties of the set \mathcal{E} listed above it is clear that the unique unknown part of the boundary of \mathcal{E} is the curve \mathcal{C} connecting the points $P_B = (\lambda_1(B), \lambda_2(B))$ and $P_{B_- \cup B_+} = (\lambda_1(B_- \cup B_+), \lambda_2(B_- \cup B_+))$. The convexity of \mathcal{E} is then guaranteed as soon as \mathcal{C} can be parametrized by a convex function. For this reason it is important to have more informations on the curve \mathcal{C} . In two dimensions, Keller and Wolf in [27] showed that the tangent of \mathcal{C} at the point P_B corresponding to a ball B is vertical. They constructed a nearly spherical perturbation of B , as recalled in Remark 3.7, and then they computed the slope of the tangent to \mathcal{C} as $\epsilon \rightarrow 0$. Moreover, in all dimensions, Brasco, Nitsch and Pratelli showed that the tangent of \mathcal{C} at the point $P_{B_- \cup B_+}$

corresponding to two balls $B_- \cup B_+$ is horizontal. In this case the limit as $\epsilon \rightarrow 0$ was computed by overlapping the two balls B_- and B_+ of a quantity measured in terms of the parameter ϵ . In the following proposition we recover these limits relying on the minimality condition of the minimizers of convex combinations (3.1) without any explicit construction. Notice that the strategy that we adopt holds in all dimensions.

Theorem 4.2. *For every dimension $N \geq 2$ and $t \in (0, 1)$, let Ω_t be a minimizer of problem (1.4). Then we have:*

i) *the tangent of \mathcal{C} at the point P_B corresponding to one ball is vertical, namely*

$$\lim_{t \rightarrow 1} \frac{\lambda_2(\Omega_t) - \lambda_2(B)}{\lambda_1(\Omega_t) - \lambda_1(B)} = -\infty \tag{4.1}$$

ii) *the tangent of \mathcal{C} at the point $P_{B_- \cup B_+}$ corresponding to two identical balls is horizontal, namely*

$$\lim_{t \rightarrow 0} \frac{\lambda_2(\Omega_t) - \lambda_2(B_- \cup B_+)}{\lambda_1(\Omega_t) - \lambda_1(B_- \cup B_+)} = 0. \tag{4.2}$$

Moreover, the following limits holds

$$\lim_{t \rightarrow 0} \lambda_2(\Omega_t) = \lambda_2(B_- \cup B_+) \quad \text{and} \quad \lim_{t \rightarrow 1} \lambda_1(\Omega_t) = \lambda_1(B). \tag{4.3}$$

Proof. From the Faber–Krahn inequality (1.1) and Theorem 1.3 we find that

$$\lambda_1(B) < \lambda_1(\Omega_t), \tag{4.4}$$

which plugged into (3.1) with $\Omega = B$ yields

$$\lambda_2(\Omega_t) < \lambda_2(B). \tag{4.5}$$

Taking $\Omega = B$ in (3.2) and dividing therein by $\lambda_2(B) - \lambda_2(\Omega_t)$ (which from (4.5) is a strictly positive value) yields

$$\frac{\lambda_1(\Omega_t) - \lambda_1(B)}{\lambda_2(B) - \lambda_2(\Omega_t)} + 1 \leq \frac{1}{t}.$$

From (4.4) and (4.5) one can see that the ratio on the left-hand side of this inequality is a positive number, therefore, letting $t \uparrow 1$, necessarily, it holds the limit in (4.1). Moreover, repeating the computations made in the proof of Theorem 1.2 and letting $t \downarrow 0$ in (3.10), it follows the limit in (4.2).

Finally, the limits in (4.3) are a consequence of (4.1), (4.2) and of the boundedness of the denominator in (4.2) (because of (3.9)) and of the numerator in (4.1) (because of (4.5)). □

Remark 4.3. The strategy of looking at the boundary of the attainable set through convex combinations can be applied to other attainable sets with different constraints, for instance to the attainable set with a perimeter constraint [3]

$$\mathcal{E}_p := \{(x, y) \in \mathbb{R}^2 : x = \lambda_1(\Omega), y = \lambda_2(\Omega), \Omega \subset \mathbb{R}^N, \Omega \text{ open}, \mathcal{P}(\Omega) = 1\},$$

or to the attainable set with a convexity constraint [2]

$$\mathcal{E}_c := \{(x, y) \in \mathbb{R}^2 : x = \lambda_1(\Omega), y = \lambda_2(\Omega), \Omega \subset \mathbb{R}^N, \Omega \text{ open and convex}, |\Omega| = 1\}.$$

In particular, as in Theorem 4.2, it is possible to show that the tangent of $\partial\mathcal{E}_p$ (or of $\partial\mathcal{E}_c$) at the point P_B is vertical.

We finish this discussion formulating an *isospectral* conjecture on the minimizers of problem (1.4), which could be used to prove the convexity of the attainable set \mathcal{E} .

Conjecture 4.4. *Let $t \in (0, 1)$ and assume $X, Y \subset \mathbb{R}^N$ to be minimizers of problem (1.4) with $F_t(X) = F_t(Y)$. Then, the lowest eigenvalues of X and Y coincide, namely*

$$\lambda_1(X) = \lambda_1(Y) \quad \text{and} \quad \lambda_2(X) = \lambda_2(Y).$$

Proposition 4.5. *The validity of Conjecture 4.4 implies that Conjecture 4.1 holds true.*

Proof. If \mathcal{E} is not convex, then we can find two points $P_X, P_Y \in \mathcal{C}$, corresponding, respectively, to X, Y , and a straight line l passing through these points such that the curve \mathcal{C} lies above l . Therefore, it is clear that l will be of the form $tx + (1 - t)y = a$ for some fixed $t \in (0, 1)$ and a real number a . Hence the sets X, Y are minimizers in (1.4) for such a t , but $\lambda_1(X) \neq \lambda_1(Y)$ and $\lambda_2(X) \neq \lambda_2(Y)$, a contradiction with Conjecture 4.4. \square

APPENDIX A. EXPLICIT CONSTANTS IN QUANTITATIVE INEQUALITIES

In the following we compute explicit constants in some quantitative inequalities. We focus in particular, on the quantitative Krahn–Szegő inequality (2.4), having in mind its application in Corollary 3.3. We start with a brief overview on the most important quantitative inequalities, without pretending of being exhaustive.

The *quantitative isoperimetric inequality for the De Giorgi perimeter \mathcal{P}* in the sharp version was proved by Fusco *et al.* in [21]: there exist a constant $C_I > 0$ (depending on the dimension) such that for all open sets $\Omega \subset \mathbb{R}^N$ of finite measure $|\Omega| = |B|$,

$$\frac{\mathcal{P}(\Omega)}{\mathcal{P}(B)} - 1 \geq C_I \mathcal{A}(\Omega)^2, \tag{A.1}$$

where \mathcal{A} is as in (2.1). In this setting, a quantitative inequality is *sharp* when it has the least exponent on the Fraenkel asymmetry, and a constant is *optimal* when it has the largest possible value. Moreover, the quantity estimated from below through the asymmetry is the *deficit* of the functional.

The same authors later proved a *quantitative Faber–Krahn inequality*, with a proof based on (A.1): there exists a constant $C_{FK} > 0$ (depending on the dimension) such that for all open sets $\Omega \subset \mathbb{R}^N$ of finite measure $|\Omega| = |B|$,

$$\frac{\lambda_1(\Omega)}{\lambda_1(B)} - 1 \geq C_{FK} \mathcal{A}(\Omega)^4. \tag{A.2}$$

Recently Brasco *et al.* proved a sharp quantitative Faber–Krahn inequality [8] with the exponent 2 instead of 4 in (A.2). Unfortunately, their proof relies on the so called *selection principle*, which does not allow to get an explicit constant.

On the other hand a *quantitative Krahn–Szegő inequality* (2.4) was proved by Brasco and Pratelli in [7], based on (A.2). Their proof has the good feature of being easily adaptable once a better quantitative Faber–Krahn inequality is available.

Finding optimal constants in quantitative inequalities is a quite difficult task. Indeed, up to our knowledge, even the optimal constant for the quantitative isoperimetric inequality (A.1) in the plane is not explicitly known (see, for instance [5, 17]). The optimal constant for the quantitative isoperimetric inequality (A.1) is explicitly known only among *convex* sets of the plane, as was proved in [1]. Moreover, it is also difficult to find explicit non-optimal constants, since often the techniques used in the proofs do not allow to identify constants. Nevertheless, a non-optimal constant for the quantitative isoperimetric inequality (A.1) was obtained in [19]. The constant for the quantitative isoperimetric inequality is important when looking for explicit constants in the quantitative Faber–Krahn and Krahn–Szegő inequalities, since both proofs of (A.2) and (2.4) rely on the quantitative isoperimetric inequality (A.1). With the path just outlined, it is then possible to find an explicit constant for (2.4) for all $N \geq 2$, although the computations are rather unpleasant and the constants are far from being optimal.

However, in two dimensions, we can follow a different strategy. In order to get an explicit constant for the quantitative Krahn–Szegő inequality, we use the quantitative Faber–Krahn inequality provide by Bhattacharya in [6]. The key point of this inequality is that it does not rely on a quantitative isoperimetric inequality. Therefore, it can be obtained a better explicit constant β_{FK} for the quantitative Faber–Krahn inequality, which plugged into the proof of the quantitative Krahn–Szegő inequality (2.4) allows to improve the constant β_{KS} as well. To prove this result we go through the papers [6, 7] step-by-step and we highlight the points where an explicit constant is needed, using, in particular, the same notation of the paper involved.

A.1. An explicit constant in the quantitative Faber–Krahn inequality

The quantitative Faber–Krahn inequality in the form proved by Bhattacharya [6] reads as: there exists a constant $\beta_{\text{FK}} > 0$ such that for all open sets $\Omega \subset \mathbb{R}^2$ of finite measure $|\Omega| = |B|$,

$$fk(\Omega) := \frac{\lambda_1(\Omega)}{\lambda_1(B)} - 1 \geq \beta_{\text{FK}} \mathcal{A}(\Omega)^3. \tag{A.3}$$

We show that the constant β_{FK} in (A.3) can be chosen as follows:

$$\beta_{\text{FK}} = \frac{1}{10^5 \times 2^3 \times j_0^2} \approx 2.161 \times 10^{-7}. \tag{A.4}$$

We use the same notation of [6], noticing that the asymmetry α used there differs from (2.1) for a factor 2. Indeed using (2.3) for an open set $\Omega \subset \mathbb{R}^2$ of unit measure and $E = B$ we have that

$$\alpha(\Omega) = \frac{\mathcal{A}(\Omega)}{2}.$$

It is enough to study the case when $fk(\Omega) \leq 1$ since once

$$fk(\Omega) \geq K\alpha(\Omega)^3, \quad \text{if } fk(\Omega) \leq 1, \tag{A.5}$$

is established, then immediately (A.3) holds true with $\beta_{\text{FK}} = \min\{K, 1\}/2^3$ (recall that $\alpha(\Omega) \leq 1$ by definition). Therefore, let us assume the deficit $fk(\Omega) \leq 1$. The first point of [6] is Lemma 3.2: it is shown a bound on the L^∞ -norm of the first eigenfunction u_1 , that is

$$\|u_1\|_{L^\infty} \leq \frac{\lambda_1(\Omega)}{2\pi} \leq \frac{\lambda_1(B)}{\pi} = j_0^2, \tag{A.6}$$

where the second inequality holds thanks to the assumption $fk(\Omega) \leq 1$, while the last equality comes from the explicit value of $\lambda_1(B)$ recalled in (1.1).

Now we can directly pass to analyze Section 4 of [6], where the proof of the main result is carried out. We recall the constants that will be used:

$$p = 2, \quad k = \frac{1}{625} = \frac{1}{5^4}. \tag{A.7}$$

Only Case 1 is of our interest, since Case 2 deals with $p < 2$. In subcase (i) one gets

$$fk(\Omega) \geq \frac{1}{10^5 j_0^2} \alpha(\Omega)^3, \tag{A.8}$$

where we used (A.7) and M therein defined as an upper bound for $\|u_1\|_{L^\infty}$ and so, according to (A.6), can be taken as j_0^2 . On the other hand, subcase (ii) gives

$$fk(\Omega) \geq \frac{61}{200} \alpha(\Omega) \geq \frac{61}{200} \alpha(\Omega)^3, \tag{A.9}$$

where the second inequality holds since $\alpha(\Omega) \leq 1$. Then, combining (A.8) with (A.9) we get (A.5) with $K = 1/(10^5 \times j_0^2) < 1$, and therefore, from the previous observation $\beta_{\text{FK}} = K/2^3$, providing (A.4).

A.2. An explicit constant in the quantitative Krahn–Szegö inequality

We can now derive an explicit constant $\beta_{\text{KS}} > 0$ for (2.5). We go through the proof of the quantitative Krahn–Szegö inequality of [7] and give the explicit value of the constants that are needed in the proofs, using the quantitative Faber–Krahn (A.3) instead of (A.2), as it was originally done in that paper. First of all we need to give an explicit constant for Lemma 3.3 of [7], such that (using their notations, but numerating the constants $\vartheta_1, \vartheta_2, \dots$ in order to keep track of them in all steps)

$$\mathcal{A}_2(\Omega) \leq \vartheta_1 \left(\mathcal{A}(\Omega_+) + \left| \frac{1}{2} - \frac{|\Omega_+|}{|\Omega|} \right| + \mathcal{A}(\Omega_-) + \left| \frac{1}{2} - \frac{|\Omega_-|}{|\Omega|} \right| \right)^{2/3}. \tag{A.10}$$

This constant is deduced by putting together three main inequalities.

- (i) First of all we call $\varepsilon_{\pm} = \frac{1}{2} - \frac{|\Omega_{\pm}|}{|\Omega|}$ and call B_{\pm} two balls centered in the origin such that $|\Omega_{\pm}| = |B_{\pm}|$. Hence there exist two points x_{\pm} such that

$$\mathcal{A}(\Omega_{\pm}) = \frac{2|(B_{\pm} + x_{\pm}) \setminus \Omega_{\pm}|}{|\Omega_{\pm}|}.$$

We then rescale the balls to measure $|\Omega|/2$ each: $\tilde{B}_{\pm} = (1 - 2\varepsilon_{\pm})^{-1/2}B_{\pm}$. We have now to translate the new balls in the direction $x_+ - x_-$ so that they are disjoint. It is easy to see that the width l of the set $(\tilde{B}_+ + x_+) \cap (\tilde{B}_- + x_-)$ can be estimated by

$$l^{3/2}|\Omega|^{1/4} \leq \vartheta_2 |(\tilde{B}_+ + x_+) \cap (\tilde{B}_- + x_-)|,$$

with the choice of $\vartheta_2 = (8\pi)^{1/4}$.

- (ii) The second intermediate inequality is

$$|(\tilde{B}_+ + x_+) \cap (\tilde{B}_- + x_-)| \leq \vartheta_3 |\Omega| (\mathcal{A}(\Omega_+) + \mathcal{A}(\Omega_-) + |\varepsilon_+| + |\varepsilon_-|),$$

and it is possible to see immediately that $\vartheta_3 = 1$ works.

- (iii) We now have to translate \tilde{B}_- so that it is tangent to $\tilde{B}_+ + x_+$, and we will call \bar{x} the new center. It is possible to prove that

$$|(\tilde{B}_- + x_-) \cap (\tilde{B}_- + \bar{x})| \leq \vartheta_4 l |\Omega|^{1/2},$$

with the constant $\vartheta_4 = \frac{\pi+2}{(2\pi)^{1/2}}$.

At the end it is possible to put together the above inequalities and get (A.10) with $\vartheta_1 = \vartheta_4(\vartheta_2\vartheta_3)^{2/3} + 2 = \frac{\pi+2}{\pi^{1/3}} + 2$.

In order to conclude the proof of the quantitative Krahn–Szegö inequality by Brasco and Pratelli we have to prove a last intermediate inequality, when $ks(\Omega) \leq 1$:

$$ks(\Omega) \geq \frac{1}{\vartheta_5} \max \left\{ \mathcal{A}(\Omega_+)^3 + \left| \frac{1}{2} - \frac{|\Omega_+|}{|\Omega|} \right|, \mathcal{A}(\Omega_-)^3 + \left| \frac{1}{2} - \frac{|\Omega_-|}{|\Omega|} \right| \right\},$$

which works for $\vartheta_5 = \frac{3}{\beta_{\text{FK}}}$, where β_{FK} is the constant (A.4) of the quantitative Faber–Krahn inequality that we computed in the paragraph above. We note that in [7] the exponent of the asymmetries is actually 4 instead of 3, since they use the quantitative Faber–Krahn (A.2) instead of the one proved by Bhattacharya, which is only two dimensional. On the other hand, if $ks(\Omega) \geq 1$ it is enough to take $\vartheta_5 = 2^3 + 1 = 9$ since $\mathcal{A}(\Omega_{\pm}) \leq 2$ and $|\varepsilon_{\pm}| \leq 1/2$. So we have that $\vartheta_5 = \max \{9, 3/\beta_{\text{FK}}\} = 3/\beta_{\text{FK}}$. Putting all the inequalities together one arrives to

$$\begin{aligned} \beta_{\text{KS}} &= \frac{1}{\vartheta_5} \frac{1}{j_1^{9/2}} = \frac{\beta_{\text{FK}}}{3} \frac{\pi^{3/2}}{(\pi + 2 + 2\pi^{1/3})^{9/2}} \\ &= \frac{\pi^{3/2}}{24(\pi + 2 + 2\pi^{1/3})^{9/2} \cdot j_0^2} \times 10^{-5} \approx 3.331 \times 10^{-11}. \end{aligned}$$

Acknowledgements. We thank Bozhidar Velichkov for pointing out a link between Theorem 1.2 with ([14], Sect. 5.3).

REFERENCES

- [1] A. Alvino, V. Ferone and C. Nitsch, A sharp isoperimetric inequality in the plane. *J. Eur. Math. Soc.* **13** (2011) 185–206.
- [2] P. Antunes and A. Henrot, On the range of the first two Dirichlet and Neumann eigenvalues of the Laplacian. *Proc. R. Soc. Lond. Ser. A* **467** (2011) 1577–1603.
- [3] P. Antunes and A. Henrot, On the range of the first two Dirichlet eigenvalues of the Laplacian with volume and perimeter constraints. *Geometry of Solutions of Partial Differential Equations*. RIMS Publ. (2013) 1850.
- [4] M. Ashbaugh and R. Benguria, Proof of the Payne–Pólya–Weinberger conjecture. *Bull. Amer. Math. Soc.* **25** (1991) 19–29.
- [5] C. Bianchini, G. Croce and A. Henrot, On the quantitative isoperimetric inequality in the plane. *ESAIM: COCV* **23** (2017) 517–549.
- [6] T. Bhattacharya, Some observations on the first eigenvalue of the p-Laplacian and its connections with asymmetry. *Electron. J. Differ. Eq.* **2001** (2001) 1–15.
- [7] L. Brasco and A. Pratelli, Sharp stability of some spectral inequalities. *Geom. Funct. Anal.* **22** (2012) 107–135.
- [8] L. Brasco, G. De Philippis and B. Velichkov, Faber–Krahn inequalities in sharp quantitative form. *Duke Math. J.* **164** (2015) 1777–1831.
- [9] L. Brasco, C. Nitsch and A. Pratelli, On the boundary of the attainable set of the Dirichlet spectrum. *Z. Angew. Math. Phys.* **64** (2013) 591–597.
- [10] D. Bucur, Minimization of the k th eigenvalue of the Dirichlet Laplacian. *Arch. Ration. Mech. Anal.* **206** (2012) 1073–1083.
- [11] D. Bucur and G. Buttazzo, Variational methods in shape optimization problems. *Progress in Nonlinear Differential Equations and their Applications*. Birkhäuser Verlag, Boston (2005).
- [12] D. Bucur, G. Buttazzo and I. Figueredo, On the attainable eigenvalues of the Laplace operator. *SIAM J. Math. Anal.* **30** (1999) 527–536.
- [13] D. Bucur, G. Buttazzo and A. Henrot, Minimization of $\lambda_2(\Omega)$ with a perimeter constraint. *Indiana Univ. Math. J.* **58** (2009) 2709–2728.
- [14] D. Bucur, G. Buttazzo and B. Velichkov, Spectral optimization problems with internal constraint. *Ann. Inst. Henri Poincaré - Anal. Nonlinéaire* **30** (2013) 477–495.
- [15] D. Bucur, D. Mazzoleni, A. Pratelli and B. Velichkov, Lipschitz regularity of the eigenfunctions on optimal domains. *Arch. Rational Mech. Anal.* **216** (2015) 117–151.
- [16] G. Buttazzo and G. Dal Maso, An existence result for a class of shape optimization problems. *Arch. Rational Mech. Anal.* **122** (1993) 183–195.
- [17] M. Cicalese and G.P. Leonardi, Best constants for the isoperimetric inequality in quantitative form. *J. Eur. Math. Soc.* **15** (2013) 1101–1129.
- [18] G. De Philippis and B. Velichkov, Existence and regularity of minimizers for some spectral functionals with perimeter constraint. *Appl. Math. Optim.* **69** (2014) 199–231.
- [19] A. Figalli, F. Maggi and A. Pratelli, A mass transportation approach to quantitative isoperimetric inequalities. *Invent. Math.* **182** (2010) 167–211.
- [20] I. Fragalà and F. Gazzola, Partially overdetermined elliptic boundary value problems. *J. Differ. Eq.* **245** (2008) 1299–1322.
- [21] N. Fusco, F. Maggi and A. Pratelli, The sharp quantitative isoperimetric inequality. *Ann. Math.* **168** (2008) 941–980.
- [22] N. Fusco, F. Maggi and A. Pratelli, Stability estimates for certain Faber–Krahn, Isocapacitary and Cheeger inequalities. *Ann. Sc. Norm. Super. Pisa Cl. Sci.* **8** (2009) 51–71.
- [23] A. Henrot, Extremum Problems for Eigenvalues of Elliptic Operators. *Frontiers in Mathematics*. Birkhäuser Verlag, Basel (2006).
- [24] A. Henrot and E. Oudet, Minimizing the second eigenvalue of Laplace operator with Dirichlet boundary conditions. *Arch. Ration. Mech. Anal.* **169** (2003) 73–89.
- [25] A. Henrot and M. Pierre, Variation et optimisation de formes. Vol. 48 of *Mathématiques et Applications*. Springer (2005).
- [26] C.-Y. Kao and B. Osting, Minimal convex combinations of three sequential Laplace–Dirichlet eigenvalues. *Appl. Math. Optim.* **69** (2014) 123–139.
- [27] J.B. Keller and S.A. Wolf, Range of the First Two Eigenvalues of the Laplacian. *Proc. Roy. Soc. London Ser. A* **447** (1994) 397–412.
- [28] M. Iversen and D. Mazzoleni, Minimising convex combinations of low eigenvalues. *ESAIM: COCV* **20** (2014) 442–459.
- [29] D. Mazzoleni and A. Pratelli, Existence of minimizers for spectral problems. *J. Math. Pures Appl.* **100** (2013) 433–453.
- [30] E. Oudet, Numerical minimization of eigenmodes of a membrane with respect to the domain. *ESAIM: COCV* **10** (2004) 315–330.
- [31] J. Serrin, A symmetry problem in potential theory. *Arch. Ration. Mech. Anal.* **43** (1971) 304–318.