

MINIMAL 2-DOMINATING SETS IN TREES *

MARCIN KRZYWKOWSKI¹

Abstract. We provide an algorithm for listing all minimal 2-dominating sets of a tree of order n in time $\mathcal{O}(1.3248^n)$. This implies that every tree has at most 1.3248^n minimal 2-dominating sets. We also show that this bound is tight.

Mathematics Subject Classification. 05C05, 05C69, 05C85, 68R10, 68W05.

1. INTRODUCTION

Let $G = (V, E)$ be a graph. The order of a graph is the number of its vertices. By the neighborhood of a vertex v of G we mean the set $N_G(v) = \{u \in V(G) : uv \in E(G)\}$. The degree of a vertex v , denoted by $d_G(v)$, is the cardinality of its neighborhood. By a leaf we mean a vertex of degree one, while a support vertex is a vertex adjacent to a leaf. The distance between two vertices of a graph is the number of edges in a shortest path connecting them. The eccentricity of a vertex is the greatest distance between it and any other vertex. The diameter of a graph G , denoted by $\text{diam}(G)$, is the maximum eccentricity among all vertices of G . By P_n we denote a path on n vertices. By a star we mean a connected graph in which exactly one vertex has degree greater than one.

A subset $D \subseteq V(G)$ is a dominating set of G if every vertex of $V(G) \setminus D$ has a neighbor in D , while it is a 2-dominating set of G if every vertex of $V(G) \setminus D$ has at least two neighbors in D . A dominating (2-dominating, respectively) set D is minimal if no proper subset of D is a dominating (2-dominating, respectively) set of G .

Keywords and phrases. Domination, 2-domination, minimal 2-dominating set, tree, counting, exact exponential algorithm, listing algorithm.

* *The research was supported by the Polish National Science Centre grant 2011/02/A/ST6/00201.*

¹ Faculty of Electronics, Telecommunications and Informatics, Gdańsk University of Technology, Narutowicza 11/12, 80–233 Gdańsk, Poland. marcin.krzywkowski@gmail.com

A minimal 2-dominating set is abbreviated as m2ds. Note that 2-domination is a type of multiple domination in which each vertex, which is not in the dominating set, is dominated at least k times for a fixed positive integer k . Multiple domination was introduced by Fink and Jacobson [7], and further studied for example in [2, 10, 17]. For a comprehensive survey of domination in graphs, see [11, 12].

Observation 1.1. *Every leaf of a graph G is in every 2-dominating set of G .*

One of the typical questions in graph theory is how many subgraphs of a given property can a graph on n vertices have. For example, the famous Moon and Moser theorem [16] says that every graph on n vertices has at most $3^{n/3}$ maximal independent sets.

Combinatorial bounds are of interest not only on their own, but also because they are used for algorithm design as well. Lawler [15] used the Moon-Moser bound on the number of maximal independent sets to construct an $(1 + \sqrt[3]{3})^n \cdot n^{\mathcal{O}(1)}$ time graph coloring algorithm, which was the fastest one known for twenty-five years. In 2003 Eppstein [6] reduced the running time of a graph coloring to $\mathcal{O}(2.4151^n)$. In 2006 the running time was reduced [1, 14] to $\mathcal{O}(2^n)$. For an overview of the field, see [9].

Fomin *et al.* [8] constructed an algorithm for listing all minimal dominating sets of a graph on n vertices in time $\mathcal{O}(1.7159^n)$. There were also given graphs ($n/6$ disjoint copies of the octahedron) having $15^{n/6} \approx 1.5704^n$ minimal dominating sets. This establishes a lower bound on the running time of an algorithm for listing all minimal dominating sets of a given graph.

The number of maximal independent sets in trees was investigated in [18]. Coururier *et al.* [5] considered minimal dominating sets in various classes of graphs. The authors of [13] investigated the enumeration of minimal dominating sets in graphs.

Bród and Skupień [3] gave bounds on the number of dominating sets of a tree. They also characterized the extremal trees. The authors of [4] investigated the number of minimal dominating sets in trees containing all leaves.

We provide an algorithm for listing all minimal 2-dominating sets of a tree of order n in time $\mathcal{O}(1.3248^n)$. This implies that every tree has at most 1.3248^n minimal 2-dominating sets. We also show that this bound is tight.

2. LISTING ALGORITHM

In this section we describe a recursive algorithm which lists all minimal 2-dominating sets of a given input tree T . The iterator of the solutions is denoted by $\mathcal{F}(T)$.

Algorithm

Notice that the diameter of a tree can be easily determined in a polynomial time.

Let T be a tree. If $\text{diam}(T) = 0$, then $T = P_1 = v_1$. Let $\mathcal{F}(T) = \{\{v_1\}\}$. If $\text{diam}(T) = 1$, then $T = P_2 = v_1v_2$. Let $\mathcal{F}(T) = \{\{v_1, v_2\}\}$. If $\text{diam}(T) = 2$, then T is a star. By x we denote the support vertex of T . Let $\mathcal{F}(T) = \{V(T) \setminus \{x\}\}$.

Now consider trees T with $\text{diam}(T) \geq 3$. Thus the order n of the tree T is at least four.

If some support vertex of T , say x , is adjacent to at least three leaves (we denote one of them by y), then let $T' = T - y$ and

$$\mathcal{F}(T) = \{D' \cup \{y\} : D' \in \mathcal{F}(T')\}.$$

Now consider trees T , in which every support vertex is adjacent to at most two leaves. The tree T can easily be rooted at a vertex r of maximum eccentricity $\text{diam}(T)$ in polynomial time. A leaf, say t , at maximum distance from r , can also be easily computed in polynomial time. Let v denote the parent of t and let u denote the parent of v in the rooted tree. If $\text{diam}(T) \geq 4$, then let w denote the parent of u . By T_x we denote the subtree induced by a vertex x and its descendants in the rooted tree T .

If $d_T(v) = 3$, then by a we denote the leaf adjacent to v and different from t . Let $T' = T - T_v$ and $T'' = T - t - a$, and let $\mathcal{F}(T)$ be as follows,

$$\begin{aligned} & \{D' \cup \{t, a\} : D' \in \mathcal{F}(T')\} \\ & \cup \{D'' \cup \{t, a\} : D'' \in \mathcal{F}(T'') \text{ and } D'' \setminus \{v\} \notin \mathcal{F}(T')\}. \end{aligned}$$

If $d_T(v) = 2$ and $d_T(u) \geq 3$, then let $T' = T - T_v$, $T'' = T - T_u$, and

$$\mathcal{F}(T) = \{D' \cup \{t\} : u \in D' \in \mathcal{F}(T')\} \cup \{D'' \cup V(T_u) \setminus \{u\} : D'' \in \mathcal{F}(T'')\}.$$

If $d_T(v) = d_T(u) = 2$, then let $T' = T - T_v$, $T'' = T - T_u$, and

$$\mathcal{F}(T) = \{D' \cup \{t\} : D' \in \mathcal{F}(T')\} \cup \{D'' \cup \{v, t\} : w \in D'' \in \mathcal{F}(T'')\}.$$

3. BOUNDING THE NUMBER OF MINIMAL 2-DOMINATING SETS

Now we prove that the running time of the algorithm from the previous section is $\mathcal{O}(1.3248^n)$.

Theorem 3.1. *For every tree T of order n , the algorithm from the previous section lists all minimal 2-dominating sets in time $\mathcal{O}(1.3248^n)$.*

Proof. We prove that the running time of the algorithm is $\mathcal{O}(1.3248^n)$. Moreover, we prove that the number of minimal 2-dominating sets of T is at most α^n , where $\alpha \approx 1.3248$ is the positive solution of the equation $x^3 - x - 1 = 0$.

We proceed by induction on the number n of vertices of a tree T . If $\text{diam}(T) = 0$, then $T = P_1 = v_1$. Obviously, $\{v_1\}$ is the only m2ds of the path P_1 . We have $n = 1$ and $|\mathcal{F}(T)| = 1$. Obviously, $1 < \alpha$. If $\text{diam}(T) = 1$, then $T = P_2 = v_1v_2$. It is easy to see that $\{v_1, v_2\}$ is the only m2ds of the path P_2 . We have $n = 2$ and

$|\mathcal{F}(T)| = 1$. Obviously, $1 < \alpha^2$. If $\text{diam}(T) = 2$, then T is a star. By x we denote the support vertex of T . It is easy to observe that $V(T) \setminus \{x\}$ is the only m2ds of the tree T . We have $n \geq 3$ and $|\mathcal{F}(T)| = 1$. Obviously, $1 < \alpha^n$.

Now assume that $\text{diam}(T) \geq 3$. Thus the order n of the tree T is at least four. The results we obtain by the induction on the number n . Assume that they are true for every tree T' of order $n' < n$.

First assume that some support vertex of T , say x , is adjacent to at least three leaves. Let y be a leaf adjacent to x . Let $T' = T - y$. Let D' be a m2ds of the tree T' . Obviously, $D' \cup \{y\}$ is an m2ds of T . Thus all elements of $\mathcal{F}(T)$ are minimal 2-dominating sets of the tree T . Now let D be any m2ds of the tree T . By Observation 1.1 we have $y \in D$. Let us observe that $D \setminus \{y\}$ is an m2ds of the tree T' as the vertex x is still dominated at least twice. By the inductive hypothesis we have $D \setminus \{y\} \in \mathcal{F}(T')$. Therefore $\mathcal{F}(T)$ contains all minimal 2-dominating sets of the tree T . Now we get $|\mathcal{F}(T)| = |\mathcal{F}(T')| \leq \alpha^{n-1} < \alpha^n$. Henceforth, we can assume that every support vertex of T is adjacent to at most two leaves.

We now root T at a vertex r of maximum eccentricity $\text{diam}(T)$. Let t be a leaf at maximum distance from r , v be the parent of t , and u be the parent of v in the rooted tree. If $\text{diam}(T) \geq 4$, then let w be the parent of u . By T_x we denote the subtree induced by a vertex x and its descendants in the rooted tree T .

Assume that $d_T(v) = 3$. By a we denote the leaf adjacent to v and different from t . Let $T' = T - T_v$ and $T'' = T - t - a$. Let us observe that all elements of $\mathcal{F}(T)$ are minimal 2-dominating sets of the tree T . Now let D be any m2ds of the tree T . By Observation 1.1 we have $t, a \in D$. If $v \notin D$, then observe that $D \setminus \{t, a\}$ is an m2ds of the tree T' . By the inductive hypothesis we have $D \setminus \{t, a\} \in \mathcal{F}(T')$. Now assume that $v \in D$. Let us observe that $D \setminus \{t, a\}$ is an m2ds of the tree T'' . By the inductive hypothesis we have $D \setminus \{t, a\} \in \mathcal{F}(T'')$. The set $D \setminus \{v, t, a\}$ is not an m2ds of the tree T' , otherwise $D \setminus \{v\}$ is a 2-dominating set of the tree T , a contradiction to the minimality of D . By the inductive hypothesis we have $D \setminus \{v, t, a\} \notin \mathcal{F}(T')$. Therefore $\mathcal{F}(T)$ contains all minimal 2-dominating sets of the tree T . Now we get $|\mathcal{F}(T)| = |\mathcal{F}(T')| + |D'' \in \mathcal{F}(T'') : D'' \setminus \{v\} \notin \mathcal{F}(T')| \leq |\mathcal{F}(T')| + |\mathcal{F}(T'')| \leq \alpha^{n-3} + \alpha^{n-2} = \alpha^{n-3}(\alpha + 1) = \alpha^{n-3} \cdot \alpha^3 = \alpha^n$.

Now assume that $d_T(v) = 2$. Assume that $d_T(u) \geq 3$. Let $T' = T - T_v$ and $T'' = T - T_u$. Let us observe that all elements of $\mathcal{F}(T)$ are minimal 2-dominating sets of the tree T . Now let D be any m2ds of the tree T . By Observation 1.1 we have $t \in D$. If $v \notin D$, then $u \in D$ as the vertex v has to be dominated twice. Observe that $D \setminus \{t\}$ is an m2ds of the tree T' . By the inductive hypothesis we have $D \setminus \{t\} \in \mathcal{F}(T')$. Now assume that $v \in D$. We have $u \notin D$, otherwise $D \setminus \{v\}$ is a 2-dominating set of the tree T , a contradiction to the minimality of D . Observe that $D \cap V(T'')$ is an m2ds of the tree T'' . By the inductive hypothesis we have $D \cap V(T'') \in \mathcal{F}(T'')$. Therefore $\mathcal{F}(T)$ contains all minimal 2-dominating sets of the tree T . Now we get $|\mathcal{F}(T)| \leq |\mathcal{F}(T')| + |\mathcal{F}(T'')| \leq \alpha^{n-2} + \alpha^{n-3} = \alpha^{n-3}(\alpha + 1) = \alpha^{n-3} \cdot \alpha^3 = \alpha^n$.

Now assume that $d_T(u) = 2$. Let $T' = T - T_v$ and $T'' = T - T_u$. Let us observe that all elements of $\mathcal{F}(T)$ are minimal 2-dominating sets of the tree T . Now let D be any m2ds of the tree T . By Observation 1.1 we have $t \in D$. If $v \notin D$, then observe that $D \setminus \{t\}$ is an m2ds of the tree T' . By the inductive hypothesis we have $D \setminus \{t\} \in \mathcal{F}(T')$. Now assume that $v \in D$. We have $u \notin D$, otherwise $D \setminus \{v\}$ is a 2-dominating set of the tree T , a contradiction to the minimality of D . Moreover, we have $w \in D$ as the vertex u has to be dominated twice. Observe that $D \setminus \{v, t\}$ is an m2ds of the tree T'' . By the inductive hypothesis we have $D \setminus \{v, t\} \in \mathcal{F}(T'')$. Therefore $\mathcal{F}(T)$ contains all minimal 2-dominating sets of the tree T . Now we get $|\mathcal{F}(T)| \leq |\mathcal{F}(T')| + |\mathcal{F}(T'')| \leq \alpha^{n-2} + \alpha^{n-3} = \alpha^{n-3}(\alpha + 1) = \alpha^{n-3} \cdot \alpha^3 = \alpha^n$. \square

It follows from the proof of the above theorem that any tree of order n has at most 1.3248^n minimal 2-dominating sets.

Corollary 3.2. *Every tree of order n has at most α^n minimal 2-dominating sets, where $\alpha \approx 1.3248$ is the positive solution of the equation $x^3 - x - 1 = 0$.*

Now we show that the bound from the previous corollary is tight. Let a_n denote the number of minimal 2-dominating sets of the path P_n . The next remark follows from the Proof of Theorem 3.1.

Remark 3.3. For every positive integer n we have

$$a_n = \begin{cases} 1 & \text{if } n \leq 3; \\ a_{n-3} + a_{n-2} & \text{if } n \geq 4. \end{cases}$$

We have $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = \alpha$, where $\alpha \approx 1.3247$ is the positive solution of the equation $x^3 - x - 1 = 0$. This implies that the bound from Corollary 3.2 is tight.

REFERENCES

- [1] A. Björklund and T. Husfeldt, Inclusion-exclusion algorithms for counting set partitions. *Proc. of FOCS* (2006) 575–582.
- [2] M. Blidia, O. Favaron and R. Lounes, Locating-domination, 2-domination and independence in trees. *Australas. J. Combin.* **42** (2008) 309–316.
- [3] D. Bród and Z. Skupień, Trees with extremal numbers of dominating sets. *Australas. J. Combin.* **35** (2006) 273–290.
- [4] D. Bród, A. Włoch and I. Włoch, On the number of minimal dominating sets including the set of leaves in trees. *Internat. J. Contemporary Math. Sci.* **4** (2009) 1739–1748.
- [5] J.-F. Couturier, P. Heggernes, P. van ’t Hof and D. Kratsch, Minimal dominating sets in graph classes: combinatorial bounds and enumeration. *Proc. of SOFSEM 2012, Lect. Notes Comput. Sci.*, vol. 7147. Springer-Verlag, Berlin (2012) 202–213.
- [6] D. Eppstein, Small maximal independent sets and faster exact graph coloring, *J. Graph Algorithms Appl.* **7** (2003) 131–140.
- [7] J. Fink and M. Jacobson, n-domination in graphs, *Graph Theory with Applications to Algorithms and Computer Science*. Wiley, New York (1985) 282–300.
- [8] F. Fomin, F. Grandoni, A. Pyatkin and A. Stepanov, Combinatorial bounds via measure and conquer: bounding minimal dominating sets and applications. *ACM Transactions on Algorithms*, vol. 5, article 9 (2009).

- [9] F. Fomin and D. Kratsch, *Exact Exponential Algorithms*. Springer, Berlin (2010).
- [10] J. Fujisawa, A. Hansberg, T. Kubo, A. Saito, M. Sugita and L. Volkmann, Independence and 2-domination in bipartite graphs. *Australas. J. Combin.* **40** (2008) 265–268.
- [11] T. Haynes, S. Hedetniemi and P. Slater, *Fundamentals of Domination in Graphs*. Marcel Dekker, New York (1998).
- [12] T. Haynes, S. Hedetniemi and P. Slater, *Domination in Graphs: Advanced Topics*. Marcel Dekker, New York (1998).
- [13] M. Kanté, V. Limouzy, A. Mary and L. Nourine, Enumeration of minimal dominating sets and variants, Fundamentals of computation theory. *Lect. Notes Comput. Sci.*, vol. 6914. Springer, Heidelberg (2011) 298–309.
- [14] M. Koivisto, An $\mathcal{O}(2^n)$ algorithm for graph coloring and other partitioning problems via inclusion-exclusion. *Proc. 47th Annual IEEE Symposium on Foundations of Comput. Sci.* (FOCS 2006), IEEE (2006) 583–590.
- [15] E. Lawler, A note on the complexity of the chromatic number problem. *Information Proc. Lett.* **5** (1976) 66–67.
- [16] J. Moon and L. Moser, On cliques in graphs. *Israel J. Math.* **3** (1965) 23–28.
- [17] R. Shaheen, Bounds for the 2-domination number of toroidal grid graphs. *Internat. J. Comput. Math.* **86** (2009) 584–588.
- [18] H. Wilf, The number of maximal independent sets in a tree. *SIAM J. Algebr. Discrete Methods* **7** (1986) 125–130.

Communicated by B. Vallée.

Received May 7, 2012. Accepted April 30, 2013.