

A SHORT PROOF THAT SHUFFLE SQUARES ARE 7-AVOIDABLE

GUILLAUME GUÉGAN¹ AND PASCAL OCHEM²

Abstract. A shuffle square is a word that can be partitioned into two identical words. We obtain a short proof that there exist exponentially many words over the 7 letter alphabet containing no shuffle square as a factor. The method is a generalization of the so-called power series method using ideas of the entropy compression method as developed by Gonçalves *et al.* [Entropy compression method applied to graph colorings. [arXiv:1406.4380](https://arxiv.org/abs/1406.4380)]

Mathematics Subject Classification. 68R15.

1. INTRODUCTION

Entropy compression has been used to avoid squares [5] and patterns [9] in infinite words over a small alphabet. The proofs require many features (an algorithm, a record, an analysis of the size the record, ...). Gonçalves, Montassier, and Pinlou [4] have recently obtained a generic way of using the entropy compression method in the context of graph coloring that avoids a lot of these technicalities.

In a recent paper [8], we have used ideas from the entropy compression method to generalize the power series method as used in combinatorics on words by Bell and Goh [1], Rampersad [10], and Blanchet–Sadri and Woodhouse [2]. We describe this method in Section 2 to make the paper self-contained.

A shuffle square is a word that can be partitioned into two identical words. For example, every square is a shuffle square, **aabbcc** and **abacbc** are shuffle squares of *abc*, and **cbcbbaca** is a shuffle square of *cbca*.

Recently, Currie [3] has answered a question of Karhumäki by showing that there exist infinite words over a finite (but large) alphabet containing no shuffle square as a factor using the Lovász local lemma. Then Müller has lowered the alphabet size to 10 in his thesis [7] and has also proved that shuffle cubes are avoidable over the 6 letter alphabet. We apply the method in Section 2 to obtain the following result in Section 3.

Theorem 1.1. *There exist at least 5.59^n words of length n over the 7 letter alphabet containing no shuffle square as a factor.*

Grytczuk *et al.* [6] have an independent proof of the list version of Theorem 1.1 using another flavor of entropy compression and different parameters. Notice that words avoiding shuffle squares avoid in particular the patterns *AA* and *ABACBC*. We have checked that words over 3 letters avoiding *AA* and *ABACBC* have

Keywords and phrases. Combinatorics on words, shuffle square, entropy compression.

¹ University Montpellier 2, LIRMM, France. guegan@lirmm.fr

² CNRS, LIRMM, Montpellier, France. ochem@lirmm.fr

finite length, so at least 4 letters are needed to avoid shuffle squares. Thus, the minimum alphabet size for an infinite word avoiding shuffle squares remains an open problem and is between 4 and 7.

2. DESCRIPTION OF THE METHOD

Let $\Sigma_m = \{0, 1, \dots, m-1\}$ be the m -letter alphabet and let $L \subset \Sigma_m^*$ be a factorial language defined by a set F of forbidden factors of length at least 2. We denote the factor complexity of L by $n_i = L \cap \Sigma_m^i$. We define L' as the set of words w such that w is not in L and the prefix of length $|w| - 1$ of w is in L . For every forbidden factor $f \in F$, we choose a number $1 \leq s_f \leq |f|$. Then, for every $i \geq 1$, we define an integer a_i such that

$$a_i \geq \max_{u \in L} |\{v \in \Sigma_m^i \mid uv \in L', uv = bf, f \in F, s_f = i\}|. \quad (2.1)$$

We consider the formal power series $P(x) = 1 - mx + \sum_{i \geq 1} a_i x^i$. If $P(x)$ has a positive real root x_0 , then $n_i \geq x_0^{-i}$ for every $i \geq 0$.

Let us rewrite that $P(x_0) = 1 - mx_0 + \sum_{i \geq 1} a_i x_0^i = 0$ as:

$$m - \sum_{i \geq 1} a_i x_0^{i-1} = x_0^{-1}. \quad (2.2)$$

Since $n_0 = 1$, we will prove by induction that $\frac{n_i}{n_{i-1}} \geq x_0^{-1}$ in order to obtain that $n_i \geq x_0^{-i}$ for every $i \geq 0$. By using (2.2), we obtain the base case: $\frac{n_1}{n_0} = n_1 = m \geq x_0^{-1}$. Now, for every length $i \geq 1$, there are:

- m^i words in Σ_m^i ;
- n_i words in L ;
- at most $\sum_{1 \leq j \leq i} n_{i-j} a_j$ words in L' ;
- $m(m^{i-1} - n_{i-1})$ words in $\Sigma_m^i \setminus \{L \cup L'\}$.

This gives $n_i + \sum_{1 \leq j \leq i} n_j a_{i-j} + m(m^{i-1} - n_{i-1}) \geq m^i$, that is, $n_i \geq mn_{i-1} - \sum_{1 \leq j \leq i} n_{i-j} a_j$.

$$\begin{aligned} \frac{n_i}{n_{i-1}} &\geq m - \sum_{1 \leq j \leq i} a_j \frac{n_{i-j}}{n_{i-1}} \\ &\geq m - \sum_{1 \leq j \leq i} a_j x_0^{j-1} && \text{By induction} \\ &\geq m - \sum_{j \geq 1} a_j x_0^{j-1} \\ &= x_0^{-1} && \text{By (2.2).} \end{aligned}$$

3. AVOIDING SHUFFLE SQUARES

We apply the method of the previous section to the avoidance of shuffle squares. The q -*prefix* (resp. q -*suffix*) of a word is its prefix (resp. suffix) of length q . A shuffle square is *minimal* if it does not contain a smaller shuffle square as a factor. A shuffle square is *small* if its length is two and is *large* otherwise. The set F of forbidden factors contains every minimal shuffle square. We set $s_f = 1$ if $f \in F$ is small and $s_f = |f| - 2$ otherwise.

We set $a_1 = 1$ because $s_f = 1$ only for small shuffle squares and there is only one way to extend a prefix by one letter to obtain a suffix xx with $x \in \Sigma_m$. To obtain reasonable upper bounds a_t for $t \geq 2$, we need to bound the number of large minimal shuffle squares. To every shuffle square f of a word w of length i , we associate the *height function* $h: [0, \dots, 2i] \rightarrow \mathbb{Z}$ defined as follows:

- $h(0) = 0$.
- For $0 < j \leq 2i$, $h(j) = h(j-1) + 1$ if the j th letter of f belongs to the subword w containing the first letter of f , and $h(j) = h(j-1) - 1$ otherwise.

Since f is a shuffle square, we have $h(2i) = 0$. Moreover, if $h(j) = 0$ for some $0 < j < 2i$, then the prefix of length j of f is a shuffle square. So, if h is the height function of a minimal shuffle square, then $h(j) > 0$ for every $0 < j < 2i$. Thus, every height function of a minimal shuffle square is associated to a unique Dyck word of length $2i - 2$. The number of height functions is thus at most $\frac{(2i-2)!}{i!(i-1)!}$. According to (2.1), we need to bound the number of solutions to $uv = bf$ such that u is fixed and $|v| = s_f = |f| - 2 = 2i - 2$. The 2-prefix of f is fixed since it corresponds to the 2-suffix of u . Notice that the 2-prefix of a large minimal shuffle square of a word w is equal to the 2-prefix of w , so the 2-prefix of w is also fixed. Thus, there are at most m^{i-2} possibilities for w . Since f is determined by w and its height function, there are at most $m^{i-2} \frac{(2i-2)!}{i!(i-1)!}$ possibilities for f . So we set $a_{2i-2} = m^{i-2} \frac{(2i-2)!}{i!(i-1)!}$ and consider the polynomial

$$\begin{aligned} P(x) &= 1 - mx + x + \sum_{i \geq 2} m^{i-2} \frac{(2i-2)!}{i!(i-1)!} x^{2i-2} \\ &= 1 - (m-1)x + \left(\frac{2x}{1 + \sqrt{1-4mx^2}} \right)^2. \end{aligned}$$

For $m = 6$, $P(x)$ has no positive root. For $m = 7$, we have $P(x_0) = 0$ with $x_0 = 0.1788487593\dots$. So there exists at least α^n words of length n over Σ_7 that avoid shuffle squares, where $\alpha = x_0^{-1} = 5.5913163944\dots$

REFERENCES

- [1] J. Bell and T.L. Goh, Exponential lower bounds for the number of words of uniform length avoiding a pattern. *Inform. Comput.* **205** (2007) 1295–1306.
- [2] F. Blanchet-Sadri and B. Woodhouse, Strict bounds for pattern avoidance. *Theor. Comput. Sci.* **506** (2013) 17–27.
- [3] J. Currie, Shuffle squares are avoidable. Manuscript.
- [4] D. Gonçalves, M. Montassier and A. Pinlou, Entropy compression method applied to graph colorings. Preprint [arXiv:1406.4380](https://arxiv.org/abs/1406.4380) (2015).
- [5] J. Grytczuk, J. Kozik and P. Micek, A new approach to nonrepetitive sequences. *Random Structures & Algorithms* **42** (2013) 214–225.
- [6] J. Grytczuk, J. Kozik and B. Zaleski, Avoiding tight twins in sequences by entropy compression. Available at <http://ssdm.mimuw.edu.pl/pliki/prace-studentow/st/pliki/bartosz-zaleski-3.pdf>
- [7] M. Müller, *Avoiding and enforcing repetitive structures in words*. Ph.D. thesis (2014).
- [8] P. Ochem, Doubled patterns are 3-avoidable. Preprint [arXiv:1510.01753](https://arxiv.org/abs/1510.01753) (2015).
- [9] P. Ochem and A. Pinlou, Application of entropy compression in pattern avoidance. *Electron. J. Combin.* **21** (2014) #RP2.7.
- [10] N. Rampersad, Further applications of a power series method for pattern avoidance. *Electron. J. Combin.* **18** (2011) #P134.

Communicated by D. Jamet.

Received March 24, 2016. Accepted March 24, 2016.