

quatrième série - tome 49 fascicule 3 mai-juin 2016

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Vincent HUMILIÈRE & Rémi LECLERCQ & Sobhan SEYFADDINI

Reduction of symplectic homeomorphisms

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Antoine CHAMBERT-LOIR

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE
de 1883 à 1888 par H. DEBRAY
de 1889 à 1900 par C. HERMITE
de 1901 à 1917 par G. DARBOUX
de 1918 à 1941 par É. PICARD
de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} janvier 2016

N. ANANTHARAMAN I. GALLAGHER
P. BERNARD B. KLEINER
E. BREUILLARD E. KOWALSKI
R. CERF M. MUSTAȚĂ
A. CHAMBERT-LOIR L. SALOFF-COSTE

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.
annales@ens.fr

Édition / *Publication*

Société Mathématique de France
Institut Henri Poincaré
11, rue Pierre et Marie Curie
75231 Paris Cedex 05
Tél. : (33) 01 44 27 67 99
Fax : (33) 01 40 46 90 96

Abonnements / *Subscriptions*

Maison de la SMF
Case 916 - Luminy
13288 Marseille Cedex 09
Fax : (33) 04 91 41 17 51
email : smf@smf.univ-mrs.fr

Tarifs

Europe : 515 €. Hors Europe : 545 €. Vente au numéro : 77 €.

© 2016 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593

Directeur de la publication : Stéphane Seuret
Périodicité : 6 n^{os} / an

REDUCTION OF SYMPLECTIC HOMEOMORPHISMS

BY VINCENT HUMILIÈRE, RÉMI LECLERCQ
AND SOBHAN SEYFADDINI

ABSTRACT. – In [9], we proved that symplectic homeomorphisms preserving a coisotropic submanifold C , preserve its characteristic foliation as well. As a consequence, such symplectic homeomorphisms descend to the reduction of the coisotropic C .

In this article we show that these reduced homeomorphisms continue to exhibit certain symplectic properties. In particular, in the specific setting where the symplectic manifold is a torus and the coisotropic is a standard subtorus, we prove that the reduced homeomorphism preserves spectral invariants and hence the spectral capacity.

To prove our main result, we use Lagrangian Floer theory to construct a new class of spectral invariants which satisfy a non-standard triangle inequality.

RÉSUMÉ. – Nous avons démontré dans [9], qu'un homéomorphisme symplectique qui laisse invariante une sous-variété coïsothrope C , préserve également son feuilletage caractéristique. Il induit donc un homéomorphisme sur la réduction symplectique de C .

Dans cet article, nous démontrons que l'homéomorphisme ainsi obtenu exhibe certaines propriétés symplectiques. En particulier, dans le cas où la variété symplectique ambiante est un tore et la sous-variété coïsothrope est un sous-tore standard, nous démontrons que l'homéomorphisme réduit préserve les invariants spectraux et donc aussi la capacité spectrale.

Pour démontrer notre résultat principal, nous construisons, à l'aide de l'homologie de Floer lagrangienne, une nouvelle famille d'invariants spectraux qui satisfont un nouveau type d'inégalité triangulaire.

1. Introduction

1.1. Context and main result

The main objects under study in this paper are symplectic homeomorphisms. Given a symplectic manifold (M, ω) , a homeomorphism $\phi : M \rightarrow M$ is called a symplectic homeomorphism if it is the C^0 -limit of a sequence of symplectic diffeomorphisms. This definition is motivated by a celebrated theorem due to Gromov and Eliashberg which asserts

that if a symplectic homeomorphism ϕ is smooth, then it is a symplectic diffeomorphism in the usual sense: $\phi^*\omega = \omega$.

Understanding the extent to which symplectic homeomorphisms behave like their smooth counterparts constitutes the central theme of C^0 -symplectic geometry. A recent discovery of Buhovsky and Opshtein suggests that these homeomorphisms are capable of exhibiting far more flexibility than symplectic diffeomorphisms: In [5], they construct an example of a symplectic homeomorphism of the standard \mathbb{C}^3 whose restriction to the symplectic subspace $\mathbb{C} \times \{0\} \times \{0\}$ is the contraction $(z, 0, 0) \mapsto (\frac{1}{2}z, 0, 0)$. Such behavior is impossible for a symplectic diffeomorphism but of course very typical for a volume-preserving homeomorphism. On the other hand, it is well-known that symplectic homeomorphisms are surprisingly rigid in comparison to volume-preserving maps. The following example of rigidity is the starting point of this article: Recall that a coisotropic submanifold is a submanifold $C \subset M$ whose tangent space, at every point of C , contains its symplectic orthogonal: $TC^\omega \subset TC$. Moreover, the distribution TC^ω is integrable and the foliation it spans is called the characteristic foliation of C .

THEOREM 1 ([9]). – *Let C be a smooth coisotropic submanifold of a symplectic manifold (M, ω) . Let ϕ denote a symplectic homeomorphism. If $C' = \phi(C)$ is smooth, then it is coisotropic. Furthermore, ϕ maps the characteristic foliation of C to that of C' .*

Prior to the discovery of the above theorem, the special cases of Lagrangian submanifolds and hypersurfaces have been treated, respectively, by Laudénbach-Sikorav [11] and Opshtein [17].

We are now in position to describe the problem we are interested in. Denote by \mathcal{F} and \mathcal{F}' , respectively, the characteristic foliations of the coisotropics C and C' from the above theorem. The reduced spaces $\mathcal{R} = C/\mathcal{F}$ and $\mathcal{R}' = C'/\mathcal{F}'$ are defined as the quotients of the coisotropic submanifolds by their characteristic foliations. These spaces are, at least locally, smooth manifolds and they can be equipped with natural symplectic structures induced by ω . Since $\phi(\mathcal{F}) = \mathcal{F}'$, the homeomorphism ϕ induces a homeomorphism $\phi_R : \mathcal{R} \rightarrow \mathcal{R}'$ of the reduced spaces. It is a classical fact that when ϕ is smooth, and hence symplectic, the reduced map ϕ_R is a symplectic diffeomorphism as well. It is therefore natural to ask whether the homeomorphism ϕ_R remains symplectic, in any sense, when ϕ is not assumed to be smooth. This is the question we seek to answer in this article.

We begin by first supposing that the reduction ϕ_R is smooth. It turns out that this scenario can be resolved rather easily using a result of [9].

PROPOSITION 2. – *Let C be a coisotropic submanifold whose reduction \mathcal{R} is a symplectic manifold⁽¹⁾, and ϕ be a symplectic homeomorphism. Assume that $C' = \phi(C)$ is smooth and therefore is coisotropic and admits a reduction \mathcal{R}' . Denote by $\phi_R : \mathcal{R} \rightarrow \mathcal{R}'$ the map induced by ϕ . Then, if ϕ_R is smooth, it is symplectic.*

We would like to point out that a similar result, with a similar proof, has already appeared in [5] (See Proposition 6).

⁽¹⁾ This is always locally true.

Proof. – We will prove that for any smooth function f_R on \mathcal{R}' , the Hamiltonian flow generated by the function $f_R \circ \phi_R$ is $\phi_R^{-1} \phi_{f_R}^t \phi_R$, where $\phi_{f_R}^t$ is the Hamiltonian flow generated by f_R . It is not hard to conclude from this that ϕ_R is symplectic: For example, it can easily be checked that ϕ_R preserves the Poisson bracket, i.e., $\{h_R \circ \phi_R, g_R \circ \phi_R\} = \{h_R, g_R\} \circ \phi_R$ for any two smooth functions h_R, g_R on \mathcal{R}' .

Let $f_R: \mathcal{R}' \rightarrow \mathbb{R}$ be smooth. We denote by $g_R: \mathcal{R} \rightarrow \mathbb{R}$ the function defined by $g_R = f_R \circ \phi_R$. Let f and g be any smooth lifts to M of f_R and g_R , respectively.

First, notice that by definition the restrictions to C of $f \circ \phi$ and g coincide. Since g is constant on the characteristic leaves of C , its Hamiltonian flow ϕ_g^t preserves C . Thus $H = (f \circ \phi - g) \circ \phi_g^t$ vanishes on C for all t . By [9, Theorem 3], the flow of the continuous Hamiltonian $(2)H$ follows the characteristic leaves of C . On the other hand we know that this flow is given by the formula $\phi_H^t = (\phi_g^t)^{-1} \phi^{-1} \phi_f^t \phi$. This isotopy descends to the reduction \mathcal{R} where it induces the isotopy $(\phi_{g_R}^t)^{-1} \phi_R^{-1} \phi_{f_R}^t \phi_R$. But since ϕ_H^t follows characteristics it must descend to the identity. Hence $(\phi_{g_R}^t)^{-1} \phi_R^{-1} \phi_{f_R}^t \phi_R = \text{Id}$ as claimed. \square

When ϕ_R is not assumed to be smooth, the situation becomes far more complicated. The question of whether or not ϕ_R is a symplectic homeomorphism seems to be very difficult and, at least currently, completely out of reach. Given the difficulty of this question, one could instead ask if there exist symplectic invariants which are preserved by reduced homeomorphisms. In this spirit, and since symplectic homeomorphisms are capacity preserving, Opshtein formulated the following a priori easier problem:

QUESTION 3. – *Is the reduction ϕ_R of a symplectic homeomorphism ϕ preserving a coisotropic submanifold always a capacity preserving homeomorphism?*

Partial positive results have been obtained by Buhovsky and Opshtein [5]. They proved in particular that in the case where C is a hypersurface, the map ϕ_R is a “non-squeezing map” in the sense that for every open set U containing a symplectic ball of radius r , the image $\phi_R(U)$ cannot be embedded in a symplectic cylinder over a 2-disk of radius $R < r$. This does not resolve Opshtein’s question, but since capacity preserving maps are non-squeezing it does provide positive evidence for it. In the case of general coisotropic submanifolds, they conjecture that the same holds and indicate as to how one might approach this conjecture.

In this article, we work in the specific setting where M is the torus $\mathbb{T}^{2(k_1+k_2)}$ equipped with its standard symplectic structure and $C = \mathbb{T}^{2k_1+k_2} \times \{0\}^{k_2}$. The reduction of C is \mathbb{T}^{2k_1} with its usual symplectic structure. Our main theorem shows that, in this setting, the reduced homeomorphism ϕ_R preserves certain symplectic invariants referred to as spectral invariants. This answers Opshtein’s question positively, as it follows immediately that the spectral capacity is preserved by ϕ_R .

More precisely, for a time-dependent Hamiltonian H , denote by $c_+(H)$ the spectral invariant, defined by Schwarz [20], associated to the fundamental class of M . Roughly speaking, $c_+(H)$ is the action value at which the fundamental class $[M]$ appears in the Floer homology of H ; see Equation (12) in Section 2.3 for the precise definition. (We should caution the reader that our notations and conventions are different than those of [20]. For example, $c_+(H)$ in this article corresponds to $c(1; H)$ in [20] where 1 is the generator

⁽²⁾ The continuous function H generates a continuous flow in the sense defined by Müller and Oh [16].

of $H^0(M)$.)⁽³⁾ For degenerate or continuous functions one defines $c_+(H) = \lim_{i \rightarrow \infty} c_+(H_i)$ where H_i is a sequence of smooth non-degenerate Hamiltonians converging uniformly to H . This limit is well-defined because c_+ satisfies a well-known Lipschitz estimate. We refer the reader to Section 2.3 for further details. Here is our main result:

THEOREM 4. – *Let ϕ be a symplectic homeomorphism of the torus $\mathbb{T}^{2(k_1+k_2)}$ equipped with its standard symplectic form. Assume that ϕ preserves the coisotropic submanifold $C = \mathbb{T}^{2k_1+k_2} \times \{0\}^{k_2}$. Denote by ϕ_R the induced homeomorphism on the reduced space $\mathcal{R} = \mathbb{T}^{2k_1}$. Then, for every time-dependent continuous function H on $[0, 1] \times \mathcal{R}$, we have:*

$$c_+(H \circ \phi_R) = c_+(H),$$

where $H \circ \phi_R(t, x) := H(t, \phi_R(x))$.

Note that Theorem 4 implies that other related symplectic invariants which are constructed using spectral invariants are also preserved by ϕ_R . Here is one example of such invariants: Following Viterbo [[22], Definition 4.11], we define the spectral capacity of an open set U , denoted by $c(U)$, by

$$c(U) = \sup\{c_+(H) \mid H \in C^0([0, 1] \times M), \text{ support}(H_t) \subset U \ \forall t \in [0, 1]\},$$

where $C^0([0, 1] \times M)$ denotes the space of time-dependent continuous functions on M . The following is an immediate corollary of Theorem 4.

COROLLARY 5. – *The map ϕ_R , from Theorem 4, preserves the spectral capacity, i.e., $c(\phi_R(U)) = c(U)$ for any open set U .*

In Definition 5.15 of [20], Schwarz defines a very similar capacity which he denotes by c_γ . It can easily be checked that ϕ_R preserves c_γ as well.

1.2. Main Tools: Lagrangian Floer theory and spectral invariants

For proving Theorem 4, we will use the theory of Lagrangian spectral invariants. These invariants were first introduced by Viterbo [22] in the setting of cotangent bundles and using generating functions. In [15], Oh reconstructed the same invariants using Lagrangian Floer homology. There have been many developments in the theory since then; see Section 2 for specific references.

In this article, we will use Lagrangian Floer homology, in the specific setting where the symplectic manifold and the Lagrangians are all tori, to construct a new class of spectral invariants. Below, we describe our settings and give a brief overview of the construction and properties of the particular spectral invariants which will be used in the proof of Theorem 4.

The symplectic manifold we will be working on is the product

$$M = \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2} \times \mathbb{T}^{2k_2}.$$

⁽³⁾ In [20], after constructing $c(1; H)$ the author proceeds to normalize the Hamiltonian H by requiring that $\int_M H(t, x)\omega^n = 0$ for each $t \in [0, 1]$. This leads to an invariant of Hamiltonian diffeomorphisms, $c(1; \phi_H^1)$.

We denote by (q_1, p_1) and (Q_1, P_1) the coordinates on the first and second \mathbb{T}^{2k_1} factors in the above product, respectively. The coordinates (q_2, p_2) and (Q_2, P_2) are defined similarly. We equip M with the standard symplectic structure given by

$$\omega_{\text{std}} = dq_1 \wedge dp_1 + dQ_1 \wedge dP_1 + dq_2 \wedge dp_2 + dQ_2 \wedge dP_2 .$$

The Lagrangian submanifolds of M whose Floer homology we will be studying are

$$\begin{aligned} L_0 &= \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\} , \\ L_1 &= \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\} \times \{0\} \times \mathbb{T}^{k_2} . \end{aligned}$$

Notice that both L_0 and L_1 decompose as products of smaller Lagrangians, i.e., $L_i = L \times L'_i$, where

$$\begin{aligned} L &= \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\} \subset \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2} , \\ L'_0 &= \mathbb{T}^{k_2} \times \{0\} \subset \mathbb{T}^{2k_2}, \text{ and } L'_1 = \{0\} \times \mathbb{T}^{k_2} \subset \mathbb{T}^{2k_2} . \end{aligned}$$

Observe that $L_0 \cap L_1 = L \times \{0\}$. In Section 3.1, we will construct an isomorphism between the Morse homology of L , denoted by $HM(L)$, and the Floer homology group $HF(L_0, L_1)$; see Theorem 18 for a precise statement. We will then use this isomorphism to associate a critical value of the Lagrangian action functional $\mathcal{A}_H^{L_0, L_1}$ to a non-zero class $a \in HM(L)$ and a Hamiltonian $H : [0, 1] \times M \rightarrow \mathbb{R}$. We will denote this critical value by

$$\ell(a; L_0, L_1; H).$$

This is the spectral invariant associated to a and H . Roughly speaking, $\ell(a; L_0, L_1; H)$ is the action value at which the Morse homology class a appears in the Floer homology group $HF(L_0, L_1)$.

Main properties of spectral invariants. – We now list some of the main properties of the spectral invariant $\ell(a; L_0, L_1; H)$.

1. Spectrality: Let $\text{Spec}(H)$ denote the set of critical values of the action functional $\mathcal{A}_H^{L_0, L_1}$. Then, for any Hamiltonian H and $a \in HM(L) \setminus \{0\}$,

$$\ell(a; L_0, L_1; H) \in \text{Spec}(H).$$

For further details, see Section 2.2.1.

2. Continuity: The following inequality holds for any Hamiltonians H, H'

$$\begin{aligned} \int_0^1 \min_M (H_t - H'_t) dt &\leq |\ell(a; L_0, L_1; H) - \ell(a; L_0, L_1; H')| \\ &\leq \int_0^1 \max_M (H_t - H'_t) dt . \end{aligned}$$

For further details, see Sections 2.2 and 3.1.

3. Splitting Formula: Let F and F' denote two Hamiltonians on $\mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}$ and \mathbb{T}^{2k_2} , respectively. Define the Hamiltonian $F \oplus F'$ on M by $F \oplus F'(z_1, z_2) = F(z_1) + F'(z_2)$, for $z_1 \in \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}$ and $z_2 \in \mathbb{T}^{2k_2}$. In Section 3.3, we obtain the following ‘‘splitting’’ formula:

$$\ell(a; L_0, L_1; F \oplus F') = \ell(a; L, L; F) + \ell([\text{pt}]; L'_0, L'_1; F'),$$

where $\ell(a; L, L; F)$ denotes the standard Lagrangian spectral invariant associated to $a \in HM(L)$ and $\ell([\text{pt}]; L'_0, L'_1; F')$ denotes the spectral invariant associated to the only non-zero class in $HF(L'_0, L'_1)$. See Sections 2.2 and 2.2.3 for the definitions of $\ell([\text{pt}]; L'_0, L'_1; F')$ and $\ell(a; L, L; F)$, respectively. Section 2.1.2 provides further details on $HF(L'_0, L'_1)$.

4. Triangle Inequalities: Given two Hamiltonians H, H' we denote by $H \# H'$ the Hamiltonian whose flow is the concatenation of the flows of H and H' ; see Equation (18) for a precise definition of $H \# H'$. Consider two Morse homology classes $a, b \in HM(L)$ such that the intersection product $a \cdot b \neq 0$. Lastly, for $i = 0, 1$, denote by $[L'_i]$ the fundamental class in $HM(L'_i)$. Then, the following triangle inequalities hold:

$$\begin{aligned} \ell(a \cdot b; L_0, L_1; H \# H') &\leq \ell(a; L_0, L_1; H) + \ell(b \otimes [L'_1]; L_1, L_1; H'), \\ \ell(a \cdot b; L_0, L_1; H \# H') &\leq \ell(a \otimes [L'_0]; L_0, L_0; H) + \ell(b; L_0, L_1; H'). \end{aligned}$$

The first three of the above properties are more or less standard, and in fact, we prove these in a more general setting in Section 2.1. The fourth property, which is perhaps the most interesting one, is specific to our settings and is different than the triangle inequality which appears in the standard setting where only one Lagrangian is considered.

Proofs of triangle inequalities of this nature consist of two main steps. First, one must prove a purely Floer theoretic version of the triangle inequality where Morse homology classes and the Morse intersection product are replaced with their Floer theoretic analogues. We do this, in a more general setting than what is described here in the introduction, in Section 2.5; see Theorem 17. The second step involves establishing a correspondence between the Morse and Floer theoretic versions of the intersection product, the latter being usually referred to as the pair-of-pants product. It is well-known that when L_0 and L_1 coincide (and some technical assumptions are satisfied) the two versions of the intersection product coincide up to a PSS-type isomorphism; see Equation (2.5.2). In our case, however, such a direct correspondence does not exist; the pair-of-pants product is not even defined on the tensor product of a single ring! In Theorem 21 and Remark 22, we fully describe the relation between the intersection product on $HM(L)$ and the pair-of-pants products $* : HF(L_0, L_1) \otimes HF(L_1, L_1) \rightarrow HF(L_0, L_1)$ and $* : HF(L_0, L_0) \otimes HF(L_0, L_1) \rightarrow HF(L_0, L_1)$.

Comparing the two forms of spectral invariants. – Using the aforementioned properties of the spectral invariants $\ell(a; L_0, L_1; H)$, one can deduce several other interesting properties of these invariants. Here, we will mention a comparison result which plays a significant role in our proof of Theorem 4.

Denote by $\ell([L_i]; L_i, L_i; H)$ the standard Lagrangian spectral invariant associated to the fundamental class $[L_i] \in HM(L_i)$; see Section 2.2.3 for the definition. The triangle inequality allows us to compare the two forms of spectral invariants. More precisely, we prove the following in Section 3.2.1.

PROPOSITION 6. – *For $i = 0, 1$, denote by $[L_i]$ the fundamental classes in $HM(L_i)$. Then, for any non-zero $a \in HM(L)$ and any Hamiltonian H :*

$$\ell(a; L_0, L_1; H) \leq \ell([L_i]; L_i, L_i; H).$$

In particular, $\ell([L]; L_0, L_1; H) \leq \ell([L_i]; L_i, L_i; H)$, where $[L] \in HM(L)$ is the fundamental class.

REMARK 7. – In defining the above spectral invariants $\ell(a; L_0, L_1; H)$, we were inspired by the construction of “conormal spectral invariants” defined in a cotangent bundle T^*N via consideration of the Lagrangian Floer homology of the zero section 0_N and the conormal ν^*V of a submanifold $V \subset N$ (see e.g., [15]). Indeed, if we heuristically think of the torus $\mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2} \times \mathbb{T}^{2k_2}$ as a compact version of the cotangent bundle to $\mathbb{T}^{k_1} \times \mathbb{T}^{k_1} \times \mathbb{T}^{k_2} \times \mathbb{T}^{k_2}$, then the Lagrangian L_0 corresponds to the zero section and L_1 corresponds to the conormal bundle of the submanifold $V = \mathbb{T}^{k_1} \times \mathbb{T}^{k_1} \times \mathbb{T}^{k_2} \times \{0\}$.

Of the above four properties of the spectral invariants $\ell(a; L_0, L_1; H)$, the first three also hold for conormal spectral invariants. We believe that, by readjusting the techniques used in this paper, one could obtain an appropriately reformulated version of the triangle inequality for conormal spectral invariants. This would then lead to the following comparison inequalities, corresponding to Proposition 6: For every homology class $a \in HM(V)$, and every Hamiltonian H on T^*N ,

$$\ell(a; 0_N, \nu^*V; H) \leq \ell([N]; 0_N, 0_N; H).$$

As far as we know, the triangle inequality has not yet been proven for conormal spectral invariants. However, the above comparison inequalities were proven, via generating-function techniques, in [22].

The idea that conormal spectral invariants could be useful in studying the behavior of spectral invariants under symplectic reduction has been present in works based on generating function theory (e.g., [21, 8], [19]) and goes back to Viterbo [22]. To the best of our knowledge, this article is the first place where this idea is implemented in Floer theory. We found this implementation to be necessary for our purposes as Floer theory is better suited for working on compact manifolds.

Organization of the paper

In Sections 2.1–2.4 we recall Floer theoretic preliminaries, define Lagrangian and Hamiltonian spectral invariants, and prove some of their essential properties. In Section 2.5, we define the pair-of-pants product and prove a purely Floer theoretic version of the triangle inequality in a fairly general setting. In Section 2.6, we prove a Künneth formula for Lagrangian Floer homology and use it to derive a splitting formula for spectral invariants. In Section 3, we specialize the Floer theory of Section 2 to the specific settings introduced above. In Section 3.2, we prove the aforementioned triangle inequality. Lastly, in Section 4, we use the results from Sections 2 and 3 to prove Theorem 4.

Acknowledgements. – We are grateful to Claude Viterbo for several helpful conversations.

This work is partially supported by ANR Grants ANR-11-JS01-010-01 and ANR-13-JS01-0008-01. The research leading to these results has received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement 307062.

2. Floer homology and spectral invariants

2.1. Lagrangian Floer homology

In this section, we review the construction of Floer homology. Throughout the section, we fix a closed symplectic manifold (M, ω) , two closed non-disjoint connected Lagrangian submanifolds L_0, L_1 and $p \in L_0 \cap L_1$ an intersection point. Recall that

- (M, ω) is symplectically aspherical if $\omega|_{\pi_2(M)} = 0$,
- a Lagrangian L of (M, ω) is weakly exact, or the pair (M, L) is relatively symplectically aspherical, if $\omega|_{\pi_2(M, L)} = 0$.

We say that the pair (L_0, L_1) is *weakly exact with respect to p* , if any disk in M whose boundary is on $L_0 \cup L_1$ and is “pinched” at p has vanishing symplectic area. More precisely, denote by D the unit disk in \mathbb{C} centered at 0. Denote by ∂D^+ the upper half of ∂D , $\partial D^+ = \{z \in \mathbb{C} : |z| = 1, \text{Im}(z) \geq 0\}$, and by ∂D^- its lower half.

DEFINITION 8. – *The pair of Lagrangians (L_0, L_1) is weakly exact with respect to $p \in L_0 \cap L_1$ if for any map $u : (D, \partial D^+, \partial D^-, \{-1, 1\}) \rightarrow (M, L_0, L_1, \{p\})$, $\int_D u^* \omega = 0$.*

Notice that in this case both L_0 and L_1 are weakly exact and thus M is symplectically aspherical.

EXAMPLE 9. – The Lagrangians we will consider in Sections 3 and 4 form weakly exact pairs with respect to any point in their intersections. Recall from Section 1.2 in the introduction that for $i = 1$ and 2 , $L_i = L \times L'_i$ are Lagrangians of $(\mathbb{T}^k \times \mathbb{T}^l, \omega_{\mathbb{T}^k} \oplus \omega_{\mathbb{T}^l})$ so that $\mathbb{T}^l = L'_0 \times L'_1$ and $L'_0 \cap L'_1 = \{0\}$. (In this example only, k and l denote the respective integers $4k_1 + 2k_2$ and $2k_2$ to ease the reading.) We fix a point $p = (p_k, 0)$ in $L_0 \cap L_1$.

First notice that since L is a subtorus of \mathbb{T}^k , $\pi_2(\mathbb{T}^k, L) = 0$ so that (L, L) is a weakly exact pair with respect to p_k .

Next, consider a pinched disk

$$u : (D, \partial D^+, \partial D^-, \{-1, 1\}) \rightarrow (\mathbb{T}^l, L'_0 \times \{0\}, \{0\} \times L'_1, \{0\}).$$

Denote by γ_0 and γ_1 the loops respectively in L'_0 and L'_1 , defined by $u(\partial D^-) = \gamma_0 \times \{0\}$ and $u(\partial D^+) = \{0\} \times \gamma_1$. Since $[u(\partial D^-)] = [u(\partial D^+)] \in \pi_1(\mathbb{T}^l) = \pi_1(L'_0) \times \pi_1(L'_1)$, γ_0 and γ_1 are null-homotopic in L'_0 and L'_1 respectively. By gluing to u two disks $v_i \subset L'_i$ bounding γ_i , we obtain a sphere in \mathbb{T}^l whose symplectic area necessarily vanishes. Since the v_i 's are included in Lagrangians we deduce that $\omega_{\mathbb{T}^l}(u) = 0$. Therefore (L'_0, L'_1) is weakly exact with respect to the single intersection point 0.

Finally, since the product of weakly exact pairs is weakly exact, we deduce that (L_0, L_1) is weakly exact with respect to p . ◀

Let $H : [0, 1] \times M \rightarrow \mathbb{R}$ be a smooth Hamiltonian function. We will denote by X_H the 1-parameter family of vector fields induced by H by $\omega(X_H^t, \cdot) = -dH_t$ for all t and by ϕ_H^t its flow satisfying: $\phi_H^0 = \text{Id}$ and for all t , $\partial_t \phi_H^t = X_H^t(\phi_H^t)$. We first consider a *non-degenerate* Hamiltonian, which means in this case that the intersection $\phi_H^1(L_0) \cap L_1$ is transverse. A generic Hamiltonian is non-degenerate.

We denote by $\Omega(L_0, L_1; p)$ the set of paths x from L_0 to L_1 which are in the connected component of the constant path p . Such a path admits a capping $\bar{x} : [0, 1] \times [0, 1] \rightarrow M$

so that: For all $t \in [0, 1]$, $\bar{x}(0, t) = p$ and $\bar{x}(1, t) = x(t)$, $[0, 1] \times \{0\}$ is mapped to L_0 and $[0, 1] \times \{1\}$ to L_1 .

Two cappings \bar{x}_1 and \bar{x}_2 of $x \in \Omega(L_0, L_1; p)$ have the same symplectic area since $\bar{x}_1 \# (-\bar{x}_2)$ is a pinched disk as defined above so that it has area 0 by assumption. (Recall that $-\bar{x}_2$ stands for \bar{x}_2 with reverse orientation.) Thus we can define the action functional by the formula:

$$(1) \quad \mathcal{Q}_H^{L_0, L_1} : \Omega(L_0, L_1; p) \rightarrow \mathbb{R}, \quad x \mapsto - \int \bar{x}^* \omega + \int_0^1 H_t(x(t)) dt .$$

The critical points of $\mathcal{Q}_H^{L_0, L_1}$ are paths $x \in \Omega(L_0, L_1; p)$ which are orbits of H that is for all t , $x(t) = \phi_H^t(x(0))$. These orbits are in one-to-one correspondence with $\phi_H^1(L_0) \cap L_1$ so that their number is finite since H is non-degenerate and M compact. One defines the Floer complex as the \mathbb{Z}_2 -vector space $CF(L_0, L_1; p; H) = \langle \text{Crit}(\mathcal{Q}_H^{L_0, L_1}) \rangle_{\mathbb{Z}_2}$. The set of critical values of $\mathcal{Q}_H^{L_0, L_1}$ is called its spectrum and is denoted by $\text{Spec}(H)$.

Now Floer’s differential is defined thanks to perturbed pseudo-holomorphic strips: we pick a 1-parameter family of tame, ω -compatible, almost complex structures J . We define the set of Floer trajectories between two orbits of H , x_- and x_+ , as

$$\widehat{\mathcal{M}}^{L_0, L_1}(x_-, x_+; H, J) = \left\{ u : \mathbb{R} \times [0, 1] \rightarrow M \left| \begin{array}{l} \partial_s u + J_t(u)(\partial_t u - X_H^t(u)) = 0 \\ \forall t, u(\pm\infty, t) = x_{\pm}(t) \\ u(\mathbb{R} \times \{0\}) \subset L_0 \\ u(\mathbb{R} \times \{1\}) \subset L_1 \end{array} \right. \right\}$$

where the limits $u(\pm\infty, t)$ are uniform in t . There is an obvious \mathbb{R} -action by reparametrization $s \mapsto s + \tau$ and we define $\mathcal{M}^{L_0, L_1}(x_-, x_+; H, J)$ as the quotient $\widehat{\mathcal{M}}^{L_0, L_1}(x_-, x_+; H, J)/\mathbb{R}$.

Requiring that the pair (H, J) is *regular*, that is the linearization of the operator $\bar{\partial}_{J, H} : u \mapsto \partial_s u + J_t(u)(\partial_t u - X_H(u))$ is surjective for all $u \in \widehat{\mathcal{M}}^{L_0, L_1}(x_-, x_+; H, J)$, ensures that $\mathcal{M}^{L_0, L_1}(x_-, x_+; H, J)$ is a smooth manifold. We denote its 0- and 1-dimensional components respectively by $\mathcal{M}_{[0]}^{L_0, L_1}(x_-, x_+; H, J)$ and $\mathcal{M}_{[1]}^{L_0, L_1}(x_-, x_+; H, J)$.

REMARK 10. – It turns out that we do not need to consider *graded* complexes. As a consequence, we do not mention the different standard indices usually entering into play in such theories. In particular, we do not require additional assumptions concerning these indices.

Nonetheless, let us recall that the $(i+1)$ -dimensional component of $\widehat{\mathcal{M}}^{L_0, L_1}(x_-, x_+; H, J)$ consists of those Floer trajectories whose Maslov-Viterbo index equals $i+1$, see e.g [3]. When there are no such trajectories, we put $\mathcal{M}_{[i]}^{L_0, L_1}(x_-, x_+; H, J)$ to be the empty set.

The 0-dimensional component of the moduli space of all Floer trajectories running between any two orbits, $\mathcal{M}_{[0]}^{L_0, L_1}(H, J) = \bigcup_{x_-, x_+} \mathcal{M}_{[0]}^{L_0, L_1}(x_-, x_+; H, J)$ is compact. Floer’s differential is defined by linearity on $CF(L_0, L_1; p; H)$ after setting the image of a generator as

$$\partial_{H, J}^{L_0, L_1}(x_-) = \sum_{x_+} \# \mathcal{M}_{[0]}^{L_0, L_1}(x_-, x_+; H, J) \cdot x_+$$

where $\#\mathcal{M}$ is the mod 2 cardinal of \mathcal{M} and the sum runs over all orbits x_+ . Since the asphericity assumption prevents bubbling of disks and spheres, by Gromov's compactness Theorem and standard gluing $\mathcal{M}_{[1]}^{L_0, L_1}(H, J)$, the 1-dimensional component of $\mathcal{M}^{L_0, L_1}(H, J)$, can be compactified, and this fact ensures that $(\partial_{H, J}^{L_0, L_1})^2 = 0$ that is, $\partial_{H, J}^{L_0, L_1}$ is a differential.

The Floer homology of the pair (L_0, L_1) is the homology of this complex $HF(L_0, L_1; p; H, J) = H(CF(L_0, L_1; p; H), \partial_{H, J}^{L_0, L_1})$. Because it is often useful to keep in mind the specific Floer data which we used to define the complex, we will keep (H, J) in the notation, however the homology does not depend on the choice of the regular pair (H, J) .⁽⁴⁾ Indeed, there are morphisms

$$\Psi_{H, J}^{H', J'} : CF(L_0, L_1; p; H) \rightarrow CF(L_0, L_1; p; H')$$

inducing isomorphisms in homology which are called continuation isomorphisms. Roughly, $\Psi_{H, J}^{H', J'}$ is defined thanks to a regular homotopy between (H, J) and (H', J') , (\tilde{H}, \tilde{J}) , by considering the 0-dimensional component of the moduli space of Floer trajectories for the pair (\tilde{H}, \tilde{J}) running from an orbit of H to an orbit of H' with boundary condition on L_0 and L_1 respectively. It is also standard—and the proof is based on the same principle by considering a homotopy between homotopies—that $\Psi_{H, J}^{H', J'}$ does not depend on the choice of the homotopy (\tilde{H}, \tilde{J}) . From these facts, one gets that they are “canonical,” that is they satisfy

$$(2) \quad \Psi_{H, J}^{H, J} = \text{Id} \quad \text{and} \quad \Psi_{H, J}^{H', J'} \circ \Psi_{H', J'}^{H'', J''} = \Psi_{H, J}^{H'', J''}$$

for any three regular pairs (H, J) , (H', J') , and (H'', J'') .

We now present two situations which will be of particular interest to us and in which one can actually compute Floer homology.

2.1.1. *The case of a single Lagrangian.* – Assume that L_0 and L_1 coincide and denote $L = L_0 = L_1$. Assume moreover that L is connected. In that case, the assumption that the pair (L, L) is weakly exact with respect to any given point $p \in L$ is equivalent to requiring the Lagrangian L to be weakly exact.

It is well-known that there exists an isomorphism between the Floer homology of (L, L) and the Morse homology of L , called PSS isomorphism. It was defined in the Hamiltonian setting by Piunikhin-Salamon-Schwarz [18], then adapted to Lagrangian Floer homology by Katić-Milinković [10] for cotangent bundles, and by Barraud-Cornea [4] and Albers [2] for compact manifolds. For details, we refer to Leclercq [12] which deals with weakly exact Lagrangians in compact manifolds which is the situation we are interested in here.

The PSS morphism requires an additional choice of a Morse-Smale pair (f, g) , consisting of a Morse function $f: L \rightarrow \mathbb{R}$ and a metric g on L . It is defined at the chain level $\Phi_{H, J}^L: CM(L; f, g) \rightarrow CF(L, L; p; H, J)$ by counting the number of elements of suitable moduli spaces. It commutes with the differential and thus induces a morphism in homology:

$$\Phi_{H, J}^L: HM(L) \rightarrow HF(L, L; p; H, J)$$

⁽⁴⁾ This being said, when there is no risk of confusion we will denote $HF(L_0, L_1; p; H, J)$ by $HF(L_0, L_1; p)$ to simplify the notation.

(as the notation suggests, we will omit the Morse data). It is an isomorphism and its inverse $\Phi_L^{H,J} : HF(L, L; p; H, J) \rightarrow HM(L)$ is defined in the same fashion. The main properties of the PSS morphism which will be needed are the following two:

1. PSS morphism commutes with continuation morphisms, that is

$$(3) \quad \begin{array}{ccc} HM(L) & \xrightarrow{\Phi_{H,J}^L} & HF(L, L; p; H, J) \\ & \searrow \Phi_{H',J'}^L & \downarrow \Psi_{H,J}^{H',J'} \\ & & HF(L, L; p; H', J') \end{array}$$

commutes for any two regular pairs (H, J) and (H', J') .

2. PSS morphism intertwines the Morse and Floer theoretic versions of the intersection product in homology, the latter being known as pair-of-pants product, see Subsection 2.5.2 for the precise statement.

2.1.2. *The case of two Lagrangians intersecting transversely at a single point.* – When L_0 and L_1 intersect transversely at a single point p , the Hamiltonian $H = 0$ is non-degenerate. The associated Floer complex obviously has a single generator, p itself. Moreover, for any choice of almost complex structure J such that $(0, J)$ is regular, the boundary map is trivial since the 0-dimensional component of the moduli space of Floer trajectories from p to itself is empty. It follows that $HF(L_0, L_1; p; 0, J)$, and hence $HF(L_0, L_1; p; H, J)$ for any regular (H, J) , is isomorphic to the group with two elements. We will refer to this isomorphism, which is uniquely defined, as a PSS-type morphism and will denote it by

$$\Phi_{H,J}^{L_0,L_1} : \mathbb{Z}_2 \rightarrow HF(L_0, L_1; p; H, J).$$

The only non-zero class in $HF(L_0, L_1; p; H, J)$ will be denoted by [pt].

Since there exists only one isomorphism between two given groups with two elements, the following diagram commutes for any two regular pairs (H, J) and (H', J') :

$$(4) \quad \begin{array}{ccc} \mathbb{Z}_2 & \xrightarrow{\Phi_{H,J}^{L_0,L_1}} & HF(L_0, L_1; p; H, J) \\ & \searrow \Phi_{H',J'}^{L_0,L_1} & \downarrow \Psi_{H,J}^{H',J'} \\ & & HF(L_0, L_1; p; H', J'). \end{array}$$

2.2. Lagrangian spectral invariants

Spectral invariants for Lagrangians in cotangent bundles were introduced by Viterbo [22] using generating functions. This was adapted to Floer homology by Oh [15]. Since then there have been several extensions to other settings. See in particular Leclercq [12] for a single weakly-exact Lagrangian and Zapolsky [23] for a weakly exact pair of Lagrangians intersecting at a single point.

We provide below a new extension of the definition for a general weakly-exact pair (L_0, L_1) with a given intersection point p .

To give this definition, the starting observation is the standard fact that for every Floer trajectory $u \in \mathcal{M}^{L_0, L_1}(x_-, x_+; H, J)$,

$$\mathcal{A}_H^{L_0, L_1}(x_-) - \mathcal{A}_H^{L_0, L_1}(x_+) = \int_{\mathbb{R} \times [0, 1]} \|\partial_s u\|^2 ds dt \geq 0,$$

where $\|\cdot\|$ is the norm associated to the metric $\omega(\cdot, J\cdot)$. Thus the action decreases along Floer trajectories. Now let H be a non-degenerate Hamiltonian and let $a \in \mathbb{R}$ be a regular value of the action functional, i.e., $a \notin \text{Spec}(H)$. It follows from this observation that if $CF^a(L_0, L_1; p; H)$ denotes the \mathbb{Z}_2 -vector space generated by Hamiltonian chords of action $< a$, then $CF^a(L_0, L_1; p; H)$ is a subcomplex of $CF(L_0, L_1; p; H)$. We denote $i^a : HF^a(L_0, L_1; p; H, J) \rightarrow HF(L_0, L_1; p; H, J)$ the map induced in homology by the inclusion. For every non-zero Floer homology class $\alpha \in HF(L_0, L_1; p; H, J)$, we define the spectral invariant associated to α to be the number

$$(5) \quad \ell(\alpha; L_0, L_1; p; H) = \inf\{a \in \mathbb{R} : \alpha \in \text{im}(i^a)\}.$$

We wish to have the ability to compare the spectral invariants of different Hamiltonians. For this purpose it will be convenient to fix a reference Floer homology group: pick a regular pair $(H_{\text{ref}}, J_{\text{ref}})$ and set

$$HF_{\text{ref}}(L_0, L_1; p) = HF(L_0, L_1; p; H_{\text{ref}}, J_{\text{ref}}).$$

DEFINITION 11. – *For every non-degenerate Hamiltonian function H , the spectral invariant associated to $\alpha \in HF_{\text{ref}}(L_0, L_1; p)$, $\alpha \neq 0$, is the number*

$$\ell(\alpha; L_0, L_1; p; H) := \ell(\Psi_{H_{\text{ref}}, J_{\text{ref}}}^{H, J}(\alpha); L_0, L_1; p; H).$$

As the notation suggests, the number $\ell(\alpha; L_0, L_1; p; H)$, both in the above definition and in Equation (5), does not depend on the necessary choice of an almost complex structure J so that the pair (H, J) is regular. This follows from the following inequality which holds for every two regular pairs (H, J) , (H', J') :

$$(6) \quad \begin{aligned} \ell(\Psi_{H_{\text{ref}}, J_{\text{ref}}}^{H', J'}(\alpha); L_0, L_1; p; H') &\leq \ell(\Psi_{H_{\text{ref}}, J_{\text{ref}}}^{H, J}(\alpha); L_0, L_1; p; H) \\ &\quad + \int_0^1 \max_M (H'_t - H_t) dt. \end{aligned}$$

We now sketch a proof of the above inequality. Since the continuation morphism is injective, the image of any non-zero class $\alpha \in HF(L_0, L_1; p; H, J)$ is non-zero. By definition of $\Psi_{H, J}^{H', J'}$ there exist Floer trajectories for a homotopy (\tilde{H}, \tilde{J}) between the generators of $CF(L_0, L_1; p; H)$ whose linear combination represents α and the generators of $CF(L_0, L_1; p; H')$ representing $\Psi_{H, J}^{H', J'}(\alpha)$. Computing the energy of such a trajectory and using the fact that the result is positive yields:

$$\ell(\Psi_{H, J}^{H', J'}(\alpha); L_0, L_1; p; H') \leq \ell(\alpha; L_0, L_1; p; H) + \int_0^1 \max_M (H'_t - H_t) dt.$$

Thus, in particular Inequality (6) follows.

Furthermore, Inequality (6) implies that for every non-degenerate H, H' ,

$$(7) \quad \int_0^1 \min_M(H_t - H'_t) dt \leq |\ell(\alpha; L_0, L_1; p; H) - \ell(\alpha; L_0, L_1; p; H')| \leq \int_0^1 \max_M(H_t - H'_t) dt.$$

As a consequence, the number $\ell(\alpha; L_0, L_1; p; H)$ is Lipschitz continuous with respect to the Hamiltonian H . Moreover, it follows that $\ell(\alpha; L_0, L_1; p; H)$ can be defined by continuity for every continuous function $H : [0, 1] \times M \rightarrow \mathbb{R}$.

2.2.1. *Spectrality.* – It is rather easy in our situation (where one does not need to keep track of cappings) to prove the *spectrality property* of the invariants $\ell(\alpha; L_0, L_1; p; H)$ regardless the non-degeneracy of H . Namely, for all non-zero $\alpha \in HF_{\text{ref}}(L_0, L_1; p)$,

$$\ell(\alpha; L_0, L_1; p; H) \in \text{Spec}(H).$$

We will need the following consequence of this property (we refer to [14, Lemma 2.2] for a proof).

COROLLARY 12. – *Let L be a weakly exact closed Lagrangian of (M, ω) and $H : [0, 1] \times M \rightarrow \mathbb{R}$ be continuous. If $H|_L = c$ for some $c \in \mathbb{R}$, then $\ell(\alpha; L, L; p; H) = c$ for all $\alpha \neq 0$ in $HF(L, L; p)$.*

We end this subsection by recalling that $\text{Spec}(H)$, for any Hamiltonian H , is a measure zero subset of \mathbb{R} . This fact will be used in Section 4.

2.2.2. *Naturality.* – Lagrangian Floer homology is *natural* in the sense that for any symplectomorphism $\psi : (M, \omega) \rightarrow (M', \omega')$ and any two Lagrangians L_0 and L_1 of (M, ω) , the following Floer homologies are isomorphic

$$(8) \quad HF(L_0, L_1; p; H, J) \simeq HF(\psi(L_0), \psi(L_1); \psi(p); H \circ \psi^{-1}, (\psi^{-1})^* J)$$

since the respective complexes as well as the respective moduli spaces involved in the definition of the differential are in one-to-one correspondence. This one-to-one correspondence is given on the generators of the complex by the obvious identification

$$x \in \text{Crit}(\mathcal{A}_H^{L_0, L_1}) \Leftrightarrow \psi(x) \in \text{Crit}(\mathcal{A}_{H \circ \psi^{-1}}^{\psi(L_0), \psi(L_1)})$$

where $\psi(x)$ denotes the orbit of $H \circ \psi^{-1}$ given as $t \mapsto \psi(x(t))$. Furthermore, the above bijection preserves the action, namely

$$\forall x \in \text{Crit}(\mathcal{A}_H^{L_0, L_1}), \quad \mathcal{A}_H^{L_0, L_1}(x) = \mathcal{A}_{H \circ \psi^{-1}}^{\psi(L_0), \psi(L_1)}(\psi(x)).$$

From this, it is easy to see that the respective Lagrangian spectral invariants coincide: For any non-zero Floer homology class α in $HF(L_0, L_1; p; H, J)$ and its image via (8), α_ψ in $HF(\psi(L_0), \psi(L_1); \psi(p); H \circ \psi^{-1}, (\psi^{-1})^* J)$, we have

$$(9) \quad \ell(\alpha; L_0, L_1; p; H) = \ell(\alpha_\psi; \psi(L_0), \psi(L_1); \psi(p); H \circ \psi^{-1}).$$

2.2.3. *The case of a single Lagrangian.* – In the particular situation of Section 2.1.1 where we consider a single Lagrangian $L (= L_0 = L_1)$, one can easily associate spectral invariants not only to Floer homology classes of (L, L) but also to (Morse) homology classes of L via the PSS isomorphism. For convenience, we denote these invariants in the same way: To any $a \neq 0$ in $HM(L)$, we associate

$$(10) \quad \ell(a; L, L; H) = \ell(\Phi_{H,J}^L(a); L, L; p; H)$$

with p any point in L and the right-hand side defined by (5).

As in the general case, this quantity requires the additional choice of an almost complex structure J such that (H, J) is regular, it is Lipschitz continuous with respect to H , so that it is independent of the choice of J and its definition naturally extends to any continuous $H: [0, 1] \times M \rightarrow \mathbb{R}$.

The naturality (9) of spectral invariants also holds in this case. More precisely, it reads

$$(11) \quad \ell(a; L, L; p; H) = \ell(\psi_*(a); \psi(L), \psi(L); \psi(p); H \circ \psi^{-1})$$

for all non-zero homology classes a of L and all symplectomorphisms ψ . Here $\psi_*(a)$ denotes the image of a by the morphism induced by $\psi|_L$ on $HM(L)$. Indeed, to see that this holds one should pick a Morse-Smale pair (f, g) on L , and use $(f \circ \psi^{-1}, (\psi^{-1})^*g)$ as Morse-Smale pair on $\psi(L)$. For these choices, the following diagram commutes:

$$\begin{array}{ccc} HM(L; f, g) & \xrightarrow{\psi_* = (\psi|_L)_*} & HM(\psi(L); f \circ \psi^{-1}, (\psi^{-1})^*g) \\ \Phi_{H,J}^L \downarrow & & \downarrow \Phi_{H \circ \psi^{-1}, (\psi^{-1})^*J}^{\psi(L)} \\ HF(L, L; p; H, J) & \xrightarrow{\psi_*} & HF(\psi(L), \psi(L); \psi(p); H \circ \psi^{-1}, (\psi^{-1})^*J) \end{array}$$

even at the chain level (this is a mild generalization of [7, Lemma 5.1] where ψ was assumed to be a Hamiltonian diffeomorphism preserving L). The fact that spectral invariants do not depend on the Morse data then leads to (11).

Finally, in the case of a single Lagrangian one spectral invariant will be of particular interest to us, namely the one associated to $[L]$, the fundamental class of L : $\ell([L]; L, L; H)$.

2.3. Hamiltonian Floer theory, spectral invariants, and capacity

2.3.1. *Hamiltonian Floer homology.* – We work in a symplectically aspherical manifold (M, ω) . Formally, this case is very similar to the Lagrangian case of Section 2.1.1.

Namely, we pick a Hamiltonian H which is non-degenerate in the sense that the graph of $\phi_H^1, \Gamma_{\phi_H^1}$, intersects transversely the diagonal $\Delta \subset M \times M$.

Instead of $\Omega(L, L; p)$, we consider the set of contractible free loops in M . We denote this set by $\Omega(M)$ and a typical element by γ . The action functional $\mathcal{A}_H: \Omega(M) \rightarrow \mathbb{R}$ is defined by the same formula as (1) except that for $\gamma \in \Omega(M)$, $\bar{\gamma}$ denotes a capping of γ , that is a disk in M whose boundary is mapped to the image of γ . Again, the asphericity condition ensures that \mathcal{A}_H is well-defined.

Its critical points are the contractible 1-periodic orbits of H which form a finite set by genericity of H and generate a \mathbb{Z}_2 -vector space which we denote $CF(M; H)$.

We again pick a 1-parameter family of tame, ω -compatible, almost complex structures J and consider the moduli spaces:

$$\widehat{\mathcal{M}}(\gamma_-, \gamma_+; H, J) = \left\{ u: \mathbb{R} \times S^1 \rightarrow M \left| \begin{array}{l} \partial_s u + J_t(u)(\partial_t u - X_H^t(u)) = 0 \\ \forall t, u(\pm\infty, t) = \gamma_{\pm}(t) \end{array} \right. \right\}$$

and their quotient by the obvious \mathbb{R} -reparametrization in s which we denote $\mathcal{M}(\gamma_-, \gamma_+; H, J)$. These moduli spaces share the same properties as their Lagrangian counterpart and the differential is defined accordingly:

$$\partial_{H,J}(\gamma_-) = \sum_{\gamma_+} \# \mathcal{M}_{[0]}(\gamma_-, \gamma_+; H, J) \cdot \gamma_+$$

on generators and extended by linearity. Again, the sum runs over all contractible 1-periodic orbits of H and $\mathcal{M}_{[0]}$ is the 0-dimensional component of the moduli space \mathcal{M} .

The Floer homology of (M, ω) is defined as the homology of this complex $HF(M) = H(CF(M; H), \partial_{H,J})$ and does not depend on the choice of the regular pair (H, J) in the sense that there are continuation isomorphisms defined in the exact same fashion as in the Lagrangian case and built via regular homotopies of the data. When H is C^2 -small enough, the Floer complex coincides with the Morse complex of M .

Finally, there is also a PSS morphism constructed from a regular pair (H, J) and a Morse-Smale pair (f, g) on M similarly to its Lagrangian counterpart. As for the latter, we will omit the Morse data and denote it $\Phi_{H,J}: HM(M) \rightarrow HF(M; H, J)$.

2.3.2. Hamiltonian spectral invariants. – This case corresponds to the one studied by Schwarz in [20]. As in the Lagrangian case described above, the Floer complex is naturally filtered by action values since the action functional decreases along Floer trajectories. So any regular value of the Hamiltonian action \mathcal{U}_H gives rise to a subcomplex $CF^a(M; H) \xrightarrow{i^a} CF(M; H)$ and the Hamiltonian spectral invariants are defined for any non-zero Floer homology class of M as in (5). Thanks to the PSS isomorphism, one can also associate spectral invariants to any non-zero (Morse) homology class of $\alpha \in HM(M)$ as in Section 2.2.3. We will temporarily use the notation $c(\alpha; H, J)$ to denote these invariants.

These invariants share similar properties with their Lagrangian counterparts. In particular they satisfy a Lipschitz estimate similar to (7). It follows that they are independent of the choice of almost complex structure and hence we will denote them $c(\alpha; H)$. Furthermore, being Lipschitz continuous, $c(\alpha; H)$ extends to continuous functions on $[0, 1] \times M$, i.e., for a continuous $H \in C^0([0, 1] \times M)$ we can define $c(\alpha; H) = \lim_{i \rightarrow \infty} c(\alpha; H_i)$ where H_i is any sequence of smooth non-degenerate Hamiltonians converging to H .

As in 2.2.3, one of these invariants will be of greatest interest to us, $c_+(H) = c([M]; H)$, the Hamiltonian spectral invariant associated to the fundamental class of M . It follows from the above discussion that for non-degenerate H , it is defined via the following expression:

$$(12) \quad c_+(H) = \inf \{ a \in \mathbb{R} : \text{PSS}([M]) \in \text{im}(HF^a(M; H) \xrightarrow{i_*^a} HF(M; H)) \}.$$

2.3.3. *The spectral capacity c .* – Following Viterbo [22], we extract from c_+ the spectral capacity c mentioned in the introduction. Namely, for any open set U in M , we define

$$c(U) = \sup\{c_+(H) : H \in C^0([0, 1] \times M), \text{supp}(H_t) \subset U \forall t \in [0, 1]\}.$$

This quantity satisfies the properties defining a capacity, see [22].

2.4. Comparison between Lagrangian and Hamiltonian spectral invariants

There is an action-preserving isomorphism between the Hamiltonian Floer complex of (M, ω) associated to a regular pair (H, J) and the Lagrangian Floer complex of the diagonal $\Delta \simeq M$ seen as a Lagrangian in $(M \times M, (-\omega) \oplus \omega)$ and associated to an appropriate regular pair (\hat{H}, \hat{J}) . The goal of this section is to prove that the respective spectral invariants coincide (see also [12, Section 3.4]). Given Hamiltonians H and G on M , we will denote by $H \oplus G$ the Hamiltonian given for every $(x, y) \in M \times M$ by $H \oplus G(x, y) = H(x) + G(y)$.

PROPOSITION 13. – *Let (M, ω) be a symplectically aspherical manifold. Let $\alpha \neq 0$ in $HM(M)$ and denote $\hat{\alpha}$ the corresponding class in $HM(\Delta)$. For any continuous time-dependent Hamiltonian H on M , $c(\alpha; H) = \ell(\hat{\alpha}; \Delta, \Delta; 0 \oplus H)$. In particular, $c_+(H) = \ell([\Delta]; \Delta, \Delta; 0 \oplus H)$.*

Notice that we are in the case of a single Lagrangian submanifold Δ , so that $\ell(\hat{\alpha}; \Delta, \Delta; 0 \oplus H)$ refers to this particular setting, see Section 2.2.3.

At several points in this paper, and to begin with in the proof of the proposition above, we will need to work with Hamiltonians H such that $H_t = 0$ for t near 0 and 1. This can always be achieved, without affecting the spectral invariants of H , by time reparametrization. This is the content of the following remark.

REMARK 14. – Let (H, J) be a regular pair. Pick a smooth increasing function $\sigma: [0, 1] \rightarrow \mathbb{R}$ so that $\sigma(t) = 0$ for all $t \in [0, \varepsilon]$ and $\sigma(t) = 1$ for all $t \in [1 - \varepsilon', 1]$ for some $\varepsilon, \varepsilon'$ so that $0 < \varepsilon < 1 - \varepsilon' < 1$. Then define $H_t^\sigma(x) = \sigma'(t)H_{\sigma(t)}(x)$. There is an obvious bijection between the sets of orbits of H and H^σ which leads to a bijection on the Floer complexes as vector spaces:

$$(13) \quad CF(M; H) \rightarrow CF(M; H^\sigma), \quad \gamma \mapsto [\gamma^\sigma : t \mapsto \gamma(\sigma(t))]$$

which preserves the action, namely $\mathcal{A}_{H^\sigma}(\gamma^\sigma) = \mathcal{A}_H(\gamma)$ (since geometrically the orbits are the same, a capping of γ also caps γ^σ).

Then define J^σ as $J_t^\sigma(x) = J_{\sigma(t)}(x)$. Notice that (H^σ, J^σ) is regular, and that there is a bijection between the moduli spaces $\mathcal{M}(\gamma_-, \gamma_+; H, J)$ and $\mathcal{M}(\gamma_-^\sigma, \gamma_+^\sigma; H^\sigma, J^\sigma)$ so that (13) induces an action-preserving isomorphism of the differential complexes. Notice that geometrically the main objects (orbits and Floer's strips) remain the same. It is thus easy to see that geometrically the representatives of a given Floer homology class remain unchanged along the process so that, together with the fact that the action is preserved, the associated (Hamiltonian) spectral invariants coincide.

For the same reason, given a Lagrangian L , the Lagrangian spectral invariants associated to H also remain unchanged along such reparametrization.

We now prove Proposition 13.

Proof. – First notice that if (M, ω) is symplectically aspherical, then the diagonal Δ is a weakly exact Lagrangian of $(M \times M, (-\omega) \oplus \omega)$.

We first prove the proposition for non-degenerate Hamiltonians. So we start with a regular pair (H, J) and apply Remark 14 with $\sigma: [0, 1] \rightarrow \mathbb{R}$ so that $\sigma(t) = 0$ for all $t \in [0, 1/2]$. Then $H^\sigma = 0$ and $J_t^\sigma = J_0$ for all $t \in [0, 1/2]$.

Now we consider for $t \in [0, \frac{1}{2}]$ the Hamiltonian $\hat{H}_t = H_{\frac{1}{2}-t}^\sigma \oplus H_{\frac{1}{2}+t}^\sigma$ on $M \times M$. There is a bijection:

$$(14) \quad \begin{aligned} &CF(M; H^\sigma) \rightarrow CF(\Delta, \Delta; \hat{H}), \\ &[\gamma: \mathbb{S}^1 \rightarrow M] \mapsto [x: [0, \frac{1}{2}] \rightarrow M \times M] \text{ with } x(t) = (\gamma(\frac{1}{2} - t), \gamma(\frac{1}{2} + t)) \end{aligned}$$

since x is an orbit of \hat{H} if and only if γ is an orbit of H^σ . Notice that by definition of σ , $\hat{H}_t = 0 \oplus H_{\frac{1}{2}+t}^\sigma$ so that by Remark 14 above

$$(15) \quad \ell([\Delta]; \Delta, \Delta; \hat{H}) = \ell([\Delta]; \Delta, \Delta; 0 \oplus H)$$

and $x(t) = (\gamma(0), \gamma(\frac{1}{2} + t))$.

Again, we need an appropriate family of almost complex structures \hat{J} on $M \times M$ which we obtained by putting $\hat{J}_t(x, y) = -J_{\frac{1}{2}-t}^\sigma(x) \times J_{\frac{1}{2}+t}^\sigma(y)$ for $t \in [0, \frac{1}{2}]$. It is easy to see that (H^σ, J^σ) is regular and that the bijection (14) above is compatible with the differentials of the complexes. Indeed, pick any two generators of $CF(M; H^\sigma)$, γ_- and γ_+ and any cylinder $u: \mathbb{R} \times \mathbb{S}^1$ which uniformly converges to γ_\pm when s converges to $\pm\infty$. Denote respectively by x_\pm the generators of $CF(\Delta, \Delta; \hat{H})$ given by (14) from γ_\pm and consider

$$\hat{u}: \mathbb{R} \times [0, \frac{1}{2}] \rightarrow M \times M, \quad \hat{u}(s, t) = (u(s, \frac{1}{2} - t), u(s, \frac{1}{2} + t)).$$

When s goes to $\pm\infty$, \hat{u} uniformly converges to $(\gamma_\pm(\frac{1}{2} - t), \gamma_\pm(\frac{1}{2} + t)) = x_\pm(t)$. The boundary conditions are: $\hat{u}(s, 0) = (u(s, \frac{1}{2}), u(s, \frac{1}{2}))$ and $\hat{u}(s, \frac{1}{2}) = (u(s, 0), u(s, 1))$ which both lie in Δ for any $s \in \mathbb{R}$. Finally, projecting Floer's equation

$$\forall t \in [0, \frac{1}{2}], \quad \partial_s \hat{u} + \hat{J}_t(\hat{u})(\partial_t \hat{u} - X_{\hat{H}}^t(\hat{u})) = 0$$

to both components of the product shows that it is satisfied if and only if

$$\forall t \in [0, 1], \quad \partial_s u + J_t^\sigma(u)(\partial_t u - X_{H^\sigma}^t(u)) = 0.$$

Thus $\hat{u} \in \mathcal{M}^{\Delta, \Delta}(x_-, x_+; \hat{H}, \hat{J})$ if and only if $u \in \mathcal{M}(\gamma_-, \gamma_+; H^\sigma, J^\sigma)$ and (14) induces an isomorphism of complexes.

Finally, remark that there is an obvious correspondence between the cappings of a 1-periodic orbit γ and the half-cappings of its associated orbit x . In particular, a capping $\bar{\gamma}$ of γ can be thought of as a half-capping for x , by putting $\bar{x} = (\bar{x}_1, \bar{x}_2): D^2 \rightarrow M \times M$, with \bar{x}_1 the constant half-capping mapping D^2 to $\gamma(0)$ and x_2 the half-capping mapping ∂D^+ to the image of γ and ∂D^- to $\gamma(0)$. By doing so, not only \bar{x} maps ∂D^+ to the image of x in $M \times M$ and ∂D^- to $(\gamma(0), \gamma(0)) \in \Delta$, but the symplectic area of $\bar{\gamma}$ with respect to ω and the symplectic area of \bar{x} with respect to $(-\omega) \oplus \omega$ coincide. It easily follows that the action is preserved along the above transformation, namely $\mathcal{A}_{H^\sigma}(\gamma) = \mathcal{A}_{\hat{H}}^{\Delta, \Delta}(x)$.⁽⁵⁾

⁽⁵⁾ To be perfectly precise, we should have used an additional time-reparametrization to define \hat{H} on the whole interval $[0, 1]$. Since such a reparametrization is harmless in terms of spectral invariants as explained in Remark 14, we omitted it.

Now pick a Morse-Smale pair (f, g) on M and define (\hat{f}, \hat{g}) on Δ by putting $\hat{f}(x, x) = f(x)$ and $\hat{g}_{(x,x)}((\xi, \xi), (\eta, \eta)) = g_x(\xi, \eta)$ for all x in M and all ξ and η in $T_x M$. Then the pair (\hat{f}, \hat{g}) is a Morse-Smale pair for Δ and it is easy to show that the moduli spaces involved in the definition of the Hamiltonian PSS morphism in M correspond to those defining the Lagrangian PSS morphism in $M \times M$ with respect to Δ along the above process. Thus, for any non-zero homology class $\alpha \in HM(M)$, which we denote $\hat{\alpha}$ when seen as a homology class in $HM(\Delta)$, $c(\alpha; H^\sigma) = \ell(\hat{\alpha}; \Delta, \Delta; \hat{H})$. In particular, when $\alpha = [M]$, $\hat{\alpha} = [\Delta]$ so that $c_+(H^\sigma) = \ell([\Delta]; \Delta, \Delta; \hat{H})$.

Combined with (15), this concludes the proof for smooth non-degenerate Hamiltonians H . In view of the extension of both c and ℓ to $C^0([0, 1] \times M)$, Proposition 13 easily follows from the non-degenerate case. \square

2.5. Products in Lagrangian Floer theory and the triangle inequality

Let L_0, L_1 , and L_2 denote three Lagrangian submanifolds of (M, ω) . We fix three intersection points $p_{01} \in L_0 \cap L_1$, $p_{12} \in L_1 \cap L_2$, $p_{02} \in L_0 \cap L_2$ and suppose that each pair (L_i, L_j) is weakly exact, in the sense of Definition 8, with respect to the intersection point $p_{ij} \in L_i \cap L_j$. In this section, we describe the usual product structure on Lagrangian Floer homology.

We will be closely following the construction of this product as described in [1, Section 3]. There exist several other ways of defining the same product; see for example [3]. Let Σ denote the Riemann surface obtained by removing three points from the boundary of the closed unit disk in \mathbb{C} . We view Σ as a strip with a slit:

$$\Sigma = (\mathbb{R} \times [-1, 0] \sqcup \mathbb{R} \times [0, 1]) / \sim,$$

where $(s, 0^-) \sim (s, 0^+)$ for all $s \geq 0$. This is indeed a Riemann surface whose interior is naturally identified with $\mathbb{R} \times (-1, 1) \setminus (-\infty, 0] \times \{0\}$ and whose boundary consists of the three components $\mathbb{R} \times \{-1\}$, $\mathbb{R} \times \{1\}$, and $(-\infty, 0] \times \{0^-, 0^+\}$. At any point, other than $(0, 0)$, the inclusion of Σ into \mathbb{C} induces the standard complex structure $(s, t) \mapsto s + it$. At the point $(0, 0)$ the complex structure is given by the map $\{z \in \mathbb{C} : \operatorname{Re}(z) \geq 0\} \rightarrow \Sigma$, $z \mapsto z^2$.




FIGURE 1. Abbondandolo-Schwarz's strip with a slit, Σ

For $0 \leq i < j \leq 2$, denote by (H_{ij}, J_{ij}) a regular pair (of a Hamiltonian and a compatible time-dependent almost complex structure) for the weakly exact pair of Lagrangians (L_i, L_j) . Without loss of generality, we may assume that $H_{ij}(t, x) = 0$ for t near 0 and 1; see

Remark 14. To define the product structure, we need some auxiliary data: For $s \in (-\infty, \infty)$ and $t \in [-1, 1]$ let $J_{(s,t)}$ denote a family of almost complex structures on M such that

$$J_{(s,t)} = \begin{cases} J_{01}^{t+1} & \text{if } s \leq -1, t \in [-1, 0], \\ J_{12}^t & \text{if } s \leq -1, t \in [0, 1], \\ J_{02}^{\frac{t+1}{2}} & \text{if } s \geq 1, t \in [-1, 1]. \end{cases}$$

Furthermore, Choose a function $K : \mathbb{R} \times [-1, 1] \times M \rightarrow \mathbb{R}$ such that

$$K(s, t, x) = \begin{cases} H_{01}(t + 1, x) & \text{if } s \leq -1, t \in [-1, 0], \\ H_{12}(t, x) & \text{if } s \leq -1, t \in [0, 1], \\ \frac{1}{2}H_{02}(\frac{t+1}{2}, x) & \text{if } s \geq 1, t \in [-1, 1]. \end{cases}$$

For any three Hamiltonian chords $x_{ij} \in CF(L_i, L_j; p_{ij}; H_{ij})$, consider the moduli space $\mathcal{M}(x_{01}, x_{12}; x_{02})$ of maps $u : \Sigma \rightarrow M$ solving the Floer-type equation $\partial_s u + J_{(s,t)}(u)(\partial_t u - X_K^{s,t}(u)) = 0$ and subject to the following asymptotic and boundary conditions

$$\begin{cases} \forall t \in [-1, 0], u(-\infty, t) = x_{01}(t + 1) \text{ and } \forall t \in [0, 1], u(-\infty, t) = x_{12}(t), \\ \forall t \in [-1, 1], u(+\infty, t) = x_{02}(\frac{t+1}{2}), \\ u(\mathbb{R} \times \{-1\}) \subset L_0, u(\mathbb{R} \times \{1\}) \subset L_2, u((-\infty, 0] \times \{0^-, 0^+\}) \subset L_1. \end{cases}$$

For generic choices of K and J , the moduli space $\mathcal{M}(x_{01}, x_{12}; x_{02})$ is a smooth finite dimensional manifold. Its 0-dimensional component, denoted by $\mathcal{M}_{[0]}(x_{01}, x_{12}; x_{02})$, is compact and thus finite. We denote by $\#\mathcal{M}_{[0]}(x_{01}, x_{12}; x_{02})$ its cardinality modulo 2. We can now define a bilinear map

$$\begin{aligned} CF(L_0, L_1; p_{01}; H_{01}) \times CF(L_1, L_2; p_{12}; H_{12}) &\rightarrow CF(L_0, L_2; p_{02}; H_{02}) \\ (x_{01}, x_{12}) &\mapsto \sum_{x_{02}} \#\mathcal{M}_{[0]}(x_{01}, x_{12}; x_{02}) \cdot x_{02}. \end{aligned}$$

This map depends on the auxiliary data (K, J) . However, it can be shown that it induces a well-defined associative product at the level of homology:

$$\begin{aligned} HF(L_0, L_1; p_{01}; H_{01}, J_{01}) \otimes HF(L_1, L_2; p_{12}; H_{12}, J_{12}) \\ \longrightarrow HF(L_0, L_2; p_{02}; H_{02}, J_{02}). \end{aligned}$$

We will refer to this product as the pair-of-pants product. Given Floer homology classes α, β , we will denote their pair-of-pants product by $\alpha * \beta$.

2.5.1. *Compatibility of the pair-of-pants product with continuation maps.* – Denote by $H'_{ij}, 0 \leq i < j \leq 2$, three additional Hamiltonians which are non-degenerate with respect to the pairs (L_i, L_j) and pick three almost complex structures J'_{ij} so that the pairs (H'_{ij}, J'_{ij}) are regular. Let $\alpha \in HF(L_0, L_1; p_{01}; H_{01}, J_{01})$ and $\beta \in HF(L_1, L_2; p_{12}; H_{12}, J_{12})$. The pair-of-pants product $*$ is compatible with continuation maps in the following sense:

$$(16) \quad \Psi_{H_{01}, J_{01}}^{H'_{01}, J'_{01}}(\alpha) * \Psi_{H_{12}, J_{12}}^{H'_{12}, J'_{12}}(\beta) = \Psi_{H_{02}, J_{02}}^{H'_{02}, J'_{02}}(\alpha * \beta).$$

One can prove this formula by considering 0- and 1-dimensional components of suitable moduli spaces of objects combining continuation Floer strips and pair-of-pants strips with slits.

Note that this compatibility between pair-of-pants and continuation maps allows one to consider the pair-of-pants product as a product on Lagrangian Floer homology, independently of the auxiliary data:

$$HF(L_0, L_1; p_{01}) \otimes HF(L_1, L_2; p_{12}) \longrightarrow HF(L_0, L_2; p_{02}).$$

2.5.2. *The pair-of-pants product when $L_0 = L_1$.* – As mentioned in Section 2.1.1, in the case of a single Lagrangian L , the PSS isomorphism intertwines the Morse and Floer theoretic versions of the intersection product. Namely,

$$(17) \quad \Phi_{H,J}^L(a \cdot b) = \Phi_{H,J}^L(a) * \Phi_{H,J}^L(b)$$

for any regular pair (H, J) and any two classes a and b in $HM(L)$. So in the case of a single Lagrangian, the pair-of-pants product turns $HF(L, L; p)$ into a ring with unit $\Phi_{H,J}^L([L])$, where $[L]$ is the fundamental class of L .

2.5.3. *The triangle inequality.* – We continue to work with the Lagrangians L_0, L_1, L_2 from the previous sections. We call a triple (L_0, L_1, L_2) of Lagrangians weakly exact if any disk with boundary on $L_0 \cup L_1 \cup L_2$ and “corners” at p_{01}, p_{12}, p_{02} has vanishing symplectic area. More precisely, denote by D the closed unit disk in \mathbb{C} , and fix the three points $z_0 = 1, z_1 = e^{\frac{2\pi}{3}i}, z_2 = e^{-\frac{2\pi}{3}i}$ on the boundary of D . Let γ_0 denote the segment on the boundary of D between z_0 and z_1 , and similarly define γ_1, γ_2 .

DEFINITION 15. – *The triple (L_0, L_1, L_2) is weakly exact with respect to the intersection points (p_{01}, p_{12}, p_{02}) if $\int_D v^* \omega = 0$ for every disk*

$$v : (D, \gamma_0, \gamma_1, \gamma_2, z_0, z_1, z_2) \rightarrow (M, L_0, L_1, L_2, p_{01}, p_{12}, p_{02}).$$

Our main motivation for introducing the above definition is to establish sharp estimates needed to prove the triangle inequality satisfied by spectral invariants.

EXAMPLE 16. – Weakly exact pairs of Lagrangians in the sense of Definition 8 provide examples of weakly exact triples. Namely, if (L_0, L_1) is weakly exact with respect to $p \in L_0 \cap L_1$, then (L_0, L_0, L_1) and (L_0, L_1, L_1) are weakly exact with respect to (p, p, p) since a disk as in the definition above is a particular case of pinched disks as in Definition 8. This, combined with Example 9, shows that the triples in which we will be interested in the course of the proof of Theorem 4 (more precisely, in Theorem 23 below) are weakly exact with respect to (p, p, p) for any $p \in L_0 \cap L_1$. ◀

Let H_{01}, H_{12} denote any two time-dependent Hamiltonians on M . Define

$$(18) \quad H_{01} \# H_{12}(t, x) = \begin{cases} 2H_{01}(2t, x) & \text{if } t \in [0, \frac{1}{2}] \\ 2H_{12}(2t - 1, x) & \text{if } t \in [\frac{1}{2}, 1]. \end{cases}$$

Once again, without loss of generality we may assume that both H_{01} and H_{12} vanish for t near 0 and 1, see Remark 14. Hence, $H_{01} \# H_{12}$ is a smooth Hamiltonian. Observe that $\phi_{H_{01} \# H_{12}}^1 = \phi_{H_{12}}^1 \circ \phi_{H_{01}}^1$. The main goal of this section is to prove the following triangle inequality:

THEOREM 17. – *Let (L_0, L_1, L_2) be a triple of Lagrangians which is weakly exact with respect to (p_{01}, p_{12}, p_{02}) , where $p_{ij} \in L_i \cap L_j$. Denote by α, β homology classes in the reference Floer homology groups $HF_{\text{ref}}(L_0, L_1; p_{01})$ and $HF_{\text{ref}}(L_1, L_2; p_{12})$, respectively. The following inequality holds:*

$$\ell(\alpha * \beta; L_0, L_2; p_{02}; H_{01} \# H_{12}) \leq \ell(\alpha; L_0, L_1; p_{01}; H_{01}) + \ell(\beta; L_1, L_2; p_{12}; H_{12}).$$

Note that, the compatibility of the pair-of-pants product with continuation maps, as described in Section 2.5.1, allows us to view $\alpha * \beta$ in the reference Floer homology group $HF_{\text{ref}}(L_0, L_2; p_{02})$. We will now prove the triangle inequality.

Proof. – Recall that, by Inequality (7), the spectral invariant $\ell(\cdot; L_i, L_j; p_{ij}; H)$ depends continuously on H . Hence by replacing H_{01} , and H_{12} with nearby non-degenerate Hamiltonians if needed, we may assume that H_{01}, H_{12} and $H_{01} \# H_{12}$ are all regular.

Write $H_{02} = H_{01} \# H_{12}$. As in the previous section, take Hamiltonian chords $x_{ij} \in CF(L_i, L_j; p_{ij}; H_{ij})$ and consider the moduli space appearing in the definition of the pair-of-pants product, $\mathcal{M}(x_{01}, x_{12}; x_{02})$. For any $\epsilon > 0$, it is possible to pick the function $K : \mathbb{R} \times [-1, 1] \times M \rightarrow \mathbb{R}$ in the auxiliary data (K, J) such that

$$(19) \quad \left| \frac{\partial K_{s,t}}{\partial s} \right| \leq \frac{\epsilon}{4} \text{ if } s \in [-1, 1], \text{ and } \frac{\partial K_{s,t}}{\partial s} = 0 \text{ otherwise.}$$

Indeed, this can be achieved by making a small perturbation of the following function

$$K'(s, t, x) = \begin{cases} H_{01}(t + 1, x) & \text{for } t \in [-1, 0], \\ H_{12}(t, x) & \text{for } t \in [0, 1]. \end{cases}$$

We leave it to the reader to verify that proving the triangle inequality amounts to showing that $\mathcal{O}_{H_{02}}^{L_0, L_2}(x_{02}) \leq \mathcal{O}_{H_{01}}^{L_0, L_1}(x_{01}) + \mathcal{O}_{H_{12}}^{L_1, L_2}(x_{12})$. We will now prove this last inequality. For any $u \in \mathcal{M}(x_{01}, x_{12}; x_{02})$, the following holds:

$$\begin{aligned} 0 &\leq \int_{\Sigma} \|\partial_s u(s, t)\|^2 ds dt = \int_{\Sigma} \omega(\partial_s u, J_{(s,t)}(u) \partial_s u) ds dt \\ &= \int_{\Sigma} \omega(\partial_s u, \partial_t u - X_K^{s,t}(u)) ds dt = \int_{\Sigma} u^* \omega - \int_{\Sigma} dK_{s,t}(\partial_s u) ds dt. \end{aligned}$$

Now, let \bar{x}_{ij} denote homotopies from the chords x_{ij} to the constant paths p_{ij} , i.e., cappings for x_{ij} . Since, the triple (L_0, L_1, L_2) is weakly exact with respect to (p_{01}, p_{12}, p_{02}) , the disk $\bar{x}_{01} \# \bar{x}_{12} \# u \# (-\bar{x}_{02})$ has symplectic area zero. Hence, we see that

$$\int_{\Sigma} u^* \omega = - \int_D (\bar{x}_{01})^* \omega - \int_D (\bar{x}_{12})^* \omega + \int_D (\bar{x}_{02})^* \omega.$$

On the other hand, Equation (19) implies that $\int_{\Sigma} \partial_s K_{s,t}(u) ds dt \leq \epsilon$ and hence we obtain the following:

$$\begin{aligned} - \int_{\Sigma} dK_{s,t}(\partial_s u) ds dt &= - \int_{\Sigma} \partial_s (K_{s,t} \circ u) ds dt + \int_{\Sigma} \partial_s K_{s,t}(u) ds dt \\ &\leq \int_0^1 H_{01}(t, x_{01}(t)) dt + \int_0^1 H_{12}(t, x_{12}(t)) dt - \int_{-1}^1 H_{02}(t, x_{02}(t)) dt + \epsilon. \end{aligned}$$

We conclude from the above that

$$0 \leq \int_{\Sigma} \|\partial_s u(s, t)\|^2 ds dt \leq \mathcal{A}_{H_{01}}^{L_0, L_1}(x_{01}) + \mathcal{A}_{H_{12}}^{L_1, L_2}(x_{12}) - \mathcal{A}_{H_{02}}^{L_0, L_2}(x_{02}) + \epsilon$$

which finishes the proof of the triangle inequality. \square

2.6. A Künneth formula for Lagrangian Floer homology and a splitting formula for spectral invariants

Let (M', ω') and (M'', ω'') denote two closed symplectically aspherical symplectic manifolds. Let (L'_0, L'_1) denote a pair of Lagrangians in M' which is weakly exact with respect to a fixed intersection point $p' \in L'_0 \cap L'_1$. Take (H', J') to be a regular pair (of a Hamiltonian and an almost complex structure), as defined in Section 2.1, for the weakly exact pair of Lagrangians (L_0, L_1) . Similarly, we define (L''_0, L''_1) , $p'' \in L''_0 \cap L''_1$, and (H'', J'') in M'' .

Consider the product Lagrangians $L_0 = L'_0 \times L''_0$, $L_1 = L'_1 \times L''_1$ in $(M' \times M'', \omega' \oplus \omega'')$, the Hamiltonian $H_t(x, y) = H' \oplus H''(t, (x, y)) := H'_t(x) + H''_t(y)$, and the almost complex structure $J = J' \oplus J''$. Note that the pair (L_0, L_1) is weakly exact with respect to the intersection point $(p', p'') \in L_0 \cap L_1$. It is easy to see that the Hamiltonian H is non-degenerate, and moreover, the Hamiltonian chords of H are of the form $x = (x', x'')$ where x', x'' are Hamiltonian chords of H' and H'' .

The pair (H, J) is regular for (L_0, L_1) : This is because the linearization of the operator $u \mapsto \partial_s u + J_t(u)(\partial_t u - X_H^t(u))$ splits into a product of the corresponding linearizations for (H', J') and (H'', J'') ; see, for example, [13] for further details. It follows that, for any two chords $x_- = (x'_-, x''_-)$ and $x_+ = (x'_+, x''_+)$ of H , the moduli space $\widehat{\mathcal{M}}^{L_0, L_1}(x_-, x_+; H, J)$, used in the definition of the Floer boundary map, coincides with the product

$$\widehat{\mathcal{M}}^{L'_0, L'_1}(x'_-, x'_+; H', J') \times \widehat{\mathcal{M}}^{L''_0, L''_1}(x''_-, x''_+; H'', J'').$$

We leave it to the reader to conclude from the discussion in the preceding paragraph that

$$CF(L_0, L_1; p; H) = CF(L'_0, L'_1; p'; H') \otimes CF(L''_0, L''_1; p''; H''),$$

where the boundary map ∂ is defined by $\partial x = \partial' x' \otimes x'' + x' \otimes \partial'' x''$, with ∂' and ∂'' denoting the boundary maps for the Floer complexes of H' and H'' , respectively. Recall that we are working over \mathbb{Z}_2 and thus applying the standard Künneth formula we obtain

$$(20) \quad HF(L_0, L_1; p; H, J) = HF(L'_0, L'_1; p'; H', J') \otimes HF(L''_0, L''_1; p''; H'', J'').$$

2.6.1. *A splitting formula for spectral invariants.* – We present in this section a splitting formula⁽⁶⁾ for spectral invariants in the situation described above. Consider a Floer homology class $\alpha = \alpha' \otimes \alpha'' \neq 0$ in $HF(L'_0, L'_1; p'; H', J') \otimes HF(L''_0, L''_1; p''; H'', J'')$. By the discussion above, α is a homology class in $HF(L_0, L_1; p; H, J)$. The following splitting formula holds:

$$(21) \quad \ell(\alpha; L_0, L_1; p; H) = \ell(\alpha'; L'_0, L'_1; p'; H') + \ell(\alpha''; L''_0, L''_1; p''; H'').$$

⁽⁶⁾ This is sometimes called the “product formula” in the literature. We have chosen this alternative terminology in order to avoid any possible confusion with the triangle inequality coming from product of homology classes.

In [6, Section 5], a more abstract and general version of the above formula is proven for spectral invariants of “decorated \mathbb{Z}_2 -graded complexes,” see [6, Theorem 5.2]. Formula (21) is an immediate corollary of this theorem.

2.6.2. *Compatibility of the Künneth Formula with the pair-of-pants product.* – In this section, we describe the compatibility of the Künneth Formula (20) with the pair-of-pants product as defined in Section 2.5.

Let $L_0 = L'_0 \times L''_0$, $L_1 = L'_1 \times L''_1 \subset M' \times M''$ be as in the previous section, and consider additionally a third Lagrangian $L_2 = L'_2 \times L''_2$. For $0 \leq i < j \leq 2$, take three intersection points $p_{ij} = (p'_{ij}, p''_{ij}) \in L_i \cap L_j$ and suppose that (L'_i, L'_j) , (L''_i, L''_j) are weakly exact with respect to the intersection points p'_{ij} , p''_{ij} , respectively. Lastly, let (H'_{ij}, J'_{ij}) and (H''_{ij}, J''_{ij}) denote regular pairs for (L'_i, L'_j) and (L''_i, L''_j) , respectively.

As in the previous section, we consider the split Hamiltonians and almost complex structures $H_{ij} = H'_{ij} \oplus H''_{ij}$, $J_{ij} = J'_{ij} \oplus J''_{ij}$. By the Künneth Formula (20), $HF(L_i, L_j; p_{ij}; H_{ij}, J_{ij})$ is generated by elements of the form $\alpha' \otimes \alpha''$, where $\alpha' \in HF(L'_i, L'_j; p'_{ij}; H'_{ij}, J'_{ij})$ and $\alpha'' \in HF(L''_i, L''_j; p''_{ij}; H''_{ij}, J''_{ij})$. Therefore, describing the pair-of-pants product, in this setting, reduces to describing the product for such elements.

Consider $\alpha' \otimes \alpha'' \in HF(L_0, L_1; p_{01}; H_{01}, J_{01})$ and $\beta' \otimes \beta'' \in HF(L_1, L_2; p_{12}; H_{12}, J_{12})$. Then, the following equality holds in $HF(L_0, L_2; p_{02}; H_{02}, J_{02})$:

$$(22) \quad (\alpha' \otimes \alpha'') * (\beta' \otimes \beta'') = (\alpha' * \beta') \otimes (\alpha'' * \beta'').$$

The reasoning as to why the above holds is very similar to the reasoning for the Künneth Formula (20): Let $x_{ij} = (x'_{ij}, x''_{ij})$ denote Hamiltonian chords for $H_{ij} = H'_{ij} \oplus H''_{ij}$. Recall from Section 2.5 the moduli space $\mathcal{M}(x_{01}, x_{12}; x_{02})$ which is used to define the pair-of-pants product. Such moduli spaces split into products:

$$\mathcal{M}(x_{01}, x_{12}; x_{02}) = \mathcal{M}(x'_{01}, x'_{12}; x'_{02}) \times \mathcal{M}(x''_{01}, x''_{12}; x''_{02}).$$

2.6.3. *Compatibility of the Künneth formula with the PSS isomorphism and the splitting formula.*– Consider the case of a single Lagrangian $L = L' \times L''$. The PSS morphism as described in this particular case in Section 2.1.1 is compatible with the Künneth Formula (20). This was the content of [13, Claim 3.4] in the more general case of monotone manifolds. More precisely, the Morse theoretic version of Künneth’s formula is satisfied, that is

$$HM(L) = HM(L') \otimes HM(L'').$$

As in the Floer theoretic case this can be proven, even at the chain level, by choosing a Morse-Smale pair (f, g) on L which splits, that is $f = f' \oplus f''$ and $g = g' \oplus g''$ where (f', g') and (f'', g'') are Morse-Smale pairs for L' and L'' , respectively.

Now, for such split Morse and Floer data, respectively (f, g) and (H, J) , one can easily prove that for all $a' \in HM(L'; f', g')$ and $a'' \in HM(L''; f'', g'')$,

$$(23) \quad \Phi_{H,J}^L(a' \otimes a'') = \Phi_{H',J'}^{L'}(a') \otimes \Phi_{H'',J''}^{L''}(a'')$$

again, even at the chain level, since the moduli spaces involved in the construction of the PSS isomorphism themselves split along the product. (The fact that the PSS isomorphism is compatible with Morse and Floer continuation morphisms then allows one to consider, at the homological level, non necessarily split data.)

It follows that the splitting Formula (21) restricts to the following:

$$(24) \quad \ell(a \otimes b; L, L; H' \oplus H'') = \ell(a; L', L'; H') + \ell(b; L'', L''; H'')$$

for all non-zero Morse homology classes $a \in HM(L')$ and $b \in HM(L'')$. Notice that this corresponds to [14, Theorem 2.14] in the case of Hamiltonians with complete flows on cotangent bundles.

3. Lagrangian Floer theory of tori

In this section, we specialize the theory developed in Section 2 to the settings introduced in Section 1.2. Recall that, $M = \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2} \times \mathbb{T}^{2k_2}$ and that it is equipped with the standard symplectic form ω_{std} . Furthermore, recall that the Lagrangians we are interested in, L_0 and L_1 , are defined as follows

$$\begin{aligned} L_0 &= \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\}, \\ L_1 &= \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\} \times \{0\} \times \mathbb{T}^{k_2}. \end{aligned}$$

As noted in Section 1.2, $L_i = L \times L'_i$, where

$$\begin{aligned} L &= \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\} \subset \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}, \\ L'_0 &= \mathbb{T}^{k_2} \times \{0\} \subset \mathbb{T}^{2k_2}, \text{ and } L'_1 = \{0\} \times \mathbb{T}^{k_2} \subset \mathbb{T}^{2k_2}. \end{aligned}$$

The pairs of Lagrangians (L, L) , (L'_0, L'_1) , and $(L_0, L_1) = (L \times L'_0, L \times L'_1)$ are all weakly exact with respect to any point in their corresponding intersections; see Example 9. We fix, for the rest of this article, in the intersection of each of the above pairs the point all of whose coordinates are zero, and carry out the constructions of Floer homology and spectral invariants (as described in Section 2) with respect to this intersection point. We will omit the intersection point from our notation.

3.1. $HF(L_0, L_1)$ and the associated spectral invariants

In this section, we construct an isomorphism between Morse homology $HM(L)$ and Floer homology $HF(L_0, L_1; H, J)$. We will then use this isomorphism to associate spectral invariants to Morse homology classes.

THEOREM 18. – *There exists a PSS-type isomorphism*

$$\Phi_{H,J}^{L_0,L_1} : HM(L) \rightarrow HF(L_0, L_1; H, J),$$

associated to every regular pair (H, J) . Furthermore, the isomorphism $\Phi_{H,J}^{L_0,L_1}$ is compatible with continuation morphisms in the following sense:

$$(25) \quad \Phi_{H,J}^{L_0,L_1} = \Psi_{H',J'}^{H,J} \circ \Phi_{H',J'}^{L_0,L_1},$$

where (H', J') is any other regular pair.

Proof. – Pick a Hamiltonian F and an almost complex structure j , on $\mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}$, such that (F, j) is regular for the pair of Lagrangians (L, L) . Similarly, we pick (F', j') , on \mathbb{T}^{2k_2} , such that (F', j') is regular for the pair (L'_0, L'_1) . Define the Hamiltonian $F \oplus F'$ on M by $F \oplus F'(z_1, z_2) = F(z_1) + F'(z_2)$, for $z_1 \in \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}$ and $z_2 \in \mathbb{T}^{2k_2}$.

We know from Sections 2.1.1 and 2.1.2 that there exist a PSS isomorphism

$$\Phi_{F,j}^L : HM(L) \rightarrow HF(L, L; F, j)$$

and a PSS-type isomorphism between $\Phi_{F',j'}^{L'_0,L'_1} : \mathbb{Z}_2 \rightarrow HF(L'_0, L'_1; F', j')$.

We define

$$\Phi_{F \oplus F', j \oplus j'}^{L_0, L_1} : HM(L) \rightarrow HF(L, L; F, j) \otimes HF(L'_0, L'_1; F', j')$$

to be the tensor product of these two isomorphisms, i.e., $\forall a \in HM(L)$ we have:

$$(26) \quad \Phi_{F \oplus F', j \oplus j'}^{L_0, L_1}(a) = \Phi_{F,j}^L(a) \otimes [\text{pt}],$$

where $[\text{pt}]$ denotes the non-trivial homology class in $HF(L'_0, L'_1; F', j')$. On the other hand, the Künneth Formula (20) tells us that

$$HF(L_0, L_1; F \oplus F', j \oplus j') = HF(L, L; F, j) \otimes HF(L'_0, L'_1; F', j').$$

Hence, $\Phi_{F \oplus F', j \oplus j'}^{L_0, L_1}$ gives an isomorphism between $HM(L)$ and $HF(L_0, L_1; F \oplus F', j \oplus j')$. For an arbitrary regular pair (H, J) we define $\Phi_{H,J}^{L_0, L_1} : HM(L) \rightarrow HF(L_0, L_1; H, J)$ by the formula

$$\Phi_{H,J}^{L_0, L_1} = \Psi_{F \oplus F', j \oplus j'}^{H, J} \circ \Phi_{F \oplus F', j \oplus j'}^{L_0, L_1}.$$

The proof of the second half of the theorem, concerning the compatibility of $\Phi_{H,J}^{L_0, L_1}$ with continuation morphisms, is an immediate consequence of the definition above and the fact that continuation morphisms are canonical in the sense of Equation (2). □

As an immediate consequence of Theorem 18, we can associate spectral invariants to elements of $HM(L)$.

DEFINITION 19. – *For every regular Hamiltonian H , the spectral invariant associated to a non-zero element $a \in HM(L)$ is the number*

$$\ell(a; L_0, L_1; H) := \ell(\Phi_{H,J}^{L_0, L_1}(a); L_0, L_1; H),$$

where the right-hand side is defined via Equation (5).

REMARK 20. – The above definition is compatible with Definition 11. In effect, what we have done in the above definition is to take the reference pair $(H_{\text{ref}}, J_{\text{ref}})$ of Definition 11 to be the pair $(F \oplus F', j \oplus j')$ from the proof of Theorem 18.

It follows immediately from Inequality (7) that for non-degenerate H, H' , we have:

$$(27) \quad \int_0^1 \min_M (H_t - H'_t) dt \leq |\ell(a; L_0, L_1; H) - \ell(a; L_0, L_1; H')| \leq \int_0^1 \max_M (H_t - H'_t) dt.$$

We conclude that $\ell(a; L_0, L_1; H)$ can be defined by continuity for every continuous function $H : [0, 1] \times M \rightarrow \mathbb{R}$.

We end this subsection by mentioning that, as in Section 2.2.1, one can easily verify the spectrality property, $\ell(a; L_0, L_1; H) \in \text{Spec}(H)$.

3.2. Product structure and the triangle inequality

Recall that the Morse homology of any manifold X carries a ring structure where the product of $a, b \in HM(X)$ is given by the intersection product $a \cdot b$.

Consider the Lagrangian submanifolds $L_0 = L \times L'_0$, $L_1 = L \times L'_1$. As a consequence of the Künneth formula for Morse homology, the homology ring $HM(L_i)$ can be written as the tensor product of the rings $HM(L)$ and $HM(L'_i)$, i.e.,

$$HM(L_i) = HM(L) \otimes HM(L'_i).$$

For $1 \leq j \leq 3$, let (H_j, J_j) denote three regular pairs. Recall that in Section 2.5 we defined the pair-of-pants product. Here, we will consider the following two instances of the pair-of-pants product

$$\begin{aligned} * : HF(L_0, L_1; H_1, J_1) \otimes HF(L_1, L_1; H_2, J_2) &\rightarrow HF(L_0, L_1; H_3, J_3), \\ * : HF(L_0, L_0; H_1, J_1) \otimes HF(L_0, L_1; H_2, J_2) &\rightarrow HF(L_0, L_1; H_3, J_3). \end{aligned}$$

The next theorem describes the relation between the above two pair-of-pants products and the intersection product on $HM(L)$.

THEOREM 21. – *Denote by $[L'_i]$, for $i = 0, 1$, the fundamental class in $HM(L'_i)$ and by $a, b \in HM(L)$ any two Morse homology classes. The intersection and pair-of-pants products satisfy the following relations:*

1. $\Phi_{H_1, J_1}^{L_0, L_1}(a) * \Phi_{H_2, J_2}^{L_1}(b \otimes [L'_1]) = \Phi_{H_3, J_3}^{L_0, L_1}(a \cdot b)$,
2. $\Phi_{H_1, J_1}^{L_0}(a \otimes [L'_0]) * \Phi_{H_2, J_2}^{L_0, L_1}(b) = \Phi_{H_3, J_3}^{L_0, L_1}(a \cdot b)$.

Proof. – We will only prove the first of the above two identities. The second is proven in a similar fashion.

Recall that $L \subset \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}$ and $L'_0, L'_1 \subset \mathbb{T}^{2k_2}$. Let F and F' denote two non-degenerate Hamiltonians on $\mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}$ and \mathbb{T}^{2k_2} , respectively. We claim that it is sufficient to prove the theorem in the special case where $H_1 = H_2 = H_3 = F \oplus F'$. Indeed, this can be deduced using the following two ingredients: First, the compatibility of PSS and PSS-type isomorphisms with continuation isomorphisms, as described by Diagram (3) and Equation (25). Second, the compatibility of continuation isomorphisms with the pair-of-pants product as described by Equation (16). We will prove the theorem in this special case and leave it to the reader to verify that this indeed does imply the general case.

Pick almost complex structures J, J' such that the pairs (F, J) and (F', J') are both regular. Now, it follows from Equation (26) that

$$\Phi_{F \oplus F', J \oplus J'}^{L_0, L_1}(a) = \Phi_{F, J}^L(a) \otimes [\text{pt}],$$

where $[\text{pt}]$ denotes the non-trivial homology class in $HF(L'_0, L'_1; F', j')$. We also know, from Equation (23), that

$$\Phi_{F \oplus F', J \oplus J'}^{L_1}(b \otimes [L'_1]) = \Phi_{F, J}^L(b) \otimes \Phi_{F', J'}^{L'_1}([L'_1]).$$

We will be needing the following identity, whose proof we postpone for the time being:

$$(28) \quad [\text{pt}] * \Phi_{F', J'}^{L'_1}([L'_1]) = [\text{pt}].$$

Using the above, we obtain the following:

$$\begin{aligned}
 & \Phi_{F \oplus F', J \oplus J'}^{L_0, L_1}(a) * \Phi_{F \oplus F', J \oplus J'}^{L_1}(b \otimes [L'_1]) \\
 &= (\Phi_{F, J}^L(a) \otimes [\text{pt}]) * (\Phi_{F, J}^L(b) \otimes \Phi_{F', J'}^{L'_1}([L'_1])) \\
 &= (\Phi_{F, J}^L(a) * \Phi_{F, J}^L(b)) \otimes ([\text{pt}] * \Phi_{F', J'}^{L'_1}([L'_1])) && \text{by Equation (22)} \\
 &= \Phi_{F, J}^L(a \cdot b) \otimes ([\text{pt}] * \Phi_{F', J'}^{L'_1}([L'_1])) && \text{by Equation (17)} \\
 &= \Phi_{F, J}^L(a \cdot b) \otimes [\text{pt}] && \text{by Equation (28)} \\
 &= \Phi_{F \oplus F', J \oplus J'}^{L_0, L_1}(a \cdot b) && \text{by Equation (26)}.
 \end{aligned}$$

It remains to prove Equation (28). Recall that

$$L'_0 = \mathbb{T}^{k_2} \times \{0\} \subset \mathbb{T}^{2k_2}, L'_1 = \{0\} \times \mathbb{T}^{k_2} \subset \mathbb{T}^{2k_2}.$$

Let

$$\Lambda_0 = \mathbb{T}^1 \times \{0\} \subset \mathbb{T}^2, \Lambda_1 = \{0\} \times \mathbb{T}^1 \subset \mathbb{T}^2,$$

and observe that (up to a symplectomorphism)

$$L'_0 = \underbrace{\Lambda_0 \times \cdots \times \Lambda_0}_{k_2 \text{ times}}, L'_1 = \underbrace{\Lambda_1 \times \cdots \times \Lambda_1}_{k_2 \text{ times}}.$$

Equations (3) and (16) tell us, respectively, that continuation morphisms are compatible with the PSS isomorphism and the pair-of-pants product. From this, one can conclude that it is sufficient to verify Equation (28) for any specific choice of a regular pair (F', J') . Furthermore, the regular pair used for defining $HF(L'_0, L'_1)$ can indeed be different from the one used for defining $HF(L'_1, L'_1)$. We will verify the formula for the choices described in the next two paragraphs.

For $HF(L'_1, L'_1)$ we pick a regular pair of the form

$$(\underbrace{f \oplus \cdots \oplus f}_{k_2 \text{ times}}, \underbrace{j \oplus \cdots \oplus j}_{k_2 \text{ times}}),$$

where (f, j) denotes a regular pair on the 2-torus \mathbb{T}^2 , the almost complex structure $j \oplus \cdots \oplus j$ denotes the obvious split almost complex structure on \mathbb{T}^{k_2} , and $f \oplus \cdots \oplus f(z_1, \dots, z_{k_2}) = f(z_1) + \cdots + f(z_{k_2})$.

For $HF(L'_0, L'_1)$ we pick a regular pair of the form

$$(0, j_0 \oplus \cdots \oplus j_0),$$

where 0 denotes the zero Hamiltonian. Recall that since L'_0 and L'_1 intersect transversely the zero Hamiltonian is non-degenerate for this pair.

Now, by the Künneth Formula (20) we have the following splittings

$$\begin{aligned}
 HF(L'_0, L'_1; 0, j_0 \oplus \cdots \oplus j_0) &= HF(\Lambda_0, \Lambda_1; 0, j_0)^{\otimes k_2}, \text{ and} \\
 HF(L'_1, L'_1; f \oplus \cdots \oplus f, j \oplus \cdots \oplus j) &= HF(\Lambda_1, \Lambda_1; f, j)^{\otimes k_2}.
 \end{aligned}$$

Furthermore, the above splittings are compatible with the pair-of-pants product as described by Equation (22). This implies that Equation (28) is an immediate consequence of the following claim:

CLAIM. – Denote by p the unique intersection point of Λ_0 and Λ_1 and by $[p]$ the Floer homology class represented by this point. Note that this is the unique non-zero class in $HF(\Lambda_0, \Lambda_1; 0, j_0)$. The pair-of-pants product $* : HF(\Lambda_0, \Lambda_1; 0, j_0) \otimes HF(\Lambda_1, \Lambda_1; f, j) \rightarrow HF(\Lambda_0, \Lambda_1; 0, j_0)$ satisfies the following identity:

$$(29) \quad [p] * \Phi_{f,j}^{\Lambda_1}([\Lambda_1]) = [p].$$

The proof of the above claim boils down to computing one of the simplest instances of the pair-of-pants product. This is well-known to experts, and thus we only present a sketch of the proof while avoiding the technical details.

Proof of Claim. – Once again, by Equations (3) and (16), it is sufficient to verify the above claim for a given choice of a regular pair (f, j) . We pick f as follows: begin with a C^2 -small Morse function on the circle Λ_1 and extend it trivially to the product $\mathbb{T}^2 = \Lambda_0 \times \Lambda_1$. Furthermore, we pick f such that $f|_{\Lambda_1}$ has only two critical points. Denote the maximum by Q and the minimum by q .

Let $\Lambda'_1 = \phi_f^1(\Lambda_1)$. Since f is C^2 -small, Λ'_1 intersects Λ_0 transversely at a single point. We denote this point by p' . See Figure 2.




FIGURE 2. Pair-of-pants product in \mathbb{T}^2

It is well-known that there exists a natural identification of $HF(\Lambda_1, \Lambda_1; f, j)$ with $HF(\Lambda_1, \Lambda'_1; 0, \tilde{j})$, where $\tilde{j}_t = (\phi_f^t)_*^{-1} j_t$; see for example [12, Section 2.2.2] or [3, Remark 1.10]. The point Q represents a homology class $[Q] \in HF(\Lambda_1, \Lambda'_1; 0, \tilde{j})$, which corresponds to the fundamental class $[\Lambda_1] \in HM(\Lambda_1)$. Also, $HF(\Lambda_0, \Lambda'_1; 0, j_0)$ is generated by $[p']$, the homology class represented by the point p' .

Again, using Diagram (3) and Equation (16), we see that Equation (29) is equivalent to

$$(30) \quad [p] * [Q] = [p'].$$

This last equality can be verified without much difficulty.

First, note that $[p] \in HF(\Lambda_0, \Lambda_1; 0, j_0)$, $[Q] \in HF(\Lambda_1, \Lambda'_1; 0, \tilde{j})$, and $[p'] \in HF(\Lambda_0, \Lambda_1; 0, j_0)$, and thus, the Hamiltonians in question are all zero. Furthermore, we can take j_0 to be the standard complex structure on \mathbb{T}^2 and j such that $\tilde{j} = j_0$. Hence, to verify that $[p] * [Q] = [p']$, we must count the number of holomorphic disks on \mathbb{T}^2 with boundary on $\Lambda_0 \cup \Lambda_1 \cup \Lambda'_0$ and corners at the points p , Q , and p' . We leave it to the reader to verify that there exists only one such disk: the one highlighted in Figure 2. This proves Equation (30). \square

This completes the proof of Theorem 21. □

REMARK 22. – For $i = 0, 1$, denote by $H_k(L'_i)$ the Morse homology group of degree k of L'_i . Suppose that $x \in H_k(L'_i)$ where $k \neq \dim(L'_i)$. Then, one can modify the proof of Theorem 21 to obtain the following additional identities:

1. $\Phi_{H_1, J_1}^{L_0, L_1}(a) * \Phi_{H_2, J_2}^{L_1}(b \otimes x) = 0,$
2. $\Phi_{H_1, J_1}^{L_0}(a \otimes x) * \Phi_{H_2, J_2}^{L_0, L_1}(b) = 0.$

The above combined with Theorem 21, give us a full description of the relation between the intersection product on $HM(L)$ and the pair-of-pants products

$$\begin{aligned} * : HF(L_0, L_1) \otimes HF(L_1, L_1) &\rightarrow HF(L_0, L_1), \\ * : HF(L_0, L_0) \otimes HF(L_0, L_1) &\rightarrow HF(L_0, L_1). \end{aligned}$$

We will not prove these additional identities, as they are not needed for the proof of Theorem 4 and their proofs are very similar to the proof of the previous theorem. We mention here that, in the same way that the proof of Theorem 21 was reduced to establishing Equation (30), proving these identities reduces to showing the following:

$$[p] * [q] = 0,$$

where p and q are defined as in Figure 2.

3.2.1. *The triangle inequality.* – In this section, we use Theorem 21 to prove the two triangle inequalities mentioned in the introduction. Recall that given two Hamiltonians H, H' , their concatenation $H \# H'$ is defined by

$$H \# H'(t, x) = \begin{cases} 2H(2t, x) & \text{if } t \in [0, \frac{1}{2}], \\ 2H'(2t - 1, x) & \text{if } t \in [\frac{1}{2}, 1]. \end{cases}$$

THEOREM 23. – Denote by $[L'_i]$, for $i = 0, 1$, the fundamental class in $HM(L'_i)$ and by $a, b \in HM(L)$ any two Morse homology classes such that $a \cdot b \neq 0$. The following inequalities hold:

1. $\ell(a \cdot b; L_0, L_1; H \# H') \leq \ell(a; L_0, L_1; H) + \ell(b \otimes [L'_1]; L_1, L_1; H'),$
2. $\ell(a \cdot b; L_0, L_1; H \# H') \leq \ell(a \otimes [L'_0]; L_0, L_0; H) + \ell(b; L_0, L_1; H').$

Proof. – We will only prove the first of the two inequalities, as the second one is proven in a very similar fashion.

By continuity of spectral invariants (27), it is sufficient to prove the inequality in the special case where H, H' and $H \# H'$ are all non-degenerate. Pick an almost complex structure J such that the pairs $(H, J), (H', J)$, and $(H \# H', J)$ are all regular. Now, the triangle inequality becomes a simple consequence of Theorem 21 and the triangle inequality of Theorem 17. Indeed,

$$\begin{aligned} \ell(a \cdot b; L_0, L_1; H \# H') &= \ell(\Phi_{H \# H', J}^{L_0, L_1}(a \cdot b); L_0, L_1; H \# H') \\ &= \ell(\Phi_{H, J}^{L_0, L_1}(a) * \Phi_{H', J}^{L_1}(b \otimes [L'_1]); L_0, L_1; H \# H') \\ &\leq \ell(\Phi_{H, J}^{L_0, L_1}(a); L_0, L_1; H) + \ell(\Phi_{H', J}^{L_1}(b \otimes [L'_1]); L_1, L_1; H') \\ &= \ell(a; L_0, L_1; H) + \ell(b \otimes [L'_1]; L_1, L_1; H'). \end{aligned}$$

Note that the triangle inequality of Theorem 17 can be applied here because by Example 16 since (L_0, L_1) is a weakly exact pair, (L_0, L_1, L_1) is a weakly exact triple. \square

We now use the triangle inequality to prove Proposition 6.

Proof of Proposition 6. – We will only provide a proof in the case $i = 1$. The other case is proven in a similar fashion.

Applying Theorem 23 in the special case where $b = [L]$ and $H = 0$ we obtain

$$\ell(a; L_0, L_1; 0 \# H') \leq \ell(a; L_0, L_1; 0) + \ell([L_1]; L_1, L_1; H').$$

Now, $\ell(a; L_0, L_1; 0) = 0$ because the spectrum of the zero Hamiltonian is the singleton $\{0\}$. The Hamiltonian $0 \# H'$ is a reparametrization of H' and so, by Remark 14, $\ell(a; L_0, L_1; 0 \# H') = \ell(a; L_0, L_1; H')$. This finishes the proof. \square

3.3. A splitting formula

In this subsection, we will use the splitting formula (21) to obtain a similar formula in our current setting. This will be used in the proof of Theorem 4.

Recall that, $M = \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2} \times \mathbb{T}^{2k_2}$. Let F and F' denote two Hamiltonians on $\mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}$ and \mathbb{T}^{2k_2} , respectively. Define the Hamiltonian $F \oplus F'$ on M by $F \oplus F'(z_1, z_2) = F(z_1) + F'(z_2)$, for $z_1 \in \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}$ and $z_2 \in \mathbb{T}^{2k_2}$. This Hamiltonian appeared in the proof of Theorem 18.

THEOREM 24. – *Let $F \oplus F'$ denote any Hamiltonian of the form described in the previous paragraph. Let $a \in HM(L)$ denote a non-zero class. The following formula holds:*

$$\ell(a; L_0, L_1; F \oplus F') = \ell(a; L, L; F) + \ell([\text{pt}]; L'_0, L'_1; F'),$$

where $[\text{pt}]$ denotes the non-zero class in $HF(L'_0, L'_1)$.


Proof. – By continuity of spectral invariants (27), it is sufficient to prove the theorem for non-degenerate F and F' . Pick almost complex structures J and J' such that the pairs (F, J) and (F', J') are regular. We have the following chain of equalities:

$$\begin{aligned} \ell(a; L_0, L_1; F \oplus F') &= \ell(\Phi_{F \oplus F', J \oplus J'}^{L_0, L_1}(a); L_0, L_1; F \oplus F') && \text{by Definition 19} \\ &= \ell(\Phi_{F, J}^L(a) \otimes [\text{pt}]; L_0, L_1; F) && \text{by Equation (26)} \\ &= \ell(\Phi_{F, J}^L(a); L, L; F) + \ell([\text{pt}]; L'_0, L'_1; F') && \text{by Equation (21)} \\ &= \ell(a; L, L; F) + \ell([\text{pt}]; L'_0, L'_1; F') && \text{by Definition 10. } \square \end{aligned}$$

4. Proof of the main theorem (Theorem 4)

We start this section by introducing the notations needed for the proof. As in Theorem 4, we consider a symplectic homeomorphism ϕ of $(\mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}, \omega_{\text{std}})$ which preserves the coisotropic submanifold $C = \mathbb{T}^{2k_1} \times \mathbb{T}^{k_2} \times \{0\}^{k_2}$. Observe that the characteristic foliation \mathcal{F} is parallel to the subtorus $\{0\}^{2k_1} \times \mathbb{T}^{k_2} \times \{0\}^{k_2}$. The map ϕ induces a homeomorphism ϕ_R on the reduced space $\mathcal{R} = C/\mathcal{F} = \mathbb{T}^{2k_1}$. Throughout the proof, given a homeomorphism θ between two spaces X and Y , and a time-dependent function ρ on Y , i.e., a function $\rho : [0, 1] \times Y \rightarrow \mathbb{R}$, the composition $\rho \circ \theta$ will denote, with a slight abuse of notation, the time dependent function on X defined by $\rho \circ \theta(t, x) = \rho(t, \theta(x))$ for all $t \in [0, 1]$ and $x \in X$. We want to show that ϕ_R preserves the spectral invariant c_+ , i.e., for every time-dependent continuous function f_R on \mathcal{R} , $c_+(f_R \circ \phi_R) = c_+(f_R)$.

Let f_R be a time-dependent continuous function on the reduced space \mathcal{R} and denote $g_R = f_R \circ \phi_R$. We denote by f and g , respectively, the standard lifts of f_R and g_R to $\mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2}$, given by $f(z_1, z_2) = f_R(z_1)$ and $g(z_1, z_2) = g_R(z_1)$, for all $z_1 \in \mathbb{T}^{2k_1}$, $z_2 \in \mathbb{T}^{2k_2}$. Note that by construction, f coincides with $g \circ \phi^{-1}$ on the coisotropic submanifold C . The situation is summarized in the following diagram:



Our proof will be based on the use of Lagrangian spectral invariants applied to graphs of symplectic maps. Given a Hamiltonian function H on a standard symplectic torus \mathbb{T}^{2n} , the graph of its time-1 map ϕ_H^1 is a Lagrangian submanifold of $\overline{\mathbb{T}^{2n}} \times \mathbb{T}^{2n}$. This graph is the image of the diagonal by the time-1 map of the Hamiltonian function $0 \oplus H$ on $\overline{\mathbb{T}^{2n}} \times \mathbb{T}^{2n}$ given by $(0 \oplus H)_t(q, p; Q, P) = H_t(Q, P)$. It will be convenient for us to see these Lagrangians as deformations of a standard ‘‘coordinate’’ Lagrangian subtorus rather than as deformations of the diagonal in $\overline{\mathbb{T}^{2n}} \times \mathbb{T}^{2n}$. Therefore we introduce the following two symplectic identifications:

$$\begin{aligned}
 \Psi : \overline{\mathbb{T}^{2k_1+2k_2}} \times \mathbb{T}^{2k_1+2k_2} &\rightarrow \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2} \times \mathbb{T}^{2k_2}, \\
 (q_1, p_1, q_2, p_2; Q_1, P_1, Q_2, P_2) &\mapsto \\
 &(q_1, P_1 - p_1; P_1, q_1 - Q_1; q_2, P_2 - p_2; P_2, q_2 - Q_2), \\
 \text{and } \Psi_R : \overline{\mathbb{T}^{2k_1}} \times \mathbb{T}^{2k_1} &\rightarrow \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1}, \\
 (q_1, p_1; Q_1, P_1) &\mapsto (q_1, P_1 - p_1; P_1, q_1 - Q_1).
 \end{aligned}$$

We see that Ψ sends the diagonal of $\overline{\mathbb{T}^{2k_1+2k_2}} \times \mathbb{T}^{2k_1+2k_2}$ to the Lagrangian subtorus

$$L_0 = \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\},$$

already introduced in Section 1.2, and Ψ_R sends the diagonal of $\overline{\mathbb{T}^{2k_1}} \times \mathbb{T}^{2k_1}$ to the Lagrangian subtorus

$$L_R = \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_1} \times \{0\}.$$

The proof of Theorem 4 will consist of a series of equalities and inequalities between spectral invariants. These identities are organized in four claims that we now give.

The first claim follows immediately from Proposition 13 and the naturality of Lagrangian spectral invariants (11). For example, (31) below is due to the fact that

$$c_+(f_R) = \ell([\Delta]; \Delta, \Delta; 0 \oplus f_R) = \ell([L_R]; L_R, L_R; (0 \oplus f_R) \circ \Psi_R^{-1})$$

with Δ the diagonal of $\overline{\mathbb{T}^{2k_1}} \times \mathbb{T}^{2k_1}$.

CLAIM. – *We have*

$$(31) \quad c_+(f_R) = \ell([L_R]; L_R, L_R; (0 \oplus f_R) \circ \Psi_R^{-1}),$$

$$(32) \quad c_+(g_R) = \ell([L_R]; L_R, L_R; (0 \oplus g_R) \circ \Psi_R^{-1}),$$

$$(33) \quad c_+(g) = \ell([L_0]; L_0, L_0; (0 \oplus g) \circ \Psi^{-1}),$$

$$(34) \quad c_+(g \circ \phi^{-1}) = \ell([L_0]; L_0, L_0; (0 \oplus g \circ \phi^{-1}) \circ \Psi^{-1}).$$

The next statement gives a relation between the spectral invariants of Lagrangians and functions defined on different spaces. This will be based on the splitting formula (21). As in Section 1.2, we denote

$$L_1 = \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_1} \times \{0\} \times \mathbb{T}^{k_2} \times \{0\} \times \{0\} \times \mathbb{T}^{k_2}.$$

We also recall that L_0 and L_1 split in the following form (see Section 1.2):

$$L_0 = L \times L'_0 \quad \text{and} \quad L_1 = L \times L'_1.$$

CLAIM. – *We have*

$$(35) \quad \ell([L_R]; L_R, L_R; (0 \oplus f_R) \circ \Psi_R^{-1}) = \ell([L]; L_0, L_1; (0 \oplus f) \circ \Psi^{-1}),$$

$$(36) \quad \ell([L_R]; L_R, L_R; (0 \oplus g_R) \circ \Psi_R^{-1}) = \ell([L_0]; L_0, L_0; (0 \oplus g) \circ \Psi^{-1}).$$

Proof. – The Lagrangians L_0 and L_1 both contain L_R and moreover can be decomposed in the form

$$L_i = \underbrace{L_R \times \Lambda}_{= L \subset \mathbb{T}^{2k_2} \times \mathbb{T}^{2k_2} \times \mathbb{T}^{2k_1}} \times \underbrace{L'_i}_{\subset \mathbb{T}^{2k_2}} \quad \text{for } i = 0 \text{ and } 1$$

where $\Lambda = \mathbb{T}^{k_2} \times \{0\} \subset \mathbb{T}^{2k_2}$. Then, if we denote $F = (0 \oplus f) \circ \Psi^{-1}$ and $F_R = (0 \oplus f_R) \circ \Psi_R^{-1}$, we see that we can decompose F according to this splitting: $F = F_R \oplus 0 \oplus 0$, where both 0's are seen as functions on \mathbb{T}^{2k_2} . By Theorem 24,

$$\ell([L]; L_0, L_1; F) = \ell([L]; L, L; F_R \oplus 0) + \ell([pt]; L'_0, L'_1; 0).$$

The second term on the right hand side vanishes. We may then apply the splitting formula (24):

$$\begin{aligned} \ell([L]; L_0, L_1; F) &= \ell([L]; L, L; F_R \oplus 0) \\ &= \ell([L_R]; L_R, L_R; F_R) + \ell([\Lambda]; \Lambda, \Lambda; 0), \end{aligned}$$

where again the second term vanishes. This proves Equation (35).

To prove Equation (36), one only needs to replace the pair (L_0, L_1) by (L_0, L_0) , the pair (L'_0, L'_1) by (L'_0, L'_0) , the function f by g and the function f_R by g_R , and repeat the same argument. \square

We will also need the following equality which is essentially a manifestation of the fact that at any fixed time, the functions f and $g \circ \phi^{-1}$ are constant on the leaves of the coisotropic submanifold C and coincide on it.

CLAIM. – *We have*

$$(37) \quad \ell([L]; L_0, L_1; (0 \oplus f) \circ \Psi^{-1}) = \ell([L]; L_0, L_1; (0 \oplus g \circ \phi^{-1}) \circ \Psi^{-1}).$$

Proof. – Denote $F = (0 \oplus f) \circ \Psi^{-1}$ and $G = (0 \oplus g \circ \phi^{-1}) \circ \Psi^{-1}$. Since f and $g \circ \phi^{-1}$ coincide on C , the continuous functions F and G coincide on the coisotropic submanifold

$$W = \Psi(C \times C) \subset \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2} \times \mathbb{T}^{2k_2}.$$

Observe that the Lagrangian L_1 is contained in W .

Since f and g are the respective lifts of f_R and g_R , their restriction to each leaf of C only depends on the time variable. Since ϕ^{-1} preserves the characteristic foliation of C , the function $g \circ \phi^{-1}$ is also a function of time on each leaf of C . From this we deduce that F and G are functions of time on each characteristic leaf of W .

Now let $(F_k)_{k \in \mathbb{N}}, (G_k)_{k \in \mathbb{N}}$ be sequences of smooth Hamiltonians which uniformly converge to F and G , respectively, with the additional property that for all $k \in \mathbb{N}$, F_k and G_k are functions of time on each leaf of W and $F_k - G_k = 0$ on W . Such sequences can be constructed as follows. Let F'_k, G'_k be two sequences of Hamiltonians that converge uniformly to F, G . Note that the restrictions of F and G coincide on W and are functions of time on each leaf hence admit the same reduced function H . Let H_k be a sequence of Hamiltonians on the reduced space of W which converges to H uniformly. Each function H_k can be lifted to a function H'_k defined on W . By construction the functions $F'_k - H'_k$ and $G'_k - H'_k$ converge to 0 on W . Denote by F''_k and G''_k their trivial extensions to $\mathbb{T}^{2k_1} \times \mathbb{T}^{2k_1} \times \mathbb{T}^{2k_2} \times \mathbb{T}^{2k_2}$, which also converge to 0. The functions $F_k = F'_k - F''_k$ and $G_k = G'_k - G''_k$ suit our needs: They converge respectively to F and G and they both coincide with the leafwise function H'_k on W .

We will next show that $\ell([L]; L_0, L_1; F_k) = \ell([L]; L_0, L_1; G_k)$ for all k . The claim would then follow by taking the limit of both sides as $k \rightarrow \infty$. Fix k and let $H_r = rG_k + (1 - r)F_k$ where $r \in [0, 1]$. We will in fact prove the stronger statement that $\ell([L]; L_0, L_1; H_r)$ is a constant function of the variable r .

For any $r, r' \in [0, 1]$ the Hamiltonians H_r and $H_{r'}$ are functions of time on each leaf of W and $H_r = H_{r'}$ on W . This is because the same statement is true for F_k and G_k . It is not hard to check that this implies that for any point $p \in W$ and any $t \in [0, 1]$ we have:

$$(38) \quad \phi_{H_r}^t(p) \text{ and } \phi_{H_{r'}}^t(p) \text{ belong to the same characteristic leaf of } W.$$

Now, consider a critical point of the action functional $\mathcal{O}_{H_r}^{L_0, L_1}$: It is a Hamiltonian chord $\phi_{H_r}^t(p)$ where $p \in L_0$ and $\phi_{H_r}^1(p) \in L_1$. The Hamiltonian H_r is a function of time on characteristic leaves and so its flow $\phi_{H_r}^t$ preserves W . Since $\phi_{H_r}^1(p) \in L_1 \subset W$, we conclude that $\phi_{H_r}^t(p) \in W$ for all $t \in [0, 1]$. Using (38), we see that $\phi_{H_{r'}}^t(p) \in W$ for any $t, r' \in [0, 1]$. Furthermore, (38) implies that $\phi_{H_{r'}}^1(p) \in L_1$: This is because the Lagrangian $L_1 \subset W$ and hence any characteristic leaf of W which intersects L_1 is entirely contained in L_1 . We conclude from the above that $t \mapsto \phi_{H_{r'}}^t(p)$ is a critical point of the action functional $\mathcal{O}_{H_{r'}}^{L_0, L_1}$ and so there exists a bijection between the critical points of the two action functionals.

Next, we will show that the two chords $\phi_{H_r}^t(p)$ and $\phi_{H_{r'}}^t(p)$ have the same action. Since H_r and $H_{r'}$ coincide on the leaves of W we see, using (38), that $H_r(\phi_{H_r}^t(p)) = H_{r'}(\phi_{H_{r'}}^t(p))$. Hence, to show that the two chords have the same action we must prove that any two cappings u_r of $\phi_{H_r}^t(p)$ and $u_{r'}$ of $\phi_{H_{r'}}^t(p)$ have the same symplectic area. We will prove this using (38) as well. Suppose that $r < r'$. Fix any choice of u_r and define $u_{r'} = u_r \# v$, where $v : [r, r'] \times [0, 1] \rightarrow \mathbb{T}^{2n} \times \mathbb{T}^{2n}$ is defined by: $v(s, t) = \phi_{H_s}^t(p)$. We must show that the symplectic area of v is zero: Note that (38) implies that for any fixed t the path $s \mapsto \phi_{H_s}^t(p)$ is contained in the same characteristic leaf of W . Therefore, $\frac{\partial v}{\partial s}$ is always tangent to the characteristic leaves of W . This combined with the fact that the image of v is contained in W yields that $\omega_{\text{std}}(\frac{\partial v}{\partial s}, \frac{\partial v}{\partial t}) = 0$. Hence, v has zero symplectic area and the symplectic area of u_r coincides with that of $u_{r'}$.

We conclude from the previous two paragraphs that $\text{Spec}(H_r) = \text{Spec}(H_{r'})$ for any $r, r' \in [0, 1]$. Recall that the spectrum of any Hamiltonian has measure zero. We see that $r \mapsto \ell([L]; L_0, L_1; H_r)$ is a continuous function taking values in a measure zero set and thus it must be constant. This finishes the proof. \square

Finally, the following claim is a direct application of Proposition 6.

CLAIM. –

$$(39) \quad \ell([L]; L_0, L_1; (0 \oplus g \circ \phi^{-1}) \circ \Psi^{-1}) \leq \ell([L_0]; L_0, L_0; (0 \oplus g \circ \phi^{-1}) \circ \Psi^{-1}).$$

End of the proof of Theorem 4. We now gather the identities collected in the above claims. Using the fact that ϕ^{-1} preserves c_+ , we obtain:

$$\begin{aligned} c_+(f_R) &\stackrel{(31)}{=} \ell([L_R]; L_R, L_R; (0 \oplus f_R) \circ \Psi_R^{-1}) \\ &\stackrel{(35)}{=} \ell([L]; L_0, L_1; (0 \oplus f) \circ \Psi^{-1}) \\ &\stackrel{(37)}{=} \ell([L]; L_0, L_1; (0 \oplus g \circ \phi^{-1}) \circ \Psi^{-1}) \\ &\stackrel{(39)}{\leq} \ell([L_0]; L_0, L_0; (0 \oplus g \circ \phi^{-1}) \circ \Psi^{-1}) \\ &\stackrel{(34)}{=} c_+(g \circ \phi^{-1}) \\ &= c_+(g) \\ &\stackrel{(33)}{=} \ell([L_0]; L_0, L_0; (0 \oplus g) \circ \Psi^{-1}) \\ &\stackrel{(36)}{=} \ell([L_R]; L_R, L_R; (0 \oplus g_R) \circ \Psi_R^{-1}) \\ &\stackrel{(32)}{=} c_+(g_R). \end{aligned}$$

Switching the roles played by f_R and g_R yields the reverse inequality $c_+(g_R) \leq c_+(f_R)$. Hence, $c_+(f_R) = c_+(g_R)$.

BIBLIOGRAPHY

- [1] A. ABBONDANDOLO, M. SCHWARZ, Floer homology of cotangent bundles and the loop product, *Geom. Topol.* **14** (2010), 1569–1722.

- [2] P. ALBERS, A Lagrangian Piunikhin-Salamon-Schwarz morphism and two comparison homomorphisms in Floer homology, *Int. Math. Res. Not.* **2008** (2008), Art. ID rnm134, 56.
- [3] D. AUROUX, A beginner's introduction to Fukaya categories, in *Contact and symplectic topology* (F. Bourgeois et al., eds.), Bolyai Soc. Math. Stud. **26**, 2014, 85–136.
- [4] J.-F. BARRAUD, O. CORNEA, Homotopic dynamics in symplectic topology, in *Morse theoretic methods in nonlinear analysis and in symplectic topology*, NATO Sci. Ser. II Math. Phys. Chem. **217**, Springer, Dordrecht, 2006, 109–148.
- [5] L. BUHOVSKY, E. OPSHTEIN, Some quantitative results in C^0 symplectic geometry, preprint arXiv:1404.0875, to appear in *Invent. math.*
- [6] M. ENTOV, L. POLTEROVICH, Rigid subsets of symplectic manifolds, *Compos. Math.* **145** (2009), 773–826.
- [7] S. HU, F. LALONDE, R. LECLERCQ, Homological Lagrangian monodromy, *Geom. Topol.* **15** (2011), 1617–1650.
- [8] V. HUMILIÈRE, On some completions of the space of Hamiltonian maps, *Bull. Soc. Math. France* **136** (2008), 373–404.
- [9] V. HUMILIÈRE, R. LECLERCQ, S. SEYFADDINI, Coisotropic rigidity and C^0 -symplectic geometry, *Duke Math. J.* **164** (2015), 767–799.
- [10] J. KATIĆ, D. MILINKOVIĆ, Piunikhin-Salamon-Schwarz isomorphisms for Lagrangian intersections, *Differential Geom. Appl.* **22** (2005), 215–227.
- [11] F. LAUDENBACH, J.-C. SIKORAV, Hamiltonian disjunction and limits of Lagrangian submanifolds, *Int. Math. Res. Not.* **1994** (1994), 161–168.
- [12] R. LECLERCQ, Spectral invariants in Lagrangian Floer theory, *J. Mod. Dyn.* **2** (2008), 249–286.
- [13] R. LECLERCQ, The Seidel morphism of Cartesian products, *Algebr. Geom. Topol.* **9** (2009), 1951–1969.
- [14] A. MONZNER, N. VICHERY, F. ZAPOLSKY, Partial quasimorphisms and quasistates on cotangent bundles, and symplectic homogenization, *J. Mod. Dyn.* **6** (2012), 205–249.
- [15] Y.-G. OH, Symplectic topology as the geometry of action functional. I. Relative Floer theory on the cotangent bundle, *J. Differential Geom.* **46** (1997), 499–577.
- [16] Y.-G. OH, S. MÜLLER, The group of Hamiltonian homeomorphisms and C^0 -symplectic topology, *J. Symplectic Geom.* **5** (2007), 167–219.
- [17] E. OPSHTEIN, C^0 -rigidity of characteristics in symplectic geometry, *Ann. Sci. Éc. Norm. Supér.* **42** (2009), 857–864.
- [18] S. PIUNIKHIN, D. SALAMON, M. SCHWARZ, Symplectic Floer-Donaldson theory and quantum cohomology, in *Contact and symplectic geometry (Cambridge, 1994)*, Publ. Newton Inst. **8**, Cambridge Univ. Press, 1996, 171–200.
- [19] J. M. SABLOFF, L. TRAYNOR, Obstructions to the existence and squeezing of Lagrangian cobordisms, *J. Topol. Anal.* **2** (2010), 203–232.
- [20] M. SCHWARZ, On the action spectrum for closed symplectically aspherical manifolds, *Pacific J. Math.* **193** (2000), 419–461.
- [21] D. THÉRET, A lagrangian camel, *Comment. Math. Helv.* **74** (1999), 591–614.

- [22] C. VITERBO, Symplectic topology as the geometry of generating functions, *Math. Annalen* **292** (1992), 685–710.
- [23] F. ZAPOLSKY, On the Hofer geometry for weakly exact Lagrangian submanifolds, *J. Symplectic Geom.* **11** (2013), 475–488.

(Manuscrit reçu le 25 juillet 2014 ;
accepté, après révision, le 23 mars 2015.)

Vincent HUMILIÈRE
Sorbonne Universités UPMC Paris 06
Institut de Mathématiques de Jussieu-Paris Rive Gauche
Univ. Paris Diderot, Sorbonne Paris Cité
UMR CNRS 7586
4 place Jussieu
75005 Paris, France
E-mail: vincent.humiliere@imj-prg.fr

Rémi LECLERCQ
Laboratoire de Mathématiques d'Orsay
Université Paris-Sud & CNRS
Université Paris-Saclay
Bâtiment 425
91405 Orsay, France
E-mail: remi.leclercq@math.u-psud.fr

Sobhan SEYFADDINI
Department of Mathematics
Massachusetts Institute of Technology
77 Massachusetts Avenue
Cambridge, MA 02139, USA
E-mail: sobhan@mit.edu