

quatrième série - tome 50 fascicule 4 juillet-août 2017

*ANNALES
SCIENTIFIQUES
de
L'ÉCOLE
NORMALE
SUPÉRIEURE*

Adrián GONZÁLEZ-PÉREZ & Marius JUNGE & Javier PARCET

Smooth Fourier multipliers in group algebras via Sobolev dimension

SOCIÉTÉ MATHÉMATIQUE DE FRANCE

Annales Scientifiques de l'École Normale Supérieure

Publiées avec le concours du Centre National de la Recherche Scientifique

Responsable du comité de rédaction / *Editor-in-chief*

Emmanuel KOWALSKI

Publication fondée en 1864 par Louis Pasteur

Continuée de 1872 à 1882 par H. SAINTE-CLAIRE DEVILLE
de 1883 à 1888 par H. DEBRAY
de 1889 à 1900 par C. HERMITE
de 1901 à 1917 par G. DARBOUX
de 1918 à 1941 par É. PICARD
de 1942 à 1967 par P. MONTEL

Comité de rédaction au 1^{er} janvier 2017

P. BERNARD A. NEVES
S. BOUCKSOM J. SZEFTEL
E. BREUILLARD S. VŨ NGỌC
R. CERF A. WIENHARD
G. CHENEVIER G. WILLIAMSON
E. KOWALSKI

Rédaction / *Editor*

Annales Scientifiques de l'École Normale Supérieure,
45, rue d'Ulm, 75230 Paris Cedex 05, France.
Tél. : (33) 1 44 32 20 88. Fax : (33) 1 44 32 20 80.
annales@ens.fr

Édition / *Publication*

Société Mathématique de France
Institut Henri Poincaré
11, rue Pierre et Marie Curie
75231 Paris Cedex 05
Tél. : (33) 01 44 27 67 99
Fax : (33) 01 40 46 90 96

Abonnements / *Subscriptions*

Maison de la SMF
Case 916 - Luminy
13288 Marseille Cedex 09
Fax : (33) 04 91 41 17 51
email : smf@smf.univ-mrs.fr

Tarifs

Europe : 519 €. Hors Europe : 548 €. Vente au numéro : 77 €.

© 2017 Société Mathématique de France, Paris

En application de la loi du 1^{er} juillet 1992, il est interdit de reproduire, même partiellement, la présente publication sans l'autorisation de l'éditeur ou du Centre français d'exploitation du droit de copie (20, rue des Grands-Augustins, 75006 Paris).

All rights reserved. No part of this publication may be translated, reproduced, stored in a retrieval system or transmitted in any form or by any other means, electronic, mechanical, photocopying, recording or otherwise, without prior permission of the publisher.

ISSN 0012-9593

Directeur de la publication : Stéphane Seuret
Périodicité : 6 n^{os} / an

SMOOTH FOURIER MULTIPLIERS IN GROUP ALGEBRAS VIA SOBOLEV DIMENSION

BY ADRIÁN GONZÁLEZ-PÉREZ, MARIUS JUNGE
AND JAVIER PARCET

ABSTRACT. – We investigate Fourier multipliers with smooth symbols defined over locally compact Hausdorff groups. Our main results in this paper establish new Hörmander-Mikhlin criteria for spectral and non-spectral multipliers. The key novelties which shape our approach are three. First, we control a broad class of Fourier multipliers by certain maximal operators in noncommutative L_p spaces. This general principle—exploited in Euclidean harmonic analysis during the last 40 years—is of independent interest and might admit further applications. Second, we replace the formerly used cocycle dimension by the Sobolev dimension. This is based on a noncommutative form of the Sobolev embedding theory for Markov semigroups initiated by Varopoulos, and yields more flexibility to measure the smoothness of the symbol. Third, we introduce a dual notion of polynomial growth to further exploit our maximal principle for non-spectral Fourier multipliers. The combination of these ingredients yields new L_p estimates for smooth Fourier multipliers in group algebras.

RÉSUMÉ. – Nous étudions des multiplicateurs de Fourier à symboles réguliers sur des groupes localement compacts. De nouveaux critères de Hörmander-Mikhlin pour des multiplicateurs spectraux et non spectraux sont établis. Notre approche se base sur trois nouveaux résultats clés. Premièrement, nous utilisons certains opérateurs maximaux dans des espaces L_p non commutatifs pour obtenir un contrôle sur de larges classes de multiplicateurs. Ce principe général — exploité en analyse harmonique euclidienne ces 40 dernières années — présente un intérêt indépendant et pourrait admettre de nouvelles applications. Deuxièmement, en établissant une version non commutative de la théorie de plongement de Sobolev pour les semigroupes de Markov initiée par Varopoulos, la dimension de cocycle utilisée auparavant est remplacée par la dimension de Sobolev. Ceci permet plus de flexibilité sur la régularité du symbole. Enfin, nous introduisons une notion duale de la croissance polynomiale pour exploiter davantage notre principe du maximum sur des multiplicateurs de Fourier non spectraux. La combinaison de ces ingrédients produit de nouvelles estimations L_p pour des multiplicateurs de Fourier réguliers dans des algèbres de groupe.

Introduction

The aim of this paper is to study Fourier multipliers on group von Neumann algebras for locally compact Hausdorff groups. More precisely, the relation between the smoothness of

their symbols and L_p -boundedness. This is a central topic in Euclidean harmonic analysis. In the context of nilpotent groups, it has also been intensively studied in the works of Cowling, Müller, Ricci, Stein and others. In this paper we will consider the dual problem, placing our nonabelian groups in the frequency side. Today it is well understood that the dual of a nonabelian group can only be described as a quantum group, its underlying algebra being the group von Neumann algebra. The interest of Fourier multipliers over such group algebras was recognized early in the pioneering work of Haagerup [11], as well as in the research carried out thereafter. It was made clear how Fourier multipliers on these algebras can help in their classification, through the use of certain approximation properties which become invariants of the algebra. Unfortunately, the literature on this topic does not involve the L_p -theory—with a few exemptions like [21] and the very recent paper of Lafforgue and de la Salle [25]—as it is mandatory from a harmonic analysis viewpoint. In this respect, our work is a continuation of [18, 19] where 1-cocycles into finite-dimensional Hilbert spaces were used to lift multipliers from the group into a more Euclidean space. This yields Hörmander-Mikhlin type results depending of the dimension of the Hilbert space involved. Here, we shall follow a different approach through the introduction of new notions of dimension allowing more room for the admissible class of multipliers. These notions rely on noncommutative forms of the Sobolev embedding theory for Markov semigroups, which carry an ‘encoded geometry’ in the commutative setting. Prior to that, we need to investigate new maximal bounds whose Euclidean analogs are central in harmonic analysis. In this paper we shall limit ourselves to unimodular groups to avoid technical issues concerning modular theory.

This text is divided into three blocks which are respectively devoted to maximal bounds, Sobolev dimension and polynomial co-growth. Let us first put in context our maximal estimates for Fourier multipliers. Given a symbol $m : \mathbb{R}^n \rightarrow \mathbb{C}$ with corresponding Fourier multiplier T_m , there is a long tradition in identifying maximal operators \mathcal{M} which satisfy the weighted L_2 -norm inequality below for all admissible input functions f and weights w

$$(WL_2) \quad \int_{\mathbb{R}^n} |T_m f|^2 w \lesssim \int_{\mathbb{R}^n} |f|^2 \mathcal{M} w.$$

It goes back to the work of Córdoba and Fefferman in the 70’s. This general principle has deep connections with Bochner-Riesz multipliers and also with A_p weight theory. The Introduction of [2] gives a very nice historical summary and new results in this direction. The main purpose of this estimate is that elementary duality arguments yield for $p > 2$ that

$$\|T_m : L_p(\mathbb{R}^n) \rightarrow L_p(\mathbb{R}^n)\| \lesssim \|\mathcal{M} : L_{(p/2)'}(\mathbb{R}^n) \rightarrow L_{(p/2)'}(\mathbb{R}^n)\|^{1/2}.$$

The most general noncommutative form of this inequality would require too much terminology for this Introduction. Instead, let us just introduce the basic concepts to give a reasonable but weaker statement. Stronger results will be given in the body of the paper. Let G be a locally compact Hausdorff group. If we write μ for the left Haar measure of G and λ for the left regular representation $\lambda : G \rightarrow \mathcal{B}(L_2 G)$, the group von Neumann algebra $\mathcal{L}G$ is the weak operator closure in $\mathcal{B}(L_2 G)$ of $\lambda(L_1(G))$. We refer to Section 1 for a construction of the Plancherel weight τ on $\mathcal{L}G$, a noncommutative substitute of the Haar measure. Note that τ is tracial iff G is unimodular—which we assume—and it coincides with the finite trace given by $\tau(x) = \langle \delta_e, x \delta_e \rangle$ when G is discrete. In the unimodular case, $(\mathcal{L}G, \tau)$ is a semifinite von Neumann algebra with a trace and it is easier to construct the noncommutative

L_p -spaces $L_p(\mathcal{L}G, \tau)$ with norm $\|x\|_p = \tau(|x|^p)^{1/p}$, where $|x|^p = (x^*x)^{p/2}$ by functional calculus on the (unbounded) operator x^*x . Given a bounded symbol $m : G \rightarrow \mathbb{C}$, the corresponding Fourier multiplier is densely defined by $T_m\lambda(f) = \lambda(mf)$. Alternatively, it will be useful to understand these operators as convolution maps in the following way

$$T_m(x) = \lambda(m) \star x = (\tau \otimes \text{Id})(\delta\lambda(m) (\sigma x \otimes \mathbf{1})),$$

where $\delta : \mathcal{L}G \rightarrow \mathcal{L}G \overline{\otimes} \mathcal{L}G$ is determined by $\delta(\lambda_g) = \lambda_g \otimes \lambda_g$ and $\sigma : \mathcal{L}G \rightarrow \mathcal{L}G$ is the anti-automorphism given by linear extension of $\sigma(\lambda_g) = \lambda_{g^{-1}}$. The first map is called the comultiplication map for $\mathcal{L}G$, whereas σ is the corresponding coinvolution. Our next ingredient is the L_p -norm of maximal operators. Given a family of noncommuting operators $(x_\omega)_\omega$ affiliated to a semifinite von Neumann algebra \mathcal{M} , their supremum is not well-defined. We may however consider their L_p -norms through

$$\left\| \sup_{\omega \in \Omega}^+ x_\omega \right\|_{L_p(\mathcal{M})} = \left\| (x_\omega)_{\omega \in \Omega} \right\|_{L_p(\mathcal{M}; L_\infty(\Omega))},$$

where the mixed-norm $L_p(L_\infty)$ -space has a nontrivial definition obtained by Pisier for hyperfinite \mathcal{M} in [31] and later generalized in [15, 20]. This definition recovers the norm in $L_p(\Sigma; L_\infty(\Omega))$ for abelian $\mathcal{M} = L_\infty(\Sigma)$, further details in Section 1. Finally, conditionally negative lengths $\psi : G \rightarrow \mathbb{R}_+$ are symmetric functions vanishing at the identity e which satisfy $\sum_{g,h} \overline{a_g} a_h \psi(g^{-1}h) \leq 0$ for any family of coefficients with $\sum_g a_g = 0$. Due to its one-to-one relation to Markov convolution semigroups, they will play a crucial role in this paper. In the classical multiplier theorems, the symbols m are cut out with functions $\eta(|\xi|)$ for some compactly supported $\eta \in C^\infty(\mathbb{R}_+)$. Our techniques do not allow us to use compactly supported functions in \mathbb{R}_+ . Instead, we are going to use analytic functions decaying fast near 0 and near ∞ . We will call such η an \mathcal{H}_0^∞ -cut-off, see Section 1 for the precise definitions. The archetype of such functions will be $\eta(z) = z e^{-z}$. When such function is real-valued for every $z \in \mathbb{R}_+$ we will say that η is real. We will denote by $L_p^\circ(\mathcal{L}G)$ the noncommutative L_p -space $L_p(\mathcal{L}G)$ modulo the functions supported in $G_0 = \{g \in G : \psi(g) = 0\}$, see Section 1.1.7 for details.

THEOREM A. – *Let G be a unimodular group equipped with any conditionally negative length $\psi : G \rightarrow \mathbb{R}_+$. Let η be a real \mathcal{H}_0^∞ -cut-off and $m : G \rightarrow \mathbb{R}$ an essentially bounded symbol satisfying that $m(g^{-1}) = m(g)$ for every $g \in G$. Assume $B_t = \lambda(m\eta(t\psi))$ admits a decomposition $B_t = \Sigma_t M_t$ with M_t positive and satisfying $M_t = \sigma M_t$, and consider the convolution map $\mathcal{R}(x) = (|M_t|^2 \star x)_{t \geq 0}$. Then the following inequality holds for $2 < p < \infty$*

$$\|T_m\|_{\mathcal{B}(L_p^\circ(\mathcal{L}G))} \lesssim_{(p)} \left(\sup_{t \geq 0} \|\Sigma_t\|_2 \right) \left\| \mathcal{R} : L_{(p/2)'}(\mathcal{L}G) \rightarrow L_{(p/2)'}(\mathcal{L}G; L_\infty) \right\|^{\frac{1}{2}},$$

By duality, a similar statement holds for $1 < p < 2$. Moreover, the assumptions of η being real, of m satisfying that $m(g^{-1}) = m(g)$ and of M_t being positive and satisfying that $M_t = \sigma M_t$ can be removed if we allow slight modifications in the statement of the theorem, like framing the conclusion in terms of noncommutative Hardy spaces. Theorem A combines in a very neat way noncommutative generalizations of (WL_2) with square function estimates. In the particular case of Hörmander-Mikhlin symbols—as we shall see along this paper—the decomposition splits the assumptions. Namely, the L_2 -norm of Σ_t is bounded using the smoothness condition while the maximal \mathcal{R} is bounded through the geometrical

assumptions regarding the dimensional behavior of ψ . Apart from the direct consequences given in the present paper, this result is of independent interest and admits potential applications in other directions to be explored in a forthcoming publication.

Given a conditionally negative length $\psi : G \rightarrow \mathbb{R}_+$, the infinitesimal generator of the semigroup $\lambda_g \mapsto \exp(-t\psi(g))\lambda_g$ is the map determined by $A(\lambda_g) = \psi(g)\lambda_g$. In particular, ψ -radial Fourier multipliers fall in the category of spectral operators of the form $m(A)$. These maps are known as spectral multipliers and play a central role in the theory. Our aim in this second block is to find smoothness criteria on m which implies L_p -boundedness of the spectral multiplier $T_{m \circ \psi}$.

It is well understood—specially after [6, 38]—that if we want to obtain L_p boundedness for $m(A)$ from the smoothness of m , for every semigroup, we need to impose analyticity on m . To obtain a smoothness condition with a finite number of derivatives our space needs to be finite-dimensional. Indeed, it is known that the optimal smoothness order may growth with the dimension. This indicates the necessity of defining a notion of dimension in the non-commutative setting. We will take as dimension the value $d > 0$ for which a Sobolev type embedding holds for A . Recall that there is a Sobolev embedding theory for Markov semigroups introduced by Varopoulos [42]. More precisely, given a measure space (Ω, μ) and certain elliptic operator A generating the Markov process $S_t = \exp(-tA)$, one can introduce the Sobolev dimension d for which the equivalence below holds

$$\|f\|_{L^{\frac{2d}{d-2}}(\Omega)} \lesssim \|A^{\frac{1}{2}}f\|_{L_2(\Omega)} \iff \|S_t f\|_{L_\infty(\Omega)} \lesssim t^{-\frac{d}{2}} \|f\|_{L_1(\Omega)}.$$

The property of the right hand side is known as ultracontractivity. When it holds for the semigroup generated by an invariant Laplacian on a Lie group, it forces $\mu(B_t(e)) \sim t^d$. Thus, we can understand ultracontractivity as a way of describing the growth of balls. With that motivation we introduce general ultracontractivity properties

$$\|S_t : L_1(\mathcal{M}) \rightarrow \mathcal{M}\|_{\text{cb}} \lesssim \frac{1}{\Phi(\sqrt{t})},$$

where cb stands for completely bounded. The function Φ will measure the “growth of the balls”. Since doubling measure spaces are widely recognized as a natural setting for performing harmonic analysis, we will impose Φ to be doubling, i.e.,:

$$\sup_{t>0} \left\{ \frac{\Phi(2t)}{\Phi(t)} \right\} < \infty,$$

and our doubling dimension will be given by

$$D_\Phi = \log_2 \sup_{t>0} \left\{ \frac{\Phi(2t)}{\Phi(t)} \right\}.$$

In the classical abelian setting, apart from the ultracontractivity—or on-diagonal behavior of S_t —we need to impose off-diagonal decay on S_t , typically Gaussian bounds. Let (G, ψ, X) be a triple formed by a locally compact Hausdorff unimodular group G , a conditionally negative length $\psi : G \rightarrow \mathbb{R}_+$ and an element X in the extended positive cone $\mathcal{L}G_+^\wedge$, see [12, 13] for precise definitions. We will say that the triple satisfies the *standard assumptions* when:

i) *Doublingness*

$\Phi_X(s) = \tau(\chi_{[0,s]}(X))$ is doubling.

ii) *L_2 Gaussian upper bounds*

$$\tau\left(\chi_{[r,\infty)}(X)|\lambda(e^{-s\psi})|^2\right) \lesssim \frac{e^{-\beta \frac{r^2}{s}}}{\Phi_X(\sqrt{s})} \quad \text{for some } \beta > 0.$$

iii) *Hardy-Littlewood maximality*

$$\left\| \sup_{s \geq 0} \frac{\chi_{[0,s]}(X)}{\Phi_X(s)} \star x \right\|_{L_p(\mathcal{L}G)} \lesssim_{(p)} \|x\|_{L_p(\mathcal{L}G)} \quad \text{for every } 1 < p < \infty.$$

We will also require the inequality iii) to hold uniformly for matrix amplifications. As we shall see, inequality ii) implies ultracontractivity with Φ_X as growth function. We will omit the dependency of X from Φ_X when it can be understood from the context. It is also interesting to point out that, in the classical case, Gaussian bounds can be deduced from the ultracontractivity in the presence of geometrical assumptions like locality or finite speed of propagation for the wave equation, see [35, 36] and [34, Section 3]. Generalizing such results to the noncommutative setting will be the object of forthcoming research. The connection of standard assumptions with smooth ψ -radial Fourier multipliers is nearly optimal.

THEOREM B. – *Let (G, ψ, X) be any triple satisfying the standard assumptions which we considered above. Given an \mathcal{H}_0^∞ -cut-off function η and a symbol $m : \mathbb{R}_+ \rightarrow \mathbb{C}$, the following inequalities hold for $1 < p < \infty$:*

i) *If $s > (D_\Phi + 1)/2$*

$$\|T_{m \circ \psi}\|_{\mathcal{CB}(L_p^s(\mathcal{L}G))} \lesssim \sup_{t \geq 0} \|m(t) \eta(\cdot)\|_{W^{2,s}(\mathbb{R}_+)}.$$

ii) *If $s > D_\Phi/2$ and $\psi \in \text{CBPlan}_q^\Phi$ for some $q \geq 2$*

$$\|T_{m \circ \psi}\|_{\mathcal{CB}(L_p^s(\mathcal{L}G))} \lesssim \sup_{t \geq 0} \|m(t) \eta(\cdot)\|_{W^{q,s}(\mathbb{R}_+)}.$$

The last inequality holds with $q = \infty$ under the sole assumption of $s > D_\Phi/2$.

The term \mathcal{CB} also stands for “completely bounded” and the property CBPlan_q^Φ plays the role of the q -Plancherel property introduced by Duong-Ouhabaz-Sikora [41], see the body of the paper for concrete definitions. The proof of Theorem B is the most technical in this paper. It will explain the decoupling nature of Theorem A. The Σ_t are controlled using the Sobolev smoothness (via the Phragmen-Lindelöf theorem) for any degree $s > 0$, whereas the maximal bound determines optimal restrictions in terms of the Sobolev dimension D_Φ .

Theorem B should be illustrated with interesting examples. The existence of natural triples satisfying the standard assumptions for nonabelian groups is the subject of current research, which will appear elsewhere. In this paper we shall construct such triples out of finite-dimensional cocycles. This permits to recover the results in [18, 19] for ψ -radial multipliers. In fact, we should emphasize at this point that the notion of dimension in the previous approach was limited to the Hilbert space dimension of the cocycle determined by the length ψ . Working with finite-dimensional cocycles is an unfortunate limitation which we could remove for noncommutative Riesz transforms in [19]. Theorem B allows to go even further for smooth radial multipliers.

In our third and last block of this paper, we consider general (non-spectral) Fourier multipliers. Apart for the semigroup over $\mathcal{L}G$ generated by ψ we will endow G with two semigroups $\mathcal{S}_1/\mathcal{S}_2 : L_\infty(G) \rightarrow L_\infty(G)$ of left/right invariant operators. The intuition here is that \mathcal{S}_j will describe the geometry of G while the semigroup generated by ψ will describe the geometry of its dual. If A denotes the infinitesimal generator of a semigroup over $L_\infty(G)$, we use the standard notation for its nonhomogeneous Sobolev spaces

$$\|f\|_{W_A^{p,s}(G)} = \|(\mathbf{1} + A)^{\frac{s}{2}} f\|_{L_p(G)}.$$

When \mathcal{S} is left invariant there exists a positive densely defined operator \widehat{A} affiliated to $\mathcal{L}G$ such that $\lambda(Af) = \lambda(f)\widehat{A}$ for all $f \in \text{dom}_2(A)$. In a similar way we obtain $\lambda(Af) = \widehat{A}\lambda(f)$ when \mathcal{S} is right invariant, see Proposition 3.3 for the proof. Then we define the *polynomial co-growth of \widehat{A}* as follows

$$\text{cogrowth}(\widehat{A}) = \inf \left\{ r > 0 : (\mathbf{1} + \widehat{A})^{-\frac{r}{2}} \in L_1(\mathcal{L}G) \right\}.$$

Our choice for the term “polynomial co-growth” sits on the intuition that \widehat{A} behaves like $|\xi|^2$ in the case of the Laplacian Δ on \mathbb{R}^D and therefore $\text{cogrowth}(\widehat{\Delta}) = D$ follows from the fact that large balls grow like r^D . Further in Section 3 we will characterize polynomial co-growth by relating the behavior of small balls in G with “large balls” in $\mathcal{L}G$, see Remark 3.8 for further explanations. It is also worth mentioning the close relation between polynomial growth and Sobolev dimension as it will be analyzed in the body of the paper. Our main result in this direction is the following criterium for non-spectral multipliers.

THEOREM C. – *Let G be a unimodular group equipped with a conditionally negative length ψ . Let $\mathcal{S}_1/\mathcal{S}_2$ be respectively left/right invariant submarkovian semigroups on $L_\infty(G)$ whose generators A_j satisfy $\text{cogrowth}(\widehat{A}_j) = D_j$ for $j = 1, 2$. Consider an \mathcal{H}_0^∞ -cut-off function η and a symbol $m : G \rightarrow \mathbb{C}$. Then, if $s_j > D_j/2$ for $j = 1, 2$, the following inequality holds for $1 < p < \infty$*

$$\|T_m\|_{\mathcal{CB}(L_p^s(\mathcal{L}G))} \lesssim_{(p)} \sup_{t \geq 0} \max \left\{ \|\eta(t\psi) m\|_{W_{A_1}^{2,s_1}(G)}, \|\eta(t\psi) m\|_{W_{A_2}^{2,s_2}(G)} \right\}.$$

Theorem C establishes a link between the, a priori unrelated, geometries which determine ψ and \mathcal{S}_j . Indeed, we use the length ψ to cut m —determining the size of the support—and use A_j to measure the smoothness of m . It is interesting to note that passing to the dual requires a size condition on \widehat{A} , reinforcing the intuition that duality switches size and smoothness. The main difference with Theorem B is that in this general context we have been forced to place the dilation in the cut-off function η instead of the multiplier m . We conclude the paper illustrating Theorem C for Lie groups of polynomial growth by means of the subriemannian metrics determined by sublaplacians, see Corollary 3.9.

1. Maximal bounds

1.1. Preliminaries

Although the material here exposed is probably well-known to experts, let us review some notions and results in the interface between harmonic analysis and operator algebra that we will need throughout this section. We will start with a brief exposition of noncommutative

integration theory, including the construction of noncommutative L_p spaces. Our main example will be the group von Neumann algebra of a unimodular Lie group equipped with its canonical Plancherel trace. Then we will review some basics of operator space theory as well as the construction of certain mixed-norm spaces. Finally we will consider Markov semigroups with an special emphasis on semigroups of convolution type. We will revisit Hardy spaces and square function estimates associated with a semigroup.

1.1.1. *Noncommutative L_p spaces.* – Part of von Neumann algebra theory has evolved as the noncommutative form of measure theory and integration. A von Neumann algebra \mathcal{M} [24, 39, 40], is a unital weak-operator closed C^* -subalgebra of $\mathcal{B}(\mathcal{H})$, the algebra of bounded linear operators on a Hilbert space \mathcal{H} . We will write $\mathbf{1}_{\mathcal{M}}$, or simply $\mathbf{1}$, for the unit. The positive cone \mathcal{M}_+ is the set of positive operators in \mathcal{M} and a trace $\tau : \mathcal{M}_+ \rightarrow [0, \infty]$ is a linear map satisfying $\tau(x^*x) = \tau(xx^*)$. Such map is said to be normal if $\sup_{\alpha} \tau(x_{\alpha}) = \tau(\sup_{\alpha} x_{\alpha})$ for bounded increasing nets (x_{α}) ; it is semifinite if for $x \in \mathcal{M}_+ \setminus \{0\}$ there exists $0 < x' \leq x$ with $\tau(x') < \infty$; and it is faithful if $\tau(x) = 0$ implies $x = 0$. The trace τ plays the role of the integral in the classical case. A von Neumann algebra \mathcal{M} is semifinite when it admits a normal semifinite faithful (n.s.f. in short) trace τ . Any $x \in \mathcal{M}$ is a linear combination $x_1 - x_2 + ix_3 - ix_4$ of four positive operators. Thus, τ extends as an unbounded operator to nonpositive elements and the tracial property takes the familiar form $\tau(xy) = \tau(yx)$. The pairs (\mathcal{M}, τ) composed by a von Neumann algebra and a n.s.f. trace will be called *noncommutative measure spaces*. Note that commutative von Neumann algebras correspond to classical measurable spaces.

By the GNS construction, the noncommutative analog of measurable sets (characteristic functions) are orthogonal projections. Given $x \in \mathcal{M}_+$, its support is the least projection q in \mathcal{M} such that $qx = x = xq$ and is denoted by $\text{supp } x$. Let $\mathcal{S}_{\mathcal{M}}^+$ be the set of all $x \in \mathcal{M}_+$ such that $\tau(\text{supp } x) < \infty$ and set $\mathcal{S}_{\mathcal{M}}$ to be the linear span of $\mathcal{S}_{\mathcal{M}}^+$. If we write $|x| = \sqrt{x^*x}$, we can use the spectral measure dE of $|x|$ to observe that

$$x \in \mathcal{S}_{\mathcal{M}} \Rightarrow |x|^p = \int_{\mathbb{R}_+} s^p dE(s) \in \mathcal{S}_{\mathcal{M}}^+ \Rightarrow \tau(|x|^p) < \infty.$$

If we set $\|x\|_p = \tau(|x|^p)^{\frac{1}{p}}$, we obtain a norm in $\mathcal{S}_{\mathcal{M}}$ for $1 \leq p < \infty$. By the strong density of $\mathcal{S}_{\mathcal{M}}$ in \mathcal{M} , the *noncommutative L_p space* $L_p(\mathcal{M})$ is the corresponding completion for $p < \infty$ and $L_{\infty}(\mathcal{M}) = \mathcal{M}$. Many basic properties of classical L_p spaces like duality, real and complex interpolation, Hölder inequalities, etc hold in this setting. Elements of $L_p(\mathcal{M})$ can be described as measurable operators affiliated to (\mathcal{M}, τ) , we refer to Pisier/Xu's survey [33] for more information and historical references. Note that classical L_p spaces $L_p(\Omega, \mu)$ are denoted in this terminology as $L_p(\mathcal{M})$ where \mathcal{M} is the commutative von Neumann algebra $L_{\infty}(\Omega, \mu)$.

1.1.2. *Group algebras and comultiplication formulae.* – Our main example of noncommutative measure space in this paper is that of group von Neumann algebra. Let G be a locally compact and Hausdorff group (LCH group in short) equipped with its left Haar measure μ , throughout this text we will assume G to have a numerable base for the topology. Let $\lambda : G \rightarrow \mathcal{B}(L_2G)$ be the left regular representation. We will also use λ to denote the linear extension of λ to the space $L_1(G)$. We will denote by C_{λ}^*G the norm closure of $\lambda(L_1(G))$

and by $\mathcal{L}G$ the closure of C_λ^*G in the weak operator topology. $\mathcal{L}G$ is usually referred to as the *group von Neumann algebra* associated to G . There is a distinguished normal faithful weight $\tau : \mathcal{L}G_+ \rightarrow \mathbb{R}_+$ such that $\lambda : L_1(G) \cap L_2(G) \rightarrow \mathcal{L}G$ extends to an isometry from $L_2(G)$ to $L_2(\mathcal{L}G, \tau)$, the GNS construction associated to τ . Such weight is unique and it is called the Plancherel weight. When the function f belongs to the dense class $C_c(G) * C_c(G)$ we have $\tau(\lambda(f)) = f(e)$. The Plancherel weight is tracial if and only if G is unimodular. In this case it is called the Plancherel trace. From now on we will focus on unimodular groups. We will often work with the spaces $L_p(\mathcal{L}G, \tau)$ although the dependency on τ will be dropped in our terminology.

$\mathcal{L}G$ has a natural comultiplication given by linear extension of $\delta(\lambda_g) = \lambda_g \otimes \lambda_g$ which extends to a $*$ -homomorphism $\delta : C_\lambda^*G \rightarrow C_\lambda^*G \otimes_{\min} C_\lambda^*G$. There is a unique normal extension $\delta : \mathcal{L}G \rightarrow \mathcal{L}G \overline{\otimes} \mathcal{L}G$. This is a consequence of the fact that if δ is normal then $\delta_* : \mathcal{L}G_* \widehat{\otimes} \mathcal{L}G_* \rightarrow \mathcal{L}G_*$. Here \otimes_{\min} and $\widehat{\otimes}$ represent respectively the minimal and projective o.s. tensor products [32] and $\overline{\otimes}$ denotes the weak operator closure of the algebraic tensor product. Identifying $\mathcal{L}(G \times G)_*$ with $\mathcal{L}G_* \widehat{\otimes} \mathcal{L}G_*$ we have

$$\delta_* \left(\int_{G \times G} f(g_1, g_2) \lambda_{(g_1, g_2)} d\mu(g_1) d\mu(g_2) \right) = \int_G f(g, g) \lambda_g d\mu(g),$$

for every $f \in C_c(G \times G) * C_c(G \times G)$. The boundedness of δ_* is then a consequence of the Herz restriction theorem [14]. It is interesting to note that the Plancherel weight can be characterized as the unique normal, nontrivial and δ -invariant weight, where δ -invariant means that

$$(\tau \otimes \text{Id})\delta x = \tau(x)\mathbf{1}.$$

Analogously, Fourier multipliers are characterized as δ -equivariant maps

$$\delta T = (T \otimes \text{Id})\delta = (\text{Id} \otimes T)\delta.$$

We will denote by $\sigma : \mathcal{L}G \rightarrow \mathcal{L}G$ the anti-automorphism given by linear extension of $\sigma(\lambda_g) = \lambda_{g^{-1}}$. The *quantized convolution* of two elements x, y affiliated to $\mathcal{L}G$ is defined by

$$x \star y = (\tau \otimes \text{Id})(\delta x (\sigma y \otimes \mathbf{1})).$$

Observe that given $m \in L_\infty(G)$, the corresponding Fourier multiplier has the form

$$T_m(x) = \lambda(m) \star x = (\tau \otimes \text{Id})(\delta \lambda(m) (\sigma x \otimes \mathbf{1})).$$

1.1.3. *Operator space background.* – The theory of operator spaces is regarded as a noncommutative or quantized form of Banach space theory. An *operator space* E is a closed subspace of $\mathcal{B}(\mathcal{H})$. Let $M_m(E)$ be the space of $m \times m$ matrices with entries in E and impose on it the norm inherited from $M_m(E) \subset \mathcal{B}(\mathcal{H}^m)$. The morphisms in this category are the *completely bounded* linear maps (c.b. in short) $u : E \rightarrow F$, i.e., those satisfying

$$\|u\|_{\mathcal{CB}(E, F)} = \|u : E \rightarrow F\|_{\text{cb}} = \sup_{m \geq 1} \|\text{Id}_{M_m} \otimes u\|_{\mathcal{B}(M_m(E), M_m(F))} < \infty.$$

Similarly, given C^* -algebras A and B , a linear map $u : A \rightarrow B$ is called *completely positive* (c.p. in short) when $\text{Id}_{M_m} \otimes u$ is positivity preserving for $m \geq 1$. When a c.p. map $u : A \rightarrow B$ is

contractive (resp. unital) we will say it is a c.c.p. (resp. u.c.p.) map. The Kadison-Schwartz inequality for a c.c.p. map $u : \mathcal{M} \rightarrow \mathcal{M}$ claims that

$$u(x)^*u(x) \leq u(x^*x) \quad \text{for all } x \in \mathcal{M}.$$

Ruan's axioms describe axiomatically those sequences of matrix norms which can occur from an isometric embedding into $\mathcal{B}(\mathcal{H})$. Admissible sequences of matrix norms are called operator space structures (o.s.s. in short) and become crucial in the theory. Given a Banach space X and an isometric embedding $\rho : X \rightarrow \mathcal{B}(\mathcal{H})$ we will denote by X^ρ the corresponding operator space. Central branches from the theory of Banach spaces like duality, tensor norms or complex interpolation have been successfully extended to the category of operator spaces. Rather complete expositions are given in [8, 30, 32]. Two particular aspects of operator space theory which are relevant in this paper are the following:

A. Vector-valued Schatten classes. – We will denote by S_p the Schatten p -class given by $S_p = L_p(\mathcal{B}(\ell_2), \text{Tr})$ with Tr the standard trace in $\mathcal{B}(\ell_2)$. Similarly, S_p^m stands for the same space over $m \times m$ matrices. Vector-valued forms of these spaces can be defined as long as we define an o.s.s. over the space where we take values. Given an operator space E , we may define the E -valued Schatten classes $S_p^m[E]$ as the operator spaces given by interpolation

$$S_p^m[E] := [S_\infty^m[E], S_1^m[E]]_{\frac{1}{p}} := [S_\infty^m \otimes_{\min} E, S_1^m \widehat{\otimes} E]_{\frac{1}{p}}.$$

These classes provide a useful characterization of complete boundedness

$$\|u\|_{\mathcal{L}_{\mathcal{B}(E,F)}} = \sup_{m \geq 1} \|\text{Id}_{M_m} \otimes u\|_{\mathcal{B}(S_p^m(E), S_p^m(F))} \quad \text{for } 1 \leq p \leq \infty.$$

For a general hyperfinite von Neumann algebra \mathcal{M} the construction of $L_p(\mathcal{M}; E)$ is carried out by direct limits of E -valued Schatten classes. We refer to Pisier's book [31] for more on vector-valued noncommutative L_p spaces. The space $L_p(\mathcal{M}; E)$ for nonhyperfinite \mathcal{M} cannot be constructed without losing fundamental properties like projectivity/injectivity of the functor $E \mapsto L_p(\mathcal{M}; E)$. Fortunately, this drawback is solvable for the vector-valued L_p space we shall be working with.

B. Operator space structure of L_p . – Given an operator space E , its opposite E_{op} is the operator space which comes equipped with the operator space structure determined by the o.s.s. of E as follows

$$\left\| \sum_{j,k=1}^m a_{jk} \otimes e_{jk} \right\|_{M_m(E_{\text{op}})} = \left\| \sum_{j,k=1}^m a_{kj} \otimes e_{jk} \right\|_{M_m(E)},$$

where e_{jk} stand for the matrix units in M_m . Alternatively, if $E \subset \mathcal{B}(\mathcal{H})$, then $E_{\text{op}} = E^\top \subset \mathcal{B}(\mathcal{H})$, where \top is the transpose. The op construction plays a role in the construction of a “natural” o.s.s. for noncommutative L_p spaces. If \mathcal{M} is a von Neumann algebra we will denote by \mathcal{M}_{op} its opposite algebra, the original algebra with the multiplication reversed. It is a well-known result that \mathcal{M}_{op} and \mathcal{M} need not be isomorphic [5]. For every operator space E the natural inclusion $j : E \rightarrow E^{**}$ is a complete isometry. This allows us to build an operator space structure in the predual \mathcal{M}_* as the only operator space structure that makes the inclusion $j : \mathcal{M}_* \rightarrow \mathcal{M}^*$ completely isometric. The operator space structure

of $L_p(\mathcal{M})$ is given by operator space complex interpolation between $L_1(\mathcal{M}) = (\mathcal{M}_{\text{op}})^*$ and \mathcal{M} . In particular, it turns out that

$$L_p(\mathcal{M})^* \simeq L_{p'}(\mathcal{M}_{\text{op}})$$

is a complete isometry for $1 \leq p < \infty$, see [32, pp. 120-121] for further details.

1.1.4. *L_∞ -valued L_p spaces.* – Maximal inequalities are a cornerstone in harmonic analysis. Unfortunately, the supremum of a family of noncommuting operators is not well-defined, so that we do not have a proper noncommutative analog of maximal functions. Nevertheless, this difficulty can be overcome if all we want is to bound the maximal function in noncommutative L_p , as usually happens in harmonic analysis for commutative spaces. In that case we exploit the fact that the p -norm of a maximal function can always be written as a mixed $L_p(L_\infty)$ -norm of the corresponding entries. This reduces the problem to construct the vector-valued spaces $L_p(\mathcal{M}; L_\infty(\Omega))$. This construction can be carried out without requiring \mathcal{M} to be hyperfinite, relying in the commutativity of $L_\infty(\Omega)$. $L_p(\mathcal{M}; L_\infty(\Omega))$ is defined as the subspace of functions $x \in L_\infty(\Omega; L_p(\mathcal{M}))$ which admit a factorization of the form $x_\omega = \alpha y_\omega \beta$ with $\alpha, \beta \in L_{2p}(\mathcal{M})$ and $y \in L_\infty(\Omega; \mathcal{M})$. The norm in such space is then given by

$$\|(x_\omega)_{\omega \in \Omega}\|_{L_p(\mathcal{M}; L_\infty(\Omega))} = \inf \left\{ \|\alpha\|_{2p} \left(\operatorname{ess\,sup}_{\omega \in \Omega} \|y_\omega\|_{\mathcal{M}} \right) \|\beta\|_{2p} : x = \alpha y \beta \right\}.$$

When $x_\omega \geq 0$ the norm coincides with

$$(1.1) \quad \|(x_\omega)_{\omega \in \Omega}\|_{L_p(\mathcal{M}; L_\infty(\Omega))} = \inf \left\{ \|y\|_{L_p(\mathcal{M})} : x_\omega \leq y \text{ for a.e. } \omega \in \Omega \right\}.$$

Its operator space structure satisfies

$$S_p^m[L_p(\mathcal{M}; L_\infty(\Omega))] = L_p(M_m \otimes \mathcal{M}; L_\infty(\Omega)).$$

It is standard to use the following notation for the noncommutative $L_p(L_\infty)$ -norm

$$\left\| \sup_{\omega \in \Omega}^+ x_\omega \right\|_{L_p(\mathcal{M})} = \|(x_\omega)_{\omega \in \Omega}\|_{L_p(\mathcal{M}; L_\infty(\Omega))},$$

where the sup is just a symbolic notation without an intrinsic meaning. In the proof of Theorem B we will use the fact that if $(\mu_{\omega_2})_{\omega_2 \in \Omega_2}$ is a family of finite positive measures in Ω_1 and $(R_{\omega_1})_{\omega_1 \in \Omega_1}$ is a family of positivity preserving operators, then the following bound holds for $x \in L_p(\mathcal{M})_+$

$$(1.2) \quad \left\| \sup_{\omega_2 \in \Omega_2}^+ \left\{ \int_{\Omega_1} R_{\omega_1}(x) d\mu_{\omega_2}(\omega_1) \right\} \right\|_p \leq \left(\sup_{\omega_2 \in \Omega_2} \|\mu_{\omega_2}\|_{M(\Omega)} \right) \left\| \sup_{\omega_1 \in \Omega_1}^+ R_{\omega_1}(x) \right\|_p.$$

When \mathcal{M} is hyperfinite, this definition of $L_p(\mathcal{M}; L_\infty(\Omega))$ coincides with the corresponding vector-valued space as defined by Pisier [31]. This approach to handle maximal inequalities in von Neumann algebras has been successfully used in [15] to find noncommutative forms of Doob's maximal inequality for martingales and the maximal ergodic inequalities for Markov semigroups [23]. The predual can be explicitly described as the L_1 -valued space $L_{p'}(\mathcal{M}; L_1(\Omega))$. Indeed, let $S_p(\Omega)$ be the Schatten class associated to the Hilbert space $L_2(\Omega)$. Note that there is a hermitian form $q : L_{2p}(\mathcal{M}) \otimes S_2^c(\Omega) \times L_{2p}(\mathcal{M}) \otimes S_2^c(\Omega) \rightarrow L_p(\mathcal{M}) \otimes L_1(\Omega)$ given by

$$q(x \otimes m, y \otimes n) = x^* y \otimes \operatorname{diag}(m^* n),$$

where $\text{diag} : S_1(\Omega) \rightarrow L_1(\Omega)$ is the restriction to the diagonal. Define

$$\|x\|_{L_p(\mathcal{M}; L_1(\Omega))} = \inf \left\{ \|a\|_{L_{2p}(\mathcal{M}; S_2^c(\Omega))} \|b\|_{L_{2p}(\mathcal{M}; S_2^c(\Omega))} : q(a, b) = x \right\}.$$

This space satisfies that $L_p(\mathcal{M}; L_1(\Omega))^* = L_{p'}(\mathcal{M}_{\text{op}}; L_\infty(\Omega))$ for $1 \leq p < \infty$.

1.1.5. *Hilbert-valued L_p spaces.* – For certain operator spaces whose underlying Banach space is a Hilbert space we can define vector-valued noncommutative L_p spaces for general von Neumann algebras. Indeed, let \mathcal{H} be a Hilbert space and $P_e \xi = \langle e, \xi \rangle e$ for some $e \in \mathcal{H}$ of unit norm. We define the following two Hilbert-valued forms of $L_p(\mathcal{M})$

$$\begin{aligned} L_p(\mathcal{M}; \mathcal{H}^c) &= L_p(\mathcal{M} \overline{\otimes} \mathcal{B}(\mathcal{H}))(\mathbf{1}_{\mathcal{M}} \otimes P_e), \\ L_p(\mathcal{M}; \mathcal{H}^r) &= (\mathbf{1}_{\mathcal{M}} \otimes P_e)L_p(\mathcal{M} \overline{\otimes} \mathcal{B}(\mathcal{H})), \end{aligned}$$

called the L_p spaces with \mathcal{H} -column (resp. \mathcal{H} -row) values. Their o.s.s. are the ones inherited from $L_p(\mathcal{M} \overline{\otimes} \mathcal{B}(\mathcal{H}))$. If $\mathcal{H} = \ell_2^n$, then we can identify $L_p(\mathcal{B}(\mathcal{H}) \overline{\otimes} \mathcal{M})$ with $L_p(\mathcal{M})$ -valued $n \times n$ matrices. Under that identification $L_p(\mathcal{M}; \mathcal{H}^c)$ (resp. $L_p(\mathcal{M}; \mathcal{H}^r)$) corresponds to the matrices with zero entries outside the first column (resp. row) and we have that

$$\begin{aligned} \left\| \sum_{j=1}^n x_j \otimes e_{j1} \right\|_{L_p(\mathcal{M} \overline{\otimes} \mathcal{B}(\ell_2^n))} &= \left\| \left(\sum_{j=1}^n x_j^* x_j \right)^{\frac{1}{2}} \right\|_{L_p(\mathcal{M})}, \\ \left\| \sum_{j=1}^n x_j \otimes e_{1j} \right\|_{L_p(\mathcal{M} \overline{\otimes} \mathcal{B}(\ell_2^n))} &= \left\| \left(\sum_{j=1}^n x_j x_j^* \right)^{\frac{1}{2}} \right\|_{L_p(\mathcal{M})}. \end{aligned}$$

The same formulas hold after replacing the finite sums by infinite ones of even by integrals. For every $1 \leq p \leq \infty$ we can embed \mathcal{H} isometrically in S_p by sending $c_p(e_j) = e_{1j}$ or $r_p(e_j) = e_{j1}$, where $\{e_j\}$ is an orthonormal basis of \mathcal{H} . Such maps are called the p -column/ p -row embeddings. These isometries endow \mathcal{H} with several o.s. structures. Observe that, as an o.s., $L_p(\mathcal{M}; \mathcal{H}^c)$ (resp. $L_p(\mathcal{M}; \mathcal{H}^r)$) coincides with Pisier’s vector-valued L_p -space $L_p(\mathcal{M}; \mathcal{H}^{c_p})$ (resp. $L_p(\mathcal{M}; \mathcal{H}^{r_p})$) for \mathcal{M} hyperfinite. For $1 \leq p < \infty$ the duals are given by $L_p(\mathcal{M}; \mathcal{H}^c)^* = L_{p'}(\overline{\mathcal{M}}; \mathcal{H}^c)$ and $L_p(\mathcal{M}; \mathcal{H}^r)^* = L_{p'}(\overline{\mathcal{M}}; \mathcal{H}^r)$. The duality pairing can be expressed as

$$\left\langle \sum_j x_j \otimes e_j, \sum_k y_k \otimes e_k \right\rangle = \sum_j \tau(x_j^* y_j).$$

The spaces $L_p(\mathcal{M}; \mathcal{H}^r)$ and $L_p(\mathcal{M}; \mathcal{H}^c)$ form complex interpolation scales for $p \geq 1$

$$\begin{aligned} [L_\infty(\mathcal{M}; \mathcal{H}^r), L_p(\mathcal{M}; \mathcal{H}^r)]_\theta &= L_{\frac{p}{\theta}}(\mathcal{M}; \mathcal{H}^r), \\ [L_\infty(\mathcal{M}; \mathcal{H}^c), L_p(\mathcal{M}; \mathcal{H}^c)]_\theta &= L_{\frac{p}{\theta}}(\mathcal{M}; \mathcal{H}^c). \end{aligned}$$

In order to treat square functions and Hardy spaces we will need to control sums and intersections of these Hilbert valued noncommutative L_p spaces. The algebraic tensor product $L_p(\mathcal{M}) \otimes \mathcal{H}$ embeds in $L_p(\mathcal{M} \overline{\otimes} \mathcal{B}(\mathcal{H}))$ by $\text{Id} \otimes r$ and $\text{Id} \otimes c$. Taking direct sums we obtain an embedding in $X = L_p(\mathcal{M} \overline{\otimes} \mathcal{B}(\mathcal{H})) \oplus L_p(\mathcal{M} \overline{\otimes} \mathcal{B}(\mathcal{H}))$. The space $L_p(\mathcal{M}; \mathcal{H}^{r \cap c})$ is defined as the norm closure (or weak- $*$ closure if $p = \infty$) of $L_p(\mathcal{M}) \otimes \mathcal{H}$ inside X . Such space comes equipped with the norm given by

$$\|x\|_{L_p(\mathcal{M}; \mathcal{H}^{r \cap c})} = \max \left\{ \|x\|_{L_p(\mathcal{M}; \mathcal{H}^r)}, \|x\|_{L_p(\mathcal{M}; \mathcal{H}^c)} \right\}.$$

The embedding also gives $L_p(\mathcal{M}; \mathcal{H}^{r \cap c})$ an o.s.s. We will denote the dual spaces by $L_p(\mathcal{M}; \mathcal{H}^{r+c}) = L_{p'}(\mathcal{M}_{\text{op}}; \mathcal{H}^{r \cap c})^*$ for $1 < p \leq \infty$. Since the subspace

$$\begin{aligned} L_\infty(\mathcal{M}_{\text{op}}; \mathcal{H}^{r \cap c}) &\subset (\mathbf{1} \otimes P_{e_1})(\mathcal{M}_{\text{op}} \overline{\otimes} \mathcal{B}(\mathcal{H})) \oplus (\mathcal{M}_{\text{op}} \overline{\otimes} \mathcal{B}(\mathcal{H}))(\mathbf{1} \otimes P_{e_1}) \\ &\subset \mathcal{M}_{\text{op}} \overline{\otimes} \mathcal{B}(\mathcal{H}) \oplus \mathcal{M}_{\text{op}} \overline{\otimes} \mathcal{B}(\mathcal{H}) \end{aligned}$$

is weak-* closed, $L_\infty(\mathcal{M}_{\text{op}}; \mathcal{H}^{r \cap c})$ inherits a natural weak-* topology through the inclusion. The space $L_1(\mathcal{M}; \mathcal{H}^{r+c})$ is defined as the subset of weak-* continuous functionals in $L_\infty(\mathcal{M}_{\text{op}}; \mathcal{H}^{r \cap c})^*$. The sum notation comes from the fact that

$$\|x\|_{L_p(\mathcal{M}; \mathcal{H}^{r+c})} = \inf \left\{ \|y\|_{L_p(\mathcal{M}; \mathcal{H}^r)} + \|z\|_{L_p(\mathcal{M}; \mathcal{H}^c)} : x = y + z \right\}.$$

We will denote by $L_p(\mathcal{M}; \mathcal{H}^{rc})$ the spaces given by

$$L_p(\mathcal{M}; \mathcal{H}^{rc}) = \begin{cases} L_p(\mathcal{M}; \mathcal{H}^{r+c}) & \text{when } 1 \leq p < 2, \\ L_p(\mathcal{M}; \mathcal{H}^{r \cap c}) & \text{when } 2 \leq p \leq \infty. \end{cases}$$

The spaces $L_p(\mathcal{M}; \mathcal{H}^{rc})$ are crucial for the noncommutative Khintchine inequalities [27, 28], the noncommutative Burkholder-Gundy inequalities [22], noncommutative Littlewood-Paley estimates [16] and other related results.

1.1.6. *Markovian semigroups and length functions.* – A semigroup $\mathcal{S} = (S_t)_{t \geq 0}$ over a Banach space X is a family of operators $S_t : X \rightarrow X$ such that $S_0 = \text{Id}$ and $S_t S_s = S_{t+s}$. Let (\mathcal{M}, τ) be a noncommutative measure space, we will say that a semigroup \mathcal{S} over \mathcal{M} is *submarkovian* iff:

- i) each S_t is a weak-* continuous and c.c.p. map,
- ii) each S_t is a self-adjoint, i.e., $\tau(S_t x^* y) = \tau(x^* S_t y)$,
- iii) the map $t \mapsto S_t$ is pointwise weak-* continuous.

\mathcal{S} is *Markovian* if each S_t is a u.c.p. map, ie $S_t(\mathbf{1}) = \mathbf{1}$. Markovian operators satisfy $\tau \circ S_t = \tau$ while submarkovian ones satisfy $\tau \circ S_t \leq \tau$. Sometimes these semigroups are called symmetric and Markovian, where symmetric is synonym with self-adjoint. All the semigroups in this paper will be symmetric, so we will drop the adjective. Using the first two properties it is easy to see that S_t extends to a c.c.p. map on $L_1(\mathcal{M})$. By the Riesz-Thorin theorem S_t is a complete contraction over $L_p(\mathcal{M})$ for $1 \leq p \leq \infty$. The third property implies that $t \mapsto S_t$ is SOT continuous in $L_1(\mathcal{M})$. By interpolation between the pointwise norm continuity on $L_1(\mathcal{M})$ and the pointwise weak-* continuity on \mathcal{M} we obtain that $t \mapsto S_t$ is SOT continuous on $L_p(\mathcal{M})$ for $1 \leq p < \infty$. For every $1 \leq p < \infty$ there is a densely defined and closable operator A whose closed domain is given by

$$\text{dom}_p(A) = \left\{ x \in L_p(\mathcal{M}) : \exists \lim_{t \rightarrow 0^+} \frac{x - S_t x}{t} \text{ in the norm topology} \right\}.$$

When $p = 2$ we have that $S_t = e^{-tA}$ and $S_t[L_p(\mathcal{M})] \subset \text{dom}_p(A)$ for $1 \leq p < \infty$. In the case $p = \infty$ we have that A is densely defined and closable with respect to the weak-* topology with domain given by those $x \in \mathcal{M}$ such that $\lim_{t \rightarrow 0^+} (x - S_t x)/t$ exists in the weak-* topology. We will call A the infinitesimal generator of \mathcal{S} .

We are interested in those (sub)markovian semigroups over $\mathcal{M} = \mathcal{L}G$ which are of convolution type. In other words, each S_t is a Fourier multiplier. It can be proved that

$S_t = T_{e^{-t\psi}}$ for some function ψ . Let us recall a characterization of these functions. First, recall some definitions. A continuous function $\psi : G \rightarrow \mathbb{C}$ is said to be conditionally negative (c.n. in short) iff $\psi(e) = 0$ and for every finite subset $\{g_1, \dots, g_m\} \subset G$ and vector $(v_1, \dots, v_m) \in \mathbb{C}^n$ we have

$$\sum_{i=1}^m v_i = 0 \Rightarrow \sum_{i,j=1}^m \bar{v}_i \psi(g_i^{-1} g_j) v_j \leq 0.$$

When $\psi : G \rightarrow \mathbb{R}_+$ is symmetric ($\psi(g) = \psi(g^{-1})$) and c.n. we will say that ψ is a *conditionally negative length*. Let \mathcal{H} be a real Hilbert space. Given an orthogonal representation $\alpha : G \rightarrow O(\mathcal{H})$ we say that a continuous map $b : G \rightarrow \mathcal{H}$ is a 1-cocycle (with respect to α) iff it satisfies the 1-cocycle law

$$b(gh) = \alpha(g)b(h) + b(g).$$

The following characterization is proved in [1, Appendix C] or [4, Chapter 1].

THEOREM 1.1. – *Let $\mathcal{S} = (S_t)_{t \geq 0}$ be a semigroup of convolution type over the group algebra $\mathcal{L}G$. Then, the following statements are equivalent:*

- i) \mathcal{S} is a Markovian semigroup,
- ii) there is a c.n. length $\psi : G \rightarrow \mathbb{R}_+$ such that $S_t = T_{e^{-t\psi}}$,
- iii) there is a real Hilbert space \mathcal{H} , an orthogonal representation $\alpha : G \rightarrow O(\mathcal{H})$ and a 1-cocycle $b : G \rightarrow \mathcal{H}$, such that $\psi(g) = \|b(g)\|_{\mathcal{H}}^2$ and $S_t = T_{e^{-t\psi}}$.

1.1.7. *Holomorphic calculus and noncommutative Hardy spaces.* – We now introduce the Hardy spaces associated with a submarkovian semigroup on (\mathcal{M}, τ) as well as the corresponding \mathcal{H}^∞ -functional calculus. Both tools were introduced in the noncommutative setting in [16]. If \mathcal{S} is a submarkovian semigroup, the fixed point subspace $F_p = \{x \in L_p(\mathcal{M}) : S_t(x) = x \forall t \geq 0\}$ coincides with $\ker A \subset \text{dom}_p(A)$ and it is a subalgebra when $p = \infty$. It is also easily seen to be a complemented subspace for $1 < p < \infty$ with projection given by $Q_p(x) = \lim_{t \rightarrow \infty} S_t x$ where the limit converges in the weak-* topology of L_p , for $1 < p < \infty$. In that range of p we will denote by $L_p^\circ(\mathcal{M}) = L_p(\mathcal{M})/F_p$ which is also a complemented subspace with projection given by $P_p = \text{Id} - Q_p$. Note that $L_p(\mathcal{M}) \simeq L_p^\circ(\mathcal{M}) \oplus_p F_p$. When S_t are Fourier multipliers over $\mathcal{M} = \mathcal{L}G$ with symbol $e^{-t\psi}$ we define $G_0 = \{g \in G : \psi(g) = 0\}$. In that case it is easy to see that F_p coincides with the space bellow

$$\overline{\{x \in L_p(\mathcal{M}) : x = \lambda(v) \text{ with } \text{supp}(v) \subset G_0\}}^{\|\cdot\|_p}.$$

Indeed, let us denote the space above by B_p . The inclusion $B_p \subset F_p$ is trivial. For the other we are going to distinguish between two cases. The first is when $1 < p \leq 2$. In such range the operator $\lambda^* : L_p(\mathcal{L}G) \rightarrow L_{p'}(G)$ is contractive and so, if $S_t(x) = x$, we have that $x = \lambda(f)$ and that $e^{-t\psi} f = f$. Which implies that $f|_{G \setminus G_0} = 0$. In the range $2 < p < \infty$, assume that $x_0 \in F_p$ but not in B_p , then by the Hahn-Banach theorem, there is a $y_0 \in L_{p'}(\mathcal{L}G)$ such that $\langle y_0, x_0 \rangle > \epsilon$ and $\langle y_0, x \rangle = 0$ when $x \in B_p$. We have that $y_0 = \lambda(f_0)$ and by the second condition we have that

$$0 = \langle y_0, \lambda(v) \rangle = \int_G f_0 d\nu,$$

for every ν with $\text{supp}[\nu] \subset G_0$ which in turn implies that $f_0|_{G_0} = 0$. As a consequence $e^{-t\psi} f_0 \rightarrow 0$ pointwise. On the other hand, since x_0 satisfies that $S_t(x_0) = x_0$ for every t , we can change y_0 by $S_t(y_0)$ keeping the separation properties. The same holds for every element in the weak-* closed convex hull of such family

$$\mathcal{C} = \overline{\text{hull}^{\text{w}*}} \{ \lambda(e^{-t\psi} f_0) : 0 \leq t \}.$$

But, the limit $e^{-t\psi(g)} f_0(g) \rightarrow 0$ implies that $0 \in \mathcal{C}$, which contradicts $\langle y, x_0 \rangle > \epsilon$.

In a similar way we find that $\lambda(\nu) \in L_p^\circ(\mathcal{M})$ if and only if the support of the measure ν doesn't intersect G_0 . It is also worth recalling that for G_0 of measure zero $L_p^\circ(\mathcal{L}G) = L_p(\mathcal{L}G)$.

For any given $x \in \mathcal{M}$ we define the function $Tx : (0, \infty) \rightarrow L_p(\mathcal{M})$ given by $t \mapsto t \partial_t S_t x$. We can see $x \mapsto Tx$ as a map from certain domain $D \subset \mathcal{M}$ into $L_p(\mathcal{M}; \mathcal{H}^r)$, $L_p(\mathcal{M}; \mathcal{H}^c)$ or $L_p(\mathcal{M}; \mathcal{H}^{rc})$, where $\mathcal{H} = L_2(\mathbb{R}_+, dt/t)$. The induced seminorms on $D \subset \mathcal{M}$ are called the row Hardy space, column Hardy space or Hardy space seminorms. Observe that the map T has as kernel those elements fixed by \mathcal{S} . Quotient out the null space and taking the completion with respect to any of those norms when $p < \infty$ (resp. the weak-* topology for $p = \infty$) gives the Hardy spaces $H_p^r(\mathcal{M}; \mathcal{S})$, $H_p^c(\mathcal{M}; \mathcal{S})$ or $H_p(\mathcal{M}; \mathcal{S})$. We can represent such norms as follows

$$\|x\|_{H_p^c(\mathcal{M}; \mathcal{S})} = \left\| \left(\int_{\mathbb{R}_+} \left(t \frac{d}{dt} S_t x \right)^* \left(t \frac{d}{dt} S_t x \right) \frac{dt}{t} \right)^{\frac{1}{2}} \right\|_{L_p(\mathcal{M})},$$

$$\|x\|_{H_p^r(\mathcal{M}; \mathcal{S})} = \left\| \left(\int_{\mathbb{R}_+} \left(t \frac{d}{dt} S_t x \right) \left(t \frac{d}{dt} S_t x \right)^* \frac{dt}{t} \right)^{\frac{1}{2}} \right\|_{L_p(\mathcal{M})}.$$

We will drop the dependency on the semigroup and write $H_p^c(\mathcal{M})$ whenever it can be understood from the context. These spaces inherit their o.s.s. from that of $L_p(\mathcal{M}; \mathcal{H}^r)$ or $L_p(\mathcal{M}; \mathcal{H}^c)$. Therefore we have the following identities

$$S_p^n[H_p^c(\mathcal{M}; \mathcal{S})] = H_p^c(\mathcal{M} \overline{\otimes} \mathcal{B}(\ell_2^n); \mathcal{S} \otimes \text{Id}),$$

$$S_p^n[H_p^r(\mathcal{M}; \mathcal{S})] = H_p^r(\mathcal{M} \overline{\otimes} \mathcal{B}(\ell_2^n); \mathcal{S} \otimes \text{Id}).$$

The duality is obtained from that of $L_p(\mathcal{M}; \mathcal{H}^c)$ or $L_p(\mathcal{M}; \mathcal{H}^r)$, resulting in the cb-isometries $H_p^r(\mathcal{M}; \mathcal{S})^* = H_p^c(\mathcal{M}_{\text{op}}; \mathcal{S})$ for $1 \leq p < \infty$. The same holds for the column case. Finally let us recall that by [16, Chapters 7 and 10] we have that if $1 < p < \infty$ then

$$(1.3) \quad H_p(\mathcal{M}; \mathcal{S}) \simeq L_p^\circ(\mathcal{M}),$$

with the equivalence as operator spaces depending on the constant p . The result fails for $p = 1, \infty$ and $H_1(\mathcal{M}; \mathcal{S})$ is smaller in general than $L_1^\circ(\mathcal{M})$. Observe that $t \partial_t S_t x = \eta(tA)x$ where $\eta(z) = ze^{-z}$. Due to the results in [16] we can change η by other analytic functions in certain class obtaining equivalent norms. We will say that a holomorphic function ρ defined over the sector $\Sigma_\theta = \{z \in \mathbb{C} : |\arg(z)| < \theta\}$ is in $\mathcal{H}^\infty(\Sigma_\theta)$ iff it is bounded and we will say that it is in $\mathcal{H}_0^\infty(\Sigma_\theta) \subset \mathcal{H}^\infty(\Sigma_\theta)$ iff there is an $s > 0$ such that

$$|\rho(z)| \lesssim \frac{|z|^s}{(1 + |z|)^{2s}}.$$

We will denote by \mathcal{H}^∞ or \mathcal{H}_0^∞ the spaces $\bigcap_{0 < \theta < \pi/2} \mathcal{H}^\infty(\Sigma_\theta)$ or $\bigcap_{0 < \theta < \pi/2} \mathcal{H}_0^\infty(\Sigma_\theta)$ respectively. If needed, we will equip these spaces with their natural inverse limit topologies.

We have that for any $\rho \in \mathcal{H}_0^\infty$ the following holds

$$(1.4) \quad \begin{aligned} \|x\|_{H_p^c(\mathcal{M})} &\sim_{(p)} \left\| \left(\int_{\mathbb{R}_+} (\rho(tA)x)^* \rho(tA)x \frac{dt}{t} \right)^{\frac{1}{2}} \right\|_{L_p(\mathcal{M})}, \\ \|x\|_{H_p^r(\mathcal{M})} &\sim_{(p)} \left\| \left(\int_{\mathbb{R}_+} \rho(tA)x (\rho(tA)x)^* \frac{dt}{t} \right)^{\frac{1}{2}} \right\|_{L_p(\mathcal{M})}. \end{aligned}$$

The equivalence also holds after matrix amplifications. This type of identities also hold for wider classes of unbounded operators A satisfying certain resolvent estimates, see [16] for further details.

1.2. The general principle

We are now ready to prove our maximal bounds in Theorem A. In fact, we shall obtain a more general principle in Theorem 1.3 which decouples in terms of row and column Hardy spaces.

DEFINITION 1.2. – Let $(B_t)_{t \geq 0}$ be a family of operators affiliated to $\mathcal{L}G$. We say that $(B_t)_{t \geq 0}$ has an L_p -square-max decomposition when there is a decomposition $B_t = \Sigma_t M_t$ such that :

$$(SM_p) \quad \begin{aligned} \sup_{t \geq 0} \|\Sigma_t\|_2 &< \infty, \\ \left\| \sup_{t > 0}^+ \sigma |M_t|^2 \star u \right\|_p &\lesssim_{(p)} \|u\|_p. \end{aligned}$$

Similarly, $(B_t)_{t \geq 0}$ has an L_p -max-square decomposition when $B_t = M_t \Sigma_t$ with :

$$(MS_p) \quad \begin{aligned} \sup_{t \geq 0} \|\Sigma_t\|_2 &< \infty, \\ \left\| \sup_{t > 0}^+ \sigma |M_t^*|^2 \star u \right\|_p &\lesssim_{(p)} \|u\|_p. \end{aligned}$$

When we say that $(B_t)_{t \geq 0}$ has a max-square (resp. square-max) decomposition we mean that it has an L_p -max-square (resp. L_p -square-max) decomposition for every $1 < p < \infty$.

THEOREM 1.3. – Let G be a LCH group equipped with a conditionally negative length $\psi : G \rightarrow \mathbb{R}_+$. Let $\mathcal{S} = (S_t)_{t \geq 0}$ be the convolution semigroup generated by ψ and pick any $\eta \in \mathcal{H}_0^\infty$. If $m : G \rightarrow \mathbb{C}$ is a bounded function satisfying that $B_t = \lambda(m\eta(t\psi))$ has an $L_{(p/2)'}$ -square-max decomposition $B_t = \Sigma_t M_t$ for some $2 < p < \infty$, then $T_m : H_p^c(\mathcal{L}G) \rightarrow H_p^c(\mathcal{L}G)$ and

$$\|T_m\|_{\mathcal{B}(H_p^c)} \lesssim_{(p)} \left(\sup_{t \geq 0} \|\Sigma_t\|_2 \right) \left\| (R_t^c)_{t \geq 0} : L_{(p/2)' }(\mathcal{L}G) \rightarrow L_{(p/2)' }(\mathcal{L}G; L_\infty) \right\|^{\frac{1}{2}}$$

where $R_t^c(x) = \sigma |M_t|^2 \star x$. Similarly, when $(B_t)_{t \geq 0}$ admits an $L_{(p/2)'}$ -max-square decomposition $B_t = M_t \Sigma_t$ for some $2 < p < \infty$, we get $T_m : H_p^r(\mathcal{L}G) \rightarrow H_p^r(\mathcal{L}G)$ and the following estimate holds

$$\|T_m\|_{\mathcal{B}(H_p^r)} \lesssim_{(p)} \left(\sup_{t \geq 0} \|\Sigma_t\|_2 \right) \left\| (R_t^r)_{t \geq 0} : L_{(p/2)' }(\mathcal{L}G) \rightarrow L_{(p/2)' }(\mathcal{L}G; L_\infty) \right\|^{\frac{1}{2}}$$

where $R_t^r(x) = \sigma |M_t^*|^2 \star x$. By duality, similar identities also hold for $1 < p < 2$.

COROLLARY 1.4. – If G , ψ , η and m are as above and $B_t = \lambda(m\eta(t\psi))$ admits both a $L_{(p/2)'}\text{-max-square}$ and a $L_{(p/2)'}\text{-square-max decomposition}$, then it turns out that $T_m : L_p^\circ(\mathcal{L}G) \rightarrow L_p^\circ(\mathcal{L}G)$ boundedly. Furthermore, if $m \equiv c$ in $G_0 = \{g \in G : \psi(g)\}$ then T_m is a bounded map on $L_p(\mathcal{L}G)$.

Proof. – The first assertion follows trivially from (1.3). For the second we use that $L_p^\circ(\mathcal{L}G)$ is a complemented subspace, and so

$$\begin{aligned} \|T_m x\|_p &\leq \|P_p T_m x\|_p + \|Q_p T_m x\|_p \\ &= \|T_m P_p x\|_{L_p^\circ(\mathcal{L}G)} + \|T_{m|_{G_0}} Q_p x\|_p \lesssim_{(p)} \left(\|T_m\|_{\mathcal{B}(L_p^\circ(\mathcal{L}G))} + c \right) \|x\|_p. \quad \square \end{aligned}$$

Proof of Theorem 1.3. – Assume that $B_t = \lambda(m\eta(t\psi))$ has an $L_{(p/2)'}\text{-square-max decomposition}$. According to (1.4) with $\rho(z) = \eta(z)\varrho(z)$ for some $\varrho \in \mathcal{H}_0^\infty$, and using that T_m commutes with the spectral calculus of A (the generator of \mathcal{G}) we obtain

$$\begin{aligned} \|T_m(x)\|_{H_p^\zeta} &\sim_{(p)} \left\| (\eta(tA)\varrho(tA)T_m x)_{t \geq 0} \right\|_{L_p(\mathcal{L}G; L_2^\zeta)} \\ &= \left\| (\eta(tA)T_m \varrho(tA)x)_{t \geq 0} \right\|_{L_p(\mathcal{L}G; L_2^\zeta)} \\ &= \left\| (T_{m_t}(x_t))_{t \geq 0} \right\|_{L_p(\mathcal{L}G; L_2^\zeta)}, \end{aligned}$$

where $m_t(g) = m(g)\eta(t\psi(g))$ and $x_t = T_{\varrho(t\psi)}x$. Recall also that the L_2 -space involved is $L_2(\mathbb{R}_+, dt/t)$. Now we may express the term on the right hand side as follows

$$\begin{aligned} (1.5) \quad \left\| (T_{m_t}(x_t))_{t \geq 0} \right\|_{L_p(\mathcal{L}G; L_2^\zeta)}^2 &= \left\| \int_{\mathbb{R}_+} |T_{m_t} x_t|^2 \frac{dt}{t} \right\|_{\frac{p}{2}} \\ &= \tau \left(u \int_{\mathbb{R}_+} |T_{m_t} x_t|^2 \frac{dt}{t} \right) = \int_{\mathbb{R}_+} \tau(u |T_{m_t} x_t|^2) \frac{dt}{t}, \end{aligned}$$

where $u \in L_{(p/2)'}(\mathcal{L}G)_+$ is the unique element realizing the $L_{p/2}$ -norm, which exists by the weak- $*$ compactness of the unit ball of $L_{(p/2)'}(\mathcal{L}G)$. Now we have to estimate the term inside the integral. As $u \geq 0$, we may write $u = w^*w$ for some $w \in L_{2(p/2)'}$ and

$$\langle u, |T_{m_t}(x_t)|^2 \rangle = \tau(w |T_{m_t}(x_t)|^2 w^*) = \tau \left(\underbrace{w |(\tau \otimes \text{Id})(\delta B_t(\sigma x_t \otimes \mathbf{1}))|^2}_{L_t} w^* \right).$$

As $L_t \mapsto w L_t w^*$ is order preserving, any bound of L_t gives a bound of the above term. By the complete positivity of the canonical trace we can apply Proposition 1.1 in [26], i.e.,

$$\langle x, y \rangle^* \langle x, y \rangle \leq \| \langle x, x \rangle \| \langle y, y \rangle$$

to the operator-valued inner product $\langle x, y \rangle = (\tau \otimes \text{Id})(x^* y)$. This yields

$$\begin{aligned} L_t &= |(\tau \otimes \text{Id})(\delta \Sigma_t \delta M_t(\sigma x_t \otimes \mathbf{1}))|^2 \\ &\leq \|(\tau \otimes \text{Id})(\delta |\Sigma_t|^2)\|_{\mathcal{L}G} (\tau \otimes \text{Id}) \left((\sigma x_t^* \otimes \mathbf{1}) \delta M_t^* \delta M_t(\sigma x_t \otimes \mathbf{1}) \right) \\ &\leq \left(\sup_{t>0} \|\Sigma_t\|_2^2 \right) (\tau \otimes \text{Id}) (\delta |M_t|^2(\sigma(x_t^* x_t) \otimes \mathbf{1})) = \left(\sup_{t>0} \|\Sigma_t\|_2^2 \right) (|M_t|^2 \star x_t^* x_t). \end{aligned}$$

We have used the δ -invariance of the trace in the second inequality and the definition of the noncommutative convolution in the last identity. Now, substituting inside the trace and using the identity for the adjoint of the noncommutative convolution operator gives

$$\langle u, |T_{m_t}(x_t)|^2 \rangle \leq K^2 \tau(u(|M_t|^2 \star x_t^* x_t)) = K^2 \tau((\sigma |M_t|^2 \star u)x_t^* x_t),$$

where K is the supremum of the L_2 norm of Σ_t . This gives rise to

$$\begin{aligned} \|T_m(x)\|_{H_p^c}^2 &\lesssim_{(p)} K^2 \int_{\mathbb{R}_+} \tau((\sigma |M_t|^2 \star u)x_t^* x_t) \frac{dt}{t} \\ &\leq K^2 \inf_{\sigma |M_t|^2 \star u \leq A} \tau\left(A \int_{\mathbb{R}_+} x_t^* x_t \frac{dt}{t}\right) \\ &\leq K^2 \inf_{\sigma |M_t|^2 \star u \leq A} \|A\|_{L_{(p/2)'}} \left\| \int_{\mathbb{R}_+} x_t^* x_t \frac{dt}{t} \right\|_{p/2} \\ &\lesssim_{(p)} K^2 \left\| (R_t^c)_{t \geq 0} : L_{(p/2)'} \rightarrow L_{(p/2)'}(L_\infty) \right\| \|x\|_{H_p^c}^2 \end{aligned}$$

by using Fatou’s lemma in the second line and the definition of the $L_p(\mathcal{L}G; L_\infty)$ norm for positive elements in the last inequality. Taking square roots gives the desired estimate. The calculations for the row case are entirely analogous. \square

REMARK 1.5. – Throughout this paper we construct max-square and square-max decompositions of $B_t = \lambda(m \eta(t\psi))$ by choosing an smoothing positive factor M_t with $M_t = \sigma M_t = M_t^*$ and satisfying the appropriate maximal inequalities. Then we extract M_t from the left and from the right of B_t as

$$\begin{aligned} B_t &= (B_t M_t^{-1}) M_t, \\ B_t &= M_t (M_t^{-1} B_t). \end{aligned}$$

If the family $\Sigma_t = B_t M_t^{-1}$ is uniformly bounded in L_2 and B_t is self-adjoint then the other is automatically uniformly in L_2 by the traciality of τ . Most of the times it will be enough to check one of the two decompositions.

Proof of Theorem A. – The assumptions on m and η ensure that B_t is self-adjoint, similarly we know by assumption that M_t is symmetric and positive. The result follows easily from Theorem 1.3 and Remark 1.5. \square

REMARK 1.6. – The technique employed here gives complete bounds assuming that the maximal inequalities are satisfied with complete bounds. In order to prove that assertion, let us express the matrix extension $(T_m \otimes \text{Id}_{M_n})$ as a matrix-valued multiplier whose symbol takes diagonal values. Indeed

$$(T_m \otimes \text{Id}_{M_n})([x_{ij}]) = (\text{Id} \otimes \tau \otimes \text{Id}_{M_n}) \left(\underbrace{(\delta\lambda(m) \otimes \mathbf{1}_{M_n})}_K (\mathbf{1} \otimes [\sigma x_{ij}]) \right),$$

where K is the corresponding kernel affiliated with $\mathcal{L}G \overline{\otimes} \mathcal{L}G \overline{\otimes} \mathbb{C}\mathbf{1}_{M_n}$. Clearly, any square-max decomposition $B_t = \Sigma_t M_t$ of $B_t = \lambda(m \eta(t\psi))$ yields a diagonal decomposition $(\delta\Sigma_t \otimes \mathbf{1}_{M_n})(\delta M_t \otimes \mathbf{1}_{M_n})$ of $K_t = \delta B_t \otimes \mathbf{1}_{M_n}$. On the other hand recall that $T_m : H_p^c \rightarrow H_p^c$ is c.b. iff $T_m \otimes \text{Id}_{M_n} : S_p^n[H_p^c] \rightarrow S_p^n[H_p^c]$ is uniformly bounded for $n \geq 1$ and that $S_p^n[H_p^c(\mathcal{L}G; \mathcal{S})] = H_p^c(M_n \otimes \mathcal{L}G; \text{Id} \otimes \mathcal{S})$. That allows us to write the

norm of $S_p^n[H_p^c(\mathcal{L}G; \mathcal{S})]$ as an $L_{p/2}$ -norm like in (1.5). Then, using [26, Proposition 1.1] for $\langle x, y \rangle = (\text{Id} \otimes \tau \otimes \text{Id}_{M_n})(x^*y)$ as in the proof of Theorem 1.3, gives for $2 < p < \infty$

$$\|T_m\|_{\mathcal{CB}(H_p^c)} \lesssim_{(p)} \left(\sup_{t \geq 0} \|\Sigma_t\|_2 \right) \left\| (R_t^c)_{t \geq 0} : L_{(p/2)'}(\mathcal{L}G) \rightarrow L_{(p/2)'}(\mathcal{L}G; L_\infty) \right\|_{\text{cb}}^{\frac{1}{2}}.$$

The row case is similar. The discussion of Corollary 1.4 generalizes to c.b. norms.

2. Spectral multipliers

2.1. Ultracontractivity

Let (\mathcal{M}, τ) be a noncommutative measure space and consider a Markov semigroup $\mathcal{S} = (S_t)_{t \geq 0}$ defined on it. Given a positive function $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ and $1 \leq p < q \leq \infty$, we say that \mathcal{S} satisfies the $\mathbf{R}_\Phi^{p,q}$ ultracontractivity property when

$$(\mathbf{R}_\Phi^{p,q}) \quad \|S_t : L_p(\mathcal{M}) \rightarrow L_q(\mathcal{M})\| \lesssim \frac{1}{\Phi(\sqrt{t})^{\frac{1}{p} - \frac{1}{q}}} \quad \forall t > 0.$$

Similarly, \mathcal{S} has the $\text{CBR}_\Phi^{p,q}$ property when the above estimate holds for the c.b. norm of $S_t : L_p(\mathcal{M}) \rightarrow L_q(\mathcal{M})$. These inequalities have been extensively studied for commutative measure spaces [43, Chapter 1]. In the theory of Lie groups with an invariant Riemannian metric (equipped with the heat semigroup generated by the invariant Laplacian) ultracontractivity holds for the function $\Phi(t) = \mu(B_t(e))$ which assigns the volume of a ball for a given radius. Influenced by that, we will interpret the above-defined properties as a way of describing the “growth of the balls” in the noncommutative geometry determined by $\mathcal{S} = (S_t)_{t \geq 0}$. For that reason, we will work with *doubling functions* Φ . Doubling functions are increasing functions $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\Phi(0) = 0$ and satisfying

$$\sup_{t > 0} \left\{ \frac{\Phi(2t)}{\Phi(t)} \right\} < \infty.$$

The doubling condition for Φ is a natural requirement since metric measure spaces (Ω, μ, d) with $\Phi_x(t) = \mu(B_x(t))$ uniformly doubling in x constitute an adequate setting for performing harmonic analysis in commutative measure spaces. Given a Markov semigroup $\mathcal{S} = (S_t)_{t \geq 0}$ over a noncommutative measure space (\mathcal{M}, τ) , let us recall the following:

- i) If \mathcal{S} satisfies $\mathbf{R}_\Phi^{p_0, q_0}$, it satisfies $\mathbf{R}_\Phi^{p, q}$ for $1 \leq p_0 \leq p < q \leq q_0 \leq \infty$.
- ii) If Φ is doubling and \mathcal{S} satisfies $\mathbf{R}_\Phi^{p_0, q_0}$ for some $1 \leq p_0 < q_0 \leq \infty$, then it satisfies $\mathbf{R}_\Phi^{p, q}$ for $1 \leq p \leq q \leq \infty$.

The same holds for the $\text{CBR}_\Phi^{p_0, q_0}$ ultracontractivity property. The proof follows the same lines than [43, Theorem II.1.3]. In the noncommutative setting a similar result is stated in [17, Lemma 1.1.2] for $\Phi(t) = t^D$. As a consequence, all the ultracontractivity properties $\mathbf{R}_\Phi^{p, q}$ are equivalent for doubling Φ . We shall denote them simply by \mathbf{R}_Φ and similarly CBR_Φ . As a corollary, we obtain that if \mathcal{M} is an abelian von Neumann algebra $\text{CBR}_\Phi^{p, q}$ and $\mathbf{R}_\Phi^{p, q}$ are equivalent for doubling Φ since $\mathbf{R}_\Phi^{p, q}$ is equivalent to $\mathbf{R}_\Phi^{p, \infty}$ and any bounded map into an abelian C^* -algebra is completely bounded. For any doubling function Φ we may define its doubling dimension D_Φ as

$$D_\Phi = \log_2 \sup_{t > 0} \left\{ \frac{\Phi(2t)}{\Phi(t)} \right\}.$$

It is quite simple to show that any doubling $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ admits upper/lower polynomial bounds for large/small values of $t > 0$. More precisely, we have the bounds

$$(2.1) \quad \begin{aligned} \Phi(t) &\lesssim_{(D_\Phi)} t^{D_\Phi} \Phi(1) \quad \text{when } t > 1, \\ \Phi(t) &\gtrsim_{(D_\Phi)} t^{D_\Phi} \Phi(1) \quad \text{when } t \leq 1. \end{aligned}$$

Of course, the converse of this assertion is false. Whenever a Markovian semigroup \mathcal{S} satisfies \mathbf{R}_Φ (resp. \mathbf{CBR}_Φ) for doubling Φ we will call D_Φ the *Sobolev dimension* (resp. *c.b. Sobolev dimension*) of (\mathcal{M}, τ) with respect to \mathcal{S} . The reason for this name is based on the well-known relation between ultracontractivity estimates for a Markov semigroup and Sobolev embedding estimates for its infinitesimal generator. One of the first contributions to that relation is in the work of Varopoulos, who proved in [42] that when $\Phi(t) = t^D$ the property \mathbf{R}_Φ is equivalent to a whole range of Sobolev type estimates for the infinitesimal generator of the semigroup. See also [43] for more on that topic. Whenever $\Phi(t) = t^D$ we will denote the ultracontractivity properties by \mathbf{R}_D or \mathbf{CBR}_D . By adding a zero, like $\mathbf{R}_\Phi(0)$, we will mean that the inequality $\mathbf{R}_\Phi^{p,q}$ is satisfied for $t \leq 1$ for every $p < q$. This notation is borrowed from [43, II.5]. Recall that if \mathcal{S} satisfies \mathbf{R}_Φ (resp. \mathbf{CBR}_Φ) for some doubling function Φ then, by the polynomial bounds in (2.1), we have $\mathbf{R}_{D_\Phi}(0)$ (resp. $\mathbf{CBR}_{D_\Phi}(0)$).

Our characterization of co-polynomial growth in Section 3 below requires the following equivalence for Sobolev-type inequalities in term of the ultracontractivity properties $\mathbf{R}_D(0)$. We did not find the proposition below in the literature, but it could be well-known to experts. We include a sketch of the proof.

PROPOSITION 2.1. – *Let \mathcal{S} be a submarkovian semigroup acting on a noncommutative measure space (\mathcal{M}, τ) . Let A denote its infinitesimal generator. Then, the following properties are equivalent :*

- i) for every $\varepsilon > 0$, \mathcal{S} satisfies the $\mathbf{R}_{D+\varepsilon}(0)$ property,
- ii) for every $\varepsilon > 0$, we have that

$$\|(\mathbf{1} + A)^{-D/4-\varepsilon} : L_2(\mathcal{M}) \rightarrow \mathcal{M}\| \lesssim_{(\varepsilon)} 1.$$

Similarly, $\mathcal{S} \in \mathbf{CBR}_{D+\varepsilon}(0)$ for all $\varepsilon > 0$ iff $(\mathbf{1} + A)^{-s} : L_2(\mathcal{M}) \xrightarrow{\text{cb}} \mathcal{M}$ for all $\varepsilon > 0$.

Proof. – The implication i) \Rightarrow ii) follows from the identity

$$(\mathbf{1} + A)^{-s}(x) = \frac{1}{\Gamma(s)} \left(\int_{\mathbb{R}_+} t^s e^{-t} S_t(x) \frac{dt}{t} \right).$$

The integral in $[0, 1]$ may be estimated applying the $\mathbf{R}_D(0)$ property, whereas the integral for $t > 1$ is easily estimated using the semigroup law. This gives the desired implication. For the converse, we now take $s = D/4 + \varepsilon$ and use that $\|f(A)\|_{\mathcal{B}(L_2)} \leq \|f\|_\infty$

$$\begin{aligned} \|S_t : L_2(\mathcal{M}) \rightarrow \mathcal{M}\| &= \left\| (\mathbf{1} + A)^{-\frac{s}{2}} (\mathbf{1} + A)^{\frac{s}{2}} S_t \right\|_{\mathcal{B}(L_2(\mathcal{M}), \mathcal{M})} \\ &\leq \left\| (\mathbf{1} + A)^{-\frac{s}{2}} \right\|_{\mathcal{B}(L_2(\mathcal{M}), \mathcal{M})} \left\| (\mathbf{1} + A)^{\frac{s}{2}} S_t \right\|_{\mathcal{B}(L_2)} \\ &\lesssim_{(\varepsilon, s)} \left(\frac{s}{2} \right)^{\binom{s}{2}} e^{-\frac{s}{2}} \frac{e^t}{t^{\frac{s}{2}}}. \end{aligned} \quad \square$$

REMARK 2.2. – Observe that if $R_D(0)$ is satisfied then ii) also holds. Nevertheless the converse is not true since the norm $\|S_t : L_1(\mathcal{M}) \rightarrow \mathcal{M}\|$ could be comparable to, say, $t^D(1 + \log(t))$ for $0 \leq t \leq 1$. The original result proved by Varopoulos [42] established a equivalence between $R_D(0)$ and the bounds

$$(\mathbf{1} + A)^{-s} : L_p(\mathcal{M}) \rightarrow L_{\frac{pn}{n-sp}}(\mathcal{M})$$

for every $0 \leq s < n/p$. When $s > n/p$ the image space of $L_p(\mathcal{M})$ is certainly much smaller than $L_\infty(\mathcal{M})$, for example in \mathbb{R}^n with the usual Laplacian the image space lies inside spaces of Hölder functions. Therefore, by describing the behavior of $(\mathbf{1} + A)^{-s}$ in $L_\infty(\mathcal{M})$ we lose information and we can no longer recover $R_D(0)$.

We will denote by $W_A^{p,s}(\mathcal{M})$, or simply $W^{p,s}(\mathcal{M})$ when the semigroup $S_t = e^{-tA}$ can be understood from the context, the closed domain in $L_p(\mathcal{M})$ of the unbounded operator $(\mathbf{1} + A)^{s/2}$, with norm given by

$$\|x\|_{W_A^{p,s}} = \|(\mathbf{1} + A)^{s/2} f\|_p.$$

These are called the *fractional Sobolev spaces* associated with \mathcal{S}^c . They satisfy the natural interpolation identities. Namely, if we set $1/p_3 = (1 - \theta)/p_1 + \theta/p_2$ we get

$$\begin{aligned} [W_A^{p_1,s}(\mathcal{M}), W_A^{p_2,s}(\mathcal{M})]_\theta &\simeq W_A^{p_3,s}(\mathcal{M}), \\ [W_A^{p_1,s_1}(\mathcal{M}), W_A^{p_2,s_2}(\mathcal{M})]_\theta &\simeq W_A^{p_3,s_1\theta+s_2(1-\theta)}(\mathcal{M}). \end{aligned}$$

Point ii) in Proposition 2.1 may be rephrased as $W_A^{2,s}(\mathcal{M}) \subset \mathcal{M}$ for every $s > D/2$.

2.1.1. *L₂ bounds for $\mathcal{CB}(L_2(\mathcal{L}G), \mathcal{L}G)$ multipliers.* – We shall work extensively with Markovian convolution semigroups over $\mathcal{L}G$ with the CBR_Φ ultracontractivity property for doubling Φ . In general, determining the c.b. norm of a multiplier between general L_p spaces is a problem that nobody expects to be solvable with a closed formula. Despite that, we can obtain characterizations in some particular cases. One of these cases is that of the c.b. multipliers $T_m : L_2(\mathcal{L}G) \rightarrow \mathcal{L}G$. That will allow us to express the $\text{CBR}_\Phi^{2,\infty}$ property of $\mathcal{S}^c = (T_{e^{-t\psi}})_{t \geq 0}$ as a condition over ψ . The next theorem is probably known to experts. Since we could not find it in the literature, we include it here for the sake of completeness.

THEOREM 2.3. – *If T denotes the map $m \mapsto T_m :$*

- i) $T : L_2^c(G) \rightarrow \mathcal{CB}(L_2^c(\mathcal{L}G), \mathcal{L}G)$ *is a complete isometry.*
- ii) $T : L_2^r(G) \rightarrow \mathcal{CB}(L_2^r(\mathcal{L}G), \mathcal{L}G)$ *is a complete isometry.*

The image of T is the set of multipliers $T_m : L_2^\dagger(\mathcal{L}G) \xrightarrow{\text{cb}} \mathcal{L}G$ for $\dagger \in \{c, r\}$ resp.

Proof. – Let V and W be operator spaces and pick $x \otimes y \in V^* \otimes W^*$. According to [32, Theorem 4.1] the map $\mathcal{J}_{x \otimes y}(w) = x\langle y, w \rangle$ extends linearly to an isomorphism $\mathcal{J} : (V \widehat{\otimes} W)^* \rightarrow \mathcal{CB}(W, V^*)$. Using the pairing $\langle \cdot, \cdot \rangle : L_2^r(\mathcal{L}G) \times L_2^c(\mathcal{L}G) \rightarrow \mathbb{C}$ given by $\langle y, w \rangle = \tau(y \sigma w)$ we obtain as a consequence that

$$\mathcal{J}_{\delta z}(w) = (\text{Id} \otimes \tau)(\delta z (\mathbf{1} \otimes \sigma w)) = z \star w,$$


where δz denotes the comultiplication map acting on z . This yields

$$\|T_m : L_2^\dagger(\mathcal{L}G) \rightarrow \mathcal{L}G\|_{\text{cb}} = \|\delta\lambda(m)\|_{(\mathcal{L}G_* \widehat{\otimes} L_2^\dagger(\mathcal{L}G))^*},$$

where $\dagger \in \{r, c\}$ is either the row or the column o.s.s. We now claim that the natural map

$$\iota : L_\infty(\mathcal{L}G; L_2^{\dagger\text{op}}(\mathcal{L}G)) \hookrightarrow (\mathcal{L}G_* \widehat{\otimes} L_2^\dagger(\mathcal{L}G))^*$$

is a complete isometry with $\dagger^{\text{op}} = r$ for $\dagger = c$ and vice-versa. This is all what is needed to complete the argument since we have the following commutative diagram of complete isometries



Let us therefore justify our claim. According to [8]

$$(\mathcal{L}G_* \widehat{\otimes} L_2^\dagger(\mathcal{L}G))^* \simeq \mathcal{L}G \otimes_{\mathcal{F}} L_2^{\dagger\text{op}}(\mathcal{L}G)$$

where $\otimes_{\mathcal{F}}$ stands for the Fubini tensor product of dual operator spaces. Bear in mind that if V^* and W^* are dual operator spaces, there are weak-* continuous embeddings $V^* \subset \mathcal{B}(\mathcal{H}_1)$ and $W^* \subset \mathcal{B}(\mathcal{H}_2)$ and we can define the weak-* spatial tensor product $V^* \overline{\otimes} W^*$ as

$$V^* \overline{\otimes} W^* = \overline{(V^* \otimes W^*)}^{w*}.$$

Such construction is representation independent and $V^* \overline{\otimes} W^*$ embeds completely isometrically in $V^* \otimes_{\mathcal{F}} W^*$. Since the column and row embeddings of $L_2(\mathcal{L}G)$ into $\mathcal{B}(L_2(\mathcal{L}G))$ are weak-* continuous, $L_\infty(\mathcal{L}G; L_2^{\dagger\text{op}}(\mathcal{L}G)) = \mathcal{L}G \overline{\otimes} L_2^{\dagger\text{op}}(\mathcal{L}G)$. This proves that ι is a complete isometry and so is the map $m \mapsto T_m = \mathcal{J} \iota \delta \lambda(m)$. \square

REMARK 2.4. – Since $L_2^r(\mathcal{L}G)$ and $L_2^c(\mathcal{L}G)$ are isometric as Banach spaces, the norms for multipliers in $\mathcal{CB}(L_2^r(\mathcal{L}G), \mathcal{L}G)$ and $\mathcal{CB}(L_2^c(\mathcal{L}G), \mathcal{L}G)$ coincide too, even if their matrix amplifications do not. Indeed we obtain that

$$\|T_m\|_{\mathcal{CB}(L_2^r(\mathcal{L}G), \mathcal{L}G)} = \|m\|_{L_2(G)} = \|T_m\|_{\mathcal{CB}(L_2^c(\mathcal{L}G), \mathcal{L}G)}.$$

For non-hyperfinite $\mathcal{L}G$, the space of Fourier multipliers in $\mathcal{CB}(L_2(\mathcal{L}G), \mathcal{L}G)$, may be difficult to describe as an operator space. Nevertheless, as a consequence of the above identities, its underlying Banach space is the Hilbert space $L_2(G)$.

REMARK 2.5. – As a consequence of the above, if G is a group and $\mathcal{S} = (T_{e^{-t\psi}})_{t \geq 0}$ is a semigroup of Fourier multipliers satisfying $\text{CBR}_\Phi^{2, \infty}$ for some function Φ , then G is amenable. To see it just notice that $e^{-t\psi} \in L_2(G)$ and so $e^{-2t\psi} \in L_1(G)$ for all $t > 0$. But a group is amenable iff there is a sequence of integrable positive type functions converging to 1 uniformly in compacts.

2.2. Standard assumptions

Let $\mathcal{L}G_+^\wedge$ denote the extended positive cone of $\mathcal{L}G$. As it will become clear along the paper, we shall treat unbounded operators X in $\mathcal{L}G_+^\wedge$ as noncommutative or quantized metrics over $\mathcal{L}G$. Note that if G is LCH and abelian, any translation-invariant metric over its dual group can be associated with the positive function $\Delta : \chi \mapsto d(\chi, e)$. The metric conditions impose that Δ is symmetric, does not vanish outside e and $\Delta(\chi_1\chi_2) \leq \Delta(\chi_1) + \Delta(\chi_2)$. Here we will only require X to be symmetric, i.e., to satisfy $\sigma X = X$. Recall that the anti-automorphism σ extends to $\mathcal{L}G_+^\wedge$. Following the intuition relating symmetric operators in $\mathcal{L}G_+^\wedge$ to metrics, we will say that $X \in \mathcal{L}G_+^\wedge$ is doubling iff the function $\Phi_X(r) = \tau(\chi_{[0,r]}(X))$ is doubling. When the dependency on the operator X can be understood from the context we will just write Φ . In a similar fashion, we will say that X satisfies the L_p -Hardy-Littlewood maximal property when

$$(HL_p) \quad \left\| \sup_{r \geq 0}^+ \left\{ \frac{\chi_{[0,r]}(X)}{\Phi_X(r)} \star u \right\} \right\|_p \lesssim \|u\|_p.$$

If we say that X has the HL property, omitting the dependency on p , we mean that the HL property is satisfied for every $1 < p \leq \infty$, with constants depending on p . When the property HL_p holds uniformly for all matrix amplifications, we will say that X satisfies the *completely bounded Hardy-Littlewood maximal property* ($CBHL_p$ in short). Let $\psi : G \rightarrow \mathbb{R}_+$ be a conditionally negative length generating a semigroup \mathcal{S} . We will say that \mathcal{S} has L_2 Gaussian bounds with respect to X when there is some $\beta > 0$ such that

$$(L_2GB) \quad \tau \left\{ \chi_{[r,\infty)}(X) |\lambda(e^{-t\psi})|^2 \right\} \lesssim \frac{e^{-\beta \frac{r^2}{t}}}{\Phi_X(\sqrt{t})}.$$

DEFINITION 2.6. – A triple $(\mathcal{L}G, \mathcal{S}, X)$, where \mathcal{S} is a Markov semigroup of Fourier multipliers generated by $\psi : G \rightarrow \mathbb{R}_+$ and $X \in (\mathcal{L}G)_+^\wedge$, is said to satisfy the standard assumptions when

- i) X is symmetric and doubling.
- ii) \mathcal{S} has L_2GB with respect to X .
- iii) X satisfies the $CBHL$ property.

Since $\mathcal{L}G$ is determined by G and \mathcal{S} by ψ we shall often write (G, ψ, X) instead.

REMARK 2.7. – If \mathcal{S} has L_2GB then it admits $CBR_{\Phi_X}^{2,\infty}$ ultracontractivity. Namely if we take $r = 0$ in (L_2GB) , it follows from Theorem 2.3 and Remark 2.4. If X is in addition doubling, \mathcal{S} has the whole range of ultracontractivity properties CBR_{Φ_X} .

2.2.1. *Stability under Cartesian products.* – It is interesting to note that the standard assumptions are stable under certain algebraic operations, the most trivial of them is probably the Cartesian product. Stability under crossed products also holds under natural conditions, see Remark 2.10 below.

LEMMA 2.8. – Assume that

$$\mathcal{S}^j = (S_{\omega_j}^j)_{\omega_j \in \Omega_j} : L_p(\mathcal{M}_j) \rightarrow L_p(\mathcal{M}_j; L_\infty(\Omega_j))$$

is completely positive for $j \in \{1, 2\}$. Then $\mathcal{S}^1 \otimes \mathcal{S}^2$ is also c.p. and

$$\begin{aligned} \left\| \mathcal{S}^1 \otimes \mathcal{S}^2 : L_p(\mathcal{M}_1 \overline{\otimes} \mathcal{M}_2) \rightarrow L_p(\mathcal{M}_1 \overline{\otimes} \mathcal{M}_2; L_\infty(\Omega_1) \otimes_{\min} L_\infty(\Omega_2)) \right\|_{\text{cb}} \\ \lesssim \prod_{j \in \{1,2\}} \left\| \mathcal{S}^j : L_p(\mathcal{M}_j) \rightarrow L_p(\mathcal{M}_j; L_\infty(\Omega_j)) \right\|_{\text{cb}}. \end{aligned}$$

Proof. – It follows from $\mathcal{S}^1 \otimes \mathcal{S}^2 = (\mathcal{S}^1 \otimes \text{Id}) \circ (\text{Id} \otimes \mathcal{S}^2)$ and (1.1), details are omitted. \square

THEOREM 2.9. – *Let (G_j, ψ_j, X_j) be triples satisfying the standard assumptions for $j = 1, 2$ and consider the Cartesian product $G = G_1 \times G_2$. Then (G, ψ, X) also satisfies the standard assumptions with the c.n.length $\psi(g_1, g_2) = \psi_1(g_1) + \psi_2(g_2)$ and $X \in \mathcal{L}G^{\wedge}_+$ determined by the formula $X^2 = X_1^2 \otimes \mathbf{1} + \mathbf{1} \otimes X_2^2$.*

Proof. – Proving that X is doubling and that the semigroup generated by ψ has Gaussian bounds amount to a trivial calculation. Indeed, Φ_X is controlled from the inequalities $\chi_{[0,r/2]}(a)\chi_{[0,r/2]}(b) \leq \chi_{[0,r]}(a+b) \leq \chi_{[0,r]}(a)\chi_{[0,r]}(b)$, which are valid for positive and commuting operators a, b . On the other hand, the $L_2\text{GB}$ follow similarly from the inequality $\chi_{[r,\infty)}(a+b) \leq \chi_{[r/2,\infty)}(a) + \chi_{[r/2,\infty)}(b)$. Let us now justify the CBHL property. Let $m : L_p(\mathcal{L}G; L_\infty \otimes_{\min} L_\infty) \rightarrow L_p(\mathcal{L}G; L_\infty)$ be the map given by $m(x \otimes f \otimes g) = x \otimes fg$, which is c.p. By Lemma 2.8

$$\mathcal{R}^1 \otimes \mathcal{R}^2 = (R_s^1 \otimes R_t^2)_{s,t \geq 0} : L_p(\mathcal{L}G) \rightarrow L_p(\mathcal{L}G; L_\infty(ds) \otimes_{\min} L_\infty(dt)),$$

where $R_s^j(x) = \Phi_{X_j}(s)^{-1} \chi_{[0,s]}(X_j) \star x$ is c.p. As a consequence $m \circ (\mathcal{R}^1 \otimes \mathcal{R}^2)$ is also completely positive. Therefore, by the doubling property we obtain the following estimate

$$\left(\frac{\chi_{[0,r]}(X)}{\Phi_X(r)} \star x \right)_{r \geq 0} \lesssim_{(D_{\Phi_1}, D_{\Phi_2})} \left(\frac{\chi_{[0,r]}(X_1)}{\Phi_{X_1}(r)} \otimes \frac{\chi_{[0,r]}(X_2)}{\Phi_{X_2}(r)} \star x \right)_{r \geq 0} = m \circ (\mathcal{R}^1 \otimes \mathcal{R}^2)(x)$$

for $x \geq 0$. This is all what we need to reduce CBHL of X to that of X_1 and X_2 . \square

REMARK 2.10. – Let H and G be LHC unimodular groups and $\theta : G \rightarrow \text{Aut}(H)$ be a measure preserving action. Let (H, ψ_1, X_1) and (G, ψ_2, X_2) be triples satisfying the standard assumptions. It is possible to prove that, under certain invariance conditions on X_1 and ψ_1 , the semidirect product $K = H \rtimes_{\theta} G$ satisfies the standard assumptions for some $X \in \mathcal{L}K^{\wedge}_*$ and certain c.n. length function $\psi : K \rightarrow \mathbb{R}_+$ built up from X_1, X_2 and ψ_1, ψ_2 respectively. Since the techniques required to prove this result are quite involved and of independent interest, we postpone its proof to a forthcoming paper were we shall explore other applications involving Bochner-Riesz summability and related topics.

2.3. Hörmander-Mikhlin criteria

In this subsection we shall give a proof of Theorem B i) by means of a suitably chosen max-square decomposition. The key is to prove that, if $B_t = \lambda(m \eta(t\psi))$, then

$$(2.2) \quad B_t = \underbrace{B_t \left(\mathbf{1} + \frac{X^2}{t} \right)^{\frac{\gamma}{2}}}_{\Sigma_t} \underbrace{\Phi(\sqrt{t})^{\frac{1}{2}} \Phi(\sqrt{t})^{-\frac{1}{2}} \left(\mathbf{1} + \frac{X^2}{t} \right)^{-\frac{\gamma}{2}}}_{M_t}$$

is a square-max decomposition for $\gamma > D_{\Phi}/2$. Breaking the symbol m into its real and imaginary parts and using Remark 1.5, we obtain a max-square decomposition by placing

the smoothing factor $(1 + X^2/t)^{\gamma/2}$ on the left hand side of B_t . The proof of the maximal inequality consists in expressing the maximal operator as a linear combination of Hardy-Littlewood maximal operators associated to X and apply (1.2). For the square estimate we will use the smoothness condition.

LEMMA 2.11. – Assume that $F_t \in C_0(\mathbb{R}_+)$ is a family of bounded variation functions parametrized by $t > 0$. Let ∂F_t be its Lebesgue-Stieltjes derivative and $|\partial F_t(\lambda)|$ its absolute variation, then for every doubling operator X , we have:

$$\left\| \left(\sup_{t \geq 0}^+ F_t(X) \star x \right) \right\|_{L_p} \leq \left(\sup_{t > 0} \|\Phi\|_{L_1(|dF_t|)} \right) \left\| \left(\sup_{r > 0}^+ \frac{\chi_{[0,r]}(X)}{\Phi(r)} \star x \right) \right\|_{L_p}.$$

Proof. – By integration by parts we have that

$$\begin{aligned} F_t(s) &= \int_{\mathbb{R}_+} F_t(r) d\delta_s(r) = \int_{\mathbb{R}_+} F_t(r) \partial \chi_{(s,\infty)}(r) \\ &= - \int_{\mathbb{R}_+} \chi_{(s,\infty)}(r) \partial F_t(r) = - \int_{\mathbb{R}_+} \frac{\chi_{[0,r]}(s)}{\Phi(r)} \Phi(r) \partial F_t(r). \end{aligned}$$

By functional calculus, the same holds for $F_t(X)$. Applying (1.2) ends the proof. \square

According to Theorem A, the right choice for the square-max decomposition is given by $F_t(s) = |M_t|^2(s) = \Phi(\sqrt{t})^{-1} (1 + s^2/t)^{-\gamma}$. It will suffice to pick here $\gamma > D_\Phi/2$, the condition in Theorem B i) will be justified later on. In order to prove the finiteness of the maximal bound in Theorem A, we just need to verify the condition of Lemma 2.11 for this concrete function.

LEMMA 2.12. – For any doubling Φ , we find

$$\int_{\mathbb{R}_+} \Phi(s) \left| \frac{d}{ds} \left(1 + \frac{s^2}{t} \right)^{-\frac{D_\Phi + \varepsilon}{2}} \right| ds \lesssim_{(D_\Phi, \varepsilon)} \Phi(\sqrt{t}).$$

Proof. – Changing variables $s \mapsto \sqrt{t}v$, we obtain

$$\begin{aligned} \int_{\mathbb{R}_+} \Phi(s) \left| \frac{d}{ds} \left(1 + \frac{s^2}{t} \right)^{-\frac{D_\Phi + \varepsilon}{2}} \right| ds &\sim_{(D_\Phi)} \int_{\mathbb{R}_+} \Phi(s) \left(1 + \frac{s^2}{t} \right)^{-\frac{D_\Phi + 2 + \varepsilon}{2}} \frac{2s}{t} ds \\ &= \int_{\mathbb{R}_+} \Phi(\sqrt{t}\sqrt{v}) (1+v)^{-\frac{D_\Phi + 2 + \varepsilon}{2}} dv \\ &= \left(\int_0^1 + \sum_{k=0}^{\infty} \int_{4^k}^{4^{k+1}} \right) = A + \sum_{k=0}^{\infty} B_k. \end{aligned}$$

The monotonicity of Φ gives $A \leq \Phi(\sqrt{t})$, while its doublingness yields

$$\begin{aligned} B_k &\leq \Phi(\sqrt{t}) 2^{D_\Phi(k+1)} \int_{4^k}^{4^{k+1}} (1+v)^{-\frac{D_\Phi + 2 + \varepsilon}{2}} dv \\ &\sim_{(D_\Phi)} \Phi(\sqrt{t}) 2^{D_\Phi(k+1)} 2^{-(D_\Phi + \varepsilon)k} \\ &\sim_{(D_\Phi)} \Phi(\sqrt{t}) 2^{-\varepsilon k}. \end{aligned}$$

Since the sequence of B_k s is summable, we have proved the desired estimate. \square

For the estimate of the square part, let us start by extending the Gaussian bounds to the complex half-plane $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Re}(z) > 0\}$. We need the following version of the Phragmen-Lindelöf theorem, see [7] for the proof.

THEOREM 2.13. – *If F is analytic over \mathbb{H} and satisfies*

$$\begin{aligned} |F(|z|e^{i\theta})| &\lesssim (|z| \cos \theta)^{-\beta}, \\ |F(|z|)| &\lesssim |z|^{-\beta} \exp(-\alpha|z|^{-\rho}), \end{aligned}$$

for some $\alpha, \beta > 0$ and $0 < \rho \leq 1$, then we find the following estimate

$$|F(|z|e^{i\theta})| \lesssim_{(\beta)} (|z| \cos \theta)^{-\beta} \exp\left(-\frac{\alpha\rho}{2}|z|^{-\rho} \cos \theta\right).$$

We may now generalize the Gaussian L_2 -bounds to the complex half-plane.

PROPOSITION 2.14. – *Let G be a unimodular group, $\psi : G \rightarrow \mathbb{R}_+$ a c.n. length and $X \in \mathcal{L}G_\dagger$ a doubling operator satisfying $L_2\text{GB}$. If we set $h_z = \lambda(e^{-z\psi})$, the following bound holds for every $z \in \mathbb{H}$*

$$\tau\left\{\chi_{[r,\infty)}(X)|h_z|^2\right\} \lesssim \frac{1}{\Phi(\sqrt{\operatorname{Re}\{z\}})} e^{-\frac{\beta}{2}\frac{r^2}{|z|} \frac{\operatorname{Re}\{z\}}{|z|}}.$$

Proof. – Let x be an element of $L_2(\mathcal{L}G)$ with $\|x\|_2 \leq 1$. Assume in addition that $x = px$ for $p = \chi_{[r,\infty)}(X)$. Then we define G_x as the following holomorphic function

$$G_x(z) = e^{-\frac{z}{t}} \Phi(\sqrt{t}) \tau(h_z x)^2.$$

Then, the estimate below holds in \mathbb{H}

$$\begin{aligned} |G_x(z)| &= e^{-\frac{\operatorname{Re}\{z\}}{t}} \Phi(\sqrt{t}) |\tau(h_z x)|^2 \leq e^{-\frac{\operatorname{Re}\{z\}}{t}} \Phi(\sqrt{t}) \tau(|h_z|^2) \\ &= e^{-\frac{|z|\cos\theta}{t}} \Phi(\sqrt{t}) \|h_{\operatorname{Re}\{z\}}\|_{L_2(\mathcal{L}G)}^2 \lesssim e^{-\frac{\operatorname{Re}\{z\}}{t}} \Phi(\sqrt{t}) / \Phi(\sqrt{\operatorname{Re}\{z\}}). \end{aligned}$$

Note that the second identity above follows from Plancherel theorem and the last inequality from $L_2\text{GB}$ for $r = 0$. On the other hand, since Φ is doubling it satisfies $\Phi(s(1+r)) \lesssim \Phi(s)(1+r)^{D_\Phi}$ for every $r > 0$ and

$$|G_x(z)| \lesssim e^{-\frac{\operatorname{Re}\{z\}}{t}} \frac{\Phi(\sqrt{t})}{\Phi(\sqrt{\operatorname{Re}\{z\}})} \lesssim e^{-\frac{\operatorname{Re}\{z\}}{t}} \left(1 + \frac{\sqrt{t}}{\sqrt{\operatorname{Re}\{z\}}}\right)^{D_\Phi} \lesssim \left(\frac{t}{|z|\cos\theta}\right)^{\frac{D_\Phi}{2}},$$

by using that $e^{-s^2}(1+1/s)^a \lesssim (1/s)^a$ in the last inequality. We also have

$$\begin{aligned} |G_x(|z|)| &= e^{-\frac{|z|}{t}} \Phi(\sqrt{t}) |\tau(h_{|z|} x)|^2 \\ &\leq e^{-\frac{|z|}{t}} \Phi(\sqrt{t}) \tau\{p h_{|z|}^* h_{|z|}\} \lesssim e^{-\frac{|z|}{t}} \frac{\Phi(\sqrt{t})}{\Phi(\sqrt{|z|})} e^{-\beta\frac{r^2}{|z|}} \\ &\lesssim e^{-\frac{|z|}{t}} \left(1 + \frac{\sqrt{t}}{\sqrt{|z|}}\right)^{D_\Phi} e^{-\beta\frac{r^2}{|z|}} \lesssim \left(\frac{t}{|z|}\right)^{\frac{D_\Phi}{2}} e^{-\beta\frac{r^2}{|z|}}. \end{aligned}$$

The Phragmen-Lindelöf theorem allows us to combine both estimates, giving

$$|G_x(|z|e^{i\theta})| \lesssim t^{\frac{D_\Phi}{2}} (|z|\cos\theta)^{-\frac{D_\Phi}{2}} e^{-\frac{\beta r^2}{2} \frac{\cos\theta}{|z|}}.$$

Taking the supremum over all x with $\|x\|_2 \leq 1$ and $x = p x$ we get

$$\sup_x |G_x(z)| = e^{-\frac{\operatorname{Re}\{z\}}{t}} \Phi(\sqrt{t}) \tau(p|h_z|^2).$$

Our previous estimate then yields

$$e^{-\frac{\operatorname{Re}\{z\}}{t}} \Phi(\sqrt{t}) \tau(p|h_z|^2) \lesssim t^{\frac{D\Phi}{2}} (|z|\cos\theta)^{-\frac{D\Phi}{2}} e^{-\frac{\beta r^2 \cos\theta}{2|z|}}.$$

Choosing the parameter $t \geq 0$ to be $t = \operatorname{Re}\{z\}$ gives the desired estimate. □

LEMMA 2.15. – *If $X \in \mathcal{LG}_+^\wedge$ is doubling and $\psi : G \rightarrow \mathbb{R}_+$ has $L_2\text{GB}$, then*

$$\tau\left\{\left(1 + \frac{X^2}{t}\right)^\kappa |h_{t(1-i\xi)}|^2\right\}^{\frac{1}{2}} \lesssim_{(\kappa)} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}} (1 + |\xi|)^\kappa \quad \text{for all } \kappa > 0.$$

Proof. – Writing $z = t(1 - i\xi)$ in Proposition 2.14 gives

$$\tau\left(\chi_{[r,\infty)}(X)|h_z|^2\chi_{[r,\infty)}(X)\right) \lesssim \frac{1}{\Phi(\sqrt{t})} e^{-\frac{\beta}{2} \frac{r^2}{t} \frac{1}{(1+|\xi|^2)}}.$$

Using the spectral measure dE_X of X and since $(1 + s^2)^\kappa \lesssim_{(\kappa)} 1 + s^{2\kappa}$

$$\begin{aligned} \tau\left\{\left(1 + \frac{X^2}{t}\right)^\kappa |h_{t(1-i\xi)}|^2\right\} &\lesssim_{(\kappa)} \tau\{|h_{t(1-i\xi)}|^2\} + \tau\{|h_{t(1-i\xi)}|^2 t^{-\kappa} X^{2\kappa}\} \\ &\lesssim \frac{1}{\Phi(\sqrt{t})} + \underbrace{\tau\{|h_{t(1-i\xi)}|^2 \int_{\mathbb{R}_+} \left(\frac{s^2}{t}\right)^\kappa dE_X(s)\}}_A. \end{aligned}$$

To estimate the term A we use integration by parts

$$\begin{aligned} A &= \int_{\mathbb{R}_+} \left(\frac{s^2}{t}\right)^\kappa \tau\{|h_{t(1-i\xi)}|^2 dE_X(s)\} \\ &= \int_{\mathbb{R}_+} \left(\frac{s^2}{t}\right)^\kappa (-\partial_s) \tau\{|h_{t(1-i\xi)}|^2 \chi_{[s,\infty)}(X)\} \\ &= \int_{\mathbb{R}_+} \frac{d}{ds} \left(\frac{s^2}{t}\right)^\kappa \tau\{|h_{t(1-i\xi)}|^2 \chi_{[s,\infty)}(X)\} ds. \end{aligned}$$

In the second line, by $-\partial_s \tau\{|h_{t(1-i\xi)}|^2 \chi_{[s,\infty)}(X)\}$, we mean the Lebesgue-Stieltjes measure associated with the increasing function $g(s) = -\tau\{|h_{t(1-i\xi)}|^2 \chi_{[s,\infty)}(X)\}$ and the third line is just an application of the integration by parts formula for Lebesgue-Stieltjes integrals. A calculation gives the desired result

$$\begin{aligned} A &\lesssim \int_{\mathbb{R}_+} \left(\frac{2\kappa s^{2\kappa-1}}{t^\kappa}\right) \frac{1}{\Phi(\sqrt{t})} e^{-\frac{\beta}{2} \frac{s^2}{t} \frac{1}{(1+|\xi|^2)}} ds \\ &\sim_{(\kappa)} \frac{(1 + |\xi|^2)^\kappa}{\Phi(\sqrt{t})} \int_{\mathbb{R}_+} s^{2\kappa-1} e^{-\frac{\beta}{2} s^2} ds \sim_{(\kappa)} \frac{(1 + |\xi|)^{2\kappa}}{\Phi(\sqrt{t})}. \quad \square \end{aligned}$$

PROPOSITION 2.16. – *Let $B_t = \lambda(m(\psi)\eta_1(t\psi))$ where $\eta_1(z) = \eta(z) e^{-z}$ for some $\eta \in \mathcal{H}_0^\infty$. Assume also that X is a doubling operator satisfying $L_2\text{GB}$, then the following estimate holds for every $\delta > 0$ and $\kappa > 0$*

$$\tau\left\{\left(1 + \frac{X^2}{t}\right)^\kappa |B_t|^2\right\}^{\frac{1}{2}} \lesssim_{(\kappa,\delta)} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}} \|m(t^{-1}\cdot)\eta(\cdot)\|_{W^{2,\kappa+\frac{1+\delta}{2}}(\mathbb{R}_+)}.$$

Proof. – By Fourier inversion formula

$$m(s)\eta_1(ts) = \underbrace{m(s)\eta(ts)}_{m_t(ts)} e^{-ts} = \left(\frac{1}{2\pi} \int_{\mathbb{R}} \widehat{m}_t(\xi) e^{i\xi ts} d\xi \right) e^{-ts}.$$

Thus, by composing with ψ and applying the left regular representation

$$B_t = \frac{1}{2\pi} \int_{\mathbb{R}} \widehat{m}_t(\xi) h_{t(1-i\xi)} d\xi.$$

Triangular inequality for the L_2 -norm with weight $(1 + X^2/t)$ and Lemma 2.15 give

$$\begin{aligned} \tau \left\{ \left(1 + \frac{X^2}{t} \right)^\kappa |B_t|^2 \right\}^{\frac{1}{2}} &= \tau \left\{ \left(1 + \frac{X^2}{t} \right)^\kappa \left| \frac{1}{2\pi} \int_{\mathbb{R}} \widehat{m}_t(\xi) h_{t(1-i\xi)} d\xi \right|^2 \right\}^{\frac{1}{2}} \\ &\leq \frac{1}{2\pi} \int_{\mathbb{R}} |\widehat{m}_t(\xi)| \tau \left\{ \left(1 + \frac{X^2}{t} \right)^\kappa |h_{t(1-i\xi)}|^2 \right\}^{\frac{1}{2}} d\xi \\ &\lesssim_{(\kappa)} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}} \int_{\mathbb{R}} |\widehat{m}_t(\xi)| (1 + |\xi|)^{\kappa + \frac{1+\delta}{2}} (1 + |\xi|)^{-\frac{1+\delta}{2}} d\xi = A. \end{aligned}$$

Höder’s inequality in conjunction with the definition of Sobolev space then yield

$$\Phi(\sqrt{t})^{\frac{1}{2}} A \leq \left(\int_{\mathbb{R}} (1 + |\xi|)^{-(1+\delta)} d\xi \right)^{\frac{1}{2}} \|m(t^{-1} \cdot) \eta(\cdot)\|_{W^{2,\kappa + \frac{1+\delta}{2}}(\mathbb{R}_+)}$$

The integral above is dominated by $(1 + \delta^{-1})^{\frac{1}{2}}$ and the assertion follows. □

Proof of Theorem B i). – Let $B_t = \lambda(m(\psi)\eta_1(t\psi))$ with $\eta_1(s) = e^{-s}\eta(s)$ and $B_t = \Sigma_t M_t$ be the decomposition (2.2) with $\gamma > D_\Phi/2$. Since we are assuming X to be symmetric, we have that $\sigma|M_t|^2 = |M_t|^2$ and, by Lemma 2.11 and Lemma 2.12, M_t satisfies the maximal inequality of (SM_p) . By Proposition 2.16 we have that

$$\sup_{t>0} \|\Sigma_t\|_{L_2(\mathcal{Z}G)} \lesssim_{(\gamma)} \sup_{t>0} \|m(t^{-1} \cdot) \eta(\cdot)\|_{W^{2,\gamma + \frac{1+\delta}{2}}(\mathbb{R}_+)}.$$

Therefore $B_t = \Sigma_t M_t$ is a square-max decomposition. By similar means we obtain a max-square decomposition $B_t = M_t \Sigma_t$. Since our maximal bounds trivially extend to matrix amplifications, we may apply Theorem 1.3 in conjunction with Remark 1.6 to deduce complete bounds of our multiplier $T_{m \circ \psi}$ in both row and column Hardy spaces. Finally, arguing as in Corollary 1.4 and noticing that $m \circ \psi \equiv m(0)$ on the subgroup $G_0 = \{g \in G : \psi(g) = 0\}$, we deduce the assertion. □

REMARK 2.17. – It is interesting to observe that the proof given here can be adapted to the classical case. Indeed, let $S_t = e^{-tA}$ be a Markovian semigroup acting on $L_\infty(X, \mu)$. Assume further that the metric measure space (X, d_Γ, μ) , where d_Γ is the gradient metric [34, Definition 3.1], is doubling, i.e.,:

$$\operatorname{ess\,sup}_{x \in X} \sup_{r>0} \left\{ \frac{\mu(B_x(2r))}{\mu(B_x(r))} \right\} < \infty$$

and that its integral kernel $k_t(x, y)$ has Gaussian bounds with respect to the gradient distance, i.e.,:

$$\|\chi_{[r,\infty)}(d_\Gamma(x, \cdot)) k_t(x, \cdot)\|_2^2 \lesssim \frac{e^{-\beta \frac{r^2}{t}}}{\mu(B_x(\sqrt{t}))}.$$

In that case we can apply the well known covering arguments for doubling spaces to prove that the Hardy-Littlewood maximal operator is of weak type $(1, 1)$ and by interpolation the HL inequalities hold. Since (X, d_Γ, μ) is a doubling metric measure space with bounded Hardy-Littlewood maximal inequalities and Gaussian Bounds we can apply the results above to reprove the classical spectral Hörmander-Mikhlin theorem as stated in [41]. We shall consider this a new proof of the classical spectral Hörmander-Mikhlin. Interestingly, some of the steps of the proof are parallel to that of [41] even when the main idea of our approach is to use maximal inequalities instead of Calderón-Zygmund estimates for the kernels.

2.4. The q -Plancherel condition

In this subsection we shall refine our results by proving Theorem B ii). Our first task is to introduce the noncommutative form of the Plancherel condition assumed in the statement.

DEFINITION 2.18. – Let (\mathcal{M}, τ) be a noncommutative measure space and let \mathcal{S} be a submarkovian semigroup generated by A . We say that \mathcal{S} satisfies the completely bounded q -Plancherel condition, denoted by CBPlan_q^Φ , where Φ is some increasing function and $q \in (2, \infty]$, whenever

$$\|F(A)\|_{\mathcal{CB}(L_2(\mathcal{Z}G), \mathcal{Z}G)} \lesssim \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}} \|F(t^{-1} \cdot)\|_{L_q(\mathbb{R}_+)},$$

for every $t > 0$ and for every function $F : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\text{supp}(F) \subset [0, t^{-1}]$.

REMARK 2.19. – In the context of this paper $\mathcal{M} = \mathcal{Z}G$ for some LCH unimodular group G endowed with its canonical trace and $\mathcal{S} = (T_{e^{-t\psi}})_{t \geq 0}$ is a semigroup of convolution type. In that case $F(A) = T_{F(\psi)}$ and by Theorem 2.3 and Remark 2.4 we have that

$$\begin{aligned} \|T_{F(\psi)}\|_{\mathcal{CB}(L_2(\mathcal{Z}G), \mathcal{Z}G)} &= \|T_{F(\psi)}\|_{\mathcal{CB}(L_2^\dagger(\mathcal{Z}G), \mathcal{Z}G)} \\ &= \|T_{F(\psi)}\|_{\mathcal{CB}(L_2^\dagger(\mathcal{Z}G), \mathcal{Z}G)} = \|F(\psi)\|_{L_2(G)}. \end{aligned}$$

Thus, the CBPlan_q^Φ condition can be restated as a bound on the $\mathcal{CB}(L_2^\dagger(\mathcal{Z}G), \mathcal{Z}G)$ norm, where \dagger is either the column or the row o.s.s. of $L_2(\mathcal{Z}G)$, or as a bound in the $L_2(G)$ -norm of the symbol $F(\psi)$. Furthermore, since ψ determines \mathcal{S} we will sometimes say that ψ has the CBPlan_q^Φ .

For every F with $\text{supp}(F) \subset [0, t^{-1}]$ we have that $F(t^{-1} \cdot)$ is supported in $[0, 1]$. Using that $L_q([0, 1]) \subset L_p([0, 1])$, with contractive inclusion, we see that $\text{CBPlan}_p^\Phi \Rightarrow \text{CBPlan}_q^\Phi$ for $p \leq q$.

PROPOSITION 2.20. – Let (G, ψ) be a pair formed by a LCH unimodular group and a c.n. length. Let Φ be a doubling function. If ψ satisfies the ultracontractivity estimates $\text{CBR}_\Phi^{2, \infty}$ then it satisfies $\text{CBPlan}_\infty^\Phi$.

Proof. – We pick $s > 0$, to be chosen later, and notice that

$$F(\psi(g)) = F(\psi(g))e^{s\psi(g)}e^{-s\psi(g)} = G_s(\psi(g))e^{-s\psi}$$

where G_s is a bounded function with $\|G_s\|_\infty \leq \|F\|_\infty e^{s/t}$. Therefore

$$\|T_{F(\psi)}\|_{\mathcal{CB}(L_2(\mathcal{Z}G), \mathcal{Z}G)} = \|T_{G_s(\psi)}S_s\|_{\mathcal{CB}(L_2(\mathcal{Z}G), \mathcal{Z}G)}$$

$$\begin{aligned} &\leq \|T_{G_s(\psi)}\|_{\mathcal{CB}(L_2(\mathcal{Z}G))} \|S_s\|_{\mathcal{CB}(L_2(\mathcal{Z}G), \mathcal{Z}G)} \\ &\lesssim \|F\|_\infty e^{s/t} \Phi(\sqrt{s})^{-\frac{1}{2}}. \end{aligned}$$

Making $s = t$ and noticing that $\|F\|_\infty = \|F(t^{-1} \cdot)\|_\infty$ gives the desired result. □

The terminology of the q -Plancherel condition comes from the so-called spectral Plancherel measures which arise in the study of spectral properties of infinitesimal generators of Markovian semigroups over some measure spaces [35, 41]. In the case of a semigroup of Fourier multipliers generated by a c.n. length we can define the Plancherel measure μ_ψ , as the only σ -finite measure over \mathbb{R}_+ satisfying that for every $F \in C_c(\mathbb{R}_+)$

$$(2.3) \quad \|T_{F(\psi)}\|_{\mathcal{CB}(L_2(\mathcal{Z}G), \mathcal{Z}G)} = \left(\int_{\mathbb{R}_+} |F(s)|^2 d\mu_\psi(s) \right)^{\frac{1}{2}}.$$

It is trivial to see that $d\mu_\psi(r) = \partial_r \mu(\{g \in G : \psi(g) \leq r\})$, where ∂_r represents the Lebesgue-Stieltjes derivative of the increasing function $g(r) = \mu(\{g \in G : \psi(g) \leq r\})$.

2.4.1. *Characterization of the q -Plancherel condition.* – By formula (2.3) the $\mathcal{CB}(L_2(\mathcal{Z}G), \mathcal{Z}G)$ norm of $T_{F(\psi)}$ can be expressed as an integral of F . The following lemma (whose proof is straightforward and we shall omit) allows to express the CBPlan_q^Φ property as a $L_{(q/2)'(\mathbb{R}_+)}$ bound on μ_ψ .

LEMMA 2.21. – *Let (Ω, Σ) be a measurable space and consider two measures μ, ν on it. Assume in addition that μ is a positive measure. Then, we have the inequality*

$$(2.4) \quad \left| \int_{\Omega} f(\omega) d\nu(\omega) \right| \leq K \|f\|_{L_p(d\mu)}$$

if and only if $\nu \ll \mu$ and $\phi = d\nu/d\mu$ satisfies $\|\phi\|_{L_{p'}(d\mu)} \leq K$. Furthermore, the optimal K in (2.4) is precisely $\|\phi\|_{L_{p'}(d\mu)}$. If ν is also positive, it is enough for (2.4) to hold only for positive functions.

PROPOSITION 2.22. – *Let G be a LCH unimodular group equipped with a c.n. length $\psi : G \rightarrow \mathbb{R}_+$. Then, this pair satisfies the CBPlan_q^Φ property with respect to some increasing function $\Phi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ if and only if $d\mu_\psi(r) = \partial_r \mu\{g \in G : \psi(g) \leq r\}$ fulfills the following conditions:*

- i) $d\mu_\psi \ll dm$.
- ii) $\left\| \frac{d\mu_\psi}{dm} \chi_{[0,R]} \right\|_{L_{(q/2)'(\mathbb{R}_+)}} \lesssim \Phi(R^{-\frac{1}{2}})^{-1} R^{-\frac{2}{q}}$ for every $R > 0$.

Proof. – Let $t = 1/R$ and $G(s) = |F(s)|^2$. By (2.3), CBPlan_q^Φ is equivalent to

$$\begin{aligned} \int_0^R G(s) d\mu_\psi(s) &\lesssim \Phi(R^{-\frac{1}{2}})^{-1} \left(\int_0^1 |F(t^{-1}s)|^q ds \right)^{\frac{2}{q}} \\ &= \Phi(R^{-\frac{1}{2}})^{-1} R^{-\frac{2}{q}} \left(\int_0^R |G(s)|^{\frac{q}{2}} ds \right)^{\frac{2}{q}}. \end{aligned}$$

Then, the result follows applying Lemma 2.21 to $(\Omega, d\nu, d\mu) = (\mathbb{R}_+, d\mu_\psi, dm)$. □

The result above uses the crucial fact that the spectrum of the semigroup \mathcal{S} generated by ψ can be identified with G . Therefore, spectral properties of the semigroup can be translated into geometrical properties of G . It is also interesting to note that the characterization in Proposition 2.22 can be expressed as a bound for the size of the spheres associated to the pseudo-metric $d_\psi(g, h) = \psi(g^{-1}h)^{1/2}$.

2.4.2. *Stability under direct products.* – Consider two pairs (G_j, ψ_j) of LCH unimodular groups equipped with c.n. lengths for $j = 1, 2$. Then it is clear that $\psi : G_1 \times G_2 \rightarrow \mathbb{R}_+$ given by $\psi(g, h) = \psi_1(g) + \psi_2(h)$ is also a c.n. length. Notice that

$$\begin{aligned} \|T_{F(\psi)}\|_{\mathcal{CB}(L_2(\mathcal{I}G), \mathcal{I}G)}^2 &= \int_{G_1 \times G_2} |F(\psi_1(g) + \psi_2(h))|^2 d\mu_{G_1}(g) d\mu_{G_2}(h) \\ &= \int_{\mathbb{R}_+} \int_{\mathbb{R}_+} |F(\xi + \zeta)|^2 d\mu_{\psi_1}(\xi) d\mu_{\psi_2}(\zeta) \\ &= \int_{\mathbb{R}_+} |F(\xi)|^2 d(\mu_{\psi_1} * \mu_{\psi_2})(\xi). \end{aligned}$$

Thus, the Plancherel measure is $\mu_\psi = \mu_{\psi_1} * \mu_{\psi_2}$ and we obtain the following result.

THEOREM 2.23. – *Assume (G_j, ψ_j) satisfy $\text{CBPlan}_{q_j}^{\Phi_j}$ for $j = 1, 2$. Then the pair $(G_1 \times G_2, \psi)$ defined above satisfies the CBPlan_q^Φ property with $\Phi = \Phi_1 \Phi_2$ and with*

$$q = \max \left\{ 2, \left(\frac{1}{q_1} + \frac{1}{q_2} \right)^{-1} \right\}.$$

Proof. – The result is a simple consequence of Young's inequality for convolutions and we shall just sketch the argument for the (slightly more involved) case where $1/q_1 + 1/q_2 > 1/2$, so that $q = 2$. According to Proposition 2.22, it suffices to see that

$$\left\| \frac{d\psi_1}{dm} * \frac{d\psi_2}{dm} \right\|_{L_\infty(0, R)} \leq \frac{1}{R \Phi_1(R^{-1/2}) \Phi_2(R^{-1/2})}.$$

The $\text{CBPlan}_{q_1}^{\Phi_1}$ property of (G_1, ψ_1) implies

$$\left\| \frac{d\psi_1}{dm} * \frac{d\psi_2}{dm} \right\|_\infty \leq \left\| \frac{d\psi_1}{dm} \right\|_{(q_1/2)'} \left\| \frac{d\psi_2}{dm} \right\|_{q_1/2} \leq \frac{1}{R^{2/q_1} \Phi_1(R^{-1/2})} \left\| \frac{d\psi_2}{dm} \right\|_{q_1/2}.$$

Now, since $1/q_1 + 1/q_2 > 1/2$ it turns out that

$$\frac{1}{q_1/2} = \frac{1}{(q_2/2)'} + \frac{1}{r} \Rightarrow \left\| \frac{d\psi_2}{dm} \right\|_{q_1/2} \leq R^{1/r} \left\| \frac{d\psi_2}{dm} \right\|_{(q_2/2)'}$$

The result follows from the characterization of $\text{CBPlan}_{q_2}^{\Phi_2}$ in Proposition 2.22. \square

REMARK 2.24. – A result along the same lines can be obtained for crossed products under invariance assumptions on ψ_1 . This goes in the same spirit as Remark 2.10.

2.4.3. *Refinement of the smoothness condition.* – Here we are going to see how we can prove the optimal smoothness order in the Hörmander-Mikhlin condition of Theorem B ii) when ψ satisfies the CBPlan_q^Φ property. We need several preparatory lemmas. In the next one we denote by $W_\eta^{p,s}(\mathbb{R}_+)$, where $\eta \in \mathcal{H}_0^\infty$, the Sobolev space given by completion with respect to the norm

$$\|f\|_{W_\eta^{p,s}(\mathbb{R}_+)} = \|(1 - \partial_x^2)^{s/2}(\eta f)\|_p.$$

LEMMA 2.25. – *Given $f, g : \mathbb{R}_+ \rightarrow \mathbb{C}$, the following holds:*

i) *For every $\varepsilon > 0$*

$$\|(1 - \partial_x^2)^{s/2}(fg)\|_2 \lesssim_{(s,\varepsilon)} \|(1 - \partial_x^2)^{(s+1+\varepsilon)/2} f\|_\infty \|(1 - \partial_x^2)^{s/2} g\|_2.$$

ii) *If $\rho(z) = z^s e^{-z}$ and $\eta \in \mathcal{H}_0^\infty$*

$$\|(1 - \partial_x^2)^{s/2}(\eta \rho f)\|_2 \lesssim_{(s,\varepsilon)} \|(1 - \partial_x^2)^{(s+1+\varepsilon)/2}(\eta f)\|_\infty.$$

Equivalently, we find the embedding $W_\eta^{\infty,s+1+\varepsilon}(\mathbb{R}_+) \subset_{(s,\varepsilon)} W_{\eta\rho}^{2,s}(\mathbb{R}_+)$.

Proof. – The second point follows immediately from the first one by noticing that $\rho(z) = z^s e^{-z}$ has finite $W^{2,s}(\mathbb{R}_+)$ norm. We are going to prove the first point for $s \in \mathbb{N}$ and use interpolation. Given $s \in \mathbb{N}$, we have

$$\begin{aligned} \|(1 - \partial_x^2)^{s/2}(fg)\|_2 &\sim \sum_{k=0}^s \|\partial_x^k(fg)\|_2 \\ &= \sum_{k=0}^s \left\| \sum_{j=0}^k \binom{k}{j} (\partial_x^j f)(\partial_x^{k-j} g) \right\|_2 \\ &\lesssim_{(s)} \left(\max_{0 \leq j \leq s} \|\partial_x^j f\|_\infty \right) \left(\sum_{k=0}^s \|\partial_x^k g\|_2 \right) \\ &\sim \left(\max_{0 \leq j \leq s} \|\partial_x^j f\|_\infty \right) \|(1 - \partial_x^2)^{s/2} g\|_2. \end{aligned}$$

Thus, all we have to see is that for every $j \in \{0, 1, 2, \dots, s\}$

$$\|\partial_x^j (1 - \partial_x^2)^{-(s+\varepsilon+1)/2} f\|_\infty \lesssim_{(s,\varepsilon)} \|f\|_\infty.$$

Recall that if the symbol of a Fourier multiplier is given by the Fourier transform of finite measure, then it is bounded in $L_\infty(\mathbb{R})$. Thus, we just need to see that there is a finite measure $\mu_{j,s}$ such that

$$\begin{aligned} \widehat{\mu}_{j,s}(\xi) &= \frac{\xi^j}{(1 + |\xi|^2)^{\frac{s+\varepsilon+1}{2}}} \\ &= \text{sgn}(\xi)^j \frac{1}{(1 + |\xi|^2)^{\frac{s+\varepsilon-j+1}{2}}} \frac{|\xi|^j}{(1 + |\xi|^2)^{\frac{j}{2}}} = (H_{[j]}(\nu_{s,j}) * m_j)^\wedge(\xi), \end{aligned}$$

where $H_{[j]}$ is the Hilbert transform for j odd and the identity map for j even. By [37, V.3, Lemma 2] m_j is a finite measure. Therefore, it is enough to see that if $\widehat{\nu}_{s,j}(\xi) = 1/(1 + |\xi|^2)^{(s+\varepsilon-j+1)/2}$, then $H_{[j]}(\nu_{s,j})$ is a finite measure. Applying the Hilbert transform or identity map to [37, V.(26)] gives the desired result. \square

LEMMA 2.26. – Assume G is a LCH unimodular group, $\psi : G \rightarrow \mathbb{R}_+$ is a c.n. length and that they satisfy the CBPlan_q^Φ property. If $\eta_1, \eta_2 \in \mathcal{H}_0^\infty(\Sigma_\theta)$, with η_1 satisfying that there is $\gamma > 0$ such that $|\eta_1(z)| \lesssim e^{-\gamma \text{Re}(z)}$ for all $z \in \Sigma_\theta$, then the following estimate holds for all $m \in L_\infty(\mathbb{R}_+)$

$$\|\lambda(m(\psi)\eta_1(t\psi)\eta_2(t\psi))\|_{L_2(\mathcal{Z}G)} \lesssim_{(D_\Phi, q, \gamma)} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}} \|m(t^{-1} \cdot)\eta_2(\cdot)\|_{L_q(\mathbb{R}_+)}.$$

Proof. – Using integration by parts we obtain

$$\begin{aligned} \|\lambda(m(\psi)\eta_1(t\psi)\eta_2(t\psi))\|_{L_2(\mathcal{Z}G)} &= \left\| \int_{\mathbb{R}_+} \lambda(m(\psi)\eta_1'(r)\eta_2(t\psi)\chi_{[0,r]}(t\psi)) dr \right\|_{L_2(\mathcal{Z}G)} \\ &\leq \int_{\mathbb{R}_+} \eta_1'(r) \|\lambda(m(\psi)\eta_2(t\psi)\chi_{[0,r]}(t\psi))\|_{L_2(\mathcal{Z}G)} dr. \end{aligned}$$

Nos, applying the CBPlan_q^Φ property, we obtain

$$\begin{aligned} &\|\lambda(m(\psi)\eta_1(t\psi)\eta_2(t\psi))\|_{L_2(\mathcal{Z}G)} \\ &\lesssim_{(q)} \int_{\mathbb{R}_+} \eta_1'(r) \frac{1}{\Phi(\sqrt{t/r})^{\frac{1}{2}}} \|m((r/t)\cdot)\eta_2(r\cdot)\|_{L_q([0,1])} dr \\ &= \left(\int_{\mathbb{R}_+} \eta_1'(r) \frac{r^{-1/q}}{\Phi(\sqrt{t/r})^{\frac{1}{2}}} dr \right) \|m(t^{-1} \cdot)\eta_2(\cdot)\|_{L_q(\mathbb{R}_+)}. \end{aligned}$$

So, we just need to estimate the integral in the right hand side term

$$\int_{\mathbb{R}_+} \eta_1'(r) \frac{r^{-1/q}}{\Phi(\sqrt{t/r})^{\frac{1}{2}}} dr = \left\{ \int_0^1 + \sum_{j=0}^\infty \int_{4^j}^{4^{j+1}} \right\} \eta_1'(r) \frac{r^{-1/q}}{\Phi(\sqrt{t/r})^{\frac{1}{2}}} dr = A + \sum_{j=0}^\infty B_j.$$

The first term is bounded as follows

$$A \leq \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}} \int_0^1 \eta_1'(r) r^{-1/q} dr \lesssim_{(q)} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}}.$$

For the rest of the terms, we apply the doubling condition to obtain

$$B_j \leq 3 \cdot 4^j \|\eta_1'\|_{L_\infty([4^j, 4^{j+1}])} \frac{2^{\frac{D_\Phi}{2}(j+1)}}{\Phi(\sqrt{t})^{\frac{1}{2}}} = \frac{3 \cdot 2^{\frac{D_\Phi}{2}}}{\Phi(\sqrt{t})^{\frac{1}{2}}} \|\eta_1'\|_{L_\infty([4^j, 4^{j+1}])} 2^{\left(\frac{D_\Phi}{2}+2\right)j}.$$

The function η_1 decreases exponentially and so does η_1' . Therefore $\eta_1'(z) \lesssim e^{-\gamma z}$ for $\text{Re}\{z\}$ large enough. That allows us to sum up all the terms in the series obtaining $\sum_j B_j \lesssim \Phi(\sqrt{t})^{-\frac{1}{2}}$ up to a constant depending on (D_Φ, γ) , as desired. \square

PROPOSITION 2.27. – Assume G is a LCH unimodular group, $\psi : G \rightarrow \mathbb{R}_+$ is a c.n. length and that they satisfy the CBPlan_q^Φ property. Assume in addition that $X \in \mathcal{L}G_+^\wedge$ is doubling and admits $L_2\text{GB}$. Then, we find for $\kappa, \delta, \varepsilon > 0$

$$\tau \left\{ \left(\mathbf{1} + \frac{X^2}{t} \right)^\kappa |B_t|^2 \right\}^{\frac{1}{2}} \lesssim_{(D_\Phi, q, \kappa, \delta, \varepsilon)} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}} \|m(t^{-1} \cdot)\eta(\cdot)\|_{W^{p, \kappa + \delta}(\mathbb{R}_+)},$$

where $B_t = \lambda(m(\psi)\eta(t\psi)e^{-2t\psi}(t\psi)^a)$, η is a \mathcal{H}_0^∞ -cut-off and $a = 2\kappa/\delta + (1 + \varepsilon)/2$.

Proof. – Fix $\kappa, \delta, \varepsilon > 0$ and $a = 2\kappa/\delta + (1 + \varepsilon)/2$. We define the linear, unbounded map $K_t : D \subset L_\infty(\mathbb{R}_+) \rightarrow L_2(\mathcal{L}G)$ by $K_t(m) = \lambda(m(t\psi)\eta(t\psi)e^{-2t\psi}(t\psi)^a)$. Using Lemma 2.26 with $\eta_1(z) = z^a e^{-2z}$ and $\eta_2(z) = \eta(z)$ gives that

$$(2.5) \quad \left\| K_t : W_\eta^{q,0}(\mathbb{R}_+) \rightarrow L_2(\mathcal{L}G) \right\| \lesssim_{(D_\Phi,q)} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}}.$$

Let us denote by $\phi_{t,\kappa}$ the family of weights given by $\phi_{t,\kappa}(x) = \tau\{(\mathbf{1} + t^{-1}X^2)^\kappa x\}$ and let $L_2(\mathcal{L}G, \phi_{t,\kappa})$ be the Hilbert spaces associated to the GNS construction of $\phi_{t,\kappa}$. We know from Proposition 2.16 that

$$\left\| K_t : W_{\eta\rho}^{2,s+\frac{1+\varepsilon}{2}}(\mathbb{R}_+) \rightarrow L_2(\mathcal{L}G, \phi_{t,s}) \right\| \lesssim_{(\kappa,\delta,\varepsilon)} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}},$$

where $s = 2\kappa/\delta$ and $\rho(z) = z^a e^{-z}$. Composing with the inclusion

$$W_\eta^{q,s+\frac{1+\varepsilon}{2}+1+\varepsilon'}(\mathbb{R}_+) \subset_{(s,\varepsilon')} W_{\eta\rho}^{2,s+\frac{1+\varepsilon}{2}}(\mathbb{R}_+),$$

which follows by interpolation from Lemma 2.25 for $q = \infty$ and the trivial inclusion for $q = 2$, gives

$$(2.6) \quad \left\| K_t : W_\eta^{q,s+\frac{1+\varepsilon}{2}+1+\varepsilon'}(\mathbb{R}_+) \rightarrow L_2(\mathcal{L}G, \phi_{t,s}) \right\| \lesssim_{(\kappa,\delta,\varepsilon,\varepsilon')} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}}.$$

Notice that the spaces obtained through GNS construction $L_2(\mathcal{L}G, \phi_{t,\kappa})$ are well behaved with respect to the complex interpolation method. In particular, the expected identity below holds

$$[L_2(\mathcal{L}G, \phi_{t,\kappa_1}), L_2(\mathcal{L}G, \phi_{t,\kappa_2})]_\theta = L_2(\mathcal{L}G, \phi_{t,(1-\theta)\kappa_1+\theta\kappa_2}).$$

Therefore, interpolating (2.5) and (2.6) with $\theta = \delta/2$ yields

$$\left\| K_t : W_\eta^{q,\kappa+\frac{\delta}{2}(\frac{1+\varepsilon}{2}+1+\varepsilon')}(\mathbb{R}_+) \rightarrow L_2(\mathcal{L}G, \phi_{t,\theta s}) \right\| \lesssim_{(D_\Phi,q,\kappa,\delta,\varepsilon,\varepsilon')} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}}.$$

Finally, choosing ε and ε' such that $((1 + \varepsilon)/2 + 1 + \varepsilon') \leq 2$ gives

$$\left\| K_t : W_\eta^{q,\kappa+\delta}(\mathbb{R}_+) \rightarrow L_2(\mathcal{L}G, \phi_{t,\kappa}) \right\| \lesssim_{(D_\Phi,q,\kappa,\delta)} \frac{1}{\Phi(\sqrt{t})^{\frac{1}{2}}}.$$

Therefore, applying this bound to the function $m(t^{-1}\cdot)$ proves the assertion. □

Proof of Theorem B ii). – Let $s > D_\Phi/2$. For any $\eta \in \mathcal{S}'_0^\infty$ and $\delta, \varepsilon > 0$ we can define $\eta_1(z) = \eta(z)e^{-2z}z^a$, where $a = 2s/\delta + (1 + \varepsilon)/2$. Set $B_t = \lambda(m(\psi)\eta_1(t\psi))$ and apply (2.2). By Proposition 2.27

$$\sup_{t>0} \|\Sigma_t\|_{L_2(\mathcal{L}G)} \lesssim_{(D_\Phi,q,s,\delta,\varepsilon)} \sup_{t>0} \|m(t^{-1}\cdot)\eta(\cdot)\|_{W^{p,s+\delta}(\mathbb{R}_+)}.$$

Once this is settled, the argument continues as in the proof of Theorem B i). □

2.5. An application for finite-dimensional cocycles

Our aim is to recover the main result in [18] for the case of radial multipliers to illustrate how the Sobolev dimension approach is, a priori, more flexible than the one used in [18]. We will start proving that c.n. lengths coming from surjective and proper finite-dimensional cocycles satisfy the standard assumptions. Then we will reduce the case of general finite-dimensional cocycles to surjective and proper ones.

Let $b : G \rightarrow \mathbb{R}^n$ be a finite-dimensional cocycle. Assume that b is surjective and proper (i.e., $b^{-1}[K]$ is a compact set for every compact K). Then the pullback of the Haar measure $b^*\mu(E) = \mu(b^{-1}[E])$ in \mathbb{R}^n is translation invariant and therefore satisfies that $d b^*\mu(\xi) = cd \xi$. Indeed, let $\alpha : G \rightarrow O(\mathbb{R}^n)$ be the orthogonal action naturally associated to b . Given a Borel compact set $E \subset \mathbb{R}^n$ with $b^{-1}(E) = A \subset G$ and since $b(gA) = \alpha_g(b(A)) + b(g)$, we conclude that

$$b^*\mu(E) = \mu(A) = \mu(gA) = b^*\mu(\alpha_g(E) + b(g)).$$

Note that $\mu(A)$ is well-defined and finite since b is continuous and proper. Applying this identity to the α -invariant sets $E = B_r(0)$ and using the surjectivity of b , we conclude the assertion. By rescaling b if necessary we will assume, without loss of generality, that $b^*\mu$ is the Lebesgue measure over \mathbb{R}^n . An important consequence of this fact is that

$$\|S_t\|_{\mathcal{CB}(L_2(\mathcal{L}G), \mathcal{L}G)}^2 = \int_G |e^{-t\psi(g)}|^2 d\mu(g) = \int_{\mathbb{R}^n} e^{-2t|\xi|^2} d(b^*\mu)(\xi) = \frac{1}{\Phi(\sqrt{t})},$$

where $\mathcal{S} = (S_t)_{t \geq 0}$ is the semigroup associated with $\psi(g) = \|b(g)\|^2$ and $\Phi(t) \sim t^n$. Therefore, the semigroup associated to any proper and surjective finite-dimensional cocycle satisfies the CBR_Φ property. In the same way, the measure μ_ψ defined in (2.3) can be expressed (using polar coordinates) as in terms of $b^*\mu$ and a trivial calculation gives that ψ has the CBPlan_2^Φ property. We need to find a suitable $X_b \in \mathcal{L}G_+^\wedge$. We shall prove that b induces a natural transference map from functions $f : \mathbb{R}^n \rightarrow \mathbb{C}$ into operators $x \in \mathcal{L}G$ given by

$$\mathcal{J}(f) = \lambda(\widehat{f} \circ b).$$

Therefore, if \mathcal{R} is a distribution in \mathbb{R}^n such that $\widehat{\mathcal{R}}(x) = |x|$, our choice will be $X_b = \lambda(\mathcal{R}(b))$. Before proving $X_b \in \mathcal{L}G_+^\wedge$ we will need the following auxiliary result.

LEMMA 2.28. – *If $\varphi_j : \mathbb{R}^n \rightarrow \mathbb{C}$ are radial L_1 -functions*

$$\lambda(\varphi_1 \circ b) \lambda(\varphi_2 \circ b) = \lambda((\varphi_1 * \varphi_2) \circ b)$$

for any group G equipped with a proper and surjective cocycle $b : G \rightarrow \mathbb{R}^n$.

Proof. – We know that $d(b^*\mu) = dm$, so that

$$\begin{aligned} (\varphi_1 \circ b) * (\varphi_2 \circ b)(g) &= \int_G \varphi_1(b(h))\varphi_2(b(h^{-1}g)) d\mu(h) \\ &= \int_G \varphi_1(b(h))\varphi_2(\alpha_{h^{-1}}(b(g) - b(h))) d\mu(h) \\ &= \int_G \varphi_1(b(h))\varphi_2(b(g) - b(h)) d\mu(h) \\ &= \int_{\mathbb{R}^n} \varphi_1(\xi)\varphi_2(b(g) - \xi) d(b^*\mu)(\xi) \end{aligned}$$

$$= \int_{\mathbb{R}^n} \varphi_1(\zeta) \varphi_2(b(g) - \zeta) d\zeta = (\varphi_1 * \varphi_2)(b(g)).$$

Taking the left regular representation at both sides yields the assertion. □

It is straightforward to restate Lemma 2.28 in terms of the transference operator \mathcal{J} . We shall be working with the following subclass of radial functions in the Euclidean space \mathbb{R}^n

$$A(\mathbb{R}^n)_{\text{rad}} = \{ \phi : \mathbb{R}^n \rightarrow \mathbb{C} \mid \phi \text{ radial, } \hat{\phi} \in L_1(\mathbb{R}^n) \}.$$

The norm on $A(\mathbb{R}^n)_{\text{rad}}$ given by $\|\phi\|_{A(\mathbb{R}^n)_{\text{rad}}} = \|\hat{\phi}\|_1$ makes such space a Banach algebra. If needed, $A(\mathbb{R}^n)_{\text{rad}}$ will be given the o.s.s. inherited from the Fourier class $A(\mathbb{R}^n)$. Observe that Lemma 2.28 implies that

$$(2.7) \quad \mathcal{J}(\phi_1 \phi_2) = \mathcal{J}(\phi_1) \mathcal{J}(\phi_2),$$

for any $\phi_j \in A(\mathbb{R}^n)_{\text{rad}}$. In fact, we have the following.

LEMMA 2.29. – *Let $b : G \rightarrow \mathbb{R}^n$ be a proper and surjective cocycle. Then,*

- i) $\mathcal{J} : A(\mathbb{R}^n)_{\text{rad}} \rightarrow \mathcal{L}G$ is contractive, a $*$ -homomorphism for the natural conjugation and positivity preserving,
- ii) for every radial function $\phi \in A(\mathbb{R}^n)_{\text{rad}}$, we have that $\|\mathcal{J}(\phi)\|_{\mathcal{L}G} = \|\phi\|_{\infty}$.

Proof. – Let us start with i). It is trivial that \mathcal{J} is bounded since

$$\|\mathcal{J}(\phi)\|_{\mathcal{L}G} = \|\lambda(\hat{\phi} \circ b)\|_{\mathcal{L}G} \leq \|\hat{\phi} \circ b\|_{L_1(G)} = \|\hat{\phi}\|_{L_1(\mathbb{R}^n)}.$$

The multiplicativity follows from (2.7) and the fact that the map is $*$ -preserving is trivial. To see that the map is positivity preserving just notice that if $0 \leq \phi \in A(\mathbb{R}^n)_{\text{rad}}$, then $\hat{\phi}$ is of positive type. But, if that is the case, then

$$\hat{\phi}(b(g^{-1}h)) = \hat{\phi}(\alpha_{g^{-1}}(b(g) - b(h))) = \hat{\phi}(b(g) - b(h))$$

and so $\hat{\phi} \circ b$ is also of positive type over G .

For ii), let $\mathcal{M} \subset \mathcal{L}G$ be the weak- $*$ closure of $\mathcal{A} = \mathcal{J}[A(\mathbb{R}^n)_{\text{rad}}]$. Clearly, since

$$\tau(\mathcal{J}(\phi)) = (\hat{\phi} \circ b)(e) = \int_{\mathbb{R}^n} \phi(\xi) dm,$$

we have that $\tau|_{\mathcal{M}}$ is also semifinite and faithful. As a consequence we get

$$(2.8) \quad \|x\|_{\mathcal{M}} = \sup_{\xi \in \text{Ball}(L_2(\mathcal{M}, \tau|_{\mathcal{M}}))} \|x \xi\|_2$$

and by the Plancherel identity $L_2(\mathcal{M}) \cong L_2(\mathbb{R}^n)_{\text{rad}}$. We are also going to use that if $\mathcal{A} \subset \mathcal{M}$ is a weak- $*$ dense $*$ -subalgebra of \mathcal{M} , then $L_2(\mathcal{M}) \cap \mathcal{A}$ is norm dense inside $L_2(\mathcal{M})$. Those two facts yield that

$$\begin{aligned} \|\mathcal{J}(\phi)\|_{\mathcal{L}G} &= \sup_{\xi \in \text{Ball}(L_2(\mathcal{M}))} \|\mathcal{J}(\phi) \xi\|_2 \\ &= \sup_{\mathcal{J}(\psi) \in \mathcal{A} \cap \text{Ball}(L_2(\mathcal{M}))} \|\mathcal{J}(\phi \psi)\|_2 \\ &= \sup_{\psi \in \text{Ball}(L_2(\mathbb{R}^n)_{\text{rad}})} \|\phi \psi\|_2 = \|\phi\|_{\infty}, \end{aligned}$$

for any radial ϕ and that concludes the proof. □

In order to define X_b as an element of $\mathcal{L}G_+^\wedge$, we need to express it as the supremum of positive operators in $\mathcal{L}G$. We use

$$1 = \int_{\mathbb{R}_+} s |\xi|^2 e^{-s|\xi|^2} \frac{d s}{s},$$

and think of $\eta_s(\xi) = |\xi|^2 s e^{-s|\xi|^2}$ as a continuous partition of the unit. Note that $\eta_s \in A(\mathbb{R}^n)_{\text{rad}}$. Hence

$$|\xi| = \int_{\mathbb{R}_+} |\xi| \eta_s(\xi) \frac{d s}{s} \rightsquigarrow \phi_{\varepsilon, R}(\xi) := \int_{\varepsilon}^R |\xi| \eta_s(\xi) \frac{d s}{s} \rightsquigarrow X_b := \sup_{0 < \varepsilon \leq R < \infty} \mathcal{J}(\phi_{\varepsilon, R}),$$

where, again, $\phi_{\varepsilon, R} \in A(\mathbb{R}^n)_{\text{rad}}$. This presents X_b as a well-defined element of the extended positive cone $\mathcal{L}G_+^\wedge$.

THEOREM 2.30. – *Let G be a LCH unimodular group and consider an n -dimensional proper and surjective cocycle $b : G \rightarrow \mathbb{R}^n$ equipped with the conditionally negative length $\psi(g) = \|b(g)\|^2$. Then (G, ψ, X_b) satisfies the standard assumptions.*

Proof. – We will start by proving the $L_2\text{GB}$. By noticing that $\zeta \mapsto \chi_{[r, \infty)}(\zeta)$ is an increasing function and the normality of the weight $x \mapsto \tau \{x |\lambda(e^{-t\psi})|^2\}$ we obtain that

$$\tau \left\{ \chi_{[r, \infty)}(X_b) |\lambda(e^{-t\psi})|^2 \right\} = \sup_{0 < \varepsilon \leq R < \infty} \tau \left\{ \chi_{[r, \infty)}(\mathcal{J}(\phi_{\varepsilon, R})) |\lambda(e^{-t\psi})|^2 \right\}.$$

If P is a polynomial, (2.7) gives $P(\mathcal{J}(\phi)) = \mathcal{J}(P(\phi))$. The function $\chi_{[r, \infty)}$ may not be a polynomial but we can approximate it by analytic functions as follows. Let F be

$$(2.9) \quad F(\zeta) = \frac{1}{2} + \frac{1}{\sqrt{\pi}} \int_0^\zeta e^{-s^2} ds.$$

We define the function $\chi_{n, r} \geq 0$ by

$$\chi_{n, r}(\zeta) = (F(n(\zeta - r)) - F(-nr))^2.$$

For $r > 0$, the positive functions $\chi_{r, n}$ converge pointwise and boundedly as $n \rightarrow \infty$ to $\chi_{[r, \infty)}$ except at r , where the value is $1/2$. Furthermore, $\chi_{n, r}(0) = 0$ and $\chi_{n, r}$ is a real analytic function with arbitrarily large convergence radius. By the analyticity it holds that, for any radial ϕ in the Schwartz class,

$$\chi_{n, r}(\mathcal{J}(\phi)) = \mathcal{J}(\chi_{r, n}(\phi)).$$

The right hand side is well-defined since $\chi_{r, n}(\phi)$ is again a Schwartz class function and so its Fourier transform is integrable. By [9, Proposition 1.48] if $\chi_{n, r}$ converges to $\chi_{[r, \infty)}$ pointwise and boundedly then $\chi_{n, r}(x)$ converges to $\chi_{[0, \infty)}(x)$ is the SOT topology for any positive $x \in \mathcal{L}G$. We have that

$$\begin{aligned} \tau \left\{ \chi_{[r, \infty)}(X_b) |\lambda(e^{-t\psi})|^2 \right\} &= \sup_{0 < \varepsilon \leq R < \infty} \tau \left\{ \text{SOT-}\lim_{n \rightarrow \infty} \chi_{r, n}(\mathcal{J}(\phi_{\varepsilon, R})) |\lambda(e^{-t\psi})|^2 \right\} \\ &= \sup_{0 < \varepsilon \leq R < \infty} \lim_{n \rightarrow \infty} \tau \left\{ \mathcal{J}(\chi_{r, n} \circ \phi_{\varepsilon, R}) |\lambda(e^{-t\psi})|^2 \right\} \\ &\leq \lim_{n \rightarrow \infty} \sup_{0 < \varepsilon \leq R < \infty} \tau \left\{ \mathcal{J}(\chi_{r, n} \circ \phi_{\varepsilon, R}) |\lambda(e^{-t\psi})|^2 \right\}. \end{aligned}$$

Moreover, $\lambda(e^{-t\psi}) = \mathcal{J}(h_t)$ for the heat kernel h_t in \mathbb{R}^n and

$$\begin{aligned} \tau\left\{\chi_{[r,\infty)}(X_b) |\lambda(e^{-t\psi})|^2\right\} &\leq \lim_{n \rightarrow \infty} \sup_{0 < \varepsilon \leq R < \infty} \tau\left\{\mathcal{J}(\chi_{r,n} \circ \phi_{\varepsilon,R}) |\mathcal{J}(h_t)|^2\right\} \\ &= \lim_{n \rightarrow \infty} \sup_{0 < \varepsilon \leq R < \infty} \tau\left\{\mathcal{J}((\chi_{r,n} \circ \phi_{\varepsilon,R}) |h_t|^2)\right\} \\ &= \lim_{n \rightarrow \infty} \sup_{0 < \varepsilon \leq R < \infty} \int_{\mathbb{R}^n} \chi_{r,n}(\phi_{\varepsilon,R}(\xi)) |h_t(\xi)|^2 d\xi \\ &\leq \lim_{n \rightarrow \infty} \int_{\mathbb{R}^n} \chi_{r,n}(|\xi|) |h_t(\xi)|^2 d\xi \\ &= \int_{\mathbb{R}^n} \chi_{[r,\infty)}(|\xi|) |h_t(\xi)|^2 d\xi \lesssim \frac{1}{\Phi(\sqrt{t})} e^{-\frac{r^2}{2t}}. \end{aligned}$$

The CBHL inequality will follow from the L_∞ Gaussian lower bounds

$$(L_\infty \text{GLB}) \quad \left\| \left(\chi_{[0,r)}(X_b) \lambda(e^{-t\psi}) \chi_{[0,r)}(X_b) \right)^{-1} \right\|_\infty^{-1} \gtrsim \frac{e^{-\beta \frac{r^2}{t}}}{\Phi(\sqrt{t})}.$$

Recall that if $x \in \mathcal{M}_+$ and p is a projection, then $p x p \in (p \mathcal{M} p)_+$ and, since any positive element can be inverted in the extended positive cone, it holds that $(p x p)^{-1} \in (p \mathcal{M} p)_+^\wedge$. We have $p \|(p x p)^{-1}\|_\infty^{-1} \leq p x p$, where the L_∞ -norm is taken in $p \mathcal{M} p$ and the left hand side may be zero. So, we can understand the right hand side of $(L_\infty \text{GLB})$ as a lower bound on $\chi_{[0,r)}(X_b) \lambda(e^{-t\psi}) \chi_{[0,r)}(X_b)$. The $L_\infty \text{GLB}$ allow to bound the noncommutative Hardy-Littlewood maximal operator by the maximal operator associated with the semigroup. Indeed, since X_b and $\lambda(e^{-t\psi})$ commute from (2.7) we deduce that $(L_\infty \text{GLB})$ yield

$$\frac{\chi_{[0,t)}(X_b)}{\Phi(t)} \lesssim \chi_{[0,t)}(X_b) \lambda(e^{-t^2\psi}) \chi_{[0,t)}(X_b) \leq \lambda(e^{-t^2\psi}).$$

This implies

$$\frac{\chi_{[0,t)}(X_b)}{\Phi(t)} \star x \lesssim S_{t^2}(x),$$

for every positive x . Now, using the maximal inequalities for semigroups of [23] gives the boundedness of the noncommutative Hardy-Littlewood maximal for every $1 < p < \infty$. The fact that $S_t \otimes \text{Id}$ is again a Markovian semigroup gives the complete bounds and so the CBHL inequality holds. To prove that $(L_\infty \text{GLB})$ holds we use that $\mathcal{J} : \mathcal{A} \rightarrow \mathcal{LG}$ is a complete contraction. Justifying the calculations like in the case of upper L_2 Gaussian bounds and using (2.7) we obtain that

$$\begin{aligned} \left\| \left(\chi_{[0,r)}(X_b) \lambda(e^{-t\psi}) \chi_{[0,r)}(X_b) \right)^{-1} \right\|_\infty &= \left\| \lambda(e^{-t\psi})^{-\frac{1}{2}} \chi_{[0,r)}(X_b) \lambda(e^{-t\psi})^{-\frac{1}{2}} \right\|_\infty \\ &\leq \left\| \lambda(e^{-t\psi})^{-\frac{1}{2}} \tilde{\chi}_r(X_b) \lambda(e^{-t\psi})^{-\frac{1}{2}} \right\|_\infty \\ &= \left\| \mathcal{J}(\tilde{\chi}_r h_t^{-1}) \right\|_\infty \\ (2.10) \quad &= \left\| \tilde{\chi}_r h_t^{-1} \right\|_{L_\infty(\mathbb{R}^n)} \lesssim t^{\frac{n}{2}} e^{\beta \frac{r^2}{t}}, \end{aligned}$$

where $\chi_{[0,r)}(|x|) \leq \tilde{\chi}_r$ is a radial and analytic function approximating $\chi_{[0,r)}(|x|)$ such that $h_t^{-1} \tilde{\chi}_r \lesssim t^{\frac{n}{2}} e^{\beta \frac{r^2}{t}}$. One can safely take $\tilde{\chi}_r(x) = F(M(1 - |x/r|^2))F(M(1 + |x/r|^2))$

for large enough M , where F is defined like in (2.9). The analyticity is used to ensure that $\tilde{\chi}_r(\mathcal{J}(\phi)) = \mathcal{J}(\tilde{\chi}_r(\phi))$. In Line (2.10) we used the second point in Lemma 2.29. \square

COROLLARY 2.31. – *Given a LCH amenable unimodular group G , let $b : G \rightarrow \mathbb{R}^n$ be a finite-dimensional cocycle with associated c.n. length $\psi(g) = |b(g)|^2$. Then, given a symbol $m : \mathbb{R}_+ \rightarrow \mathbb{C}$ and $1 < p < \infty$, the following estimate holds for any \mathcal{H}_0^∞ cut-off function η and any $s > n/2$*

$$\|T_{m \circ \psi}\|_{\mathcal{CB}(L_p^\circ(\mathcal{I}G))} \lesssim_{(p)} \sup_{t>0} \|m(t \cdot)\eta(\cdot)\|_{W^{2,s}(\mathbb{R}_+)}.$$


Proof. – If the cocycle b is surjective and proper the result follows from Theorem B. Indeed, in that case we know from Theorem 2.30 that (G, ψ, X_b) satisfies the standard assumptions with $\Phi(s) = s^n$ and Sobolev dimension $D_\Phi = n$. Moreover, the CBPlan_2^Φ property also holds as we explained before Lemma 2.28. In the general case take $G_\rtimes = \mathbb{R}^n \rtimes_\alpha G$ where $\alpha : G \rightarrow O(n)$ is the orthogonal representation that makes $g \mapsto (x \mapsto \alpha_g x + b(g))$ an affine representation. The function $b_\rtimes : G_\rtimes \rightarrow \mathbb{R}^n$ given by $b_\rtimes(\xi, g) = \xi + b(g)$ satisfies the cocycle law with cocycle action $\beta : G_\rtimes \rightarrow \mathbb{R}^n$ given by $\beta_{(\xi,g)} = \alpha_g$. Indeed, we have

$$\begin{aligned} b_\rtimes(\xi + \alpha_g \zeta, gh) &= \xi + \alpha_g \zeta + b(gh) \\ &= \xi + \alpha_g \zeta + \alpha_g b(h) + b(g) \\ &= \beta_{(\xi,g)}(b_\rtimes(\zeta, h)) + b_\rtimes(\xi, g). \end{aligned}$$

Furthermore b_\rtimes is clearly surjective but it may not be proper. In that case, we shall take the associated affine representation $\pi_\rtimes : G_\rtimes \rightarrow \mathbb{R}^n \rtimes O(n)$ and note that the quotient representation $\pi_\rtimes^\circ : G_\rtimes^\circ = G_\rtimes / \ker(\pi_\rtimes) \rightarrow \mathbb{R}^n \rtimes O(n)$ satisfies that its associated cocycle $b_\rtimes^\circ : G_\rtimes^\circ \rightarrow \mathbb{R}^n$ is always proper (even if it is not injective). To see that, let $p_1 : \mathbb{R}^n \rtimes O(n) \rightarrow \mathbb{R}^n$ be the natural projection into the first component and consider a compact set $K \subset \mathbb{R}^n$. Then

$$(b_\rtimes^\circ)^{-1}[K] = (\pi_\rtimes^\circ)^{-1}[p_1^{-1}[K]] = (\pi_\rtimes^\circ)^{-1}[K \times O(n)]$$

and the last term is compact since $K \times O(n)$ is compact and π_\rtimes° is continuous bijection between second countable locally compact groups and hence a continuous group isomorphism. Summing up, we have the following commutative diagram



According to Theorem 2.30, for the last cocycle we can use that $(G_\rtimes^\circ, \psi_\rtimes^\circ, X_{b_\rtimes^\circ})$ satisfy the standard assumptions, where ψ_\rtimes° is the c.n. length naturally associated to b_\rtimes° . By Theorem B, this implies

$$\|T_{m \circ \psi_\rtimes^\circ}\|_{\mathcal{CB}(L_p(\mathcal{I}G_\rtimes^\circ))} \lesssim_{(p)} \sup_{t>0} \|m(t \cdot)\eta(\cdot)\|_{W^{2,s}(\mathbb{R}_+)}.$$

Now, using de Leeuw's type periodization [3, Theorem 8.4 iii)] we obtain the same complete bounds for $T_{m \circ \psi_\times}$ in $L_p(\mathcal{L}G_\times)$ for every $1 < p < \infty$. In order to prove the assertion, we just need to restrict to the subgroup $\{0\} \times G \leq G_\times$. This follows from the de Leeuw's restriction type result in [3, Theorem 8.4 i)]. \square

2.6. Foreword

During the exposition of the contents of Section 2 several natural questions arise.

1. The first question is whether all finite-dimensional proper cocycles, not necessarily surjective, such that their associated c.n. length satisfy CBR_Φ have $L_2\text{GB}$ for some $X \in \mathcal{L}G_+^\wedge$. We have only been able to prove it in the easier case of surjective cocycles. To that end, our intuition is that a (probably nontrivial) generalization of (2.7) will be required.
2. The second point sprouts from the annoyance of the fact that we have not been able to produce explicit examples of infinite-dimensional cocycles with $L_2\text{GB}$. We are not confident about their existence. It will be of great interest for us to either construct infinite-dimensional cocycles having $L_2\text{GB}$ or to prove that all c.n. lengths admitting X with $L_2\text{GB}$ come from finite-dimensional cocycles. A way of relaxing such problem is to change the family of c.n. lengths arising from finite-dimensional to the family of (real) analytic c.n. lengths (in order to make sense of analyticity we will require G to be a Lie group). Note that every finite-dimensional cocycle $b : G \rightarrow \mathbb{R}^n$ over a Lie group G induces a group homomorphism of Lie groups $\pi : G \rightarrow \mathbb{R}^n \rtimes O(n)$. Such homomorphisms are automatically analytic. Therefore, the function $\psi : G \rightarrow \mathbb{R}_+$ is real analytic. It is reasonable to conjecture that every $\psi : G \rightarrow \mathbb{R}_+$ defined on a Lie group and with $L_2\text{GB}$ is analytic.
3. A possible strategy for constructing conditionally negative lengths coming from infinite-dimensional cocycles with $L_2\text{GB}$ is to extend the stability results (announced in Remark 2.10) for crossed products to non θ -invariant $\psi_1 : H \rightarrow \mathbb{R}_+$ and $X_1 \in \mathcal{L}H_+^\wedge$. If either G is amenable or $\theta : G \rightarrow \text{Aut}(H)$ is an amenable action, some sort of averaging procedure may give new c.n. lengths having $L_2\text{GB}$ if the original ones do have $L_2\text{GB}$. It will also be desirable to extend the stability of the standard assumptions to extensions of topological groups.

3. Non-spectral multipliers

3.1. Polynomial co-growth

As we have seen, elements in the extended positive cone $\mathcal{L}G_+^\wedge$ can be understood as quantized metrics over $\mathcal{L}G$. Indeed, when G is abelian, any invariant distance over its dual group is determined by the (positive unbounded) function $d(e, \chi)$ affiliated to $L_\infty(\widehat{G})$, since $d(\chi_1, \chi_2) = d(e, \chi_1^{-1}\chi_2)$. It may seem natural to require X to satisfy properties analogous to the triangular inequality, the faithfulness and the symmetry. Nevertheless, such assumptions will not be necessary here since we will need just "asymptotic" properties of X . Indeed, one of our main families of examples will come from the unbounded multiplication symbols of invariant Laplacians over G . In order to match the classical case of \mathbb{R}^n with the standard

Laplacian, whose multiplication symbol is $|\xi|^2$, we will use the convention that X behaves like $d(e, \chi)^2$. That will explain the $1/2$ exponent in some of the formulas.

DEFINITION 3.1. – Given $X \in \mathcal{L}G_+^\wedge$, we say that X has polynomial co-growth of order D iff

$$D = \inf \left\{ r > 0 : (\mathbf{1} + X)^{-r/2} \in L_1(\mathcal{L}G) \right\} < \infty.$$

The definition is motivated by the fact that if we are in an abelian group and X is the unbounded positive function given by $d(e, \chi)^2$, where d is a translation invariant metric then, defining $\Phi(r) = \tau(\chi_{[0,r^2]}(X)) = \mu(B_r(e))$, we get

$$\|(\mathbf{1} + X)^{-D/2-\varepsilon}\|_1 = \int_{\mathbb{R}_+} \frac{1}{(1+r^2)^{\frac{D}{2}+\varepsilon}} d\Phi(r) = \left(\frac{D}{2} + \varepsilon\right) \int_{\mathbb{R}_+} \frac{2r\Phi(r)}{(1+r^2)^{\frac{D}{2}+1+\varepsilon}} dr.$$

In particular the last expression is finite whenever $\mu(B_r(e)) \lesssim r^D$.

REMARK 3.2. – In the proof of Theorem C we are only going to use that the convolution operator $u \mapsto u \star (\mathbf{1} + X)^{-\beta}$ is completely bounded on $L_p(\mathcal{L}G)$ for $\beta > D$. Any element in $L_1(\mathcal{L}G)$ induces such bounded operator. Indeed we could have defined a similar notion of polynomial co-growth alternatively as

$$D = \inf \left\{ r > 0 : (\mathbf{1} + X)^{-r/2} \in \mathcal{CB}(L_1(\mathcal{L}G)) \right\} < \infty,$$

where $(\mathbf{1} + X)^{-r/2}$ is identified with the operator $x \mapsto (\mathbf{1} + X)^{-r/2} \star x$. This condition is a priori weaker than co-polynomial growth although they coincide for amenable groups. We will stick to the original since it is a condition general enough to allow us to prove Theorem C and restrictive enough to be fully characterized.

Now we are going to prove the existence of unbounded operators affiliated to $\mathcal{L}G$ behaving like multiplication symbols for left or right invariant Laplacians. Recall that a submarkovian semigroup \mathcal{S} acting on $L_\infty(G)$ is respectively called left/right invariant when $S_t \circ \lambda_g = \lambda_g \circ S_t$ or $S_t \circ \rho_g = \rho_g \circ S_t$ accordingly.

PROPOSITION 3.3. – Let G be a LCH unimodular group and consider any submarkovian semigroup \mathcal{S} over $L_\infty(G)$. Let A denote its positive generator. Then, the following properties hold:

- i) If \mathcal{S} is left invariant then there is a densely defined and closable unbounded positive operator \widehat{A} affiliated to $\mathcal{L}G$ such that, for all $f \in \text{dom}(A) \subset L_2(G)$

$$\lambda(Af) = \lambda(f)\widehat{A}.$$

- ii) If \mathcal{S} is right invariant then there is densely defined and closable unbounded positive operator \widehat{A} affiliated to $\mathcal{L}G$ such that, for all $f \in \text{dom}(A) \subset L_2(G)$

$$\lambda(Af) = \widehat{A}\lambda(f).$$

Proof. – We start by proving ii). Notice that $A : \text{dom}(A) \subset L_2(G) \rightarrow L_2(G)$ is densely defined. It is affiliated with $\mathcal{L}G$ iff for every unitary $u \in \mathcal{L}G' = \mathcal{R}G$ we have that $uA = Au$. Since S_t is ρ invariant and we can approximate in the SOT topology every element in $\mathcal{R}G$ by linear combinations of elements in $(\rho_g)_{g \in G}$, we obtain that S_t commutes with any element $x \in \mathcal{R}G$. A function $f \in L_2(G)$ is in $\text{dom}(A)$ when

$$\lim_{t \rightarrow 0^+} \frac{\text{Id} - S_t}{t} f$$

exists in $L_2(G)$ and we then have

$$\lim_{t \rightarrow 0^+} \left\| Af - \frac{\text{Id} - S_t}{t} f \right\|_2 = 0.$$

This implies $u \text{dom}(A) \subset \text{dom}(A)$ for any $U(\mathcal{R}G)$. Multiplying by u we obtain

$$\|uAf - Au f\|_2 \leq \lim_{t \rightarrow 0^+} \left\| uAf - \frac{\text{Id} - S_t}{t} u f \right\|_2 + \lim_{t \rightarrow 0^+} \left\| \frac{\text{Id} - S_t}{t} u f - Au f \right\|_2 = 0$$

for every $f \in \text{dom}(A)$. This proves that A is affiliated with $\mathcal{L}G$. Notice that $\lambda : L_2(G) \rightarrow L_2(\mathcal{L}G)$ unitarily. We will define $\widehat{A} = \lambda A \lambda^*$. By definition \widehat{A} is an unbounded operator on $L_2(\mathcal{L}G)$ affiliated with $(\lambda \mathcal{R}G \lambda^*)' = \lambda \mathcal{L}G \lambda^*$ which is also equal to the von Neumann algebra $\mathcal{L}G$ acting by left multiplication in the GNS construction associated to its trace. The operator \widehat{A} is densely defined and closable since A is densely defined and closable. The identity of ii) follows by definition. The construction for i) is somewhat analogous. We need two trivial observations:

1. The anti-automorphism $\sigma : \mathcal{L}G \rightarrow \mathcal{L}G$ extends to a unitary operator $\sigma_2 : L_2(\mathcal{L}G) \rightarrow L_2(\mathcal{L}G)$ since $\tau \circ \sigma = \tau$. If $\pi_r : \mathcal{L}G_{\text{op}} \rightarrow \mathcal{B}(L_2(\mathcal{L}G))$ and $\pi_\ell : \mathcal{L}G \rightarrow \mathcal{B}(L_2(\mathcal{L}G))$ are the right and left GNS representations, then $\sigma_2 \circ \pi_r(x) = \pi_\ell(\sigma x) \circ \sigma_2$.
2. The anti-automorphism σ extends to an automorphism of the extended positive cone $\mathcal{L}G_\pm^\wedge$. We are going to denote such extension again by σ .

Notice that, since $\pi_\ell[\mathcal{L}G]' = \pi_r[\mathcal{L}G]$, any element in $x \in \pi_\ell[\mathcal{L}G]'$ can be expressed as $\pi_r(x')$ for some $x' \in \mathcal{L}G$. By point 1, the map that sends x to x' is given, after identifying $\mathcal{L}G$ with its GNS representation $\pi_\ell[\mathcal{L}G]$, by $x' = \sigma(\sigma_2 x \sigma_2)$. Let S be given by $S = \lambda A \lambda^*$. Then S is affiliated with $(\lambda \mathcal{L}G \lambda^*)' = \pi_\ell[\mathcal{L}G]'$. If we define \widehat{A} as $\widehat{A} = \sigma(\sigma_2 S \sigma_2)$, where σ is the extension of point 2, we obtain i). □

REMARK 3.4. – *Since G is unimodular, the unitary $\iota : L_2(G) \rightarrow L_2(G)$ given by $f(g) \mapsto f(g^{-1})$ is an isometry that intertwines ρ_g and λ_g . We can characterize the pairs of left and right invariant operators A_1, A_2 whose left and right multiplication symbols, \widehat{A}_1 and \widehat{A}_2 respectively, coincide. By a trivial calculation those are the operators such that $A_1 \iota = \iota A_2$. Indeed, using that $\lambda : L_2(G) \rightarrow L_2(\mathcal{L}G)$ satisfies $\lambda \circ \iota = \sigma_2 \circ \lambda$ and that if A is the infinitesimal generator of a submarkovian semigroup then $A^\top = A$, we obtain that*

$$\sigma(\lambda A_1 \lambda^*) = \sigma_2 \lambda A_2 \lambda^* \sigma_2 = \lambda \iota A_2 \iota \lambda^*,$$

but the right hand side satisfies that $\sigma(\lambda A_1 \lambda^) = \lambda A_1^\top \lambda^* = \lambda A_1 \lambda^*$.*

Now we are going to characterize those semigroups whose infinitesimal generator has polynomial co-growth. In order to prove the characterization we will need the following two lemmas. Recall that the Fourier algebra AG is defined as those $f : G \rightarrow \mathbb{C}$ such that $\lambda(f) \in L_1(\mathcal{L}G)$ with $\|f\|_{AG} = \|\lambda(f)\|_{L_1(\mathcal{L}G)}$. We will use below the straightforward inequalities for $f \in AG$

$$(3.1) \quad |\tau(\lambda(f))| \leq \|f\|_\infty \leq \tau(|\lambda(f)|).$$

Indeed, both follow from the identity $\tau(\lambda_g^* \lambda(f)) = f(g)$ which is valid for $f \in AG$.

LEMMA 3.5. – *Let G be a LCH unimodular group and \mathcal{S} a submarkovian semigroup of right (resp. left) invariant operators satisfying that $S_t : C_0(G) \rightarrow C_0(G)$. Let A be the positive generator and assume further that \widehat{A} has polynomial cogrowth of order D . Then $W_A^{2,s}(G) \cap AG$ is dense inside $W_A^{2,s}(G)$ for every $s > D/2 + \varepsilon$.*

Proof. – We will prove only the right invariant case. Notice that AG is closed by left and right translations. The fact that $S_t : C_0(G) \rightarrow C_0(G)$, together with the Riesz representation theorem gives that for every $g \in G$ there is weak-* continuous family of unit measures on G , $(\mu_t^g)_{g \in G, t \geq 0}$ such that

$$S_t f(g) = \int_G f(h) d\mu_t^g(h).$$

Applying the right invariance gives us that $d\mu_t^g(h) = d\mu_t^e(hg^{-1})$. This yields

$$(3.2) \quad S_t f(g) = \int_G \rho_g f d\mu_t^e = \iota^* \mu_t^e * f(g),$$

where $(\iota^* \mu_t^e)(E) = \mu_t^e(E^{-1})$. It is clear that $\|S_t f - f\|_{L_2(G)} \rightarrow 0$ as $t \rightarrow 0^+$. Recall that the same is true for $f \in W_A^{2,s}(G)$ in the $W_A^{2,s}(G)$ -norm for every $s > 0$. Suppose that $f \in W_A^{2,s}(G)$, then, applying the formula (3.2) together with the polynomial co-growth, we have that

$$S_t f = \iota^* \mu_t * f = \iota^* \mu_t * (\mathbf{1} + A)^{-\frac{s}{2}} (\mathbf{1} + A)^{\frac{s}{2}} f = h_{t,s} * g,$$

where $g = (\mathbf{1} + A)^{s/2} f$. We have that $\|g\|_2 = \|f\|_{W_A^{s,2}}$ and

$$\|h_{t,s}\|_2 \leq \|\mu_t^e\|_{M(G)} \|(\mathbf{1} + \widehat{A})^{-s/2}\|_{L_2(\mathcal{L}G)} < \infty.$$

This proves that $S_t f \in AG \cap W_A^{2,s}(G)$. Making $t \rightarrow 0^+$ completes the claim. \square

THEOREM 3.6. – *Let G be a unimodular LCH group and let \mathcal{S} be a right (resp. left) invariant submarkovian semigroup over G . Let A be its infinitesimal generator and assume further that $S_t : C_0(G) \rightarrow C_0(G)$. Then, the following assertions are equivalent:*

- i) *The multiplication symbol \widehat{A} of A has polynomial co-growth of order D .*
- ii) *\mathcal{S} satisfies the following inequality for every $\varepsilon > 0$*

$$\left\| (\mathbf{1} + A)^{-(\frac{D}{4} + \varepsilon)} : L_2(G) \rightarrow L_\infty(G) \right\| \lesssim_{(\varepsilon)} 1.$$

Proof. – To prove i) \Rightarrow ii), pick $f \in AG \cap W^{2,s}(G)$ for $s = D/2 + 2\varepsilon$ and note

$$\begin{aligned} \|f\|_\infty &\leq \|\lambda((\mathbf{1} + A)^{-s/2}(\mathbf{1} + A)^{s/2} f)\|_1 \\ &= \|(\mathbf{1} + \widehat{A})^{-s/2} \lambda((\mathbf{1} + A)^{s/2} f)\|_1 \\ &\leq \|(\mathbf{1} + \widehat{A})^{-s/2}\|_2 \|\lambda((\mathbf{1} + A)^{s/2} f)\|_2 \\ &= \|(\mathbf{1} + \widehat{A})^{-s}\|_1^{1/2} \|f\|_{W_A^{2,s}(G)} \lesssim_{(\varepsilon)} \|f\|_{W_A^{2,s}(G)}. \end{aligned}$$

We have used (3.1) in the first inequality, Proposition 3.3 in the first identity and the polynomial cogrowth in the last inequality. By the density Lemma 3.5 we conclude that $W_A^{2,s}(G)$ embeds in $L_\infty(G)$ which is a rephrasal of ii). For the implication ii) \Rightarrow i) we note that from (3.1)

$$\left| \tau\left((\mathbf{1} + \widehat{A})^{-\frac{D}{4}-\varepsilon} \lambda(f)\right) \right| \leq \|(\mathbf{1} + A)^{-\frac{D}{4}-2\varepsilon} f\|_\infty \lesssim_{(\varepsilon)} \|f\|_2.$$

Taking the supremum over $f \in L_2(G)$ with norm 1 gives the desired result. □

REMARK 3.7. – *Due to Proposition 2.1 we obtain that the point ii) is equivalent to satisfying the ultracontractivity property $R_{D+\varepsilon}(0)$ for every $\varepsilon > 0$. Since $R_D(0)$ implies $R_{D+\varepsilon}(0)$ for every $\varepsilon > 0$, it is sufficient to prove $R_D(0)$ in order to have polynomial co-growth of order D .*

REMARK 3.8. – *Sobolev inequalities involving powers of $\mathbf{1} + A$ are sometimes called local [43, II.X] since they are tightly connected to the ultracontractivity estimates for $0 < t \leq 1$ and in many contexts that amounts to describing the growth of ball of small radius. Therefore Theorem 3.6 relates the behavior of the large balls of $\mathcal{L}G$ with the behavior of small balls in G . This goes along the common intuition that taking group duals exchanges local and asymptotic/coarse properties.*

Proof of Theorem C. – Let $B_t = \lambda(m \eta(t\psi))$ and let \widehat{A}_1 be the multiplication symbol associated with the generator of the right invariant semigroup \mathcal{S}_1 which is determined by Proposition 3.3. Then

$$B_t = \underbrace{(\mathbf{1} + \widehat{A}_1)^{-\frac{s_1}{2}}}_{M_t} \underbrace{(\mathbf{1} + \widehat{A}_1)^{\frac{s_1}{2}}}_{\Sigma_t} B_t$$

is a max-square decomposition. By the definition of co-polynomial growth we have that $\sigma|M_t|^2 = (\mathbf{1} + \sigma\widehat{A}_1)^{-s_1} \in L_1(\mathcal{L}G)$ and therefore it is a c.b. multiplier in every $L_p(\mathcal{L}G)$ for $1 \leq p \leq \infty$. Since M_t does not depend on t , the maximal inequality (MS_p) is satisfied trivially. By the construction of \widehat{A}_1 we have

$$\sup_{t>0} \|\Sigma_t\|_{L_2(\mathcal{L}G)} = \sup_{t>0} \|(\mathbf{1} + \widehat{A}_1)^{\frac{s_1}{2}} \lambda(m\eta(t\psi))\|_{L_2(\mathcal{L}G)} = \sup_{t>0} \|m\eta(t\psi)\|_{W_{A_1}^{2,s_1}(G)}.$$

The square-max decomposition is manufactured in exactly the same way. □

3.2. Sublaplacians over polynomial-growth Lie groups

Here we are going to work with left (resp. right) invariant submarkovian semigroups over $L_\infty(G)$ generated by sublaplacians. Let M be a smooth manifold, $\mathbb{X} = \{X_1, \dots, X_r\}$ be a family of smooth vector fields and μ a σ -finite measure over M . Let us denote by $(\sigma_j(t))_{t \in (-\varepsilon_j, \varepsilon_j)}$ the one-parameter diffeomorphism generated by X_j and assume further that μ is invariant under $(\sigma_j(t))_{t \in (-\varepsilon_j, \varepsilon_j)}$. Then, the semigroup whose infinitesimal generator is given by the sublaplacian associated to \mathbb{X}

$$\Delta_{\mathbb{X}} = - \sum_{j=1}^r X_j^2$$

is submarkovian. This is a consequence of the theory of symmetric Dirichlet forms [10]. If $M = G$ is a Lie group, μ its left Haar measure and $\mathbb{X} = \{X_1, \dots, X_r\}$ left invariant vector fields. By the invariance under the one parameter subgroup generated by X_j of μ we have that $S_t = e^{-t\Delta_{\mathbb{X}}}$ is a submarkovian semigroup of left invariant operators. The same construction can be performed using right invariant vector fields if G is unimodular. Any sublaplacian carries a natural subriemannian metric given by

$$d_{\mathbb{X}}(x, y) = \inf_{\substack{\gamma: [0,1] \rightarrow M \\ \gamma(0)=x, \gamma(1)=y}} \left\{ \left(\int_0^1 \sum_{j=1}^r |a_j(t)|^2 dt \right)^{\frac{1}{2}} \mid \gamma'(t) = \sum_{j=0}^r a_j(t) X_j(\gamma(t)) \right\}.$$

This metric coincides with the Lipschitz distance given by the gradient form, also known as Meyer's carré du champs [29]. Observe also that, if G is a connected Lie group, then its subriemannian distance is finite iff \mathbb{X} generates the whole Lie algebra. Similarly, $f \in \text{Ker}_p(\Delta_{\mathbb{X}})$ iff $f \in L_p(M)$ and $f(x) = f(y)$ whenever the subriemannian distance $d_{\mathbb{X}}(x, y)$ is finite.

The main family of illustrations of Theorem C comes from Lie groups endowed with right and left invariant sublaplacians. Indeed, let $\mathbb{V} = \{v_1, v_2, \dots, v_r\} \subset T_e G$ be a collection of, linearly independent, vectors generating the whole Lie algebra and $\mathbb{X}_1 = \{X_1, \dots, X_r\}$ and $\mathbb{X}_2 = \{Y_1, \dots, Y_r\}$ be its right and left invariant extensions respectively. Then their associated sublaplacians satisfy $\iota \Delta_{\mathbb{X}_1} = \Delta_{\mathbb{X}_2} \iota$ where we use $\iota f(g) = f(g^{-1})$. Hence, it suffices to study the polynomial co-growth for $\widehat{\Delta}_{\mathbb{X}_1}$. By Remark 3.7 we just need to show that $S_t = e^{-t\Delta_{\mathbb{X}}}$ has the $R_D(0)$ property and by [43, Theorem VIII.2.9] we know that if G is a Lie group of polynomial growth, then

$$\frac{e^{-\beta_1 \frac{d_{\mathbb{X}_1}(x,y)^2}{t}}}{\mu(B_e(\sqrt{t}))} \lesssim h_t(x, y) \lesssim \frac{e^{-\beta_2 \frac{d_{\mathbb{X}_1}(x,y)^2}{t}}}{\mu(B_e(\sqrt{t}))},$$

where h_t is the heat kernel associated with S_t , $d_{\mathbb{X}_1}$ is the subriemannian distance associated to \mathbb{X}_1 and $B_e(r)$ are the balls of radius r with respect to that metric. It is a well known fact, see [43], that

$$\mu(B_e(r)) \sim t^{D_0},$$

for t small. Here D_0 is the *local dimension* associated to \mathbb{X}_1 , given by

$$D_0 = \sum_{j=0}^{\infty} j \dim(F_{j+1}/F_j),$$

where $F_0 = \{0\}$, $F_1 = \mathbb{X}_1$ and $F_{j+1} = \text{span}\{F_j, [F_j, \mathbb{X}_1]\}$. As a consequence S_t has the $R_{D_0}(0)$ property and therefore $\widehat{\Delta}_{\mathbb{X}_1}$, and so $\widehat{\Delta}_{\mathbb{X}_2}$, have polynomial co-growth of order D_0 . As a corollary we obtain the following theorem.

THEOREM 3.9. – *Let G be a polynomial growth Lie group equipped with a c.n. length $\psi : G \rightarrow \mathbb{R}_+$. Let $\eta \in \mathcal{H}_0^\infty$ and consider a generating set $\mathbb{X} = \{X_1, X_2, \dots, X_r\}$ of independent right invariant vector fields. Let us write $\Delta_{\mathbb{X}}$ for its sublaplacian. Then, the following inequality holds for every $1 < p < \infty$ and any $s > D_0/2$*

$$\|T_m\|_{\mathcal{CB}(L_p^s(\mathfrak{z}G))} \lesssim_{(p)} \sup_{t \geq 0} \max \left\{ \|\eta(t\psi)m\|_{W_{\Delta_{\mathbb{X}}}^{2,s}(G)}, \|\eta(t\psi)\iota m\|_{W_{\Delta_{\mathbb{X}}}^{2,s}(G)} \right\}.$$

Acknowledgements. – González-Pérez is partially supported by FPI scholarship BES-2011-04419. González-Pérez and Parcet are partially supported by the ERC StG-256997-CZOSQP, Junge is partially supported by the NSF DMS-1201886 and all authors of this paper are also supported in part by the ICMAT Grant “Severo Ochoa” SEV-2011-0087 (Spain).

BIBLIOGRAPHY

- [1] B. BEKKA, P. DE LA HARPE, A. VALETTE, *Kazhdan's property (T)*, New Mathematical Monographs **11**, Cambridge Univ. Press, Cambridge, 2008.
- [2] J. BENNETT, Optimal control of singular Fourier multipliers by maximal operators, *Anal. PDE* **7** (2014), 1317–1338.
- [3] M. CASPERS, J. PARCET, M. PERRIN, E. RICARD, Noncommutative de Leeuw theorems, *Forum Math. Sigma* **3** (2015), e21.
- [4] P.-A. CHERIX, M. COWLING, P. JOLISSAINT, P. JULG, A. VALETTE, *Groups with the Haagerup property*, Progress in Math. **197**, Birkhäuser, 2001.
- [5] A. CONNES, A factor not anti-isomorphic to itself, *Ann. Math.* **101** (1975), 536–554.
- [6] M. G. COWLING, Harmonic analysis on semigroups, *Ann. of Math.* **117** (1983), 267–283.
- [7] E. B. DAVIES, Uniformly elliptic operators with measurable coefficients, *J. Funct. Anal.* **132** (1995), 141–169.
- [8] E. G. EFFROS, Z.-J. RUAN, *Operator spaces*, London Mathematical Society Monographs. New Series **23**, The Clarendon Press, Oxford Univ. Press, New York, 2000.
- [9] G. B. FOLLAND, *A course in abstract harmonic analysis*, Studies in Advanced Math., CRC Press, Boca Raton, FL, 1995.
- [10] M. FUKUSHIMA, Y. OSHIMA, M. TAKEDA, *Dirichlet forms and symmetric Markov processes*, extended ed., De Gruyter Studies in Mathematics **19**, Walter de Gruyter & Co., Berlin, 2011.
- [11] U. HAAGERUP, An example of a nonnuclear C^* -algebra, which has the metric approximation property, *Invent. math.* **50** (1978/79), 279–293.
- [12] U. HAAGERUP, Operator-valued weights in von Neumann algebras. I, *J. Funct. Anal.* **32** (1979), 175–206.

- [13] U. HAAGERUP, Operator-valued weights in von Neumann algebras. II, *J. Funct. Anal.* **33** (1979), 339–361.
- [14] C. HERZ, Problems of extrapolation and spectral synthesis on groups, in *Conference on Harmonic Analysis (Univ. Maryland, College Park, Md., 1971)*, Lecture Notes in Math. **266**, Springer, Berlin, 1972, 157–166.
- [15] M. JUNGE, Doob’s inequality for non-commutative martingales, *J. reine angew. Math.* **549** (2002), 149–190.
- [16] M. JUNGE, C. LE MERDY, Q. XU, H^∞ functional calculus and square functions on noncommutative L^p -spaces, *Astérisque* **305** (2006).
- [17] M. JUNGE, T. MEI, Noncommutative Riesz transforms—a probabilistic approach, *Amer. J. Math.* **132** (2010), 611–680.
- [18] M. JUNGE, T. MEI, J. PARCET, Smooth Fourier multipliers on group von Neumann algebras, *Geom. Funct. Anal.* **24** (2014), 1913–1980.
- [19] M. JUNGE, T. MEI, J. PARCET, Noncommutative Riesz transforms—Dimension free bounds and Fourier multipliers, preprint arXiv:1407.2475.
- [20] M. JUNGE, J. PARCET, Mixed-norm inequalities and operator space L_p embedding theory, *Mem. Amer. Math. Soc.* **203** (2010).
- [21] M. JUNGE, Z.-J. RUAN, Approximation properties for noncommutative L_p -spaces associated with discrete groups, *Duke Math. J.* **117** (2003), 313–341.
- [22] M. JUNGE, Q. XU, Noncommutative Burkholder/Rosenthal inequalities, *Ann. Probab.* **31** (2003), 948–995.
- [23] M. JUNGE, Q. XU, Noncommutative maximal ergodic theorems, *J. Amer. Math. Soc.* **20** (2007), 385–439.
- [24] R. V. KADISON, J. R. RINGROSE, *Fundamentals of the theory of operator algebras. Vol. I*, Graduate Studies in Math. **15**, Amer. Math. Soc., Providence, RI, 1997.
- [25] V. LAFFORGUE, M. DE LA SALLE, Noncommutative L^p -spaces without the completely bounded approximation property, *Duke Math. J.* **160** (2011), 71–116.
- [26] E. C. LANCE, *Hilbert C^* -modules*, London Mathematical Society Lecture Note Series **210**, Cambridge Univ. Press, Cambridge, 1995.
- [27] F. LUST-PIQUARD, Inégalités de Khintchine dans C_p ($1 < p < \infty$), *C. R. Acad. Sci. Paris Sér. I Math.* **303** (1986), 289–292.
- [28] F. LUST-PIQUARD, G. PISIER, Noncommutative Khintchine and Paley inequalities, *Ark. Mat.* **29** (1991), 241–260.
- [29] P. A. MEYER, L’opérateur carré du champ, in *Séminaire de Probabilités, X (Première partie, Univ. Strasbourg, Strasbourg, année universitaire 1974/1975)*, Lecture Notes in Math. **511**, Springer, 1976, 142–161.
- [30] V. PAULSEN, *Completely bounded maps and operator algebras*, Cambridge Studies in Advanced Math. **78**, Cambridge Univ. Press, Cambridge, 2002.
- [31] G. PISIER, Non-commutative vector valued L_p -spaces and completely p -summing maps, *Astérisque* **247** (1998).
- [32] G. PISIER, *Introduction to operator space theory*, London Mathematical Society Lecture Note Series **294**, Cambridge Univ. Press, Cambridge, 2003.

- [33] G. PISIER, Q. XU, Non-commutative L^p -spaces, in *Handbook of the geometry of Banach spaces, Vol. 2*, North-Holland, Amsterdam, 2003, 1459–1517.
- [34] L. SALOFF-COSTE, Sobolev inequalities in familiar and unfamiliar settings, in *Sobolev spaces in mathematics. I*, Int. Math. Ser. 8, Springer, 2009, 299–343.
- [35] A. SIKORA, Sharp pointwise estimates on heat kernels, *Quart. J. Math. Oxford Ser.* **47** (1996), 371–382.
- [36] A. SIKORA, Riesz transform, Gaussian bounds and the method of wave equation, *Math. Z.* **247** (2004), 643–662.
- [37] E. M. STEIN, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, No. 30, Princeton Univ. Press, Princeton, N.J., 1970.
- [38] E. M. STEIN, *Topics in harmonic analysis related to the Littlewood-Paley theory*, Annals of Math. Studies, No. 63, Princeton Univ. Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1970.
- [39] M. TAKESAKI, *Theory of operator algebras. I*, Springer, New York-Heidelberg, 1979.
- [40] M. TAKESAKI, *Theory of operator algebras. II*, Encyclopaedia of Math. Sciences **125**, Springer, Berlin, 2003.
- [41] X. THINH DUONG, E. M. OUHABAZ, A. SIKORA, Plancherel-type estimates and sharp spectral multipliers, *J. Funct. Anal.* **196** (2002), 443–485.
- [42] N. T. VAROPOULOS, Hardy-Littlewood theory for semigroups, *J. Funct. Anal.* **63** (1985), 240–260.
- [43] N. T. VAROPOULOS, L. SALOFF-COSTE, T. COULHON, *Analysis and geometry on groups*, Cambridge Tracts in Mathematics **100**, Cambridge Univ. Press, Cambridge, 1992.

(Manuscrit reçu le 5 juillet 2015 ;
 accepté, après révision, le 23 juin 2016.)

Adrián GONZÁLEZ-PÉREZ
 Instituto de Ciencias Matemáticas
 CSIC-UAM-UC3M-UCM
 Universidad Autónoma de Madrid
 C/ Nicolás Cabrera 13-15
 28049 Madrid, Spain
 E-mail: adrian.gonzalez@icmat.es

Marius JUNGE
 Department of Mathematics
 University of Illinois at Urbana-Champaign
 1409 W. Green St.
 Urbana, IL 61891, USA
 E-mail: junge@math.uiuc.edu

Javier PARCET
 Instituto de Ciencias Matemáticas
 CSIC-UAM-UC3M-UCM
 Consejo Superior de Investigaciones Científicas
 C/ Nicolás Cabrera 13-15
 28049 Madrid, Spain
 E-mail: javier.parcet@icmat.es

