

# BULLETIN DE LA S. M. F.

T.J. HEAD

## Remarks on a problem in primary abelian groups

*Bulletin de la S. M. F.*, tome 91 (1963), p. 109-112

[http://www.numdam.org/item?id=BSMF\\_1963\\_\\_91\\_\\_109\\_0](http://www.numdam.org/item?id=BSMF_1963__91__109_0)

© Bulletin de la S. M. F., 1963, tous droits réservés.

L'accès aux archives de la revue « Bulletin de la S. M. F. » (<http://smf.emath.fr/Publications/Bulletin/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

REMARKS ON A PROBLEM  
IN PRIMARY ABELIAN GROUPS ;

BY

THOMAS J. HEAD

(Ames, Iowa).

---

1. All groups considered in this note are assumed to be  $p$ -primary abelian groups. If  $A$  is a subgroup of  $G$  then  $\overline{A}$  will denote the closure of  $A$  in the usual topology of  $G$  ([2], page 114). The closure of a subgroup is a subgroup, but the closure of a pure subgroup need not be pure. It is a consequence of lemma 20 of [3] that if  $G$  is a closed  $p$ -group (for definition see [2], page 114) then the closure of each pure subgroup of  $G$  is pure.

PROBLEM. — If  $G$  is a primary abelian group without elements of infinite height in which the closure of each pure subgroup is pure does it follow that  $G$  is a closed  $p$ -group ?

We do not know the answer to this question, but we can give an affirmative answer in the case of direct sums of cyclic groups :

THEOREM. — If  $G$  is a direct sum of cyclic  $p$ -groups and the closure of each pure subgroup of  $G$  is pure in  $G$  then  $G$  is a bounded  $p$ -group.

An outline of the proof of this theorem is given in paragraph 3 below.

2. **The relation of the problem to minimal pure embeddings.** — Following B. CHARLES [1], when a subgroup  $S$  of a group  $G$  is contained in a pure subgroup  $P$  of  $G$  which has the property that no proper pure subgroup of  $P$  contains  $S$  we say that  $P$  is *minimal pure containing*  $S$ .

When such a  $P$  exists we say that  $S$  has a minimal pure embedding in  $G$ . We will denote the subgroup of elements of infinite height in a group  $G$  by  $G'$ .

In the proofs below we use the following two observations :

*If  $A$  is a subgroup of  $G$  then  $\bar{A}$  is that subgroup of  $G$  containing  $A$  for which  $\bar{A}/A = (G/A)'$ . If a subgroup  $S$  of a group  $G$  is contained in  $G'$  and if  $P$  is minimal pure containing  $S$  then  $P$  is divisible.*

The latter observation follows from the fact that if  $P$  were not divisible  $P$  would contain a finite cyclic direct summand  $\{x\}$  and if  $P = \{x\} \oplus C$  then  $S$  would be contained in  $C$  where  $C$ , being a direct summand of  $P$ , would be pure in  $G$ .

For a subgroup  $S$  of  $G$  we denote by  $S'$  the subgroup of  $G$  containing  $S$  for which  $S'/S$  is the maximal divisible subgroup of  $G/S$ .

PROPOSITION. — Let  $P$  be a pure subgroup of a primary abelian group  $G$ . Let  $H$  be a subgroup of  $G$  for which  $P \subset H \subset \bar{P}$ . Then  $H$  has a minimal pure embedding in  $G$  if and only if  $H \subset P'$ .

*Proof.* — Suppose  $P_1$  is minimal pure containing  $H$ . Then  $P_1/P$  is minimal pure containing  $H/P$  in  $G/P$ . Since

$$H/P \subset \bar{P}/P = (G/P)',$$

$P_1/P$  is a divisible subgroup of  $G/P$ . Then  $P_1 \subset P'$  and  $H \subset P'$ . Conversely, if  $H \subset P'$  then  $H/P$  is contained in the maximal divisible subgroup of  $G/P$ . Then there exists a subgroup  $P_1$  of  $G$  containing  $P$  such that  $P_1/P$  is minimal divisible containing  $H/P$  in  $G/P$ . Then  $P_1$  is minimal pure containing  $H$  in  $G$ .

It has been suggested ([1], page 224) that if  $G$  is a primary abelian group without elements of infinite height and  $S$  is a subgroup of  $G$  which is the union of an ascending chain of discrete subgroups of  $G$  then  $S$  has a minimal pure embedding in  $G$ . The proposition and theorem above are sufficient to show that this is not true even if the discrete subgroups are finite :

Let  $G$  be a countable unbounded direct sum of cyclic  $p$ -groups. Let  $P$  be a pure subgroup of  $G$  for which  $\bar{P}$  is not pure.  $\bar{P}$  is the union of an ascending chain of finite (hence discrete) subgroups of  $G$ . Since  $P'$  is pure,  $\bar{P} \neq P'$ . Consequently  $\bar{P}$  is not contained in any subgroup of  $G$  which is minimal pure containing  $\bar{P}$ .

This same example is a counter-example to part 2 of theorem 6 of [1] because  $P$  is a pure subgroup of  $G$  which is dense in  $\bar{P}$  and yet  $\bar{P}$  has no minimal pure embedding in  $G$ . Along this line we have :

COROLLARY. — For a primary abelian group  $G$  the following two conditions are equivalent :

(1) Each subgroup  $H$  of  $G$  that contains a subgroup  $P$  which is pure in  $G$  and dense in  $H$  (relative to the topology of  $G$ ) has a minimal pure embedding in  $G$ .

(2) For each pure subgroup  $P$  of  $G$ ,  $\bar{P}$  is pure in  $G$ .

*Proof.* — Assume (1) and let  $P$  be pure in  $G$ . Then  $\bar{P}$  has a minimal pure embedding in  $G$ . By the proposition  $P = P'$  and  $\bar{P}$  is pure in  $G$ .

Assume (2) and let  $H$  be a subgroup of  $G$  which contains a subgroup  $P$  which is pure in  $G$  and dense in  $H$ . We have  $P \subset H \subset \bar{P}$ . Since  $\bar{P}$  is pure in  $G$  it follows from the proposition that  $\bar{P} = P'$ . The proposition then gives the conclusion that  $H$  has a minimal pure embedding in  $G$ .

3. Outline of the proof of the theorem stated in paragraph 1. —

It is sufficient to show that if  $G = \sum_{n=1}^{\infty} Z(p^{i(n)})$  where  $i(n)$  is a strictly increasing sequence of positive integers,  $i(1) \geq 2$ , and  $Z(p^{i(n)})$  is a cyclic group of order  $p^{i(n)}$  then  $G$  contains a pure subgroup  $P$  for which  $\bar{P}$  is not pure. For each positive integer  $n$  let  $g(n)$  be a generator of  $Z(p^{i(n)})$ . Then it may be verified that the following sequence of elements of  $G$  is a linearly independent set and that the subgroup,  $P$ , generated by this set is pure in  $G$  :

$$s(n) = g(2n - 1) + p^{i(2n) - i(2n-1) + 1} g(2n) + p^{i(2n+1) - i(2n-1)} g(2n + 1),$$

$$(1 \leq n < \infty).$$

Let

$$x = p^{i(1)-1} g(1).$$

Then  $x \in \bar{P}$  since modulo  $P$  we have :

$$x = p^{i(1)-1} g(1) \equiv -p^{i(3)-1} g(3) \equiv \dots \equiv (-1)^n p^{i(2n+1)-1} g(2n + 1) \equiv \dots$$

Let  $y$  be any element of  $G$  for which  $p^{i(1)-1} y = x$ . There is an integer  $N$  such that the component of  $y$  in  $Z(p^{i(N)})$  is different from 0 and the component of  $y$  in  $Z(p^{i(n)})$  is 0 for each  $n > N$ . By proceeding from the fact the component of  $y$  in  $Z(p^{i(1)})$  is the unique component of  $y$  which is not annihilated by  $p^{i(1)-1}$ , it can be verified that the neighborhood  $y + p^{i(1)-1} G$  of  $y$  is disjoint from  $P$ . Then  $y \notin \bar{P}$  and the equation  $p^{i(1)-1} z = x$ , which has the solution  $z = g(1)$  in  $G$ , is not solvable for  $z$  in  $\bar{P}$ . Thus  $\bar{P}$  is not pure in  $G$ .

## REFERENCES.

- [1] CHARLES (Bernard). — Étude sur les sous-groupes d'un groupe abélien, *Bull. Soc. math. France*, t. 88, 1960, p. 217-227.
- [2] FUCHS (Laszlo). — *Abelian groups*. Budapest, Hungarian Academy of Sciences, 1958.
- [3] KAPLANSKY (Irving). — *Infinite abelian groups*. Ann Arbor, University of Michigan Press, 1954 (*University of Michigan Publications in Mathematics*, 2).

(Manuscrit reçu le 16 septembre 1962.)

Thomas J. HEAD,  
Iowa State University,  
Ames, Iowa (États-Unis).

