

MÉMOIRES DE LA S. M. F.

EBERHARD BECKER

The reduced Witttring

Mémoires de la S. M. F., tome 59 (1979), p. 11-12

<http://www.numdam.org/item?id=MSMF_1979__59__11_0>

© Mémoires de la S. M. F., 1979, tous droits réservés.

L'accès aux archives de la revue « Mémoires de la S. M. F. » (<http://smf.emath.fr/Publications/Memoires/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

THE REDUCED WITTRING

by

Eberhard BECKER

These notes give a brief account on a joint work of L. Bröcker and the author. Detailed proofs will appear in the Journal of Algebra.

Let K be a real (= formally real) field, $X = X(K)$ the topological space of all orderings of K [5, p.63], and $W(K)$ the Witt ring of the nondegenerated bilinearforms over K . By $W_t(K)$ we denote the torsion-subgroup of $W(K)$, which is known also to be its nilradical [6].

Let $C(X, \mathbb{Z})$ be the ring of all continuous functions $X \rightarrow \mathbb{Z}$ (\mathbb{Z} provided with the discrete topology). Then we get a homomorphism $\text{sign} : W(K) \rightarrow C(X, \mathbb{Z})$ defined by $(\text{sign}(\rho))(P) := \text{sign}_P(\rho) = \text{signature of } \rho \text{ at } P$. The following basic result is due to Pfister [6].

THEOREM 1 : The sequence $0 \rightarrow W_t(K) \rightarrow W(K) \xrightarrow{\text{sign}} C(X, \mathbb{Z})$ is exact.

So it must be considered as a main task in the theory of reduced Witt rings to characterize the elements of $\text{sign } W(K)$ among the functions $f \in C(X, \mathbb{Z})$. In order to state the main result the notion of a preorder has to be introduced. A subset $T \subset K$ is called a preorder of K iff the following conditions are satisfied :

- i) $T + T \subset K, T \cdot T \subset K$ ii) $K^2 \subset T$ iii) $T \cap -T = \{0\}$.

A preorder T is the intersection of all orderings in which it is contained. Given a preorder T , the subset $X_T := \{P \supset T \mid P \text{ ordering of } K\}$ is a closed subspace of X . Clearly, we have the restriction-homomorphism $\text{Res} : C(X, \mathbb{Z}) \rightarrow C(X_T, \mathbb{Z})$.

Denote by $W_T(K)$ the image of $\text{Res} \cdot \text{sign} : W(K) \rightarrow C(X_T, \mathbb{Z})$. Choose any ordering $P_0 \supset T$. Set $P_0^X = P \setminus \{0\}, T^X = T \setminus \{0\}$; P_0^X and T^X are subgroups of K^X . As with $W(K)$, we find an epimorphism $\mathbb{Z}[P_0^X/T^X] \rightarrow W_T(K)$. Furthermore the mapping $X_T \rightarrow \text{char}(P_0^X/T^X), P \mapsto (aT^X \mapsto \text{sgn}_P(a))$ is a topological embedding of X_T into the (Pontrjagin-) character-group of P_0^X/T^X .

PROPOSITION. For a preorder T the following statements are equivalent :

- i) $\mathbb{Z}[P_0^X/T^X] \rightarrow W_T(K)$ is an isomorphism,
 ii) $X_T \rightarrow \text{char}(P_0^X/T^X)$ is a homeomorphism,
 iii) $T + T_a = T \cup T_a$ for all $a \in K$, such that $a \notin -T$.

A preorder which satisfies the equivalent conditions of the last proposition, is called a fan (in French : éventail). Fans turn out to be of great importance in other contexts, too [1], [3].

THEOREM 2. A function $f \in C(X, \mathbb{Z})$ lies in sign $W(K)$ iff

$$\sum_{P \in T} f(P) \equiv 0 \pmod{\frac{1}{2} (K^X : T^X)}$$

for all fans T with $(K^X : T^X) < \infty$.

The description of sign $W(K)$ in $C(X, \mathbb{Z})$ was also attacked by R. Brown [4] and settled for the case that K admits only finitely many real places. For the general case he was led to a conjecture which (in his terminology) states that all formally real fields are exact. From theorem 2 one can derive:

THEOREM 3. All formally real fields are exact.

The proof of theorem 2 heavily depends on two local-global principles for reduced quadratic forms, one of which has essentially been proved in [2]. Furthermore, the generalized theory of reduced Witt rings [1] is extensively used, i.e. $W(K)$ is factorized by forms $\langle 1, -t \rangle$, where $0 \neq t$ and t belongs to an arbitrary but fixed preorder T . This point of view turns out to be fundamental even for the study of ordinary reduced Witt rings.

REFERENCES

- [1] E. BECKER and E. KÖPPING, Reduzierte quadratische Formen und Semiordnungen reeller Körper, Abh. Math. Sem. Univ. Hamburg 46 (1977), 143-177.
- [2] L. BRÖCKER, Zur Theorie der quadratischen Formen über formalreellen Körpern, Math. Ann. 210 (1974), 233-256.
- [3] L. BRÖCKER, Characterizations of fans and hereditarily pythagorean fields, Math. Z. 152 (1976), 149-163.
- [4] R. BROWN, The reduced Witt ring of a formally real field, Trans. Amer. Math. Soc. 230 (1977), 257-292.
- [5] J. MILNOR and D. HUSEMOLLER, Symmetric Bilinear Forms, Springer-Verlag (1973).
- [6] A. PFISTER, Quadratische Formen in beliebigen Körpern, Invent. Math. 1 (1966), 116-132.

Eberhard BECKER

Mathematisches Institut der Universität,
Weyertal 86-90, D-5000 Köln 41.