

MÉMOIRES DE LA S. M. F.

BENEDICT H. GROSS

DON ZAGIER

On the critical values of Hecke L-series

Mémoires de la S. M. F. 2^e série, tome 2 (1980), p. 49-54

http://www.numdam.org/item?id=MSMF_1980_2_2_49_0

© Mémoires de la S. M. F., 1980, tous droits réservés.

L'accès aux archives de la revue « Mémoires de la S. M. F. » (<http://smf.emath.fr/Publications/Memoires/Presentation.html>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE CRITICAL VALUES OF HECKE L-SERIES

by

Benedict H. GROSS

and

Don ZAGIER

Let E be the elliptic curve over \mathbb{Q} with minimal model

$$y^2 + xy = x^3 - x^2 - 2x - 1 .$$

The modular invariant, discriminant, and conductor of E are given by

$$j = -3^3 \cdot 5^3$$

$$\Delta = -7^3$$

$$N = (7^2) .$$

Let Ω denote the fundamental real period of the Néron differential $\omega = \frac{dx}{2y+x}$ on E :

$$\Omega = \int_{E(\mathbb{R})} \omega = 1.93331170561681\dots$$

Over the field $K = \mathbb{Q}(\sqrt{-7})$, E has complex multiplication by $\mathcal{O} = \mathbb{Z}\left[\frac{1+\sqrt{-7}}{2}\right]$. Hence Ω can be determined explicitly, using an identity of

Chowla and Selberg [1] :

$$\Omega = \frac{\Gamma(1/7)\Gamma(2/7)\Gamma(4/7)}{\sqrt{-7} \cdot 2\pi i} .$$

Similarly, the L-series of E is equal to the L-series of a Hecke character χ of K. The conductor of χ is the ideal $(\sqrt{-7})$; for \mathcal{Q} an ideal of K which is prime to 7 :

$$\chi(\mathcal{Q}) = \alpha \text{ where } \mathcal{Q} = (\alpha) , \alpha^3 \equiv 1 \pmod{\sqrt{-7}}$$

We have calculated the central critical values of the Hecke L-series which are associated to odd powers of the character χ . Let $n \geq 1$ be an integer; the Dirichlet series

$$L(\chi^{2n-1}, s) = \sum \frac{\chi^{2n-1}(\mathcal{Q})}{N\mathcal{Q}^s}$$

converges absolutely in the right half-plane $\text{Re}(s) > n + \frac{1}{2}$. It extends to a holomorphic function on the entire complex plane: the modified function $\Lambda(\chi^{2n-1}, s) = (7/2\pi)^s \Gamma(s) L(\chi^{2n-1}, s)$ satisfies Hecke's functional equation :

$$\Lambda(\chi^{2n-1}, s) = (-1)^{n+1} \Lambda(\chi^{2n-1}, 2n-s) .$$

It follows that the value of $L(\chi^{2n-1}, s)$ at $s = n$, the center of the critical strip, vanishes when n is even.

When n is odd, define a_n by

$$L(\chi^{2n-1}, n) = \frac{\Omega^{2n-1}}{(2\pi i/\sqrt{-7})^{n-1}} \frac{a_n}{(n-1)!} .$$

(We found this normalization by trial and error; it is consistent with the work of Katz on the interpolation of real analytic Eisenstein series [2].)

The values of a_n for $1 \leq n \leq 33$ are listed in Table 1.

HECKE L-SERIES

Table 1

n	a_n
1	1/2
3	2
5	2
7	$2(3)^2$
9	$2(7)^2$
11	$2(3^2 \cdot 5 \cdot 7)^2$
13	$2(3 \cdot 7 \cdot 29)^2$
15	$2(3 \cdot 7 \cdot 103)^2$
17	$2(3 \cdot 5 \cdot 7 \cdot 607)^2$
19	$2(3^3 \cdot 7 \cdot 4793)^2$
21	$2(3^2 \cdot 5 \cdot 7 \cdot 29 \cdot 2399)^2$
23	$2(3^3 \cdot 5 \cdot 7^2 \cdot 10091)^2$
25	$2(3^2 \cdot 7^2 \cdot 29 \cdot 61717)^2$
27	$2(3^2 \cdot 5^2 \cdot 7^2 \cdot 13 \cdot 53^2 \cdot 79)^2$
29	$2(3^4 \cdot 5^2 \cdot 7^2 \cdot 113 \cdot 127033)^2$
31	$2(3^5 \cdot 5 \cdot 7^2 \cdot 71 \cdot 1690651)^2$
33	$2(3^4 \cdot 5 \cdot 7^2 \cdot 1291 \cdot 1747169)^2$

Let $p \equiv 1 \pmod{4}$ be a prime and let χ_p be the Hecke character

$$\chi_p(\alpha) = \left(\frac{\mathbb{N}\alpha}{p}\right) \chi(\alpha) .$$

The Hecke L-series $L(\chi_p, s)$ is equal to the L-series of an elliptic curve E_p/\mathbb{Q} which becomes isomorphic to E over $\mathbb{Q}(\sqrt{p})$. Let $\Omega_p = \Omega/\sqrt{p}$ and define $a_n^{(p)}$ by

$$L(\chi_p^{2n-1}, n) = \frac{\Omega_p^{2n-1}}{(2\pi i/\sqrt{-7})^{n-1}} \frac{a_n^{(p)}}{(n-1)!}.$$

Again, $a_n^{(p)}$ vanishes when n is even. For n odd and $p = 5, 13, 17, 29, 53$

we found the values listed in Table 2.

Table 2

n	P	5	13	17	29	53
1	1	1	1	1	2	0 Frank E ₅₃ (0) = 2J
3	(2 ²) ²	(2 ² . 3) ²	(2 ² . 3) ²	(2 . 3 . 13) ²	2(2 ² . 3) ²	2(2 ³ . 7) ²
5	(2 ² . 5) ²	(2 ² . 157) ²	(2 ² . 157) ²	(2 . 271) ²	2(2 ² . 5 . 37) ²	2(2 ⁴ . 3 . 5 ² . 7) ²
7	(2 ² . 3 . 5 . 61) ²	(2 ² . 3 ² . 1847) ²	(2 ² . 3 ² . 1847) ²	(2 . 3 ² . 61) ²	2(2 ² . 3 ² . 11 . 29 . 61) ²	2(2 ³ . 3 . 5 ² . 7 ² . 13 . 17) ²
9	(2 ² . 5 . 7 . 199) ²	(2 ² . 7 . 13 . 5813) ²	(2 ² . 7 . 13 . 5813) ²	(2 . 7 . 266977) ²	2(2 ² . 7 . 17 . 80779) ²	2(2 ⁵ . 3 . 7 ² . 19 . 2699) ²
11	(2 ² . 5 ² . 271) ²	(2 ² . 3 ³ . 5 . 7 . 13 . 1021) ²	(2 ² . 3 ³ . 5 . 7 . 13 . 1021) ²	(2 . 3 ⁴ . 5 . 7 . 17 . 2081) ²	2(2 ² . 3 ³ . 5 ³ . 7 . 13 . 29 . 79) ²	2(2 ³ . 3 ² . 5 ² . 7 ² . 11 . 129893) ²
13	(2 ² . 3 . 5 ² . 7 . 3767) ²	(2 ² . 3 . 7 . 13 . 3747629) ²	(2 ² . 3 . 7 . 13 . 3747629) ²			
15	(2 ² . 3 . 5 ² . 7 . 89 . 13687) ²	(2 ² . 3 ² . 7 . 13 . 101 . 317 . 15307) ²	(2 ² . 3 ² . 7 . 13 . 101 . 317 . 15307) ²			
17	(2 ² . 3 . 5 ⁴ . 7 . 26737) ²	(2 ² . 3 . 5 . 7 . 13 . 4877 . 6510011) ²	(2 ² . 3 . 5 . 7 . 13 . 4877 . 6510011) ²			

Note : In all the cases we computed, $a_n^{(p)}$ is either a square or twice a square, depending on whether $\left(\frac{p}{n}\right)$ is -1 or $+1$. Can one prove this is general? Are there "higher Tate-Shafarevitch groups" associated to the abelian varieties $(E_p)^{2n-1}$ whose orders can be conjecturally related to $a_n^{(p)}$? Do these groups carry a natural alternating pairing?

Bibliography :

- [1] CHOWLA S., and SELBERG A., On Epstein's Zeta Function. J. Crelle 227 (1967), 96-110.
- [2] KATZ N., p -adic interpolation of real analytic Eisenstein series. Annals Math. 104 (1976), 459-571.

B. Gross
Princeton University
Fine Hall
Princeton, N. J. 08540 (U.S.A.)

D. Zagier
Mat. Inst. d. Univ. Bonn
Wegelerstrasse 10
53 Bonn (B.R.D.)