



ANNALES

DE

L'INSTITUT FOURIER

Ugo BOSCAIN & Camille LAURENT

The Laplace-Beltrami operator in almost-Riemannian Geometry

Tome 63, n° 5 (2013), p. 1739-1770.

http://aif.cedram.org/item?id=AIF_2013__63_5_1739_0

© Association des Annales de l'institut Fourier, 2013, tous droits réservés.

L'accès aux articles de la revue « Annales de l'institut Fourier » (<http://aif.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://aif.cedram.org/legal/>). Toute reproduction en tout ou partie de cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

THE LAPLACE-BELTRAMI OPERATOR IN ALMOST-RIEMANNIAN GEOMETRY

by Ugo BOSCAIN & Camille LAURENT (*)

ABSTRACT. — We study the Laplace-Beltrami operator of generalized Riemannian structures on orientable surfaces for which a local orthonormal frame is given by a pair of vector fields that can become collinear.

Under the assumption that the structure is 2-step Lie bracket generating, we prove that the Laplace-Beltrami operator is essentially self-adjoint and has discrete spectrum. As a consequence, a quantum particle cannot cross the singular set (i.e., the set where the vector fields become collinear) and the heat cannot flow through the singularity. This is an interesting phenomenon since when approaching the singular set all Riemannian quantities explode, but geodesics are still well defined and can cross the singular set without singularities.

This phenomenon also appears in sub-Riemannian structures which are not equi-regular, i.e., when the growth vector depends on the point. We show this fact by analyzing the Martinet case.

RÉSUMÉ. — On étudie l'opérateur de Laplace-Beltrami dans des structures Riemanniennes généralisées sur des surfaces orientables pour lesquelles un repère orthonormé est donné par une paire de champs de vecteurs pouvant devenir colinéaires.

Sous l'hypothèse que la structure est génératrice avec des crochets de Lie d'ordre 2, on prouve que l'opérateur de Laplace-Beltrami est essentiellement autoadjoint et a un spectre discret. Par conséquent, une particule quantique ne peut pas traverser l'ensemble singulier (i.e., là où les champs de vecteurs deviennent colinéaires) et la chaleur ne peut pas diffuser à travers la singularité. Le phénomène est intéressant puisque lorsqu'on s'approche de l'ensemble singulier, toutes les quantités Riemanniennes explosent, mais les géodésiques sont encore bien définies et peuvent traverser l'ensemble singulier sans singularité.

Ce phénomène apparaît aussi dans des structures sous-Riemanniennes qui ne sont pas équirégulières, i.e., dont le vecteur de croissance dépend du point. On montre ce fait en analysant le cas Martinet.

Keywords: Grushin, Laplace-Beltrami operator, almost-Riemannian structures.

Math. classification: 53C17,35P05,58C40.

(*) This research has been supported by the European Research Council, ERC StG 2009 GeCoMethods, contract number 239748, by the ANR Project GCM, program Blanche, project number NT09-504490 and by the DIGITEO project CONGEO.

1. Introduction

A 2-dimensional Almost Riemannian Structure (2-ARS for short) is a generalized Riemannian structure that can be defined locally by a pair of smooth vector fields on a 2-dimensional manifold M , satisfying the Hörmander condition. These vector fields play the role of an orthonormal frame.

Let us denote by $\blacktriangle(q)$ the linear span of the two vector fields at a point q . Where $\blacktriangle(q)$ is 2-dimensional, the corresponding metric is Riemannian. Where $\blacktriangle(q)$ is 1-dimensional, the corresponding Riemannian metric is not well defined, but thanks to the Hörmander condition one can still define the Carnot-Carathéodory distance between two points, which happens to be finite and continuous.

2-ARSs were introduced in the context of hypoelliptic operators [20, 21], they appeared in problems of population transfer in quantum systems [13, 14, 15], and have applications to orbital transfer in space mechanics [10, 9]. 2-ARSs are a particular case of rank-varying sub-Riemannian structures (see for instance [8, 23, 35]).

Generically (i.e., for an open and dense subset of the set of all 2-ARSs, in a suitable topology), the singular set \mathcal{Z} , where $\blacktriangle(q)$ has dimension 1, is a 1-dimensional embedded submanifold and there are three types of points: Riemannian points, Grushin points where $\blacktriangle(q)$ is 1-dimensional and $\dim(\blacktriangle(q) + [\blacktriangle, \blacktriangle](q)) = 2$ and tangency points where $\dim(\blacktriangle(q) + [\blacktriangle, \blacktriangle](q)) = 1$ and the missing direction is obtained with one more bracket. One can easily show that at Grushin points $\blacktriangle(q)$ is transversal to \mathcal{Z} . Generically, at tangency points $\blacktriangle(q)$ is tangent to \mathcal{Z} and tangency points are isolated. Normal forms at Riemannian, Grushin and tangency points were established in [4] and are described in Figure 1.1.

2-ARSs present very interesting phenomena. For instance, geodesics can pass through the singular set, with no singularities even if all Riemannian quantities (as for instance the metric, the Riemannian area, the curvature) explode while approaching \mathcal{Z} . Moreover, the presence of a singular set permits the conjugate locus to be nonempty even if the Gaussian curvature is always negative (where it is defined). See [4] and [4, 2, 17, 18] for Gauss–Bonnet-type formulas and for a classification of 2-ARSs from the point of view of Lipschitz equivalence.

In this paper we study the Laplace-Beltrami operator in a 2-ARS. The main point is that the first order terms explode as a consequence of the explosion of the area on the singular set \mathcal{Z} . In particular we are interested in its self-adjointness. This is crucial to understand the evolution of a quantum particle and of the heat flow in a 2-ARS.




Figure 1.1. The local representations established in [4]. Tangency points are the most difficult to handle due to the fact that the asymptotic of the distance is different from the two sides of the singular set (see [11, 16]).

On $M \setminus \mathcal{Z}$, this operator is defined as the divergence of the gradient. The almost-Riemannian gradient can be defined with no difficulty since it involves only the inverse metric that is well defined (by continuity) even on the singular set. More precisely, on a Riemannian manifold (M, g) , ∇ is the unique operator from $C^\infty(M)$ to $\text{Vec}(M)$ satisfying $g_q(\nabla \phi(q), v) = d\phi_q(v)$, $\forall q \in M, v \in T_qM$. In the two-dimensional case, if in an open set $\Omega \subset M$ an orthonormal frame for g is given by two vector fields X_1 and X_2 , then we have on Ω ,

$$\nabla(\phi) = X_1(\phi)X_1 + X_2(\phi)X_2, \tag{1.1}$$

where by $X_i(\phi)$ we mean the Lie derivative of ϕ in the direction of X_i ($i = 1, 2$). This last formula can be used to define the gradient even where X_1 and X_2 are not linearly independent. In this paper, we use formula (1.1) to define the gradient of a function on a 2-ARS.

The divergence of a vector field requires a notion of area and in a 2-ARS the natural area (i.e., the Riemannian one) explodes while q approaches the singular set. Let us recall that the divergence of a vector field X on an orientable Riemannian manifold (M, g) is the unique function satisfying $\operatorname{div} X d\omega = L_X d\omega$ where $d\omega$ is the Riemannian volume form. In coordinates $\operatorname{div} X = \frac{1}{\omega} \sum_i \partial_i (\omega X^i)$, where $\omega = \sqrt{\det g}$.

To see how the Riemannian quantities (and in particular $d\omega$) explode while approaching the singular set \mathcal{Z} , choose a local orthonormal frame for the 2-ARS of the form $X_1 = (1, 0)$ and $X_2 = (0, f(x, y))$, where f is some smooth function. This is always possible, see for instance [4]. On $M \setminus \mathcal{Z}$, for the metric g , the area element $d\omega$, the curvature K and the gradient of a smooth function one has:

$$\begin{aligned} g(x, y) &= \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{f(x,y)^2} \end{pmatrix}, \\ d\omega(x, y) &= \frac{1}{|f(x, y)|} dx dy, \\ K(x, y) &= \frac{f(x, y) \partial_x^2 f(x, y) - 2 (\partial_x f(x, y))^2}{f(x, y)^2}, \\ \nabla(\phi) &= \partial_x(\phi) \partial_x + f^2 \partial_y(\phi) \partial_y. \end{aligned}$$

and for a vector field $Y = (Y^1, Y^2)$ one has

$$\begin{aligned} \operatorname{div}(Y) &= |f| \left(\partial_x \left(\frac{1}{|f|} Y^1 \right) + \partial_y \left(\frac{1}{|f|} Y^2 \right) \right) \\ &= \partial_x Y^1 + \partial_y Y^2 - \frac{\partial_x f}{f} Y^1 - \frac{\partial_y f}{f} Y^2 \end{aligned} \tag{1.2}$$

which is not well defined on \mathcal{Z} even on vector fields which are linear combinations of X_1 and X_2 . In particular, the divergence is not well defined on X_1 . As a consequence, the Laplace Beltrami operator presents some singularities in the first order terms. (The choice of the area does not affect the principal symbol of the operator). More precisely, one has

$$\Delta(\phi) = \partial_x^2 \phi + f^2 \partial_y^2 \phi - \frac{\partial_x f}{f} \partial_x \phi + f(\partial_y f)(\partial_y \phi). \tag{1.3}$$

The simplest case is the well known Grushin metric, which is the 2-ARS on \mathbf{R}^2 for which an orthonormal basis is given by,

$$X_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad X_2 = \begin{pmatrix} 0 \\ x \end{pmatrix}. \tag{1.4}$$

Here the singular set \mathcal{Z} is the y axis and on $\mathbf{R}^2 \setminus \mathcal{Z}$ the Riemannian metric, the Riemannian area and the Gaussian curvature are given respectively by:

$$g = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{x^2} \end{pmatrix}, \quad d\omega = \frac{1}{|x|} dx dy, \quad K = -\frac{2}{x^2}. \quad (1.5)$$

It follows that the Laplace Beltrami operator is given by

$$\Delta\phi := \operatorname{div}(\nabla(\phi)) = (\partial_x^2 + x^2\partial_y^2 - \frac{1}{x}\partial_x)\phi, \quad (1.6)$$

which is the basic model of Laplace Beltrami operator on 2-ARSs and whose self-adjointness will be studied in this paper. Notice that the operator "sum of squares",

$$\bar{\Delta} = X_1^2 + X_2^2 = \partial_x^2 + x^2\partial_y^2 \quad (1.7)$$

has been deeply studied in the literature. This operator (which in this case is the principal symbol of Δ) can be obtained as the divergence of the gradient by taking as area, the standard Euclidean area on \mathbf{R}^2 . However, this operator does not give any information about Δ because of the diverging first order term. Moreover, in 2-ARS an intrinsic Laplacian with non-diverging terms cannot be defined naturally, since an intrinsic area which does not diverge on the singular set is not known. Here by "intrinsic" we mean an area which depends only on the 2-ARS and not on the choice of the coordinates and of the orthonormal frame.

Notice that even if all Riemannian quantities are not defined on \mathcal{Z} , classical geodesics do. See Section 3.1 for the explicit expressions of geodesics in the Grushin plane. Another interesting feature of the Grushin plane is that a bounded open set intersecting \mathcal{Z} has finite diameter but infinite area.

The main purpose of this paper is to study the following question:

[Q]: Let M be a 2-D manifold endowed with a 2-ARS, and let Δ be the corresponding Laplace-Beltrami defined on $C_0^\infty(M \setminus \mathcal{Z})$. Is Δ essentially self-adjoint?

Notice that a priori one expects a negative answer to this question, since, as explained below, a positive answer would imply that neither the heat flow nor a quantum particle can pass through \mathcal{Z} , while classical geodesics cross it with no singularities.

Our main result is an unexpected positive answer to the question **[Q]** when there are no tangency points and when M is compact. More precisely, we have the following.

THEOREM 1.1. — *Let M be a 2-D compact orientable manifold endowed with a 2-ARS. Assume that*

[HA]: the singular set \mathcal{Z} is an embedded one-dimensional submanifold of M ;

[HB]: for every $q \in M$, $\blacktriangle(q) + [\blacktriangle, \blacktriangle](q) = T_qM$.

Let $d\omega$ be the corresponding Riemannian area and Δ the corresponding Laplace-Beltrami operator, both defined on $M \setminus \mathcal{Z}$. Then we have the following facts

- (1) Δ with domain $C_0^\infty(M \setminus \mathcal{Z})$ is essentially self-adjoint on $L^2(M, d\omega)$.
- (2) The domain of Δ (the self-adjoint extension denoted by the same letter) is given by

$$D(\Delta) = \{u \in L^2(M, d\omega) \mid \Delta_{D,g}u \in L^2(M, d\omega)\}$$

where $\Delta_{D,g}u$ is Δu seen as a distribution in $\Omega = M \setminus \mathcal{Z}$.

- (3) The resolvent $(-\Delta + 1)^{-1}$ is compact and therefore its spectrum is discrete and consists of eigenvalues with finite multiplicity.

Hypothesis **HA** is generic (see [4]). Hypothesis **HB** implies that every point is either a Riemannian point or a Grushin point (see Figure 1.1). The hypothesis that there are no tangency points is technical and the same result should hold in much more general situations. Indeed, the singular set behaves like a barrier as a consequence of the explosion of the volume, and in the case of tangency points this explosion is even stronger. The orientability of the manifold can be weakened. Indeed it is only necessary that each connected component of \mathcal{Z} (which is diffeomorphic to S^1) admits an open tubular neighborhood diffeomorphic to $(-1, 1) \times S^1$. See Proposition 2.3 and 2.4.

Remark 1.2. — Notice that 2-ARSs satisfying **HA** and **HB** do exist. See for instance [4] where such a 2-ARS has been built on compact orientable manifolds of any genus. Notice moreover that a 2-ARS satisfying **HA** and **HB** are structurally stable in the sense that small perturbations of the local orthonormal frames in the C^∞ norm, do not destroy conditions **HA** and **HB**.

Theorem 1.1 has a certain number of implications.

First, it implies that a quantum particle, localized for $t = 0$ on a connected component of $M \setminus \mathcal{Z}$, remains localized in this connected component for any time $t \in \mathbf{R}$. The same phenomenon holds for the wave or the heat equation. Indeed, the essential self-adjointness means that our operator can be naturally and uniquely extended (by taking its closure) to a self-adjoint operator without adding any additional boundary condition. But, in our case, one possible extension is to take an extension for the operator defined

on each connected component (for instance, the Friedrichs extension) and to “concatenate” them. This possible extension is the one which separates the dynamics. For instance, this is what we obtain when one defines the Friedrichs extension of the usual Laplace operator on \mathbf{R} defined on function in $C_0^\infty(\mathbf{R} \setminus \{0\})$. We obtain two distinct dynamics with Dirichlet boundary condition at 0. Yet, this is of course not the only possible dynamic because the operator defined on function $C_0^\infty(\mathbf{R} \setminus \{0\})$ is not essentially self-adjoint. Hence, the essential self-adjointness means that the unique self-adjoint extension of our operator is the one that separates the dynamics on each connected components. That means that it is not necessary to add any boundary condition: the explosion of the area naturally acts as a barrier which prevents the crossing of the degeneracy zone by the particules.

More precisely, we have the following.

COROLLARY 1.3. — *With the notations of Theorem 1.1, consider the unique solution u of the Schrödinger equation (according to the self-adjoint extension defined in the previous theorem),*

$$\begin{cases} i\partial_t u + \Delta u &= 0 \\ u(0) &= u_0 \in L^2(M, d\omega) \end{cases} \tag{1.8}$$

with u_0 supported in a connected component Ω of $M \setminus \mathcal{Z}$. Then, $u(t)$ is supported in Ω for any $t \geq 0$. The same holds for the solution of the heat or for the solution of the wave equation.

Second, it is well known that in Riemannian geometry one can relate properties of the Riemannian distance to those of the corresponding heat kernels. For instance, we have that

THEOREM 1.4 (Varadhan, Neel, Stroock). — *Let M be a compact, connected, smooth Riemannian manifold and d the corresponding Riemannian distance. Let $p_t(q_1, q_2)$ be the heat kernel of the heat equation $\partial_t \phi = \frac{1}{2} \Delta \phi$, where Δ is the Laplace-Beltrami operator. Define*

$$E_t(q_1, q_2) := -t \log p_t(q_1, q_2)$$

We have:

- 1): $E_t(q_1, q_2) \rightarrow \frac{1}{2}d(q_1, q_2)^2$, uniformly on $M \times M$ as $t \rightarrow 0$,
- 2): Let $Cut(q_1) \subset M$ the cut locus from q_1 . Then $q_2 \notin Cut(q_1)$ if and only if $\lim_{t \rightarrow 0} \nabla^2 E_t(q_1, q_2) = \frac{1}{2} \nabla^2 d(q_1, q_2)^2$, while $q_2 \in Cut(q_1)$ if and only if $\limsup_{t \rightarrow 0} \|\nabla^2 E_t(q_1, q_2)\| = \infty$, where $\|\cdot\|$ is the operator norm.

The first result is due to Varadhan [34], the second one to Neel and Staprook [29, 30]. See also [7] for some extensions in sub-Riemannian geometry.

Both these results do not depend upon the fact that one is using the Laplace-Beltrami operator or the principal symbol of the operator in a fixed frame to construct the heat kernel.

The situation is very different in 2-ARS. Both these results are false for the Laplace Beltrami operator. Indeed the distance between two points belonging to two different connected components of $M \setminus \mathcal{Z}$ is finite while if q belongs to a connected component W of $M \setminus \mathcal{Z}$ then $p_t(q, \cdot)$ is supported in W .

However, a result in the spirit of the one of Varadhan (which applies to the context of 2-ARG) has been obtained by Leandre in [26] for the operator “sum of squares”. Hence, the Laplace-Beltrami operator defined above, has quite different properties with respect to the operator sum of squares. The last one is not intrinsic, but still keeps tracks of intrinsic quantities as the almost-Riemannian distance. In particular, the corresponding heat flow crosses the set \mathcal{Z} which is not the case for the Laplace-Beltrami operator. For relations among the heat kernel on the diagonal and the distance in sub-Riemannian geometry see [6].

To prove Theorem 1.1 we start by analyzing the Grushin case (1.6). We first compactify in the y variable by considering it on $\mathbf{R}_x \times \mathbb{T}_y$.

By setting $f = \sqrt{|x|}g$ we are reduced to study the essential self adjointness of the following operator considered on L^2 with the usual euclidian metric

$$L = \partial_x^2 + x^2 \partial_y^2 - \frac{3}{4} \frac{1}{x^2}. \quad (1.9)$$

If for a moment we forget the term $x^2 \partial_y^2$ which is not relevant for the self-adjointness of this operator (and becomes a non-positive potential $-k^2 x^2$ after performing Fourier transform in y), we are reduced to study the operator $\partial_x^2 - \frac{3}{4} \frac{1}{x^2}$, which is well known in the literature. Indeed we have the following result

PROPOSITION 1.5. — *The operator $-\partial_x^2 + \frac{c}{|x|^2}$ defined on $L^2(]0, +\infty[)$ with domain $C_0^\infty(]0, +\infty[)$ is essentially self-adjoint if and only if $c \geq \frac{3}{4}$.*

In the literature there are different proofs of this result (see [32] Chapter X for an introduction), often leading to some stronger statements (for instance for potential $V \geq \frac{3}{4x^2}$) and to generalizations to higher dimensions.

The rest of the proof for an almost Riemannian structure consists in generalizing this result for a normal form around a connected component

of the singular set. The main tools are Kato inequality and perturbation theory.

Remark 1.6. — Notice that, as remarked in the introduction of [22], solutions of the Schrödinger equation, with the Laplacian defined as sum of square $\partial_x^2 + x^2 \partial_y^2$, display a total lack of dispersion: one can not obtain Strichartz type estimates better than what we could expect by Sobolev embedding. It would be interesting to understand if the same holds for the Laplace-Beltrami operator defined by (1.6). Moreover it would be interesting to study if a wave packet moving towards the singular set is reflected or dispersed.

The structure of the paper is the following. In section 2 we briefly recall the notion of almost-Riemannian structure. An original result is given in Proposition 2.3, in which we globalize the normal form of type (F_2) around a connected component of \mathcal{Z} in the compact orientable case and under the assumption that there are no tangency points. In Section 3 we analyze the Grushin case, both for what concerns geodesics and the self-adjointness of the Laplace-Beltrami operator. The main result is proven in Section 4.

As a byproduct of our studies, we obtain that the Laplace-Beltrami operator is not essentially self-adjoint when an orthonormal basis is given by the vectors $(1, 0)$, $(0, |x|^\alpha)$ with $\alpha \in (0, 1)$ (see remark 3.1). This fact motivates some remarks on metrics of this kind (which do not enter in the standard framework of almost-Riemannian geometry, since the vector fields are not smooth). For instance, we show that in this case, even if the area element explodes, the area of a bounded open set intersecting \mathcal{Z} is finite (which is false for 2-ARSs). Moreover, we show that there exist regular curves passing through \mathcal{Z} with a velocity not belonging to the span of X_1 and X_2 , having finite length. See Section 3.3.

Finally, in the appendix, we discuss the Martinet case to show that this kind of phenomenon appears also in sub-Riemannian structure of constant rank but which are not equiregular, (i.e., sub-Riemannian structure whose growth vector depends on the point, see for instance [28]). In this case the role of the Laplace-Beltrami operator is played by the intrinsic sub-Riemannian Laplacian defined via the Popp measure (see [3, 28]). While approaching the Martinet surface, the Popp volume explodes and so do the first order coefficients of the sub-Riemannian Laplacian.

THEOREM 1.7. — *Consider the sub-Riemannian structure in $M = \mathbb{T}_x \times \mathbb{R}_y \times \mathbb{T}_z$ for which an orthonormal basis is given by $X_1 = (1, 0, \frac{y^2}{2})$, $X_2 = (0, 1, 0)$. Then the corresponding intrinsic sub-Riemannian Laplacian which*

is given by $\Delta_{sr} = (X_1)^2 + (X_2)^2 - \frac{1}{y}X_2$ with domain $C_0^\infty(M \setminus \{y = 0\})$ is essentially self-adjoint on $L^2(M, d\omega)$, where $d\omega = \frac{1}{|y|} dx dy dz$ is the Popp volume.

2. Basic Definitions

In this section we recall some basic definitions in the framework of 2-ARSs following [4, 2].

Let M be a smooth surface without boundary. Throughout the paper, unless specified, manifolds are smooth (i.e., C^∞) and without boundary; vector fields and differential forms are smooth. The set of smooth vector fields on M is denoted by $\text{Vec}(M)$. The circle is denoted by S^1 or \mathbb{T} depending on the context. C_0^∞ denotes the set of smooth functions with compact support.

2.1. 2-Almost-Riemannian Structures

DEFINITION 2.1. — A 2-dimensional almost-Riemannian structure (2-ARS, for short) is a triple $\mathcal{S} = (E, \mathfrak{f}, \langle \cdot, \cdot \rangle)$ where E is a vector bundle of rank 2 over M and $\langle \cdot, \cdot \rangle$ is a Euclidean structure on E , that is, $\langle \cdot, \cdot \rangle_q$ is a scalar product on E_q smoothly depending on q . Finally $\mathfrak{f} : E \rightarrow TM$ is a morphism of vector bundles, i.e., **(i)** the diagram

$$\begin{array}{ccc} E & \xrightarrow{\mathfrak{f}} & TM \\ & \searrow \pi_E & \downarrow \pi \\ & & M \end{array}$$

commutes, where $\pi : TM \rightarrow M$ and $\pi_E : E \rightarrow M$ denote the canonical projections and **(ii)** \mathfrak{f} is linear on fibers.

Denoting by $\Gamma(E)$ the $C^\infty(M)$ -module of smooth sections on E , and by $\mathfrak{f}_* : \Gamma(E) \rightarrow \text{Vec}(M)$, the map $\sigma \mapsto \mathfrak{f}_*(\sigma) := \mathfrak{f} \circ \sigma$. We require that the submodule of $\text{Vec}(M)$ given by $\blacktriangle = \mathfrak{f}_*(\Gamma(E))$ to be bracket generating, i.e., $\text{Lie}_q(\blacktriangle) = T_qM$ for every $q \in M$.

Here $\text{Lie}(\blacktriangle)$ is the smallest Lie subalgebra of $\text{Vec}(M)$ containing \blacktriangle and $\text{Lie}_q(\blacktriangle)$ is the linear subspace of T_qM whose elements are evaluation at q of elements belonging to $\text{Lie}(\blacktriangle)$. The condition that \blacktriangle satisfies the Lie bracket generating assumption is also known as the Hörmander condition.

We say that a 2-ARS $(E, \mathfrak{f}, \langle \cdot, \cdot \rangle)$ is orientable if E is orientable as a vector bundle. Notice that one can build non-orientable 2-ARSs on orientable

manifolds and orientable 2-ARSs on non-orientable manifolds. See [4] for some examples. We say that a 2-ARS $(E, \mathfrak{f}, \langle \cdot, \cdot \rangle)$ is trivializable if E is isomorphic to the trivial bundle $M \times \mathbf{R}^2$. A particular case of 2-ARSs is given by Riemannian surfaces. In this case $E = TM$ and \mathfrak{f} is the identity.

Let $\mathcal{S} = (E, \mathfrak{f}, \langle \cdot, \cdot \rangle)$ be a 2-ARS on a surface M . We denote by $\blacktriangle(q)$ the linear subspace $\{V(q) \mid V \in \blacktriangle\} = \mathfrak{f}(E_q) \subseteq T_qM$. The set of points in M such that $\dim(\blacktriangle(q)) < 2$ is called *singular set* and denoted by \mathcal{Z} . Since \blacktriangle is bracket generating, the subspace $\blacktriangle(q)$ is nontrivial for every q and \mathcal{Z} coincides with the set of points q where \blacktriangle is one-dimensional.

The Euclidean structure on E allows to define a symmetric positive definite $C^\infty(M)$ -bilinear form on the submodule \blacktriangle by

$$G : \blacktriangle \times \blacktriangle \rightarrow C^\infty(M)$$

$$G(V, W) = \langle \sigma_V, \sigma_W \rangle$$

where σ_V, σ_W are the unique⁽¹⁾ sections of E satisfying $\mathfrak{f} \circ \sigma_V = V, \mathfrak{f} \circ \sigma_W = W$.

At points $q \in M$ where $\mathfrak{f}|_{E_q}$ is an isomorphism, i.e., on $M \setminus \mathcal{Z}$, G is a tensor and the value $G(V, W)|_q$ depends only on $V(q), W(q)$. In this case G defines a Riemannian metric g via

$$g_q(v, w) = G(V, W)(q), \quad v, w \in T_qM,$$

where V and W are two vector fields such that $v = V(q)$ and $w = W(q)$.

This is no longer true at points q where $\mathfrak{f}|_{E_q}$ is not injective.

If (σ_1, σ_2) is an orthonormal frame for $\langle \cdot, \cdot \rangle$ on an open subset Ω of M , an *orthonormal frame for G* on Ω is given by $(\mathfrak{f} \circ \sigma_1, \mathfrak{f} \circ \sigma_2)$. Orthonormal frames are systems of local generators of \blacktriangle .

For every $q \in M$ and every $v \in \blacktriangle(q)$ define

$$\mathbf{G}_q(v) = \inf\{\langle u, u \rangle_q \mid u \in E_q, \mathfrak{f}(u) = v\}.$$

An absolutely continuous curve $\gamma : [0, T] \rightarrow M$ is admissible for \mathcal{S} if there exists a measurable essentially bounded function

$$[0, T] \ni t \mapsto u(t) \in E_{\gamma(t)},$$

called *control function* such that $\dot{\gamma}(t) = \mathfrak{f}(u(t))$ for almost every $t \in [0, T]$. Given an admissible curve $\gamma : [0, T] \rightarrow M$, the *length of γ* is

$$\ell(\gamma) = \int_0^T \sqrt{\mathbf{G}_{\gamma(t)}(\dot{\gamma}(t))} \, dt.$$

⁽¹⁾ the uniqueness is consequence of the fact that we assume E of rank two and \blacktriangle Lie bracket generating.

The *Carnot-Caratheodory distance* (or sub-Riemannian distance) on M associated with \mathcal{S} is defined as

$$d(q_0, q_1) = \inf\{\ell(\gamma) \mid \gamma(0) = q_0, \gamma(T) = q_1, \gamma \text{ admissible}\}.$$

It is a standard fact that $\ell(\gamma)$ is invariant under reparameterization of the curve γ . Moreover, if an admissible curve γ minimizes the so-called *energy functional* $E(\gamma) = \int_0^T \mathbf{G}_{\gamma(t)}(\dot{\gamma}(t)) dt$ with T fixed (and fixed initial and final point) then $v = \sqrt{\mathbf{G}_{\gamma(t)}(\dot{\gamma}(t))}$ is constant and γ is also a minimizer of $\ell(\cdot)$. On the other hand, a minimizer γ of $\ell(\cdot)$, such that v is constant, is a minimizer of $E(\cdot)$ with $T = \ell(\gamma)/v$.

The finiteness and the continuity of $d(\cdot, \cdot)$ with respect to the topology of M are guaranteed by the Lie bracket generating assumption on the 2-ARS (see [5]). The Carnot-Caratheodory distance endows M with the structure of metric space compatible with the topology of M as differential manifold.

When the 2-ARS is trivializable, the problem of finding a curve minimizing the energy between two fixed points $q_0, q_1 \in M$ is naturally formulated as the distributional optimal control problem with quadratic cost and fixed final time

$$\dot{q} = \sum_{i=1}^2 u_i X_i(q), \quad u_i \in \mathbf{R}, \quad \int_0^T \sum_{i=1}^2 u_i^2(t) dt \rightarrow \min, \quad q(0) = q_0, \quad q(T) = q_1.$$

where $\{X_1, X_2\}$ is an orthonormal frame.

2.2. Geodesics

A *geodesic* for \mathcal{S} is a curve $\gamma : [0, T] \rightarrow M$ such that for every sufficiently small nontrivial interval $[t_1, t_2] \subset [0, T]$, $\gamma|_{[t_1, t_2]}$ is a minimizer of $E(\cdot)$. A geodesic for which $\mathbf{G}_{\gamma(t)}(\dot{\gamma}(t))$ is (constantly) equal to one is said to be parameterized by arclength. The local existence of minimizing geodesics is a standard consequence of Filippov Theorem (see for instance [5]). When M is compact any two points of M are connected by a minimizing geodesic.

Locally, in an open set Ω , if $\{X_1, X_2\}$ is an orthonormal frame, a curve parameterized by arclength is a geodesic if and only if it is the projection on Ω of a solution of the Hamiltonian system corresponding to the Hamiltonian

$$H(q, \lambda) = \frac{1}{2}((\lambda X_1(q))^2 + (\lambda X_2(q))^2), \quad q \in \Omega, \quad \lambda \in T_q^* \Omega. \quad (2.1)$$

lying on the level set $H = 1/2$. This is the Pontryagin Maximum Principle [31] in the case of 2-ARSs. Its simple form follows from the absence of abnormal extremals in 2-ARSs, as a consequence of the Hörmander condition

see [4]. Notice that when looking for a geodesic γ minimizing the energy from a submanifold \mathcal{T} (possibly of dimension zero), one should add the transversality condition $\lambda(0)T_{\gamma(0)}\mathcal{T} = 0$.

2.3. Generic 2-ARSs and normal forms

A property (P) defined for 2-ARSs is said to be *generic* if for every rank-2 vector bundle E over M , (P) holds for every f in an open and dense subset of the set of morphisms of vector bundles from E to TM , endowed with the C^∞ -Whitney topology.

Define $\blacktriangle_1 = \blacktriangle$ and $\blacktriangle_{k+1} = \blacktriangle_k + [\blacktriangle, \blacktriangle_k]$. We say that \mathcal{S} satisfies condition **(H0)** if the following properties hold: **(i)** \mathcal{Z} is an embedded one-dimensional submanifold of M ; **(ii)** the points $q \in M$ at which $\blacktriangle_2(q)$ is one-dimensional are isolated; **(iii)** $\blacktriangle_3(q) = T_qM$ for every $q \in M$. It is not difficult to prove that property **(H0)** is generic among 2-ARSs (see [4]). This hypothesis was essential to show Gauss–Bonnet type results for ARSs in [4, 2, 18]. The following theorem recalls the local normal forms for ARSs satisfying hypothesis **(H0)** (see Figure 1.1).

THEOREM 2.2 ([4]). — *Consider a 2-ARS satisfies **(H0)**. Then for every point $q \in M$ there exist a neighborhood U of q and an orthonormal frame $\{X_1, X_2\}$ of the ARS on U such that, up to a change of coordinates, $q = (0, 0)$ and $\{X_1, X_2\}$ has one of the forms*

$$\begin{aligned} \text{(F1)} \quad X_1(x, y) &= \frac{\partial}{\partial x}, & X_2(x, y) &= e^{\phi(x, y)} \frac{\partial}{\partial y}, \\ \text{(F2)} \quad X_1(x, y) &= \frac{\partial}{\partial x}, & X_2(x, y) &= xe^{\phi(x, y)} \frac{\partial}{\partial y}, \\ \text{(F3)} \quad X_1(x, y) &= \frac{\partial}{\partial x}, & X_2(x, y) &= (y - x^2\psi(x))e^{\xi(x, y)} \frac{\partial}{\partial y}, \end{aligned}$$

where ϕ, ψ and ξ are smooth functions such that $\phi(0, y) = 0$ and $\psi(0) \neq 0$.

Let \mathcal{S} be a 2-ARS satisfying **(H0)**. A point $q \in M$ is said to be an *ordinary point* if $\blacktriangle(q) = T_qM$, hence, if \mathcal{S} is locally described by (F1). We call q a *Grushin point* if $\blacktriangle(q)$ is one-dimensional and $\blacktriangle_2(q) = T_qM$, i.e., if the local description (F2) applies. Finally, if $\blacktriangle(q) = \blacktriangle_2(q)$ has dimension one and $\blacktriangle_3(q) = T_qM$ then we say that q is a *tangency point* and \mathcal{S} can be described near q by the normal form (F3).

Notice that under hypotheses **HA** and **HB** of Theorem 1.1, **H0** is fulfilled and there are no tangency points.

In the compact case, the following proposition permits to extend the normal form (F2) to a neighborhood of a connected component of \mathcal{Z} , when there are no tangency points.

PROPOSITION 2.3. — Consider a 2-ARS on a compact orientable manifold satisfying **(H0)**. Let W be a connected component of \mathcal{Z} containing no tangency points. Then there exists a tubular neighborhood U of W and an orthonormal frame $\{X_1, X_2\}$ of the 2-ARS on U such that, up to a change of coordinates, $W = \{(0, y), y \in S^1\}$ and $\{X_1, X_2\}$ has the form

$$X_1(x, y) = \frac{\partial}{\partial x}, \quad X_2(x, y) = xe^{\phi(x, y)} \frac{\partial}{\partial y}. \quad (2.2)$$

Proof. — The proof consists in using as first coordinates the distance from W and it is very similar to the proof of Theorem 2.2, which contains a local version of Proposition 2.3 (see [4] pp. 813–814, proof of Lemma 1 and proof of Theorem 1 for Grushin points). Here we just explain which modifications are necessary.

Since M is assumed to be compact then W is diffeomorphic to S^1 . Moreover, since M is assumed to be orientable, a sufficiently small open tubular neighborhood of W is diffeomorphic to $(-1, 1) \times S^1$. Then consider a smooth regular parametrization $S^1 \ni \alpha \mapsto w(\alpha)$ of W .

Let $\alpha \mapsto \lambda_0(\alpha) \in T_{w(\alpha)}^*M$ be a smooth map satisfying $H(\lambda_0(\alpha), w(\alpha)) = 1/2$, where H is the Hamiltonian of the PMP (2.1) and $\lambda_0(\alpha) \perp T_{w(\alpha)}W$. Notice that such a map exists since we are assuming that M is orientable.

Let $E(t, \alpha)$ be the projection on the q variables of the solution at time t of the Hamiltonian system given by the Pontryagin Maximum Principle with initial condition $(q(0), \lambda(0)) = (w(\alpha), \lambda_0(\alpha))$. With the same arguments given in the proof of Lemma 1 in [4], one shows that $E(t, \alpha)$ is a local diffeomorphism around every point of the type $(0, \bar{\alpha})$, $\bar{\alpha} \in S^1$. Using the fact that E is a global diffeomorphism from $\{0\} \times S^1$ to W and by suitably reducing ε one gets that it is a diffeomorphism between $(-\varepsilon, \varepsilon) \times S^1$ (for some $\varepsilon > 0$) and a tubular neighborhood of W . This permits to use as coordinates in U the pair (t, α) . Indeed t is the distance from W which we proved to be smooth in U . As in [4], one builds an orthonormal frame in U defining the vector field X_1 by

$$X_1(t, \alpha) = \partial_t E(t, \alpha) = \partial_t.$$

As in [4] one obtains that the second vector field of the orthonormal frame has to be of the form $te^{\phi(t, \alpha)} \frac{\partial}{\partial \alpha}$. \square

Remark 2.4. — Notice that in Proposition 2.3, the hypothesis that M is orientable can be weakened by requiring that W admits an open tubular neighborhood diffeomorphic to $(-1, 1) \times S^1$

3. The Grushin case

As already mentioned the Grushin plane is the trivializable almost Riemannian metric on the (x, y) plane for which an orthonormal basis is given by

$$X_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, X_2 = \begin{pmatrix} 0 \\ x \end{pmatrix}. \tag{3.1}$$

In the sense of Definition 2.1 it can be seen as a triple $(E, \mathfrak{f}, \langle \cdot, \cdot \rangle)$ where $E = \mathbf{R}^2 \times \mathbf{R}^2$, $\mathfrak{f}((x, y), (a, b)) = ((x, y), (a, bx))$ and $\langle \cdot, \cdot \rangle$ is the standard Euclidean metric.

3.1. Geodesics of the Grushin plane

In this section we recall how to compute the geodesics for the Grushin plane, with the purpose of stressing that they can cross the singular set with no singularities.

Setting $q = (x, y)$ and $\lambda = (\lambda_1, \lambda_2)$, the Hamiltonian (2.1) is given by

$$H(x, y, \lambda_1, \lambda_2) = \frac{1}{2}(\lambda_1^2 + x^2\lambda_2^2) \tag{3.2}$$

and the corresponding Hamiltonian equations are:

$$\begin{aligned} \dot{x} &= \lambda_1, & \dot{\lambda}_1 &= -x\lambda_2^2 \\ \dot{y} &= x^2\lambda_2, & \dot{\lambda}_2 &= 0 \end{aligned} \tag{3.3}$$

Geodesics parameterized by arclength are projections on the (x, y) plane of solutions of these equations, lying on the level set $H = 1/2$. We study the geodesics starting from **i)** a Grushin point, e.g. $(0, 0)$ **ii)** an ordinary point, e.g. $(-1, 0)$.

Case $(x(0), y(0)) = (0, 0)$

In this case the condition $H(x(0), y(0), \lambda_1(0), \lambda_2(0)) = 1/2$ implies that we have two families of geodesics corresponding respectively to $\lambda_1(0) = \pm 1$, $\lambda_2(0) =: a \in \mathbf{R}$. Their expression can be easily obtained and it is given by:

$$\begin{cases} x(t) = \pm t, & y(t) = 0 & \text{if } a = 0 \\ x(t) = \pm \frac{\sin(at)}{a}, & y(t) = \frac{2at - \sin(2at)}{4a^2} & \text{if } a \neq 0 \end{cases} \tag{3.4}$$

Some geodesics are plotted in Figure 3.1 together with the “front”, i.e., the end point of all geodesics at time $t = 1$. Notice that geodesics start horizontally. The particular form of the front shows the presence of a conjugate locus accumulating to the origin.




Figure 3.1. Geodesics and the front for the Grushin plane, starting from the singular set.

Case $(x(0), y(0)) = (-1, 0)$

In this case the condition $H(x(0), y(0), \lambda_1(0), \lambda_2(0)) = 1/2$ becomes $\lambda_1^2 + \lambda_2^2 = 1$ and it is convenient to set $\lambda_1 = \cos(\theta), \lambda_2 = \sin(\theta), \theta \in S^1$. The expression of the geodesics is given by:

$$\left\{ \begin{array}{l} x(t) = t - 1, \quad y(t) = 0, \quad \text{if } \theta = 0 \\ x(t) = -t - 1, \quad y(t) = 0, \quad \text{if } \theta = \pi \\ \left. \begin{array}{l} x(t) = -\frac{\sin(\theta - t \sin(\theta))}{\sin(\theta)}, \\ y(t) = \frac{2t - 2 \cos(\theta) + \frac{\sin(2\theta - 2t \sin(\theta))}{\sin(\theta)}}{4 \sin(\theta)} \end{array} \right\} \text{if } \theta \notin \{0, \pi\}$$

Some geodesics are plotted in Figure 3.2 together with the “front” at time $t = 4.8$. Notice that geodesics pass horizontally through \mathcal{Z} , with no singularities. The particular form of the front shows the presence of a conjugate locus. Geodesics can have conjugate times only after intersecting \mathcal{Z} . Before it is impossible since they are Riemannian and the curvature is negative.




Figure 3.2. Geodesics and the front for the Grushin plane, starting from a Riemannian point.

3.2. The Laplace-Beltrami operator on the Grushin plane

In this section we illustrate in details the results of the paper in the Grushin case. For the study of the Laplace Beltrami operator it will be convenient to compactify the y direction considering $x \in \mathbf{R}$ and $y \in \mathbb{T}$.

As mentioned in the introduction, the Laplace-Beltrami operator on the Grushin plane can be written as

$$\Delta = \partial_x^2 + x^2 \partial_y^2 - \frac{1}{x} \partial_x.$$

We expect that the Laplacian is the Friedrichs extension associated to the positive quadratic form

$$Q(u, u) = \int_{\mathbf{R}^2} |\nabla u|^2 d\omega = \int_{\mathbf{R}^2} \frac{1}{|x|} (|\partial_x u|^2 + |x| |\partial_y u|^2) dx dy$$

Let us make the change of variable in Hilbert space $f = \sqrt{|x|}g$ which is unitary from $L^2(\mathbf{R} \times \mathbb{T}, d\omega)$ to $L^2(\mathbf{R} \times \mathbb{T}, dx dy)$, so that $(f_1, f_2)_{L^2(\mathbf{R} \times \mathbb{T}, d\omega)} = (g_1, g_2)_{L^2(\mathbf{R} \times \mathbb{T}, dx dy)}$.

We compute the operator in the new variable:

$$\begin{aligned} \Delta f &= \partial_x^2 f + x^2 \partial_y^2 f - \frac{1}{|x|} \partial_x f \\ &= \sqrt{|x|} \left[\partial_x^2 g + x^2 \partial_y^2 g - \frac{3}{4x^2} g \right] =: \sqrt{|x|} Lg. \end{aligned}$$

Hence we are left to study the operator $Lg = \partial_x^2 g + x^2 \partial_y^2 g - \frac{3}{4x^2} g$ on $L^2(\mathbf{R} \times \mathbb{T}, dx dy)$. By decomposing a function $g(x, y)$ in the Fourier basis in the variable y , we get the following decomposition

$$L^2(\mathbf{R} \times \mathbb{T}) \approx \bigoplus_{k \in \mathbf{Z}}^\perp H_k$$

where $H_k = L^2(\mathbf{R})$ and the operator L acts on each H_k by

$$(Lu)_k(x) = \partial_x^2 u_k - x^2 k^2 u_k - \frac{3}{4x^2} u_k = \partial_x^2 u_k - V_k u_k.$$

with $V_k(x) \geq \frac{3}{4x^2}$. But, we know that in dimension 1, the operator $-\partial_x^2 + V$ with domain $C_0^\infty(]0, +\infty[)$ is essentially self-adjoint on $L^2(]0, +\infty[)$ if $V \geq \frac{3}{4x^2}$ (see [32], Theorem X.10 for the proof of the limit point case at 0 and Theorem X.8 at $+\infty$). Hence, each operator $\partial_x^2 u_k - V_k$ is essentially self-adjoint. As a consequence, L is essentially self-adjoint as well.

Remark 3.1. — $V = \frac{3}{4x^2}$ is exactly the limit singular potential to have $-\partial_x^2 + V$ in the limit point case at zero and hence essentially self-adjoint (see [32, Theorem X.10, page 159]). It is surprising to find the same constant $\frac{3}{4}$ for the Grushin operator.

Notice that the Laplace-Beltrami operator for the generalized Riemannian structure having $X = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $Y = \begin{pmatrix} 0 \\ |x|^\alpha \end{pmatrix}$ with $\alpha > 0$ as orthonormal frame, is given by

$$\Delta u = \partial_x^2 u + |x|^{2\alpha} \partial_y^2 u - \frac{\alpha}{x} \partial_x u.$$

By making the change of variable in Hilbert space $u = |x|^{\alpha/2} v$ (which is unitary from $L^2(\mathbf{R} \times \mathbb{T}, d\omega)$ to $L^2(\mathbf{R} \times \mathbb{T}, dx dy)$) we get for the transformed operator

$$\tilde{\Delta} v = \partial_x^2 v - \frac{\alpha}{2} \left(\frac{\alpha}{2} + 1 \right) \frac{1}{x^2} + |x|^{2\alpha} \partial_y^2 v,$$

which is essentially self-adjoint if and only if $\alpha \in [1, +\infty[$.

3.3. $^{1/\alpha}\sqrt{\text{Grushin}}$

In this section, motivated by Remark 3.1, we briefly discuss the generalized Riemannian structures on \mathbf{R}^2 for which an orthonormal basis is given

by

$$X_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, X_2 = \begin{pmatrix} 0 \\ |x|^\alpha \end{pmatrix}, \alpha \in (0, 1). \quad (3.5)$$

Notice that this generalized-Riemannian structure does not fit the definition of almost-Riemannian structures, since the vector fields are not smooth. As usual, call \mathcal{Z} the set in which the two vector fields are not linearly independent. It is interesting to notice the following issues, which in some cases are very different with respect to the standard Grushin case:

- The Lie bracket between X_1 and X_2 is not well defined on \mathcal{Z} . Indeed $[X_1, X_2] = (0, \alpha|x|^{\alpha-1})^T$. However, it is not difficult to prove that the corresponding distance is continuous and endows \mathbf{R}^2 with its standard topology (note that it is not the case if $\alpha < 0$).
- The curvature is given by $K = -\frac{\alpha(1+\alpha)}{x^2}$ and the area element by $d\omega = \frac{1}{|x|^\alpha} dx dy$. Hence, the area element explodes but the area of a bounded open set intersecting \mathcal{Z} is finite (and its diameter is finite as well). This is certainly the reason why the Laplace-Beltrami operator is not essentially self-adjoint and, as a consequence, a quantum particle or the heat flow can pass through the set \mathcal{Z} . Notice that for the Grushin plane a bounded open set intersecting \mathcal{Z} has finite diameter but infinite area.
- There exists regular curves passing through \mathcal{Z} with a velocity not belonging to the span of X_1 and X_2 and having finite length. For instance take $\alpha = 1/2$ and the curve $\gamma(t) := (t, t)$, $t \in [-1, 1]$. We have $\dot{\gamma}(0) \notin \text{span}\{X_1(\gamma(0)), X_2(\gamma(0))\}$ but its length is finite. Indeed it corresponds to the controls $u_1 = 1$ and $u_2 = 1/|t|^{1/2}$. Hence, $\ell(\gamma) = \int_{-1}^1 \sqrt{1 + |t|^{-1}} dt = 2(\sqrt{2} + \text{arcsinh}(1)) \sim 4.59$. Such phenomenon does not appear on the singular set of a 2-ARS, as a consequence of the smoothness of orthonormal frames.

4. Laplace-Beltrami on an Almost-Riemannian structure

4.1. Computation in local coordinates

The aim of this subsection is the proof of the following theorem.

THEOREM 4.1. — *Set $\Omega = \{x = (x, y) \in \mathbf{R} \times \mathbb{T} \mid x \neq 0\}$ and let g a Riemannian metric on Ω of the form $\text{diag}(1, x^{-2}e^{-2\Phi(x,y)})$ where Φ is a smooth function which is constant for large $|x|$. Let Δ be the corresponding Laplace-Beltrami operator with domain $C_0^\infty(\Omega)$.*

Then, Δ is essentially self-adjoint for $L^2(\mathbf{R} \times \mathbb{T}, d\omega)$, where $d\omega$ is the corresponding Riemannian area on Ω .

The same result holds by replacing Ω with Ω_{\pm} where

$$\begin{aligned} \Omega_- &= \{x = (x, y) \in \mathbf{R} \times \mathbb{T} \mid x < 0\}, \\ \Omega_+ &= \{x = (x, y) \in \mathbf{R} \times \mathbb{T} \mid x > 0\}. \end{aligned}$$

A simple consequence of this theorem is that the only self-adjoint Laplace operator that can be constructed by extension of the natural one (i.e., the Laplace-Beltrami operator defined on $C_0^\infty(\Omega)$) preserves the decomposition $L^2(\Omega, d\omega) = L^2(\Omega_-, d\omega) \oplus^\perp L^2(\Omega_+, d\omega)$.

Indeed, if we denote Δ^- (resp Δ^+) the unique self-adjoint extension of the natural Laplace operator with domain $C_0^\infty(\Omega_-)$ (resp $C_0^\infty(\Omega_+)$), we can define the operator $\bar{\Delta}(u_- + u_+) = \Delta^-u_- + \Delta^+u_+$ with the natural domain inherited from the one of Δ^- and Δ^+ . The operator $\bar{\Delta}$ is a self-adjoint extension of the Laplace-Beltrami operator with domain $C_0^\infty(\Omega)$ and it is the unique one by Theorem 4.1. Another way of seeing this fact is from the point of view of the evolution equation.

COROLLARY 4.2. — *With the notations of Theorem 4.1, consider the unique solution u of the Schrödinger equation (according to the self-adjoint extension defined in the previous theorem),*

$$\begin{cases} i\partial_t u + \Delta u &= 0 \\ u(0) &= u_0 \in L^2(\mathbf{R} \times \mathbb{T}, d\omega) \end{cases} \tag{4.1}$$

with u_0 supported in Ω_+ . Then, $u(t)$ is supported in Ω_+ for any $t \geq 0$. The same holds for the solution of the heat or for the solution of the wave equation.

This corollary is a simple consequence of the essential self-adjointness of the Laplace-Beltrami operator as discussed above.

This is in strong contrast with the classical dynamics associated with this metric where the geodesics can "cross" the barrier $\{x = 0\}$ (see Figure 3.2). The proof of Theorem 4.1 relies on a change of variable in Hilbert space similar to the one we did for the Grushin operator which leads to an operator that can be written as "Grushin type + singular potential". Then, we follow the proof of Kalf and Walter [24] (which was himself inspired by B. Simon [33]) using Kato inequality for the main part. The other part can be treated by perturbation theory. Note that some related results for singular potential were proven in [19] and [27].

Proof. — [Proof of Theorem 4.1] In coordinates, the Laplace-Beltrami operator has the form

$$\Delta = \frac{1}{\omega} \sum_{i,j} \partial_i [\omega g^{ij} \partial_j]$$

where we have denoted $\omega = \sqrt{\det g}$ the Riemannian volume.

For $u \in L^2(\mathbf{R} \times \mathbb{T}, d\omega)$, let us make the change of variable in Hilbert space $u = \omega^{-1/2}v$ which is unitary from $L^2(\mathbf{R} \times \mathbb{T}, d\omega)$ to $L^2(\mathbf{R} \times \mathbb{T}, dx dy)$ and let us compute its action in the new variable.

$$\begin{aligned} & \omega^{1/2} \Delta \omega^{-1/2} v := \tilde{\Delta} v \\ &= \frac{1}{\omega^{1/2}} \sum_{i,j} \partial_i [\omega g^{ij} \partial_j (\omega^{-1/2} v)] \\ &= \frac{1}{\omega^{1/2}} \sum_{i,j} \partial_i [\omega g^{ij} (\partial_j \omega^{-1/2}) v + g^{ij} (\omega^{1/2}) \partial_j v] \\ &= \frac{1}{\omega^{1/2}} \sum_{i,j} [\partial_i [g^{ij} \omega (\partial_j \omega^{-1/2})] v + g^{ij} \omega (\partial_j \omega^{-1/2}) \partial_i v + (\partial_i g^{ij}) \omega^{1/2} \partial_j v \\ &\quad + g^{ij} (\partial_i \omega^{1/2}) \partial_j v + g^{ij} (\omega^{1/2}) \partial_{ij} v] \\ &= \frac{1}{\omega^{1/2}} \sum_{i,j} [\partial_i [g^{ij} \omega (\partial_j \omega^{-1/2})] v + g^{ij} \omega (\partial_i \omega^{-1/2}) \partial_j v + (\partial_i g^{ij}) \omega^{1/2} \partial_j v \\ &\quad + g^{ij} (\partial_i \omega^{1/2}) \partial_j v + g^{ij} (\omega^{1/2}) \partial_{ij} v], \end{aligned}$$

where we have used the symmetry of g .

We have: $\omega (\partial_i \omega^{-1/2}) + (\partial_i \omega^{1/2}) = -\frac{1}{2} \omega \omega^{-3/2} \partial_i \omega + \frac{1}{2} \omega^{-1/2} \partial_i \omega = 0$. Moreover let

$$\begin{aligned} \mu &:= \sum_{i,j} \frac{1}{\omega^{1/2}} \partial_i [g^{ij} \omega (\partial_j \omega^{-1/2})] = - \sum_{i,j} \frac{1}{2\omega^{1/2}} \partial_i [g^{ij} (\partial_j \omega) \omega^{-1/2}] \\ &= \sum_{i,j} \left[-\frac{1}{2} \partial_i (g^{ij}) \partial_j (\ln \omega) - \frac{1}{2\omega} g^{ij} \partial_{ij} \omega + \frac{1}{4\omega^2} g^{ij} (\partial_i \omega) (\partial_j \omega) \right]. \end{aligned}$$

Hence we get

$$\begin{aligned} \tilde{\Delta} v &= \sum_{i,j} g^{ij} \partial_{ij} v + (\partial_i g^{ij}) \partial_j v + \mu v \\ &= \operatorname{div}_e (\nabla v) + \mu v \\ &= \operatorname{div}_e (G^{-1} \nabla_e v) + \mu v. \end{aligned}$$

In the last formula we have used matrix notation where $G^{-1} = (g^{ij})$ and $\operatorname{div}_e, \nabla_e$ are the divergence and the gradient of the Euclidean space. Now,

let us specify the computation in our diagonal metric.

$$\begin{aligned}
 \omega &= \frac{1}{|x|} e^{-\Phi}, \\
 \partial_1 \omega &= -\frac{\operatorname{sgn}(x)}{|x|^2} e^{-\Phi} - \frac{\partial_1 \Phi}{|x|} e^{-\Phi}, \\
 \partial_2 \omega &= -\frac{\partial_2 \Phi}{|x|} e^{-\Phi}; \quad \partial_2^2 \omega = \frac{(\partial_2 \Phi)^2}{|x|} e^{-\Phi} - \frac{\partial_2^2 \Phi}{|x|} e^{-\Phi}, \\
 \partial_1^2 \omega &= \frac{2}{|x|^3} e^{-\Phi} + \frac{2 \operatorname{sgn}(x) \partial_1 \Phi}{|x|^2} e^{-\Phi} + \frac{(\partial_1 \Phi)^2}{|x|} e^{-\Phi} - \frac{\partial_1^2 \Phi}{|x|} e^{-\Phi}, \\
 \mu &= -\frac{1}{|x|^2} - \frac{\operatorname{sgn}(x) \partial_1 \Phi}{|x|} - \frac{(\partial_1 \Phi)^2}{2} + \frac{\partial_1^2 \Phi}{2} \\
 &\quad + \frac{1}{4|x|^2} + \frac{\operatorname{sgn}(x) \partial_1 \Phi}{2|x|} + \frac{(\partial_1 \Phi)^2}{4} \\
 &\quad + x^2 e^{2\Phi} \left[(\partial_2 \Phi)^2 + \frac{1}{2} [\partial_2^2 \Phi - (\partial_2 \Phi)^2] + \frac{1}{4} (\partial_2 \Phi)^2 \right] \\
 &= -\frac{3}{4|x|^2} - \frac{\operatorname{sgn}(x) \partial_1 \Phi}{2|x|} - \frac{(\partial_1 \Phi)^2}{4} + \frac{\partial_1^2 \Phi}{2} \\
 &\quad + x^2 e^{2\Phi} \left[\frac{1}{2} \partial_2^2 \Phi + \frac{3}{4} (\partial_2 \Phi)^2 \right] \\
 &:= -\frac{3}{4|x|^2} - \mu_2.
 \end{aligned}$$

Now, we are able to conclude by the following two Lemmas. The first one proves that the main part without μ_2 is essentially self-adjoint while the second one treats μ_2 as a perturbation.

LEMMA 4.3. — *Let G^{-1} given by $\operatorname{diag}(1, x^2 e^{2\Phi(x,y)})$ with $|\Phi| \leq C$ for $C > 0$.*

The operator $L : v \mapsto -\operatorname{div}_e(G^{-1} \nabla_e v) + \frac{3}{4|x|^2} v$ defined on $L^2(\mathbf{R} \times \mathbb{T}, dx dy)$ with domain $C_0^\infty(\Omega)$ is positive and essentially self-adjoint.

Hence, since L defines a positive symmetric operator, its unique self-adjoint extension is the Friedrichs extension (in the following still denoted by L).

LEMMA 4.4. — *The operator of multiplication by μ_2 (defined with domain $C_0^\infty(\Omega)$) is infinitesimally small with respect to L . Therefore, by the theorem of Kato-Rellich (Theorem X.12 of [32]), $L + \mu_2$ is essentially self-adjoint.*

To conclude the proof of Theorem 4.1 we are left to prove the two lemmas.

Proof. — [Proof of Lemma 4.3] We closely follow the proof of Kalf and Walter [24]. Actually, we mimick the proof of the 1-D case and we use the fact that our operator is the classical Laplacian with potential for function only depending on x : $-\partial_1^2 + \frac{3}{4|x|^2}$. It is enough to prove that $\text{Ran}(L + I)$ is dense in L^2 , see Theorem X.26 of [32]. So, let $h \in L^2(\Omega)$ such that $h \perp \text{Ran}(L + I)$ and let us prove $h = 0$ (we can assume h real valued without loss of generality).

Set $\beta(x) = \frac{|x|^{3/2} e^{-|x|}}{1+|x|^{3/2}}$. We have $\beta(x) \geq 0$ and $\beta \in L^2(\Omega)$. We obtain, for $x \neq 0$,

$$\begin{aligned} -\text{div}_e(G^{-1}\nabla_e\beta) + \frac{3}{4|x|^2}\beta + \beta &= -\partial_1^2\beta + \frac{3}{4|x|^2}\beta + \beta \\ &= \frac{3|x|^{1/2}e^{-|x|}}{4(1+|x|^{3/2})^3}(|x|^2 + 4|x|^{3/2} + 7|x|^{1/2} + 4) := X(x) \end{aligned}$$

Let $\eta, \zeta \in C^\infty([0, +\infty[)$ so that

$$\begin{aligned} \eta(s) &= 0 \text{ if } s \leq 1/2 \\ &= 1 \text{ if } s \geq 1 \\ \zeta(s) &= 1 \text{ if } s \leq 1 \\ &= 0 \text{ if } s \geq 2 \end{aligned}$$

Denote $\beta_n(x) = \eta(n|x|)\zeta(\frac{|x|}{n})\beta(x) \in C_0^\infty(\Omega)$ so that $L\beta_n$ is well defined in $L^2(\Omega)$. Moreover, we have

$$\begin{aligned} \beta(x) &= \mathcal{O}(|x|^{3/2}), & |\nabla\beta(x)| &= \mathcal{O}(|x|^{1/2}), \text{ as } x \rightarrow 0 \\ \beta(x) &= \mathcal{O}(e^{-|x|}), & |\nabla\beta(x)| &= \mathcal{O}(e^{-|x|}), \text{ as } x \rightarrow \infty \end{aligned}$$

so, $L\beta_n + \beta_n = \eta(n|x|)\zeta(\frac{|x|}{n})X - 2\partial_1 \left[\eta(n|x|)\zeta(\frac{|x|}{n}) \right] \partial_1\beta - \partial_1^2 \left[\eta(n|x|)\zeta(\frac{|x|}{n}) \right] \beta$ converges weakly in L^2 to $X \in L^2(\Omega)$ such that $X(x) \geq 0$. So, we can write

$$(X, |h|)_{L^2} = \lim_{n \rightarrow \infty} (L\beta_n + \beta_n, |h|)_{L^2} \geq 0$$

But, we will use the Kato inequality:

LEMMA 4.5 (Kato’s inequality, see Lemma A of [25]). —
 For $v \in L^1_{loc}(\Omega)$ real valued such that $\text{div}_e(G^{-1}\nabla_e v) \in L^1_{loc}(\Omega)$, we have, in the sense of distributions on Ω

$$-\text{div}_e(G^{-1}\nabla_e|v|) \leq -(\text{sgn}v)\text{div}_e(G^{-1}\nabla_e v).$$

Notice that the Kato inequality of [25] can only be applied on $\Omega_\varepsilon := \{(x, y) \in \Omega : |x| > \varepsilon\}$ where the metric is positive definite. This does

not create difficulties since we are applying this Lemma in the sense of distributions on Ω .

Since $h \perp \text{Ran}(L + I)$, we have $(L + I)h = 0$ in the distributional sense. So $\text{div}_e(G^{-1}\nabla_e h) \in L^1_{loc}(\Omega)$ and we can write $(\text{sgn}h)(-\text{div}_e(G^{-1}\nabla_e h) + \frac{3}{4|x|^2}h + h) = 0$ as an equality in $L^1_{loc}(\Omega)$ and therefore in the distributional sense. Applying the Kato Lemma to h and $\beta_n > 0$ as test function, we get

$$(L\beta_n + \beta_n, |h|)_{L^2} \leq (\beta_n, (\text{sgn}h)(Lh + h))_{L^2} \leq 0.$$

So $(X, |h|)_{L^2} = 0$ and $h \equiv 0$. □

Proof. — [Proof of Lemma 4.4] The quadratic form associated to $L + 1$ is

$$Q(u, u) = \int_{\Omega} (G^{-1}\nabla_e u \cdot \nabla_e u) + \frac{3}{4|x|^2} |u|^2 + |u|^2 \, dx \, dy$$

For $u \in C_0^\infty(\Omega)$, and by Cauchy-Schwarz inequality, we have

$$\begin{aligned} Q(u, u) = (Lu, u)_{L^2} + \|u\|_{L^2}^2 &\leq \|Lu\|_{L^2} \|u\|_{L^2} + \|u\|_{L^2}^2 \\ &\leq \varepsilon \|Lu\|_{L^2}^2 + \left(\frac{1}{4\varepsilon} + 1\right) \|u\|_{L^2}^2 \end{aligned}$$

Moreover, since Φ is smooth and constant for large x , we have

$$\begin{aligned} \|\mu_2 u\|_{L^2}^2 &\leq C \left(\left\| \frac{u}{|x|} \right\|_{L^2} + \|u\|_{L^2}^2 \right) \\ &\leq CQ(u, u) \leq C\varepsilon \|Lu\|_{L^2}^2 + (C\varepsilon + 1) \|u\|_{L^2}^2. \end{aligned}$$

Since ε is arbitrary small and the estimate can be extended to any $u \in D(L)$ by taking closure, this yields the result. □

The proof of Theorem 4.1 is concluded. □

The next Lemma is the first step to establish the compactness of the resolvent of Δ in the compact case. Of course, since here we are dealing with the case of $\mathbf{R} \times \mathbb{T}$, we can not expect compactness without adding a cut-off function.

LEMMA 4.6. — *Denote by Δ the positive self-adjoint operator defined by Theorem 4.1. Then, the truncated resolvent $\rho(x)(-\Delta + 1)^{-1}$, where $\rho \in C_0^\infty(\mathbf{R})$, is well-defined and compact from $L^2(\mathbf{R} \times \mathbb{T}, d\omega)$ to itself.*

Proof. — The fact that $(-\Delta + 1)^{-1}$ is well defined comes from the positivity of $-\Delta$. For the compactness it is equivalent to prove that the operator $\rho(x)(-\tilde{\Delta} + 1)^{-1}$ is compact when defined on $L^2(\mathbf{R} \times \mathbb{T}, dx \, dy)$ with the metric induced by the Lebesgue measure.

We begin by showing that it is the case for L defined above. Let u_n be a real valued sequence in $L^2(\mathbf{R} \times \mathbb{T}, dx dy)$ with L^2 norm bounded by 1. Denote $\widetilde{f}_n := \rho(x)(L + 1)^{-1}u_n := \rho(x)f_n$. In particular, $Q(f_n, f_n) = ((L + 1)f_n, f_n) = (u_n, f_n) = \int_{\Omega} G^{-1} \nabla_e f_n \cdot \nabla_e f_n + \frac{3}{4|x|^2} |f_n|^2 + |f_n|^2 dx dy$ is bounded in $L^2(\mathbf{R} \times \mathbb{T}, dx dy)$. Set $\varepsilon > 0$. Take $\varphi \in C_0^\infty(\mathbf{R})$, supported in $] - \varepsilon, \varepsilon[$ with $0 \leq \varphi \leq 1$ and $\varphi \equiv 1$ in a neighborhood of 0. Split $\widetilde{f}_n = \varphi(x)\widetilde{f}_n + (1 - \varphi(x))\widetilde{f}_n := f_{n,1} + f_{n,2}$. Then, we have $|x\varphi(x)|^2 \leq \varepsilon^2$ and $\|\varphi f_n\|_{L^2}^2 = \int \frac{|x\varphi(x)|^2 f_n^2}{|x|^2} \leq \varepsilon^2 \frac{4}{3} Q(f_n, f_n) \leq C\varepsilon^2$. This also gives $\|f_{n,1}\|_{L^2} \leq C\varepsilon$. Now that φ is fixed, $f_{n,2} = (1 - \varphi(x))\widetilde{f}_n = (1 - \varphi(x))\rho(x)f_n$ is supported in a fixed compact of Ω and bounded in H^1 . Indeed,

$$\begin{aligned} \|\nabla_e f_{n,2}\|_{L^2}^2 &\leq 2(\|(1 - \varphi(x))\rho(x)\nabla_e f_n\|_{L^2}^2 + \|\partial_x(\varphi(x)\rho(x))f_n\|_{L^2}^2) \\ &\leq CQ(f_n, f_n) \leq C \end{aligned}$$

where the second inequality comes from the fact that there exists a constant C such that $Id \leq CG^{-1}$ (in the sense of quadratic forms) on the support of $(1 - \varphi(x))\rho(x)$. We conclude by invoking the compact embedding of H^1 into L^2 on compact sets so that $f_{n,2}$ is convergent up to extraction. Actually, we have proved that for any $\varepsilon > 0$, we can find an extraction γ such that $f_{\gamma(n)}$ can be written $f_{\gamma(n)} = f_{\gamma(n),1} + f_{\gamma(n),2}$ with $\|f_{\gamma(n),1}\|_{L^2} \leq \varepsilon$ and $f_{\gamma(n),2}$ convergent. By choosing $\varepsilon = \frac{1}{p}$, $p \in \mathbb{N}$, and by a diagonal extraction argument, we easily get that we can find a subsequence such that $f_{\gamma(n)}$ is a Cauchy sequence and converges. This gives that $\rho(L + 1)^{-1}$ is compact. It remains to prove the same result for $L + \mu_2$.

Again, let u_n be a sequence in $L^2(\mathbf{R} \times \mathbb{T}, dx dy)$ with L^2 norm bounded by 1. Denote $\widetilde{f}_n := \rho(x)(L + \mu_2 + 1)^{-1}u_n := \rho(x)f_n$. Thanks to Lemma 4.4, we get

$$\begin{aligned} \|(L + 1)f_n\|_{L^2} &\leq \|(L + \mu_2 + 1)f_n\|_{L^2} + \|\mu_2 f_n\|_{L^2} \\ &\leq \|u_n\|_{L^2} + \varepsilon \|(L + 1)f_n\|_{L^2} + C_\varepsilon \|f_n\|_{L^2} \\ &\leq C_\varepsilon + \varepsilon \|(L + 1)f_n\|_{L^2} \end{aligned}$$

so that we get by absorption for ε small enough that $(L + 1)f_n$ is bounded. Since we have proved that $\rho(L + 1)^{-1}$ is a compact operator, we get that $\widetilde{f}_n = \rho(x)f_n = \rho(L + 1)^{-1}(L + 1)f_n$ is relatively compact. \square

4.2. Case of a compact manifold

In this section, using the previous results, we prove the three statements of Theorem 1.1. For the first statement, we have to prove the following.

PROPOSITION 4.7. — Consider a 2-ARS on a compact manifold M satisfying hypotheses **HA** and **HB**. Denote by Δ the Laplace operator defined for functions of $C_0^\infty(\Omega)$ where $\Omega = M \setminus \mathcal{Z}$. Then, Δ is essentially self-adjoint on $L^2(M, d\omega)$.

Proof. — Let $M = \bigcup_{i \in I} \Omega_i$ be a finite covering of M such that for every connected component W of \mathcal{Z} there exists $i \in I$ such that $W \subset \Omega_i$ and $W \cap \Omega_j = \emptyset$ for $j \in I, j \neq i$ and an orthonormal frame for the 2-ARS in Ω_i is given by (2.2). Moreover assume that if Ω_j ($j \in I$) does not contain any Grushin point, then an orthonormal frame for the 2-ARS in Ω_i is given by the normal form (F1). This is possible thanks to Theorem 2.2 and Proposition 2.3.

Theorem 4.1 yields the result in local coordinates around a connected component of the singular set. We only have to extend it to the whole manifold.

Let $(\Psi_i)_{i \in I}$ be a partition of unity associated to $(\Omega_i)_{i \in I}$, that is $\Psi_i \in C_0^\infty(\Omega_i)$ and $\sum_{i \in I} \Psi_i = 1$. We can also assume that $\Psi_i \equiv 0$ or $\Psi_i \equiv 1$ in a neighborhood of \mathcal{Z} , so that $\nabla \Psi_i$ and $\Delta \Psi_i$ are C^∞ compactly supported in Ω . Here we use the fact that we have global coordinates around every connected component of \mathcal{Z} .

Again, let $u \in L^2(\Omega)$ such that $u \perp \text{Ran}(-\Delta + I)$ and let us prove $u = 0$ (we can assume u real valued without loss of generality). We denote $u_i = \Psi_i u \in L^2(\Omega_i)$.

Since the domain of definition of Δ contains $C_0^\infty(\Omega)$, we get that u is solution of $-\Delta u + u = 0$ in the sense of distributions on Ω . By elliptic regularity, we get that $u \in C^\infty(\Omega)$. Therefore, the only problem is the possibility to compute integration by part around the degeneracy points.

Let Ω_i with $i \in I$ be an open set around which we can find a coordinate system so that the metric g takes the diagonal form $(1, x^{-2}e^{-2\Phi(x,y)})$ that we extend arbitrarily on $\mathbf{R} \times \mathbb{T}$ with Φ smooth constant for large x . We will denote Δ_i, ∇_i and $d\omega_i$ the Laplacian, gradient and area corresponding to this extension on $\mathbf{R} \times \mathbb{T}$. Since $u_i \in L^2(\Omega_i)$, we can consider the function u_i in this local coordinates (in what follows we will not distinguish u_i with its representant in local coordinates) and make the computation (in the sense of distributions of Ω and of $\mathbf{R}^* \times \mathbb{T}$ in local coordinates):

$$-\Delta_i u_i + u_i = -2\nabla_i u \cdot \nabla_i \Psi_i - (\Delta_i \Psi_i)u.$$

Remark that in these local coordinates u has only a true meaning for small x but we can then extend this equality on $\mathbf{R}^* \times \mathbb{T}$ since Ψ_i is compactly supported. Moreover, $\nabla_i \Psi_i$ and $\Delta_i \Psi_i$ is supported outside of the zone of

degeneracy Z where u is C^∞ . So, we get $-\Delta_i u_i + u_i \in C_0^\infty(\mathbf{R}^* \times \mathbb{T})$ and so $-\Delta_i u_i + u_i \in L^2(M, d\omega_i)$. This is true in the sense of distributions, but that means that for any $\varphi \in C_0^\infty(\mathbf{R}^* \times \mathbb{T})$, we have

$$|(u_i, -\Delta_i \varphi + \varphi)_{L^2(\mathbf{R}^* \times \mathbb{T}, d\omega_i)}| \leq C \|\varphi\|_{L^2(\mathbf{R}^* \times \mathbb{T}, d\omega_i)}.$$

In particular, that means that u_i belongs to the domain of the adjoint of the operator $-\Delta_i + I$ with domain $C_0^\infty(\mathbf{R}^* \times \mathbb{T})$. But Theorem 4.1 gives that this operator is essentially self-adjoint so, $u_i \in D(L_i)$ where L_i is the self-adjoint (Friedrichs) extension of $-\Delta_i + I$ on $\mathbf{R}^* \times \mathbb{T}$. In particular, we can write (the right hand side has to be understood as a limit for a sequence $u_{i,n}$ in $C_0^\infty(\Omega)$ converging strongly to u_i for the norm of the quadratic form)

$$(u_i, -\Delta_i u_i + u_i)_{L^2(\mathbf{R}^* \times \mathbb{T}, d\omega_i)} = \|\nabla_i u_i\|_{L^2(\mathbf{R}^* \times \mathbb{T}, d\omega_i)}^2 + \|u_i\|_{L^2(\mathbf{R}^* \times \mathbb{T}, d\omega_i)}^2.$$

This quantity also makes sense when u_i is considered on the manifold M and the extension of the metric was chosen so that $-\Delta_i u_i + u_i = -\Delta u_i + u_i$ in the sense of distributions and $\nabla_i u_i = \nabla u_i$.

The same result holds for $i \in I$ corresponding to a Riemannian zone.

Moreover, if $i \neq j \in I$, the common support of u_i and u_j do not intersect the degeneracy zone Z . Hence, these functions are C^∞ in this zone and we can write

$$(u_i, -\Delta u_j + u_j)_{L^2(M)} = (\nabla u_i, \nabla u_j)_{L^2(M)} + (u_i, u_j)_{L^2(M)}$$

By summing up, we get

$$0 = (u, -\Delta u + u)_{L^2(M, d\omega)} = (\nabla u, \nabla u)_{L^2(M, d\omega)} + (u, u)_{L^2(M, d\omega)}$$

and $u \equiv 0$. □

From now on, we will use the same notation Δ for the self-adjoint extension of the symmetric operator Δ .

In order to prove the second statement of Theorem 1.1 we have to prove the following.

PROPOSITION 4.8. — *The domain of Δ is given by*

$$D(\Delta) = \{u \in L^2(M, d\omega) \mid \Delta_{D,g} u \in L^2(M, d\omega)\} \tag{4.2}$$

where $\Delta_{D,g} u$ is Δu seen as a distribution in $\Omega = M \setminus Z$.

Proof. — Let T be the Laplace operator defined with domain given by (4.2). We easily see that

$$\begin{aligned} D(T) &= \{u \in L^2(M, d\omega) \mid \exists C : \forall \varphi \in C_0^\infty(\Omega), |(u, \Delta \varphi)_{L^2}| \leq C \|\varphi\|_{L^2}\} \\ &= D(\Delta^*) = D(\Delta). \end{aligned} \tag{□}$$

In order to prove the third statement of Theorem 1.1 we have to prove the following.

PROPOSITION 4.9. — *Denote by Δ the positive self-adjoint operator defined by Proposition 4.7, then the resolvent $(-\Delta + 1)^{-1}$ is well defined and compact from $L^2(M, d\omega)$ to itself with the measure defined by the metric.*

Therefore, the spectrum of Δ is discrete and consists of eigenvalues of finite multiplicity.

Proof. — The fact that $(-\Delta + 1)^{-1}$ is well defined comes from the positivity of $-\Delta$. Now, let u_n be a bounded sequence in $L^2(M, d\omega)$ and $f_n = (-\Delta + 1)^{-1}u_n$. By density, we can assume u_n in $C_0^\infty(M \setminus \mathcal{Z})$ and $f_n \in C^\infty(M \setminus \mathcal{Z})$ by elliptic regularity. Then, we have $\int_M |\nabla f_n|^2 + |f_n|^2 = (u_n, f_n)_{L^2}$ bounded.

Consider the partition of unity $(\Psi_i)_{i \in I}$ introduced in the proof of the previous Theorem and denote $f_{i,n} := \Psi_i f_n$.

Let $i \in I$ be an index corresponding to a “Grushin zone”. By the formula

$$(-\Delta + I)f_{n,i} = \Psi_i u_n - 2\nabla \Psi_i \cdot \nabla f_n - (\Delta \Psi_i)f_n,$$

we have that $(-\Delta + I)f_{n,i}$ is bounded in $L^2(M)$. By using Lemma 4.6, we get that $\rho(x)(-\Delta_i + 1)^{-1}(-\Delta + I)f_{n,i}$ is compact for a function $\rho \in C_0^\infty(\mathbf{R})$ defined in some local coordinate charts. Here, we have denoted Δ_i the Laplacian for an extension of the local metric to $\mathbf{R} \times \mathbb{T}$. To finish, we only have to notice that $(-\Delta_i + 1)^{-1}(-\Delta + I)f_{n,i} = (-\Delta_i + 1)^{-1}(-\Delta_i + 1)f_{n,i} = f_{n,i}$ because of the support of $f_{n,i}$ (note that it is not the case for $(-\Delta_i + 1)^{-1}$ because the resolvent depends on the extension).

So, choosing ρ such that $\rho(x)f_{n,i} = f_{n,i}$ in local coordinates, we have proved that each sequence $f_{n,i}$ is compact for a Grushin zone. The same result holds for a Riemannian zone because the equivalent of Lemma 4.6 still holds for any Riemannian extension. In sum, we get that the sequence f_n is convergent up to extraction, which yields the result. □

Appendix A. The Martinet case

In this Section, we prove Theorem 1.7. A sub-Riemannian structure is a triple $(M, \blacktriangle, \mathbf{g})$, where M is a smooth manifold, \blacktriangle is a smooth vector distribution of constant rank satisfying the Hörmander condition, and \mathbf{g} is a Riemannian metric on \blacktriangle . Let $\blacktriangle_1 = \blacktriangle$ and $\blacktriangle_{k+1} = \blacktriangle_k + [\blacktriangle, \blacktriangle_k]$. A

sub-Riemannian structure is said to be equiregular if the dimension of $\mathbf{\Delta}_k$ does not depend on the point.

Almost-Riemannian and sub-Riemannian structures can be treated in the unified setting of rank-varying sub-Riemannian structures, see [1, Chapter 3] and [2, Definition 2]. For sake of readability, we omit this point of view.

In this section, we briefly treat the Martinet sub-Riemannian manifold (see for instance [12, 28]), defined by $M = \mathbf{R}^3$, $\mathbf{\Delta}(q) = \text{span}\{X_1(q), X_2(q)\}$, and $\mathbf{g}(X_i, X_j) = \delta_{ij}$, $i, j = 1, 2$, where

$$X_1 = \begin{pmatrix} 1 \\ 0 \\ \frac{y^2}{2} \end{pmatrix}, \quad X_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}. \tag{A.1}$$

Notice that $X_3 = [X_1, X_2] = \begin{pmatrix} 0 \\ 0 \\ -y \end{pmatrix}$. Hence, X_1, X_2 and X_3 span

the tangent space at any point outside the so called Martinet plane $\mathcal{Z} = \{y = 0\}$, which is the region where the structure is not equiregular. However

$X_4 = [[X_1, X_2], X_2] = [X_3, X_2] = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$, hence, the Hörmander condition

is fulfilled on the whole space.

With this structure, $\mathbf{R}^3 \setminus \mathcal{Z}$ is a 3D sub-Riemannian contact manifold. On such a structure it is possible to define intrinsically a volume form which is given by $dX_1 \wedge dX_2 \wedge dX_3$, where $\{dX_1, dX_2, dX_3\}$ is the dual basis to $\{X_1, X_2, X_3\}$. Such a volume form is independent on the choice of the orthonormal basis which define the sub-Riemannian structure, and it is called the Popp measure (see [28] and Proposition 8 of [3]). One gets $dX_1 = dx$, $dX_2 = dy$ and $dX_3 = \frac{y}{2}dx - \frac{1}{y}dz$ so that $dX_1 \wedge dX_2 \wedge dX_3 = -\frac{1}{y}dx \wedge dy \wedge dz$ and the corresponding density is $d\omega = \frac{1}{|y|}dx dy dz$. This allows to define the sub-Riemannian Laplacian as the divergence of the sub-Riemannian gradient defined as in formula (1.1) (see Remark 14 of [3]),

$$\Delta_{sr} = (X_1)^2 + (X_2)^2 - \frac{1}{y}X_2 = (\partial_x + \frac{y^2}{2}\partial_z)^2 + \partial_y^2 - \frac{1}{y}\partial_y. \tag{A.2}$$

Notice that the singularity in the first order term of Δ_{sr} appears similarly as it does in the Grushin case. We do the same reasoning.

For simplicity, we compactify in x and z and consider the same structure on $M = \mathbb{T}_x \times \mathbf{R}_y \times \mathbb{T}_z$. We will prove that Δ_{sr} is essentially self-adjoint on $L^2(M, d\omega)$ with domain $C_0^\infty(M \setminus \mathcal{Z})$. Again, we make the change of variable

in Hilbert space $f = \sqrt{|y|}g$ so that

$$\Delta_{sr}f = \sqrt{|y|} \left((\partial_x + \frac{y^2}{2}\partial_z)^2 g + \partial_y^2 g - \frac{3}{4|y|^2} g \right) = \sqrt{|y|} \widetilde{\Delta}_{sr} g. \quad (\text{A.3})$$

So, we are left to prove that $\widetilde{\Delta}_{sr} = (\partial_x + \frac{y^2}{2}\partial_z)^2 + \partial_y^2 - \frac{3}{4y^2}$ is essentially self-adjoint on $L^2(\mathbb{T} \times \mathbf{R} \times \mathbb{T}, dx dy dz)$ with domain $C_0^\infty(M \setminus \mathcal{Z})$. We compute the Fourier transform in x and z and get the decomposition

$$L^2(\mathbb{T} \times \mathbf{R} \times \mathbb{T}) \approx \bigoplus_{(k,l) \in \mathbf{Z}^2} H_{(k,l)}$$

where $H_{(k,l)} = L^2(\mathbf{R})$ and the operator L acts on each $H_{(k,l)}$ by

$$(Lu)_{(k,l)}(y) = \partial_y^2 u_{(k,l)} - (k + \frac{y^2}{2}l)^2 u_{(k,l)} - \frac{3}{4y^2} u_{(k,l)} = \partial_y^2 u_{(k,l)} - V_{(k,l)} u_{(k,l)}$$

with $V_{(k,l)}(y) \geq \frac{3}{4y^2}$. We conclude as in the Grushin case.

This result suggests the general conjecture that for a sub-Riemannian structure which is rank-varying or not equiregular on an hypersurface, the singular set acts as a barrier for the heat flow and for a quantum particle.

BIBLIOGRAPHY

- [1] A. AGRACHEV, D. BARILARI & U. BOSCAIN, *Introduction to Riemannian and sub-Riemannian geometry (Lecture Notes)*, http://people.sissa.it/agrachev/agrachev_files/notes.html.
- [2] A. A. AGRACHEV, U. BOSCAIN, G. CHARLOT, R. GHEZZI & M. SIGALOTTI, “Two-dimensional almost-Riemannian structures with tangency points”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **27** (2010), no. 3, p. 793-807.
- [3] A. AGRACHEV, U. BOSCAIN, J.-P. GAUTHIER & F. ROSSI, “The intrinsic hypoelliptic Laplacian and its heat kernel on unimodular Lie groups”, *J. Funct. Anal.* **256** (2009), no. 8, p. 2621-2655.
- [4] A. AGRACHEV, U. BOSCAIN & M. SIGALOTTI, “A Gauss-Bonnet-like formula on two-dimensional almost-Riemannian manifolds”, *Discrete Contin. Dyn. Syst.* **20** (2008), no. 4, p. 801-822.
- [5] A. A. AGRACHEV & Y. L. SACHKOV, *Control theory from the geometric viewpoint*, Encyclopaedia of Mathematical Sciences, vol. 87, Springer-Verlag, Berlin, 2004, Control Theory and Optimization, II, xiv+412 pages.
- [6] D. BARILARI, “Trace heat kernel asymptotics in 3D contact sub-Riemannian geometry”, *Journal of Mathematical Science*, To appear.
- [7] D. BARILARI, U. BOSCAIN & R. NEEL, “Small time heat kernel asymptotics at the sub-Riemannian cut-locus”, *Journal of Differential Geometry* **92** (2012), no. 3, p. 373-416.
- [8] A. BELLAÏCHE, “The tangent space in sub-Riemannian geometry”, in *Sub-Riemannian geometry*, Progr. Math., vol. 144, Birkhäuser, Basel, 1996, p. 1-78.
- [9] B. BONNARD, J.-B. CAILLAU, R. SINCLAIR & M. TANAKA, “Conjugate and cut loci of a two-sphere of revolution with application to optimal control”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **26** (2009), no. 4, p. 1081-1098.
- [10] B. BONNARD & J. B. CAILLAU, “Metrics with equatorial singularities on the sphere”, Preprint 2011, HAL, vol. 00319299, pp. 1-29.

- [11] B. BONNARD, G. CHARLOT, R. GHEZZI & G. JANIN, “The Sphere and the Cut Locus at a Tangency Point in Two-Dimensional Almost-Riemannian Geometry”, *J. Dynam. Control Systems* **17** (2011), no. 1, p. 141-161.
- [12] B. BONNARD & M. CHYBA, *Singular trajectories and their role in control theory*, Mathématiques & Applications (Berlin) [Mathematics & Applications], vol. 40, Springer-Verlag, Berlin, 2003, xvi+357 pages.
- [13] U. BOSCAIN, T. CHAMBRION & G. CHARLOT, “Nonisotropic 3-level quantum systems: complete solutions for minimum time and minimum energy”, *Discrete Contin. Dyn. Syst. Ser. B* **5** (2005), no. 4, p. 957-990.
- [14] U. BOSCAIN & G. CHARLOT, “Resonance of minimizers for n -level quantum systems with an arbitrary cost”, *ESAIM Control Optim. Calc. Var.* **10** (2004), no. 4, p. 593-614 (electronic).
- [15] U. BOSCAIN, G. CHARLOT, J.-P. GAUTHIER, S. GUÉRIN & H.-R. JAUSLIN, “Optimal control in laser-induced population transfer for two- and three-level quantum systems”, *J. Math. Phys.* **43** (2002), no. 5, p. 2107-2132.
- [16] U. BOSCAIN, G. CHARLOT & R. GHEZZI, “Normal forms and invariants for 2-dimensional almost-Riemannian structures”, *Differential Geometry and its Applications* **31** (2013), no. 1, p. 41-62.
- [17] U. BOSCAIN, G. CHARLOT, R. GHEZZI & M. SIGALOTTI, “Lipschitz Classification of Two-Dimensional Almost-Riemannian Distances on Compact Oriented Surfaces”, *J. Geom. Anal.* **23** (2013), no. 1, p. 438-455.
- [18] U. BOSCAIN & M. SIGALOTTI, “High-order angles in almost-Riemannian geometry”, in *Actes de Séminaire de Théorie Spectrale et Géométrie. Vol. 24. Année 2005–2006*, Sémin. Théor. Spectr. Géom., vol. 25, Univ. Grenoble I, 2008, p. 41-54.
- [19] H. DONNELLY & N. GAROFALO, “Schrödinger operators on manifolds, essential self-adjointness, and absence of eigenvalues”, *Journal of Geometric Analysis* **7** (1997), no. 2, p. 241-257.
- [20] B. FRANCHI & E. LANCONELLI, “Une métrique associée à une classe d’opérateurs elliptiques dégénérés”, *Rend. Sem. Mat. Univ. Politec. Torino* (1983), no. Special Issue, p. 105-114 (1984), Conference on linear partial and pseudodifferential operators (Torino, 1982).
- [21] V. V. GRUŠIN, “A certain class of hypoelliptic operators”, *Mat. Sb. (N.S.)* **83** (125) (1970), p. 456-473.
- [22] P. GÉRARD & S. GRELLIER, “The Szegő cubic equation”, *Ann. Scient. Ec. Norm. Sup.* **43** (2010), no. 4, p. 761-809.
- [23] F. JEAN, “Uniform estimation of sub-Riemannian balls”, *J. Dynam. Control Systems* **7** (2001), no. 4, p. 473-500.
- [24] H. KALF & J. WALTER, “Note on a paper of Simon on essentially self-adjoint Schrödinger operators with singular potentials”, *Archive for Rational Mechanics and Analysis* **52** (1973), no. 3, p. 258-260.
- [25] T. KATO, “Schrödinger operators with singular potentials”, *Israel Journal of Mathematics* **13** (1972), no. 1, p. 135-148.
- [26] R. LÉANDRE, “Minoration en temps petit de la densité d’une diffusion dégénérée”, *J. Funct. Anal.* **74** (1987), no. 2, p. 399-414.
- [27] M. MAEDA, “Essential selfadjointness of Schrödinger operators with potentials singular along affine subspaces”, *Hiroshima Mathematical Journal* **11** (1981), no. 2, p. 275-283.
- [28] R. MONTGOMERY, *A tour of subriemannian geometries, their geodesics and applications*, Mathematical Surveys and Monographs, vol. 91, American Mathematical Society, Providence, RI, 2002, xx+259 pages.
- [29] R. NEEL, “The small-time asymptotics of the heat kernel at the cut locus”, *Comm. Anal. Geom.* **15** (2007), no. 4, p. 845-890.

- [30] R. NEEL & D. STROOCK, “Analysis of the cut locus via the heat kernel”, in *Surveys in differential geometry. Vol. IX*, Surv. Differ. Geom., IX, Int. Press, Somerville, MA, 2004, p. 337-349.
- [31] L. S. PONTRYAGIN, V. G. BOLTYANSKIĬ, R. V. GAMKRELIDZE & E. F. MISHCHENKO, *The Mathematical Theory of Optimal Processes*, fourth ed., “Nauka”, Moscow, 1983, 392 pages.
- [32] M. REED & B. SIMON, *Methods of modern mathematical physics*, Academic press, 1980.
- [33] B. SIMON, “Essential self-adjointness of Schrödinger operators with singular potentials”, *Archive for Rational Mechanics and Analysis* **52** (1973), no. 1, p. 44-48.
- [34] S. R. S. VARADHAN, “On the behavior of the fundamental solution of the heat equation with variable coefficients”, *Comm. Pure Appl. Math.* **20** (1967), p. 431-455.
- [35] M. VENDITTELLI, G. ORIOLO, F. JEAN & J.-P. LAUMOND, “Nonhomogeneous nilpotent approximations for nonholonomic systems with singularities”, *IEEE Trans. Automat. Control* **49** (2004), no. 2, p. 261-266.

Ugo BOSCAIN
CNRS, Centre de Mathématiques Appliquées,
École Polytechnique,
Route de Saclay, 91128 Palaiseau Cedex, France,
and
INRIA, Centre de Recherche Saclay, Team GECCO.
ugo.boscain@polytechnique.edu

Camille LAURENT
CNRS, Centre de Mathématiques Appliquées,
École Polytechnique,
Route de Saclay, 91128 Palaiseau Cedex, France
and
CNRS, UMR 7598, Laboratoire Jacques-Louis
Lions, F-75005 Paris, France
laurent@ann.jussieu.fr