



ANNALES

DE

L'INSTITUT FOURIER

Renaud LEPLAIDEUR & Dawei YANG

SRB measures for higher dimensional singular partially hyperbolic attractors

Tome 67, n° 6 (2017), p. 2703-2717.

http://aif.cedram.org/item?id=AIF_2017__67_6_2703_0



© Association des Annales de l'institut Fourier, 2017,

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Annales de l'institut Fourier »
(<http://aif.cedram.org/>), implique l'accord avec les conditions générales
d'utilisation (<http://aif.cedram.org/legal/>).

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

SRB MEASURES FOR HIGHER DIMENSIONAL SINGULAR PARTIALLY HYPERBOLIC ATTRACTORS

by Renaud LEPLAIDEUR & Dawei YANG (*)

ABSTRACT. — We prove the existence and the uniqueness of the SRB measure for any singular hyperbolic attractor in dimension $d \geq 3$. The proof does not use Poincaré sectional maps, but uses basic properties of thermodynamical formalism.

RÉSUMÉ. — Nous prouvons que tout attracteur partiellement hyperbolique de dimension finie et avec singularité(s) admet une unique mesure SRB. La preuve utilise des outils simples et généraux du formalisme thermodynamique et ne nécessite pas de recourir à une section de Poincaré.

1. Introduction

1.1. Background

Our aim is to prove that any finite dimensional singular partially hyperbolic attractor admits a unique SRB measure.

SRB stands for Sinai, Ruelle and Bowen who established in the 70's the theory of thermodynamical formalism and proved the existence and the uniqueness of some special measures for Anosov systems and Axiom A attractors (diffeomorphisms or flows), see e.g. [4, 19, 20].

SRB measures usually satisfy the two following properties. On the one hand their disintegrations along unstable manifolds are absolutely continuous with respect to the Lebesgue measures on these manifolds. On the other hand, and at least for the hyperbolic case, SRB measures are expected to be

Keywords: partially hyperbolic singular flows, thermodynamical formalism, equilibrium states, SRB and physical measures.

2010 *Mathematics Subject Classification:* 37A35, 37A60, 37D20, 37D35, 47N10.

(*) RL would like to thank Soochow University for its support and hospitality. DY would like to thank the support of NSFC 11271152 and A Project Funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions(PAPD).

physical measures, i.e., their sets of generic points have positive Lebesgue measure on the manifold (see e.g. [25] for a survey).

Since Sinai, Ruelle and Bowen, it has remained a challenge to export the SRB theory to other dynamical systems presenting a weaker form of hyperbolicity. Among them, the Lorenz-like attractors represent one of the most famous families. First introduced by Lorenz in [13], this class has several typical properties of chaotic dynamics: it is robust in the C^1 -topology, every ergodic invariant measure is hyperbolic (see Lemma A.1) but the attractors themselves are not hyperbolic.

For the 3-dimensional case, Guckenheimer was inspired by the example introduced by Lorenz to define in [14] the notion of geometric Lorenz attractor. He asked about the existence of SRB measures for these attractors. In [16], Morales, Pacifico and Pujals used the notion of *singular hyperbolicity* to characterize more general Lorenz-like dynamics. In [1], Araujo–Pacifico–Pujals–Viana proved the existence and the uniqueness of SRB measures for *three-dimensional singular hyperbolic attractors*. We also mention other related works: [2, 6].

Higher-dimensional singular hyperbolic attractors have been defined in [26] and [15]. It was thus a natural question to investigate the existence and the uniqueness of the SRB measure for these attractors. In this paper we actually prove the existence and the uniqueness of such a measure.

We emphasize that the method from [1] cannot be adapted to the higher dimensional case. Indeed, their main idea is to construct a sequence of cross-sections and consider the return maps among these sections. Then, they quotient the dynamics to piecewise expanding maps of intervals and use the existence of absolutely continuous invariant measures for these maps. That idea cannot be directly adapted to the higher-dimensional case for the following two reasons. On the one hand, it is much more difficult to construct cross sections, and on the other hand, the existence of the absolutely continuous invariant measure for general higher-dimensional piecewise expanding maps may fail (see [5, 21]).

Our strategy is different. We show here that basic properties of thermodynamical formalism allow one to show the existence and the uniqueness of SRB measures. Actually, a close strategy has already been investigated in [7] where Cowieson and Young obtained SRB measures by using the Pesin entropy formula for diffeomorphisms beyond uniform hyperbolicity.

After we had posted the paper on arXiv, we had some communication with M. Viana, and learnt that he and J. Yang had a work in progress on that topic. We were not aware on that work, which did not exist as a

preprint but as a recorded talk (see [23]). We agreed with them to mention this video. In our mind, the fact that we independently got similar ideas emphasizes how natural this strategy is.

Finally, we want to emphasize that our strategy deeply uses the absolute continuity of the strong stable foliation, which has been a very important research tool for years (see [17]).

1.2. Settings and the statement of the result

Let X be a vector field on a d -dimensional manifold M , and φ_t be the flow generated by X . We recall that a compact invariant set Λ is called a topological attractor if

- there is an open neighborhood U of Λ such that $\bigcap_{t \geq 0} \varphi_t(\overline{U}) = \Lambda$,
- Λ is transitive, i.e., there is a point $x \in \Lambda$ with a dense forward-orbit.

A compact invariant set Λ is called a *singular hyperbolic attractor* (see [15, 26]) if it is a topological attractor, with at least one *singularity* σ , which means that σ satisfies $X(\sigma) = 0$. Moreover, there is a continuous invariant splitting $T_\Lambda M = E^{ss} \oplus F^{cu}$ of $D\varphi_t$ together with constants $C > 0$ and $\lambda > 0$ such that

- Domination: for any $x \in \Lambda$ and any $t > 0$,

$$\|D\varphi_t|_{E^{ss}(x)}\| \|D\varphi_{-t}|_{F^{cu}(\varphi_t(x))}\| \leq Ce^{-\lambda t}.$$

- Contraction: for any $x \in \Lambda$ and any $t > 0$, $\|D\varphi_t|_{E^{ss}(x)}\| \leq Ce^{-\lambda t}$.
- Sectional expansion: for any x , $F^{cu}(x)$ contains two non-collinear vectors, and any $t > 0$, for every pair of non-collinear vectors \mathbf{v} and \mathbf{w} in $F^{cu}(x)$, $|\det D\varphi_t|_{\text{span}\langle \mathbf{v}, \mathbf{w} \rangle}| \geq Ce^{\lambda t}$.

We emphasize that one of the difficulties to study these attractors is that the singularity may belong to the attractor and may be accumulated by recurrent regular orbits. Since the uniformly hyperbolic case has already been well-understood since [4], we assume that the attractor does contain a singularity.

We recall that entropy for a flow is defined as being the entropy of the time-1 map $f := \varphi_1$.

DEFINITION 1.1. — *An invariant measure for the flow is said to be an SRB measure if it is an SRB measure for the time-one map f of the flow, that is:*

- (1) *it has a positive Lyapunov exponent almost everywhere,*

- (2) *the conditional measures on unstable manifolds are absolutely continuous w.r.t. Lebesgue measures on unstable manifolds.*

We refer the reader to the survey [25] on SRB measures for more details. Ledrappier–Young [12] proved that μ is an SRB measure if and only if μ has a positive Lyapunov exponent almost everywhere and the entropy of μ is the integration of the sum of all positive Lyapunov exponents.

The equality is called *the Pesin entropy formula* and is the heart of our strategy.

The goal of this paper is to prove the following theorem.

THEOREM 1.2. — *Let Λ be a singular hyperbolic attractor of a \mathcal{C}^2 vector field X of any manifold M of any dimension. Then there is a unique SRB measure μ supported on Λ . Moreover μ is ergodic and physical.*

1.3. Thermodynamical formalism

The main idea of the proof of the theorem is to use the thermodynamical formalism. We refer the reader to [22] for general results on this topic.

We recall that if $\psi : \Lambda \rightarrow \mathbb{R}$ is a continuous function, an invariant probability measure μ is said to be *an equilibrium state for ψ* if it satisfies

$$h_\mu(f) + \int \psi \, d\mu = \sup_{\nu \text{ ergodic, } \text{supp } \nu \subset \Lambda} \left\{ h_\nu(f) + \int \psi \, d\nu \right\} =: \mathcal{P}(\psi).$$

$\mathcal{P}(\psi)$ is called the *pressure* for the potential ψ . We recall that it also satisfies

$$\begin{aligned} \mathcal{P}(\psi) &= \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow +\infty} \frac{1}{n} \sup \left\{ \log \sum_{x \in \mathcal{E}_{n,\varepsilon}} e^{S_n(\psi)(x)}, \mathcal{E}_{n,\varepsilon} = \text{maximal} \right. \\ &\quad \left. (n, \varepsilon)\text{-separated set in } \Lambda \right\} \\ &= \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow +\infty} \frac{1}{n} \inf \left\{ \log \sum_{x \in \mathcal{E}_{n,\varepsilon}} e^{S_n(\psi)(x)}, \mathcal{E}_{n,\varepsilon} = \text{minimal} \right. \\ &\quad \left. (n, \varepsilon)\text{-spanning set in } \Lambda \right\}, \end{aligned}$$

where $S_n(\psi) = \psi + \dots + \psi \circ f^{n-1}$. Recall that $x, y \in M$ are (n, ε) -close if $d(f^i(x), f^i(y)) < \varepsilon$ for any $0 \leq i \leq n-1$; x and y are (n, ε) -separated if $\exists 0 \leq k \leq n-1, d(f^k(x), f^k(y)) \geq \varepsilon$. A (n, ε) -ball at x is the set of all points that are (n, ε) -close to x . A finite subset Γ of Λ is called (n, ε) -spanning if any point of Λ is contained in a (n, ε) -ball of some point in Γ .

J. Yang (see [24, Theorem A]) proved that the time-one map f is entropy expansive on the singular hyperbolic attractor Λ . This yields the upper semi-continuity for the metric entropy in the weak star topology and it is well known that it implies:

LEMMA 1.3. — *Every continuous function ψ on Λ has an equilibrium state μ supported on Λ .*

Denote by J^{cu} the Jacobian for f restricted to the F^{cu} direction, i.e., $J^{cu} = \det Df|_{F^{cu}}$. We also set $V := -\log |J^{cu}|$. It is a continuous function. Thus, by Lemma 1.3, there exists at least one equilibrium state for V . Our goal is to show that this equilibrium state is unique and is the SRB measure.

1.4. Plan of the proof

The paper proceeds as follows. In Section 2 we prove $\mathcal{P}(V) = 0$. This shows that any equilibrium state for $V = -\log |J^{cu}|$ satisfies the Pesin entropy formula, and therefore, such an equilibrium state is an SRB measure if and only if it admits a positive Lyapunov exponent (by [12]). Then, we prove that an equilibrium state for V has one positive Lyapunov exponent.

In Section 3 we prove that any SRB measure is physical and then deduce the uniqueness from transitivity.

In Appendix A we recall several classical properties and state precisely some definitions such as those of strong stable manifolds, the Oseledets splitting of invariant measures, the absolute continuity, etc.

Acknowledgements

We thank R. Scott for correcting english, J. Buzzzi, Y. Cao, S. Crovisier, Y. Hua, F. Ledrappier and the anonymous referee for useful discussions, advices and improvements for the redaction of this paper.

2. The Existence of ergodic SRB measures

In this section we prove that $\mathcal{P}(V) = 0$. As we said above, a standard argument will imply that any equilibrium state for V , is an SRB measure.

2.1. Upper bound for $\mathcal{P}(V)$

LEMMA 2.1. — *$\mathcal{P}(V) \leq 0$. Moreover, if $\mathcal{P}(V) = 0$ then, there exists an ergodic measure which satisfies the Pesin entropy formula and is an equilibrium state for V .*

Proof. — We remind the reader that $V(x) = -\log |J^{cu}(x)|$. We prove the lemma by the following steps. Let μ be an ergodic invariant probability measure.

- If μ is concentrated on a singularity, by the definition of sectional expansion, we have $\int V d\mu < 0$ and $h_\mu(f) = 0$. Then

$$h_\mu(\varphi_1) + \int V d\mu < 0.$$

- If μ is not concentrated on a singularity, by Lemma A.1, we have the following measurable Oseledec splitting

$$E^{ss} \oplus \underbrace{\langle X \rangle \oplus E_\mu^u}_{=F^{cu}}.$$

Note that $-\int V d\mu$ is the integral of the sum of all Lyapunov exponents along F^{cu} . Among them, the Lyapunov exponent along the flow direction is zero and the others are positive. Therefore, the sum of positive Lyapunov exponents equals the sum of Lyapunov exponents along F^{cu} ; i.e.,

$$\int V d\mu = -\sum \lambda_\mu^+,$$

where $\sum \lambda_\mu^+$ is the sum of all positive Lyapunov exponents of μ . By the Ruelle inequality, we have

$$h_\mu(f) \leq \sum \lambda_\mu^+ = -\int V d\mu.$$

- Now we thus have

$$\mathcal{P}(V) \leq \sup_{\mu \text{ ergodic}} \{h_\mu(\varphi_1) + \int V d\mu\} \leq 0.$$

The second statement in the Lemma now easily follows: if $\mathcal{P}(V) = 0$, let μ be an equilibrium state for V (it exists due to Lemma 1.3). Therefore $\mathcal{P}(V) = h_\mu(f) + \int V d\mu = 0$ holds. Hence, $h_\nu(f) + \int V d\nu = 0 = \mathcal{P}(V)$ holds for at least one ergodic component ν of μ (see [10, Theorem 4.1.12 and Corollary 4.3.17]). Thus, ν is ergodic, satisfies the Pesin entropy formula and is an equilibrium state for V . \square

2.2. Lower bound for $\mathcal{P}(V)$

We follow and adapt the volume lemma from Bowen (see [3, Section 4.B]) and its version from Qiu (see [18]).

PROPOSITION 2.2. — *Let Λ be the singular hyperbolic attractor with splitting $T_\Lambda M = E^{ss} \oplus F^{cu}$. Then $\mathcal{P}(V) \geq 0$.*

Proof. — We use the characterization that we recalled just after the definition for pressure:

$$\mathcal{P}(V) = \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow +\infty} \frac{1}{n} \inf \left\{ \log \sum_{x \in \mathcal{E}_{n,\varepsilon}} e^{S_n(V)(x)}, \begin{array}{l} \mathcal{E}_{n,\varepsilon} = \text{minimal} \\ (n, \varepsilon)\text{-spanning set in } \Lambda \end{array} \right\}.$$

Let $\varepsilon > 0$ be fixed and consider a minimal (n, ε) -spanning set $\mathcal{E}_{n,\varepsilon} = \{\xi_i\}$. Therefore $\bigcup B_n(\xi_i, \varepsilon)$ is an open neighborhood of Λ .

LEMMA 2.3. — *There is $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$, there is $m_\varepsilon > 0$ independent of n such that any cover of Λ by (n, ε) -balls has Lebesgue measure larger than m_ε .*

Proof. — First, we claim that Λ contains regular orbits. Indeed, if this does not hold, then Λ only consists on singularities and transitivity yields that Λ is a single singularity, say σ . The bundle F^{cu} is non-trivial and sectional expanding. Therefore the unstable space $E^u(\sigma)$ is non-trivial. Consequently, $\Lambda = \{\sigma\}$ cannot be an attractor.

By [24, Theorem A], the topological entropy on Λ is thus positive. By the variational principle, there is an f -invariant ergodic measure μ with positive metric entropy. Since μ is non-trivial and supported on a singular hyperbolic attractor, μ is hyperbolic as stated in Lemma A.1. We can use Pesin theory for that measure and construct local unstable manifolds $W_{loc}^u(x)$ for μ -almost every point x .

One can choose a point $x \in \Lambda$ such that $W_{loc}^u(x)$ contains a disk for the topology of the embedded manifold $W_{loc}^u(x)$, and whose diameter for the metric of M is $\varepsilon_1 > 0$. We denote this disk by $D^u(\varepsilon_1)$. Since Λ is a topological attractor, the unstable manifold of any point of Λ is contained in Λ . Thus, $D^{cu}(\varepsilon_1) := \bigcup_{t \in [-\varepsilon_1, \varepsilon_1]} \varphi_t(D^u(\varepsilon_1))$ is contained in Λ . This set has a Lebesgue measure of order $\varepsilon_1^{\dim F^{cu}}$. See more in Subsection A.2.3 about the metrics between the manifold M and sub-manifolds.

Take ε is smaller than ε_1 and ε_Λ (which is the size of local strong stable manifolds from Subsection A.2). All the points in $D^{cu}(\varepsilon_1)$ admit a stable local manifold W_ε^{ss} . Then, $\bigcup B_n(\xi_i, \varepsilon)$ contains the set

$$\bigcup_{y \in D^{cu}(\varepsilon_1)} W_{\varepsilon/4}^{ss}(y),$$

which has a Lebesgue measure of order $\varepsilon^{\dim E^{ss}}$ (using the absolute continuity of the stable foliation, see Subsection A.2.3). □

We assume that ε is much smaller than this ε_0 , where ε is used to define minimal (n, ε) -spanning sets. Since Λ is an attractor, the bundle E^{ss} and the local strong stable foliation W_{loc}^{ss} can be extended in a neighborhood of Λ (see [9, Theorem 5.5] and Subsection A.2).

Define $B^{cu}(x, \rho)$ as the balls of center $x \in \Lambda$ and radius $\rho < \varepsilon_0$ respectively in the local cu -plaques $W_{loc}^{cu}(x)$. Denote by

$$B_n^{cu}(x, \rho) = \{y : f^i(y) \in B^{cu}(f^i(x), \rho), \forall 0 \leq i \leq n - 1\}.$$

The metrics between M and local strong stable manifolds have uniform ratios (see the property (D1–D3) in Subsection A.2.3). More precisely, there is $\kappa > 1$ such that for all ε small enough,

$$(2.1) \quad W_{\rho/\kappa}^{ss}(B^{cu}(x, \rho/\kappa)) \subset B(x, \rho) \subset W_{\kappa\rho}^{ss}(B^{cu}(x, \kappa\rho)).$$

The continuity of Df and the continuity of F^{cu} imply that there exists $C_\varepsilon > 0$ such that for every x and y in Λ ,

$$(2.2) \quad d(x, y) < \kappa\varepsilon \implies e^{-C_\varepsilon} \leq \left| \frac{J^{cu}(x)}{J^{cu}(y)} \right| \leq e^{C_\varepsilon}.$$

Note that by continuity of F^{cu} , $C_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow 0$.

Again, the absolute continuity of the stable foliation yields:

$$\begin{aligned} m_\varepsilon &\leq \sum_i \text{Leb}(B_n(\xi_i, \varepsilon)) \leq \sum_i \text{Leb}(W_{\kappa\varepsilon}^{ss}(B_n^{cu}(\xi_i, \kappa\varepsilon))) \\ &\leq C_s(\kappa.\varepsilon)^{\dim E^{ss}} \cdot \sum_i \int_{f^{n-1}(B_n^{cu}(\xi_i, \kappa\varepsilon))} e^{S_n(V)(x)} dx \\ &\leq C_s(\kappa.\varepsilon)^{\dim E^{ss}} \cdot e^{nC_\varepsilon} \sum_i e^{S_n(V)(\xi_i)} \text{Leb}^{cu}(B^{cu}(f^{n-1}(\xi_i), \kappa.\varepsilon)) \\ &\leq C_s C_{cu}(\kappa.\varepsilon)^{\dim E^{ss}} \cdot e^{nC_\varepsilon} \sum_i e^{S_n(V)(\xi_i)}. \end{aligned}$$

Where the second inequality (thus the constant C_s) comes from (A.1) (see Subsection A.2.3) and conformity of the Lebesgue measure. C_{cu} is a uniform upper bound of the Lebesgue measure on $B^{cu}(x, \rho)$.

Taking $\frac{1}{n}$ log in the last inequality, then taking $\limsup_{n \rightarrow +\infty}$ and then $\varepsilon \rightarrow 0$ we get

$$\mathcal{P}(V) \geq 0. \quad \square$$

2.3. Proof for the existence of ergodic SRB measures

The map f is \mathcal{C}^2 , we can thus use the Pesin theory. By Lemma 1.3, V has an equilibrium state. By Proposition 2.2, $\mathcal{P}(V) \geq 0$ and by Lemma 2.1,

$\mathcal{P}(V) \leq 0$. This yields $\mathcal{P}(V) = 0$. Then, the second part of Lemma 2.1 yields the existence of an ergodic equilibrium state for V , μ , which satisfies:

$$h_\mu(f) = \sum \lambda_\mu^+.$$

The fact that F^{cu} is sectional expanded and has dimension at least 2 implies that every ergodic measure has a positive Lyapunov exponent. So does μ .

Therefore, μ is an SRB measure and the proof of the existence of ergodic SRB measures in Theorem 1.2 is complete.

3. The SRB measure is unique and physical

Almost all ergodic components of an SRB measure are also SRB measures (see [11, Corollaire 4.10]). To get uniqueness for the SRB measure, it is thus sufficient to get uniqueness for ergodic SRB measures. The uniqueness will follow from the fact that the disintegration of an SRB measure for a $C^{1+\alpha}$ diffeomorphism is more than absolutely continuous with respect to Leb^u ; it is actually equivalent to it.

LEMMA 3.1. — *Let μ be an ergodic SRB measure for $f = \varphi_1$. Let $\{\mu_u\}$ be its system of conditional measures with respect to any measurable partition subordinate to the unstable foliation. Then almost every μ_u is equivalent to Leb^u .*

Proof. — Remember (see Lemma A.1) that there exist two measurable subbundles $E^c = \langle X \rangle$ and E_μ^u defined μ -almost everywhere and invariant for the time-one map f such that

$$E^{ss} \oplus \underbrace{\langle X \rangle \oplus E_\mu^u}_{=F^{cu}}.$$

Let ξ be a partition subordinate to the (Pesin) unstable foliation W^u of the time-one map f . From the remark in [12, p. 513], for μ -a.e. x , the density ρ_x of μ_u on $\xi(x)$ with respect to Leb^u satisfies

$$\frac{\rho_x(z)}{\rho_x(y)} = \prod_{k=0}^{+\infty} \frac{\left| \det Df|_{E_\mu^u}(f^{-k}(z)) \right|}{\left| \det Df|_{E_\mu^u}(f^{-k}(y)) \right|}.$$

By the (non-uniformly) backward contraction property of the unstable manifold, the above quantity converges and is also bounded away from zero, uniformly in y and z chosen in $\xi(x)$. This shows that the density is a continuous function on $\xi(x)$ and does not vanish. □

Let us now prove the uniqueness for the SRB measure and that it is physical. For the map f and the measure μ , a point x is said to be *forward μ -generic* if $1/n \sum_{i=0}^{n-1} \delta_{f^i(x)} \rightarrow \mu$ as $n \rightarrow \infty$.

LEMMA 3.2. — *Let μ be an ergodic SRB measure. Then there exists a non-empty open set U such that Lebesgue almost every point in U is forward μ -generic.*

Proof. — Since μ is an SRB measure, we have that $h_\mu + \int V d\mu = 0$. Thus μ cannot be concentrated on a singularity (see the first part of the proof of Lemma 2.1). Then it is hyperbolic for the flow φ_t by Lemma A.1. Let ξ be any measurable partition subordinate to the unstable foliation (for μ) and choose x_1 forward μ -generic. We can assume that x_1 is in the interior of $\xi(x_1)$. By Lemma 3.1, Lebesgue a.e. every x in $\xi(x_1)$ is forward μ -generic. Clearly, if x is forward μ -generic, then for every t , $\varphi_t(x)$ is also forward μ -generic. Moreover, every y in $W^{ss}(x)$ is also forward μ -generic. Let us set

$$U := \bigcup_{t \in]-\varepsilon, \varepsilon[, x \in \xi(x_1)} W_\varepsilon^{ss}(\varphi_t(x)).$$

By the hyperbolicity of μ , ξ_1 is a $(\dim F^{cu} - 1)$ -dimensional sub-manifold. Consequently, $\varphi_{[-\varepsilon, \varepsilon]}(\xi(x_1))$ is a $\dim F^{cu}$ -dimensional sub-manifold. Hence U is an open set by the invariance of domain theorem. By the absolute continuity of the strong stable foliation, Lebesgue almost every point in U is forward μ -generic. \square

Remark 3.3. — With almost the same proof of Lemma 3.2, we have the following result. If μ is a hyperbolic SRB measure for a C^2 diffeomorphism f or a C^2 vector field X , if $\text{supp}(\mu)$ admits a partially hyperbolic splitting $E^{ss} \oplus F$ satisfying that $\dim E^{ss}$ is exactly the sum of the multiplicities of negative Lyapunov exponents of μ , then there is a non-empty open set U in the manifold such that Lebesgue almost every point in U is forward μ -generic.

We can now prove the uniqueness of the SRB measure. Assume that μ_1 and μ_2 are two different ergodic SRB measures. Let U_1 and U_2 obtained from Lemma 3.2 for μ_1 and μ_2 . Now, transitivity shows that for some T , $\varphi_T(U_2) \cap U_1$ is a non-empty open set. Lebesgue almost every point in this intersection is forward μ_1 and μ_2 -generic, which shows that $\mu_1 = \mu_2$.

Appendix A. Classical results on (partially) hyperbolicity

A.1. Oseledets splitting for ergodic measures

For an ergodic invariant measure μ , denote by E_μ^s the sub-bundle associated to all the negative Lyapunov exponents and E_μ^u the sub-bundle associated to all the positive Lyapunov exponents.

An important property for singular hyperbolic attractors is that every invariant ergodic measure is *hyperbolic* in the following sense:

LEMMA A.1. — *If an ergodic measure μ is not supported on the set of singularities⁽¹⁾, then it is hyperbolic, i.e., $D\varphi_t$ has exactly one zero Lyapunov exponent that is given by the vector field X . Moreover, $E_\mu^s = E^{ss}$, and $F^{cu} = \langle X \rangle \oplus E_\mu^u$.*

Sketch of the proof. — Since μ is ergodic and is not supported on the set of singularities, we have that almost every point x of μ is a non-singular point and recurrent. Thus, there is a sequence (t_n) such that $\lim_{n \rightarrow \infty} t_n = \infty$ and $\lim_{n \rightarrow \infty} \varphi_{t_n}(x) = x$. Consequently $\lim_{n \rightarrow \infty} X(\varphi_{t_n}(x)) = X(x) \neq 0$. The Lyapunov exponent along $X(x)$ is

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{1}{t} \log \frac{|D\varphi_t X(x)|}{|X(x)|} &= \lim_{n \rightarrow \infty} \frac{1}{t_n} \log \frac{|D\varphi_{t_n} X(x)|}{|X(x)|} \\ &= \lim_{n \rightarrow \infty} \frac{1}{t_n} \log \frac{|X(\varphi_{t_n}(x))|}{|X(x)|} = 0. \end{aligned}$$

Now we show that there is at most one zero Lyapunov exponent. Note that the non-zero vector that has zero Lyapunov exponent cannot be contained in E^{ss} . Thus one can assume that it is contained in F^{cu} . Assume by contradiction that there is a vector $v \in F^{cu}(x) \setminus \langle X(x) \rangle$, such that the Lyapunov exponent of v is zero. We consider the plane generated by $X(x)$ and v . The sectional expansion property implies that the area of this plane is expanded by $D\varphi_t$. However, this contradicts to the fact that the two vectors $X(x)$ and v all have zero Lyapunov exponents.

The above arguments also implies that $E_\mu^s = E^{ss}$, and $F^{cu} = \langle X \rangle \oplus E_\mu^u$. □

⁽¹⁾Note that μ is ergodic and it is not supported on the set of singularities. Hence no singularity can be regular for μ .

A.2. Local stable manifolds and central unstable plaques

A.2.1. Local stable manifolds

One can continuously extend the bundles E^{ss} and F^{cu} in a neighborhood U of Λ (see [8, Proposition 2.7]). Note that the extension of E^{ss} can be made Df -invariant (see the proof of [8, Corollary 2.8]⁽²⁾, but generally we do not know how to extend F^{cu} to be an invariant bundle in U . We still denote the extended bundles by E^{ss} and F^{cu} .

Then, since Λ is an attractor, one can construct a stable foliation in U . More precisely, there is $\varepsilon_\Lambda > 0$ such that for any $x \in U$, $W_{\varepsilon_\Lambda}^{ss}$ is a local embedded sub-manifold of dimension $\dim E^{ss}$ by [8, Theorem 4.3] and the proof of [9, Theorem 4.1]. There are C^1 maps $\Psi_x^{ss} : E^{ss}(x) \rightarrow F^{cu}(x)$ that are C^1 close to the zero maps (on a compact neighborhood of the zero section) uniformly w.r.t. x and vary continuously w.r.t. x in the C^1 topology such that

$$W_{\varepsilon_\Lambda}^{ss}(x) = \exp_x(\text{Graph}(\Psi_x^{ss})(\varepsilon_\Lambda))$$

where $\text{Graph}(\Psi_x^{ss})(\varepsilon_\Lambda) = \{(v, \Psi_x^{ss}(v)) : v \in E^{ss}(x), \|v\| \leq \varepsilon_\Lambda\}$. Note that C and λ are the constants in the definition of the singular hyperbolicity. One can choose ε_Λ such that every point $y \in W_{\varepsilon_\Lambda}^{ss}$ satisfies:

$$d(\varphi_t(x), \varphi_t(y)) \leq 2Ce^{-\lambda t/2}, \text{ and } \frac{d(\varphi_t(x), \varphi_t(y))}{\|D\varphi_t|_{F(x)}\|} \leq 2Ce^{-\lambda t/2}, \quad \forall t \geq 0.$$

More generally, one can define $W_\varepsilon^{ss}(x)$ for every $\varepsilon < \varepsilon_\Lambda$ by taking the restricted graph.

These (local) strong stable manifolds generate a (global) stable foliation in a neighborhood of Λ ,

$$W^{ss}(x) := \bigcup_{t \geq 0} \varphi_{-t} W_{\varepsilon_\Lambda}^{ss}(\varphi_t(x)).$$

A.2.2. cu -plaques

These are locally invariant center-unstable sub-manifolds (called plaques in [9, Theorem 5.5], or [8, Theorem 4.5]) $W_{loc}^{cu}(x)$ tangent to $F^{cu}(x)$ and containing the image by \exp_x of a disk of radius ε_Λ .

(2) The statement of [8, Corollary 2.8] does not give this statement directly. However, its proof applies. More precisely, we do not use the perturbation part of diffeomorphisms, we only use the part of the neighborhood; from the proof, we know that the dominated bundle (the weaker bundle) can be extended in a neighborhood in an invariant way for an attractor.

More precisely, for each point x , there are C^1 maps $\Psi_x^{cu} : F^{cu}(x) \rightarrow E^{ss}(x)$ that are C^1 close to the zero maps (on a compact neighborhood of the zero section) uniformly w.r.t. x and vary continuously w.r.t. x in the C^1 topology such that

$$W_{\varepsilon_\Lambda}^{cu}(x) = \exp_x(\text{Graph}(\Psi_x^{cu})(\varepsilon_\Lambda)),$$

where $\text{Graph}(\Psi_x^{cu})(\varepsilon_\Lambda) = \{(v, \Psi_x^{cu}(v)) : v \in F^{cu}(x), \|v\| \leq \varepsilon_\Lambda\}$.

A.2.3. Absolute continuity, local product structure and metrics for sub-manifolds

The stable foliation has the *absolutely continuous* property. Given $y \in W_{\varepsilon_\Lambda}^{ss}(x)$, any two transversals Σ_y at y and Σ_x at x to the stable foliation, the stable foliation induces a holonomy map h^{Σ_x, Σ_y} . When the vector field is C^2 , $(h^{\Sigma_x, \Sigma_y})_* \text{Leb}_{\Sigma_x}$ is equivalent to Leb_{Σ_y} . Furthermore, the density function can be uniformly bounded when Σ_x and Σ_y are tangent to the cone field \mathcal{C}_a^F for a small, where at each point x

$$\mathcal{C}_a^F = \{v \in T_x M : v = v^s + v^c, v^s \in E^{ss}(x), v^c \in F^{cu}(x), |v^s| \leq a|v^c|\}.$$

One can see [17, Theorem 7.1] for the details of the absolute continuity of the stable foliation. Then, the absolute continuity yields for any measurable set A in a cu -plaque $W_{loc}^{cu}(x)$,

$$(A.1) \quad \frac{1}{C_s} \varepsilon^{\dim E^{ss}} \text{Leb}_{W_{loc}^{cu}(x)}^{cu}(A) \leq \text{Leb}(W_{\kappa\varepsilon}^{ss}(A)) \leq C_s \varepsilon^{\dim E^{ss}} \text{Leb}_{W_{loc}^{cu}(x)}^{cu}(A).$$

Roughly speaking, local stable manifolds and cu -plaques generate a local product structure with

$$B(x, \varepsilon) \approx W_{\varepsilon}^{ss}(x) \times W_{\varepsilon}^{cu}(x),$$

and absolutely continuity implies a local equivalence $\text{Leb} \sim \text{Leb}_x^{ss} \otimes \text{Leb}_x^{cu}$.

There exists some universal constant $\kappa > 1$ independent of x and ε such that for every small enough $\varepsilon > 0$, and for every x ,

- (D1) $W_{\varepsilon/\kappa}^{ss}(B^{cu}(x, \varepsilon/\kappa)) \subset B(x, \varepsilon) \subset W_{\kappa\varepsilon}^{ss}(B^{cu}(x, \kappa\varepsilon))$,
- (D2) for any $y, z \in W_{\varepsilon}^{ss}(x)$, we have $d_M(y, z)/\kappa \leq d^{ss}(y, z) \leq \kappa d_M(y, z)$, where d^{ss} is the metric on $W_{\varepsilon_\Lambda}^{ss}(x)$ and d_M is the metric on M ,
- (D3) for any $y, z \in W_{\varepsilon}^{cu}(x)$, we have $d_M(y, z)/\kappa \leq d^{cu}(y, z) \leq \kappa d_M(y, z)$, where d^{cu} is the metric on $W_{\varepsilon_\Lambda}^{cu}(x)$ and d_M is the metric on M .

These properties are referred as the uniform control between the metric of M and the metrics of its sub-manifolds.

BIBLIOGRAPHY

- [1] V. ARAÚJO, M. J. PACIFICO, E. R. PUJALS & M. VIANA, “Singular-hyperbolic attractors are chaotic”, *Trans. Am. Math. Soc.* **361** (2009), no. 5, p. 2431-2485.
- [2] A. ARROYO & E. R. PUJALS, “Dynamical properties of singular-hyperbolic attractors”, *Discrete Contin. Dyn. Syst.* **19** (2007), no. 1, p. 67-87.
- [3] R. BOWEN, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, 2^e ed., Lecture Notes in Mathematics, vol. 470, Springer, 2008, viii+75 pages.
- [4] R. BOWEN & D. RUELLE, “The ergodic theory of Axiom A flows”, *Invent. Math.* **29** (1975), no. 3, p. 181-202.
- [5] J. BUZZI, “No or infinitely many a.c.i.p. for piecewise expanding C^r maps in higher dimensions”, *Commun. Math. Phys.* **222** (2001), no. 3, p. 495-501.
- [6] W. COLMENÁREZ, “SRB measures for singular hyperbolic attractors”, PhD Thesis, Universidade Federal do Rio de Janeiro (Brazil), 2002.
- [7] W. COWIESON & L.-S. YOUNG, “SRB measures as zero-noise limits”, *Ergodic Theory Dyn. Syst.* **25** (2005), no. 4, p. 1115-1138.
- [8] S. CROVISIER & R. POTRIE, “Introduction to partially hyperbolic dynamics”, www.crm.umontreal.ca/sms/2017/pdf/diapos/00-CP-Trieste-Version1.pdf, 2015.
- [9] M. W. HIRSCH, C. C. PUGH & M. SHUB, *Invariant manifolds*, Lecture Notes in Mathematics, vol. 583, Springer, 1977, ii+149 pages.
- [10] A. KATOK & B. HASSELBLATT, *Introduction to the modern theory of dynamical systems*, Encyclopedia of Mathematics and its Applications, vol. 54, Cambridge University Press, 1995, With a supplementary chapter by Katok and Leonardo Mendoza, xviii+802 pages.
- [11] F. LEDRAPPIER, “Propriétés ergodiques des mesures de Sinai”, *Publ. Math., Inst. Hautes Étud. Sci.* (1984), no. 59, p. 163-188.
- [12] F. LEDRAPPIER & L.-S. YOUNG, “The metric entropy of diffeomorphisms. I. Characterization of measures satisfying Pesin’s entropy formula”, *Ann. Math.* **122** (1985), no. 3, p. 509-539.
- [13] E. N. LORENZ, “Deterministic non-periodic flow”, *J. Atmospheric Sci.* **20** (1963), p. 130-141.
- [14] J. E. MARSDEN & M. F. MCCrackEN, *The Hopf bifurcation and its applications*, Applied Mathematical Sciences, vol. 19, Springer-Verlag, New York, 1976, xiii+408 pages.
- [15] R. J. METZGER & C. A. MORALES, “Sectional-hyperbolic systems”, *Ergodic Theory Dyn. Syst.* **28** (2008), no. 5, p. 1587-1597.
- [16] C. A. MORALES, M. J. PACIFICO & E. R. PUJALS, “Robust transitive singular sets for 3-flows are partially hyperbolic attractors or repellers”, *Ann. Math.* **160** (2004), no. 2, p. 375-432.
- [17] Y. B. PESIN, *Lectures on partial hyperbolicity and stable ergodicity*, Zurich Lectures in Advanced Mathematics, European Mathematical Society (EMS), 2004, vi+122 pages.
- [18] H. QIU, “Existence and uniqueness of SRB measure on C^1 generic hyperbolic attractors”, *Commun. Math. Phys.* **302** (2011), no. 2, p. 345-357.
- [19] D. RUELLE, “A measure associated with axiom-A attractors”, *Am. J. Math.* **98** (1976), no. 3, p. 619-654.
- [20] Y. G. SINAI, “Gibbs measures in ergodic theory”, *Usp. Mat. Nauk* **27** (1972), no. 4(166), p. 21-64.
- [21] M. TSUJII, “Piecewise expanding maps on the plane with singular ergodic properties”, *Ergodic Theory Dyn. Syst.* **20** (2000), no. 6, p. 1851-1857.

- [22] P. WALTERS, *An introduction to ergodic theory*, Graduate Texts in Mathematics, vol. 79, Springer, 1982, ix+250 pages.
- [23] J. YANG, Online video, <http://video.impa.br/index.php?page=international-conference-on-dynamical-system>.
- [24] ———, “Topological entropy of Lorenz-like flows”, <https://arxiv.org/abs/1412.1207>, 2014.
- [25] L.-S. YOUNG, “What are SRB measures, and which dynamical systems have them?”, *J. Stat. Phys.* **108** (2002), no. 5-6, p. 733-754, Dedicated to David Ruelle and Yasha Sinai on the occasion of their 65th birthdays.
- [26] S. ZHU, S. GAN & L. WEN, “Indices of singularities of robustly transitive sets”, *Discrete Contin. Dyn. Syst.* **21** (2008), no. 3, p. 945-957.

Manuscrit reçu le 14 janvier 2016,
révisé le 31 janvier 2017,
accepté le 1^{er} mars 2017.

Renaud LEPLAIDEUR
LMBA, UMR6205
Université de Brest
6, avenue Victor Le Gorgeu
C.S. 93837, France
renaud.leplaidur@univ-brest.fr

Dawei YANG
School of Mathematical Sciences,
Soochow University,
No. 1, Shizi Street,
Suzhou, 215006, P.R. China
yangdw@suda.edu.cn