

ANNALES DE L'INSTITUT FOURIER

S. K. BAJPAI

On the mean values of an entire function represented by Dirichlet series

Annales de l'institut Fourier, tome 21, n° 2 (1971), p. 31-34

<http://www.numdam.org/item?id=AIF_1971__21_2_31_0>

© Annales de l'institut Fourier, 1971, tous droits réservés.

L'accès aux archives de la revue « Annales de l'institut Fourier » (<http://annalif.ujf-grenoble.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

ON THE MEAN VALUES OF AN ENTIRE FUNCTION REPRESENTED BY DIRICHLET SERIES

by S. K. BAJPAI

1. Let $f(s)$ be an entire function of the complex variable $s = \sigma + it$ defined everywhere in the complex plane by absolutely convergent Dirichlet series

$$(1.1) \quad f(s) = \sum_{n=0}^{\infty} a_n e^{s\lambda_n}$$

where $0 \leq \lambda_0 < \lambda_1 < \dots < \lambda_n \rightarrow \infty$ as $n \rightarrow \infty$.

As usual, the symbols $M(\sigma, f)$, $\mu(\sigma, f)$ and $v(\sigma, f)$ denote the maximum modulus, the maximum term and the rank of the maximum term respectively for $f(s)$, and can be found in [8]. We define

$$A_k(\sigma) = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T |f(\sigma + it)|^k dt$$

and

$$W_{y, \delta, k}(\sigma) = \lim_{T \rightarrow \infty} \frac{e^{-\delta\sigma}}{2T} \int_y^{\sigma} \int_{-T}^T |f(x + it)|^k e^{\delta x} dx dt; \quad \delta > 0.$$

The mean values for $k = 2$ and $y = 0$ were defined by Hadamard [2] and also by Kamthan ([3], [4]) who has obtained the following main results for $k = 2$ and $y = 0$

$$(1.2) \quad \lim_{\sigma \rightarrow \infty} \inf \frac{\log \log A_2(\sigma)}{\sigma} = \frac{\rho}{\lambda}$$

$$(1.3) \quad \lim_{\sigma \rightarrow \infty} \inf \frac{\log \log W_{0, \delta, 2}(\sigma)}{\sigma} = \frac{\rho}{\lambda}$$

Juneja [6] also proved (1.3) by using a different technique for $k = 2$ and $y = 0$. In [7] another extension of (1.3)

has been obtained for $k = 2$ and $y = -\infty$. Kamthan [5] has further attempted to establish (1.2) for every $k > 0$, but his arguments of the proof are vague. Thus, in the present paper, we restrict ourselves to functions of finite Ritt-order ρ , establish (1.2) and extend (1.3) for every $k > 0$ by a different technique.

2. We have

THEOREM 1. — *If $f(s)$, defined by Dirichlet series (1.1), be an entire function of finite Ritt-order ρ , lower order λ , type T and lower type t , then, for $0 < k < \infty$*

$$\frac{\rho}{\lambda} = \lim_{\sigma \rightarrow \infty} \inf \frac{\log \log A_k(\sigma)}{\sigma} = \lim_{\sigma \rightarrow \infty} \inf \frac{\log \log W_{y, \delta, k}(\sigma)}{\sigma}$$

and

$$\frac{kT}{kt} = \lim_{\sigma \rightarrow \infty} \inf \frac{\log A_k(\sigma)}{e^{\rho \sigma}} = \lim_{\sigma \rightarrow \infty} \inf \frac{\log W_{y, \delta, k}(\sigma)}{e^{\rho \sigma}}$$

First we prove the following :

LEMMA. — *Let $f(s)$, given by (1.1), be an entire function of finite Ritt-order ρ , then, for $0 < k < \infty$*

$$\log A_k(\sigma) \sim \log W_{y, \delta, k}(\sigma) \sim k \log M(\sigma, f).$$

Proof. — It is well known

$$(2.1) \quad |a_n| e^{\sigma \lambda_n} = \left| \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T f(\sigma + it) e^{-i \lambda_n t} dt \right|$$

First, let $k \geq 1$, then by Holder's inequality, we have

$$(2.2) \quad |a_n| e^{\sigma \lambda_n} \leq C_k \left[\lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T |f(\sigma + it)|^k dt \right]^{1/k}$$

where

$$C_k = \begin{cases} 1 & \text{if } k = 1 \\ 4 \left[\frac{\Gamma \left(\frac{1}{2} + \frac{k}{2(k-1)} \right)}{\sqrt{\pi} \Gamma \left(1 + \frac{k}{2(k-1)} \right)} \right]^{\frac{k-1}{k}} & \text{if } k > 1. \end{cases}$$

Further, from (2.2) we have

$$\{|a_n|e^{x\lambda_n}\|^k \leq C_k^k \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T |f(x+it)|^k dt$$

Hence, for $y \leq 0$

$$(2.3) \quad \frac{\{|a_n|e^{\sigma\lambda_n}\|^k}{k\lambda_n + \delta} - \frac{e^{-\delta\sigma}|a_n|^k e^{(k\lambda_n+\delta)y}}{k\lambda_n + \delta} \leq C_k^k W_{y,\delta,k}(\sigma) \leq \frac{C_k^k}{\delta} M^k(\sigma, f)$$

The case when $0 < k < 1$ may be treated as following:

$$(2.4) \quad |a_n|e^{\sigma\lambda_n} \leq C_{k+1} \left\{ \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T |f(\sigma+it)|^{k+1} dt \right\}^{\frac{1}{k+1}} \\ \leq C_{k+1} \{M(\sigma, f) A_k(\sigma)\}^{\frac{1}{k+1}}$$

Also, from (2.1), for $0 < k < 1$, we have

$$(2.5) \quad \frac{\{|a_n|e^{\sigma\lambda_n}\|^{k+1}}{(k+1)\lambda_n + \delta} - \frac{e^{-\delta\sigma}|a_n|^{k+1} e^{((k+1)\lambda_n+\delta)y}}{(k+1)\lambda_n + \delta} \\ \leq C_{k+1}^{k+1} \lim_{T \rightarrow \infty} \frac{e^{-\delta\sigma}}{2T} \int_0^\sigma \int_{-T}^T e^{x\delta} |f(x+it)|^{k+1} dt dx \\ \leq C_{k+1}^{k+1} M(\sigma, f) W_{y,\delta,k}(\sigma) \leq \frac{1}{\delta} C_{k+1}^{k+1} M^{k+1}(\sigma, f)$$

Since $f(s)$ is of finite Ritt-order ρ , it follows that ([1], p. 719), ([9], p. 265)

$$(2.6) \quad \log \mu(\sigma, f) \sim \log M(\sigma, f) \\ \text{and} \quad \log \log \mu(\sigma, f) \sim \log \lambda_{y(\sigma, f)}$$

Hence (2.2), (2.3), (2.4), (2.5) and (2.6) together imply the lemma. Proof of theorem 1, follows immediately from the above lemma.

THEOREM 2. — If $f(s)$ is an entire function of finite Ritt-order ρ and is defined by (1.1) then

$$\lim_{\sigma \rightarrow \infty} \inf \frac{\log \{W_{y,\delta,k}(\sigma, f')/W_{y,\delta,k}(\sigma, f)\}}{\sigma} = \frac{k\rho}{k\lambda}$$

provided k is an even integer.

For $k = 2$, this result is due to Juneja [6] and for k even it follows with the use of Minkowski's inequality and the

fact that $W_{\gamma, \delta, k}(\sigma)$ is an increasing convex function of σ for k even.

In the end author wishes to express his sincere thanks to Professor R. S. L. Srivastava for his kind guidance. The author is also thankful to the referee for his suggestions.

REFERENCES

- [1] A. G. AZPEITIA, On the maximum modulus and the maximum term of an entire Dirichlet series, *Proc. Amer. Math. Soc.*, **12** (1961), 717-721.
- [2] E. HILLE, Analytic function theory volume II; Blaisdell Pub. Comp., 1962.
- [3] P. K. KAMTHAN, On the mean values of entire functions represented by Dirichlet series; *Acta. Math. Acad. Sci. Hung.*, **15** (1964), 133-136.
- [4] P. K. KAMTHAN, On the mean values of entire function represented by Dirichlet series, *ibidem*, **16** (1965), 89-92.
- [5] P. K. KAMTHAN, On entire functions represented by Dirichlet series (IV); *Ann. Inst. Fourier*, Grenoble **16** (1966), 209-223.
- [6] O. P. JUNEJA, On the mean values of an entire function and its derivatives represented by Dirichlet series, *Ann. Polon. Math.*, **18** (1966), 307-313.
- [7] O. P. JUNEJA and K. N. AWASTHI, On the mean values of an entire function and its derivatives represented by Dirichlet series II, *Ann. Polon. Math.*, **22** (1969), 89-96.
- [8] S. MANDEL BROJT, Dirichlet series, *Rice Inst. Pamph.*, **31**, 4 (1944).
- [9] K. SUGIMURA, Übertragung einiger Satze aus der Theorie der ganzen Functionen auf Dirichletsche Reihen, *Math. Z.*, **29** (1928-1929), 264-277.

Manuscrit reçu le 24 avril 1970.

S. K. BAJPAI,

Department of Mathematics,
Indian Institute of Technology,
Kanpur 16, India.