

Journal de l'École polytechnique

Mathématiques

Karine BEAUCHARD & Camille LAURENT

Local exact controllability of the 2D-Schrödinger-Poisson system

Tome 4 (2017), p. 287-336.

http://jep.cedram.org/item?id=JEP_2017__4__287_0

© Les auteurs, 2017.

Certains droits réservés.



Cet article est mis à disposition selon les termes de la licence
CREATIVE COMMONS ATTRIBUTION – PAS DE MODIFICATION 3.0 FRANCE.
<http://creativecommons.org/licenses/by-nd/3.0/fr/>

L'accès aux articles de la revue « Journal de l'École polytechnique — Mathématiques » (<http://jep.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jep.cedram.org/legal/>).

Publié avec le soutien
du Centre National de la Recherche Scientifique

cedram

Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>

LOCAL EXACT CONTROLLABILITY OF THE 2D-SCHRÖDINGER-POISSON SYSTEM

BY KARINE BEAUCHARD & CAMILLE LAURENT

ABSTRACT. — In this article, we investigate the exact controllability of the 2D-Schrödinger-Poisson system, which couples a Schrödinger equation on a bounded domain of \mathbb{R}^2 with a Poisson equation for the electrical potential. The control acts on the system through a Neumann boundary condition on the potential, locally distributed on the boundary of the space domain. We prove several results, with or without nonlinearity and with different boundary conditions on the wave function, of Dirichlet type or of Neumann type.

RÉSUMÉ (Contrôlabilité locale exacte du système de Schrödinger-Poisson 2D)

Dans cet article, nous étudions la contrôlabilité exacte du système de Schrödinger-Poisson 2D, qui couple une équation de Schrödinger sur un ouvert borné 2D, avec une équation de Poisson pour le potentiel électrique. Le contrôle agit sur le système via une condition de Neumann sur le potentiel, localement distribuée sur le bord du domaine spatial. Nous démontrons plusieurs résultats, avec ou sans non-linéarité, avec différents types de conditions de bord sur la fonction d'onde, de type Dirichlet ou de type Neumann.

CONTENTS

| | |
|--|-----|
| 1. Introduction..... | 288 |
| 2. Comments on our assumptions..... | 296 |
| 3. Well-posedness of the Schrödinger-Poisson system and linearization..... | 300 |
| 4. Local exact control of high frequencies..... | 309 |
| 5. Local exact control around eigenfunctions..... | 315 |
| 6. Schrödinger equation with real-valued boundary controls..... | 320 |
| 7. Nonlinear equation on a rectangle..... | 322 |
| References..... | 334 |

MATHEMATICAL SUBJECT CLASSIFICATION (2010). — 35Q40, 35Q41, 93C10, 93C20.

KEYWORDS. — Control of partial differential equations, Schrödinger-Poisson system, bilinear control.

The authors were partially supported by the “Agence Nationale de la Recherche” (ANR), Projet Blanc EMAQS number ANR-2011-BS01-017-01.

1. INTRODUCTION

Microelectronics industry has driven transistor sizes to the nanometer scale. This has led to the possibility of building nanostructures like single electron transistors or single electron memories, which involve the transport of only a few electrons. In general, such devices consist in an active region, on which the electrical potential can be tuned by an electrode (the gate). In many applications, the performance of the device will depend on the possibility of controlling the electrons by acting on the gate voltage.

At the nanometer scale, quantum effects become important and a quantum transport model is necessary. In this paper, we analyze the controllability of a simplified mathematical model, of the quantum transport of electrons trapped in a two-dimensional device. The model consists in a single Schrödinger equation, on a 2D bounded domain, coupled to the Poisson equation for the electrical potential. The control acts on the system through a Neumann boundary condition on the potential, on a part of the boundary, modeling the gate. Such a model has already been studied in [24] for controllability purposes.

It would be physically relevant to impose Dirichlet boundary conditions on the wave function and to take into account the self-consistent potential modeling interactions between electrons. However, the mathematical analysis of this configuration is quite complicated, thus we investigate two simpler configurations:

- (1) a first configuration, in which Dirichlet boundary conditions are imposed on the wave function, but we neglect the self-consistent potential,
- (2) a second configuration, in which we take into account the self-consistent potential, but Neumann boundary conditions are imposed on the wave function.

This work is a first step towards more realistic models.

1.1. LINEAR PDE WITH DIRICHLET BOUNDARY CONDITIONS. — First, we consider the system

$$(1) \quad \begin{cases} (i\partial_t + \Delta)\psi(t, x) = v(t, x)\psi(t, x), & (t, x) \in (0, T) \times \Omega, \\ \psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \psi(0, x) = \psi_0(x) & x \in \Omega, \\ (-\Delta + 1)v(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu v(t, x) = g(t, x)1_{\Gamma_c}(x), & (t, x) \in (0, T) \times \partial\Omega, \end{cases}$$

where Ω is a bounded open subset of \mathbb{R}^2 , Γ_c is an open subset of $\partial\Omega$ and 1_{Γ_c} is its characteristic function. The control is the real valued function g and we want to control the wave function ψ .

1.1.1. *Definitions and notation.* — Let Ω be either a smooth bounded open subset of \mathbb{R}^2 or the rectangle $(0, \pi) \times (0, L)$ with $L > 0$.

We denote by $\langle \cdot, \cdot \rangle$ the complex valued scalar product on $L^2(\Omega, \mathbb{C})$,

$$\langle f, g \rangle := \int_{\Omega} f(x) \overline{g(x)} \, dx,$$

by \mathcal{S} the $L^2(\Omega, \mathbb{C})$ -sphere

$$\mathcal{S} := \{ \xi \in L^2(\Omega, \mathbb{C}) ; \|\xi\|_{L^2(\Omega)} = 1 \},$$

by $(-\Delta_D)$ the Laplace operator associated with Dirichlet boundary conditions

$$D(-\Delta_D) := H^2 \cap H_0^1(\Omega, \mathbb{C}), \quad -\Delta_D \xi := -\Delta \xi,$$

by $(\lambda_k)_{k \in \mathbb{N}^*}$ the nondecreasing sequence of its eigenvalues, by $(\varphi_k)_{k \in \mathbb{N}^*}$ one associated orthonormal basis of eigenfunctions,

$$\begin{cases} -\Delta \varphi_k(x) = \lambda_k \varphi_k(x), & x \in \Omega, \\ \varphi_k(x) = 0, & x \in \partial\Omega, \\ \|\varphi_k\|_{L^2(\Omega)} = 1, \end{cases}$$

by $H_{(0)}^3(\Omega)$ the Sobolev space

$$H_{(0)}^3(\Omega) := D((-\Delta_D)^{3/2}) = \{ \xi \in H^3 \cap H_0^1(\Omega, \mathbb{C}) ; \Delta \xi \in H_0^1(\Omega, \mathbb{C}) \},$$

and by \mathbb{P}_K , for $K \in \mathbb{N}^*$, the projection

$$\begin{aligned} \mathbb{P}_K : H^{-1}(\Omega, \mathbb{C}) &\longrightarrow \text{Adh}_{H^{-1}(\Omega)}(\text{Span}(\varphi_k ; k \geq K)) \\ \xi &\longmapsto \xi - \sum_{k=1}^{K-1} \langle \xi, \varphi_k \rangle_{H^{-1}, H_0^1} \varphi_k, \end{aligned}$$

where $\langle \cdot, \cdot \rangle_{H^{-1}, H_0^1}$ is the duality product between $H^{-1}(\Omega)$ and $H_0^1(\Omega)$. We also use the weak observability of the Schrödinger equation, defined below; several configurations (Ω, Γ_c) for which it has been proved are recalled in Section 2.1.

DEFINITION 1 (Observability and weak observability of Schrödinger equation)

Let Γ be an open subset of $\partial\Omega$ and $0 \leq \tau_1 < \tau_2 < \infty$. The Schrödinger equation on Ω is observable (resp. weakly observable) on $(\tau_1, \tau_2) \times \Gamma$ if there exists $\mathcal{C}_0 > 0$ such that

$$(2) \quad \begin{aligned} \|\phi_T\|_{H_0^1(\Omega)} &\leq \mathcal{C}_0 \|\partial_\nu \phi\|_{L^2((\tau_1, \tau_2) \times \Gamma)}, \quad \forall \phi_T \in H_0^1(\Omega, \mathbb{C}) \\ (\text{resp. } \|\phi_T\|_{H_0^1(\Omega)} &\leq \mathcal{C}_0 (\|\partial_\nu \phi\|_{L^2((\tau_1, \tau_2) \times \Gamma)} + \|\phi_T\|_{H^{-1}(\Omega)}), \quad \forall \phi_T \in H_0^1(\Omega, \mathbb{C})), \end{aligned}$$

where $\phi(t) := e^{i\Delta_D(t-T)} \phi_T$, i.e., ϕ is the solution of

$$(3) \quad \begin{cases} (i\partial_t + \Delta)\phi(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \phi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \phi(T, x) = \phi_T(x), & x \in \Omega. \end{cases}$$

The Schrödinger equation on Ω is observable (resp. weakly observable) on Γ if it is observable (resp. weakly observable) on $(\tau_1, \tau_2) \times \Gamma$ for some $0 \leq \tau_1 < \tau_2 < \infty$.

In the inequality (2), the last term $\|\phi_T\|_{H^{-1}(\Omega)}$ may be equivalently replaced by any term of the form $\|\phi_T\|_{H^s(\Omega)}$ with $-1 < s < 1$, by an interpolation argument.

1.1.2. *Local exact controllability of high frequencies.* — Our first result is the local exact controllability of high frequencies of the system (1), around any trajectory of the free system.

THEOREM 1. — *Let Ω be a bounded open subset of \mathbb{R}^2 , Γ_c be a non-empty open subset of $\partial\Omega$, $T > 0$, $\psi_0 \in H_{(0)}^3(\Omega) \cap \mathcal{S}$ and $\psi_{\text{ref}}(t) := e^{i\Delta_D t}\psi_0$. We assume that*

(**H1**) *either Ω is of class C^∞ and locally on one side of $\partial\Omega$; or $\Omega = (0, \pi) \times (0, L)$ for some $L > 0$ and $\bar{\Gamma}_c$ does not contain any vertex of Ω ,*

(**H2**) *the Schrödinger equation on Ω is weakly observable on $(0, \tilde{T}) \times \Gamma_c$ for some $\tilde{T} \in (0, T)$,*

(**H3**) *$|\partial_\nu \psi_{\text{ref}}(t, x)| \geq m > 0$, $\forall (t, x) \in (T', T'') \times \Gamma_c$ for some $T', T'' \in [0, T]$ such that $T'' - T' > \tilde{T}$.*

Then, there exists $K, \delta > 0$ and a C^1 -map

$$\Upsilon : \mathcal{V} \longrightarrow L^2((0, T) \times \Gamma_c, \mathbb{R}),$$

where $\mathcal{V} := \{\psi_f \in \mathbb{P}_K[H_{(0)}^3(\Omega, \mathbb{C})]; \|\psi_f - \mathbb{P}_K[\psi_{\text{ref}}(T)]\|_{H_{(0)}^3} < \delta\}$, such that

- $\Upsilon(\mathbb{P}_K[\psi_{\text{ref}}(T)]) = 0$,
- for every $\psi_f \in \mathcal{V}$, the solution of (1) with control $g = \Upsilon(\psi_f)$ satisfies

$$\mathbb{P}_K[\psi(T)] = \psi_f.$$

As a consequence, there exists $K' \geq K$ and $g \in L^2((0, T) \times \Gamma_c, \mathbb{R})$ such that the solution of (1) satisfies $\mathbb{P}_{K'}[\psi(T)] = 0$; in particular, $x \mapsto \psi(T, x)$ is a smooth function.

Such a result may be used to prove global exact controllability of the Schrödinger-Poisson system in $H_{(0)}^3(\Omega, \mathbb{C})$ with controls $g \in L^2((0, T), \mathbb{R})$, by following the strategy of [6]. This will be at the core of future works by the authors.

In particular, Theorem 1 applies, for arbitrary $T > 0$ and $\psi_0 \in H_{(0)}^3(\Omega) \cap \mathcal{S}$, when (see Propositions 1 and 2)

- $\Omega = (0, \pi) \times (0, L)$ for some $L > 0$ and Γ_c contains both a horizontal and a vertical segment,
- Ω is a disk and Γ_c is arbitrary.

When the Geometric Control Condition is fulfilled for (Ω, Γ_c) , then Theorem 1 applies for any $T > 0$ if $\psi_0 \in H_{(0)}^3(\Omega) \cap \mathcal{S}$ satisfies (**H3**) (see Proposition 1).

Assumption (**H1**) is important for the system (1) to be well-posed in $H_{(0)}^3(\Omega)$ with control $g \in L^2((0, T) \times \Gamma_c, \mathbb{R})$ (see Section 3.4). When Ω is a rectangle, it is important to assume that its vertices do not belong to $\bar{\Gamma}_c$, in order to take advantage of the usual elliptic regularity on the potential v .

1.1.3. *Local exact controllability around an eigenstate.* — Our second result is the local exact controllability of the system (1) around an eigenstate.

THEOREM 2. — *Let Ω be a bounded open subset of \mathbb{R}^2 , Γ_c be an open subset of $\partial\Omega$, $T > 0$, $R \in \mathbb{N}^*$ and $\psi_{\text{ref}}(t) := \varphi_R(x)e^{-i\lambda_R t}$. We assume **(H1)**, **(H2)**,*

(H3') $|\partial_\nu \varphi_R(x)| \geq m > 0, \forall x \in \Gamma_c$,

(H4) λ_R is a simple eigenvalue of $(-\Delta_D)$ and for any eigenvector Φ of $(-\Delta_D)$, the solution w of

$$\begin{cases} (-\Delta + 1)w(x) = \varphi_R(x)\Phi(x), & x \in \Omega, \\ \partial_\nu w(x) = 0, & x \in \partial\Omega, \end{cases}$$

does not identically vanish on Γ_c : $w|_{\Gamma_c} \not\equiv 0$.

Then, there exists $\delta > 0$ and a C^1 -map

$$\Upsilon : \mathcal{V} \longrightarrow L^2((0, T) \times \Gamma_c, \mathbb{R}),$$

where

$$\mathcal{V} := \{(\psi_0, \psi_f) \in [H^3_{(0)}(\Omega, \mathbb{C}) \cap \mathcal{S}]^2; \|\psi_0 - \psi_{\text{ref}}(0)\|_{H^3_{(0)}} + \|\psi_f - \psi_{\text{ref}}(T)\|_{H^3_{(0)}} < \delta\},$$

such that $\Upsilon[\psi_{\text{ref}}(0), \psi_{\text{ref}}(T)] = 0$ and, for every $(\psi_0, \psi_f) \in \mathcal{V}$, the solution of (1) with control $g = \Upsilon(\psi_0, \psi_f)$ satisfies $\psi(T) = \psi_f$.

Note that, under Assumption **(H1)**, then Assumption **(H3')** systematically holds with $R = 1$. The assumptions of Theorem 2 could look quite technical, but roughly speaking the main assumptions are:

- the weak observability (Assumption **(H2)**), that ensures the controllability of high frequencies, as in Theorem 1,
- the unique continuation property (Assumption **(H4)**), that ensures the controllability of low frequencies, see Proposition 5).

If **(H2)** holds, but **(H4)** is not satisfied, then the linearized system around ψ_{ref} is not controllable: it misses a finite number of directions corresponding to the eigenfunctions Φ for which **(H4)** is not satisfied, see Remark 2. It would be interesting to recover these directions by power series expansions [17, Chap. 8].

Theorem 2 applies, in particular, when $\Omega = (0, \pi) \times (0, L)$ for some $L > 0$, Γ_c is an open subset of $\partial\Omega$ that contains both a horizontal and a vertical segment and λ_R is a simple eigenvalue of $(-\Delta_D)$ (see Proposition 3).

The validity of Theorem 2 on the disk $\Omega = \{(x, y) \in \mathbb{R}^2; x^2 + y^2 < 1\}$ is an open problem: checking Assumption **(H4)** would require some study on the product of Bessel functions, which might require lengthy computations.

Following arguments by Méhats, Privat and Sigalotti [24] (relying on methods of Privat-Sigalotti [33]), it should be possible to prove that **(H4)** holds generically with respect to the domain Ω . This becomes clearer with some equivalent ways of expressing Assumption **(H4)** described in Proposition 4.

Assumption **(H4)** is quite unusual in control theory. It looks like a classical unique continuation property for eigenfunctions but it does not seem that we can refer to some already known results to prove it in great generality. We have indeed chosen to discuss about it more precisely in Subsection 2.3.

Exact controllability as in Theorem 2 probably holds close to any reference trajectory ψ_{ref} as done in Theorem 1. Yet, Assumption **(H4)** should be replaced by a unique continuation assumption complicated to state and depending on time. We have chosen to state Theorem 2 only close to an eigenfunction because the unique continuation assumption takes the nice form of **(H4)**.

1.2. CONTROLLABILITY OF SCHRÖDINGER EQUATION WITH REAL VALUED CONTROL

For the Schrödinger-Poisson system, the control g corresponds to a potential and therefore needs to be real valued to have a physical meaning. Therefore, as an intermediary result, we will also get the exact controllability of Schrödinger equation with *real valued* controls (instead of complex valued ones in the existing literature), when the equation is weakly observable. The result that we need for the control of the Schrödinger-Poisson system will actually be more complicated. Yet, we believed that it could be useful for other contexts and we give a simpler proof in the simpler context of the free Schrödinger equation (see for instance Araruna-Cerpa-Mercado-Santos [2] where related questions are raised).

THEOREM 3. — *Let Ω be an open subset of \mathbb{R}^2 , Γ be an open subset of $\partial\Omega$ and $0 < \tilde{T} < T < \infty$. If the Schrödinger equation on Ω is weakly observable on $(0, \tilde{T}) \times \Gamma$ then, for every $\psi_f \in H^{-1}(\Omega, \mathbb{C})$, there exists a real valued control $u \in L^2((0, T) \times \partial\Omega, \mathbb{R})$ such that the solution of*

$$\begin{cases} (i\partial_t + \Delta)\psi = 0, & (t, x) \in (0, T) \times \Omega, \\ \psi(t, x) = u(t, x)1_\Gamma(x), & (t, x) \in (0, T) \times \partial\Omega, \\ \psi(0, x) = 0, & x \in \Omega, \end{cases}$$

satisfies $\psi(T) = \psi_f$.

The proof will be given in Section 6.

1.3. NONLINEAR PDE, ON A RECTANGLE, WITH NEUMANN BOUNDARY CONDITIONS

Now, we consider the nonlinear Schrödinger-Poisson system on a rectangle

$$(4) \quad \begin{cases} i\partial_t\psi(t, x) = -\Delta\psi(t, x) + \tilde{v}(t, x)\psi(t, x), & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu\psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \psi(0, x) = \psi_0(x), & x \in \Omega, \\ (-\Delta + 1)\tilde{v}(t, x) = \varepsilon|\psi(t, x)|^2, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu\tilde{v}(t, x) = g(t, x)1_{\Gamma_c}(x), & (t, x) \in (0, T) \times \partial\Omega, \end{cases}$$

where $\varepsilon \in \mathbb{R}$, $x = (x_1, x_2) \in \Omega := (0, \pi) \times (0, L)$, $L > 0$ and Γ_c is an open subset of $\partial\Omega$. The control is the real valued function g and we want to control the wave function ψ .

For the nonlinear system (4), our main result is the local exact controllability around the reference trajectory, constant in space,

$$(5) \quad \left(\psi_{\text{ref}}(t, x) = \frac{e^{-i\varepsilon t/\sqrt{\pi L}}}{\sqrt{\pi L}}, \tilde{v}_{\text{ref}}(t, x) = 0, g_{\text{ref}}(t, x) = 0 \right).$$

We denote by Δ_N the Laplace operator associated with Neumann boundary conditions

$$\begin{aligned} D(\Delta_N) &= H_N^2(\Omega, \mathbb{C}) := \{ \varphi \in H^2(\Omega, \mathbb{C}) ; \partial_\nu \varphi = 0 \text{ on } \partial\Omega \} \\ \Delta_N \varphi &:= \Delta \varphi. \end{aligned}$$

THEOREM 4. — *Let $L > 0$, $\Omega := (0, \pi) \times (0, L)$, Γ_c be an open subset of $\partial\Omega$ such that $\bar{\Gamma}_c$ does not contain any vertex of Ω , $T > 0$, $\varepsilon \in \mathbb{R}$ be such that*

$$(6) \quad \varepsilon > -\frac{\pi L}{2} m(m+1)^2, \quad \text{where } m := \min \{ 1 ; (\pi/L)^2 \}.$$

and ψ_{ref} be defined by (5). There exists $\delta > 0$ and a C^1 -map

$$\Upsilon : \mathcal{V} \longrightarrow L^2((0, T) \times \partial\Omega, \mathbb{R}),$$

where

$$\mathcal{V} := \{ (\psi_0, \psi_f) \in [H_N^2(\Omega, \mathbb{C}) \cap \mathcal{S}]^2 ; \|\psi_0 - \psi_{\text{ref}}(0)\|_{H^2} + \|\psi_f - \psi_{\text{ref}}(T)\|_{H^2} < \delta \},$$

such that $\Upsilon(\psi_{\text{ref}}(0), \psi_{\text{ref}}(T)) = 0$ and for every $(\psi_0, \psi_f) \in \mathcal{V}$, the solution of (4) with control $g = \Upsilon(\psi_0, \psi_f)$ satisfies $\psi(T) = \psi_f$.

This part with the nonlinear Schrödinger equation only deals with Ω a rectangle, where the boundary is flat. Indeed, with the Neumann boundary control, the smoothing effect, required in our proof, is not well-understood for a general open set Ω . The nonhomogeneous boundary Cauchy problem is then quite complicated. This was for instance investigated for the wave equation by Tataru [35] and earlier papers by Lasiecka and Triggiani [21]. The curvature of the boundary has important consequences in this case.

1.4. BIBLIOGRAPHY

1.4.1. *Schrödinger equation with bilinear control.* — The Schrödinger equation with bilinear control has been widely studied in the literature, and is related to the system under study in this article. This model reads

$$(7) \quad \begin{cases} (i\partial_t + \Delta - V)\psi(t, x) = u(t)\mu(x)\psi(t, x), & (t, x) \in (0, T) \times \Omega, \\ \psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \end{cases}$$

where $V, \mu : \Omega \rightarrow \mathbb{R}$ are given functions, the state ψ lives in \mathcal{S} and the control is the real valued function $u : (0, T) \rightarrow \mathbb{R}$. When V and μ solve an appropriate Poisson equation,

then the system (7) is a particular case of the Schrödinger-Poisson system (1) (take $v(t, x) = V(x) + u(t)\mu(x)$).

A negative result. — A negative control result was proved by Turinici in [37], as a consequence of a general result by Ball, Marsden and Slemrod in [3]. It states that, for $V = 0$, for a given function $\mu \in C^2(\Omega, \mathbb{R})$, for a given initial condition $\psi_0 \in (H^2 \cap H_0^1)(\Omega, \mathbb{C}) \cap \mathcal{S}$, and by using controls $u \in L_{\text{loc}}^r((0, \infty), \mathbb{R})$ with $r > 1$, one may only reach a subset of $(H^2 \cap H_0^1)(\Omega) \cap \mathcal{S}$ that has an empty interior in $(H^2 \cap H_0^1)(\Omega, \mathbb{C}) \cap \mathcal{S}$. Recently, Boussaid, Caponigro and Chambrión extended this negative result to the case of controls in $L_{\text{loc}}^1((0, \infty), \mathbb{R})$, see [11]. However, this negative results are actually due to an inappropriate choice of functional setting, as emphasized in the next paragraph.

Note that this type of negative results is specific to systems with a bilinear control, which is a function of the time variable t (and not a function of both t and x). Thus, it does not apply to the Schrödinger-Poisson system (1) studied in this article. See also Section 5.3 for some comments about this type of control in our context.

Local exact results in 1D. — Beauchard proved in [4] the exact controllability of Equation (7), locally around the ground state in H^7 , with controls $u \in H^1((0, T), \mathbb{R})$ in large time T , in the case $N = 1$, $\Omega = (-1/2, 1/2)$, $\mu(x) = x$ and $V = 0$. The proof of [4] relies on Coron's return method and Nash-Moser's theorem.

The reference [5], by the authors of this paper, improves this result and establishes the exact controllability of Equation (7), locally around the ground state in H^3 , with controls $u \in L^2((0, T), \mathbb{R})$, in arbitrary time $T > 0$, and with generic functions μ when $N = 1$, $\Omega = (0, 1)$. This result, proved with $V = 0$ in [5], can be extended to an arbitrary potential V , as explained in [29]. The proof relies on a smoothing effect, that allows one to conclude with the inverse mapping theorem (instead of Nash-Moser's one).

Then, Morancey and Nersesyan developed this strategy to control a Schrödinger equation with a polarizability term [28] and a finite number of Schrödinger equations with one control [27, 29].

The goal of this article is to extend this type of strategy (previously applied in 1D) to the 2D Schrödinger-Poisson system (1). New difficulties appear at the two levels of the proof:

- in controlling the linearized system, because a new observability inequality is required,
- in proving the appropriate smoothing effect or the 2D-system (1).

Global approximate results. — Three strategies have been developed to study approximate controllability for Equation (7)

The first strategy is a variational argument introduced by Nersesyan in [30]. It proves the global controllability to the ground state, approximately in H^3 , with smooth controls $u \in C_c^\infty((0, T), \mathbb{R})$, in large time T , for generic functions (μ, V) , in arbitrary dimension N .

Note that this global approximate control result may be coupled to the previous local exact controllability results to provide global exact controllability, see [31] for Equation (7), [28] for a Schrödinger equation with a polarizability term, [29] for finite number of Schrödinger equations with the same control.

A second strategy consists in deducing approximate controllability in regular spaces (containing H^3) from exact controllability results in infinite time by Nersisyan and Nersisyan [32]

A third strategy, due to Chambrion, Mason, Sigalotti, and Boscain [16], relies on geometric techniques for the controllability of the Galerkin approximations. It proves (under appropriate assumptions on V and μ) the approximate controllability of (7) in L^2 , with piece-wise constant controls. The hypotheses of this result were refined by Boscain, Caponigro, Chambrion, and Sigalotti in [8]. The approximate controllability is proved in higher Sobolev norms in [11] for one equation, and in [9] for a finite number of equations with one control. For more details and more references about the geometric techniques, we refer the reader to the recent survey [10].

1.4.2. *Schrödinger-Poisson system.* — In [24], Méhats, Privat and Sigalotti prove the approximate controllability in L^2 for a Schrödinger-Poisson system, with mixed boundary conditions (of Dirichlet or Neumann type, depending on the place where we are on the boundary). This result holds for generic domains Ω and generic control supports Γ_c on the boundary. The proof relies on the general result of [16] and analyticity arguments to obtain genericity.

1.5. STRUCTURE OF THE ARTICLE. — This article is organized in 7 Sections.

Section 2 is a discussion about our assumptions **(H2)**, **(H3)** and **(H4)**, in particular when Ω is a rectangle domain or a disk.

Section 3 is dedicated to the well-posedness of the system (1), in appropriate spaces for our control problem. The starting point is a smoothing effect proved by Puel in [34], that needs to be recast in our context.

In Section 4, we prove local exact controllability of high frequencies, around any trajectory of the free system, i.e., Theorem 1.

Section 5 is devoted to the proof of the local exact controllability, around any eigenstate, i.e., Theorem 2.

The goal of Section 6 is to explain how the strategy of Section 5 can be used to get controllability for the Schrödinger equation with *real-valued* boundary controls (instead of complex valued ones in the literature), i.e., Theorem 3.

Finally, in Section 7, we prove the local exact controllability of the nonlinear Schrödinger-Poisson system (4) on a rectangle, i.e., Theorem 4.

1.6. NOTATION. — Implicitly functions take values in \mathbb{C} , otherwise we specify, for instance $L^2((0, T) \times \partial\Omega, \mathbb{R})$. The volume element on Ω is denoted dx and the surface element on $\partial\Omega$ is denoted $d\sigma(x)$. When $\varphi \in \mathcal{S}$ then $T_S\varphi$ denotes the tangent space to

the sphere \mathcal{S} at point φ

$$T_{\mathcal{S}}\varphi := \left\{ \xi \in L^2(\Omega, \mathbb{C}) ; \Re \left(\int_{\Omega} \varphi(x) \overline{\xi(x)} dx \right) = 0 \right\}.$$

2. COMMENTS ON OUR ASSUMPTIONS

The goal of this section is to prove that our assumptions **(H2)**, **(H3)** and **(H4)** hold for appropriate configurations (Ω, Γ_c) and to explain why **(H4)** is related to the control of low frequencies.

2.1. WEAK OBSERVABILITY OF SCHRÖDINGER EQUATION (H2). — The goal of this section is to recall known results about Assumption **(H2)**.

PROPOSITION 1. — *Property (H2) holds in the following cases.*

(1) $\Omega = (0, \pi) \times (0, L)$ for some $L > 0$, $\tilde{T} > 0$ and Γ_c is an open subset of $\partial\Omega$ that contains both a horizontal and a vertical segment with non-zero length (necessary and sufficient condition).

(2) Ω is smooth, does not have a contact of infinite order with its tangents and (Ω, Γ_c) satisfies the Geometric Control Condition.

(3) $\Omega = \{(x, y) ; x^2 + y^2 < 1\}$ and Γ_c is an arbitrary open set of the boundary.

Statement 1 is proved by Tenenbaum and Tucsnak in [36, Th. 1.4]. Precisely, they prove that the Schrödinger equation is observable on $(0, \tilde{T}) \times \Gamma_c$ iff Γ_c contains both a horizontal and a vertical segment with non-zero length. Thus weak observability also holds in this configuration. Moreover, the same counter-example as in [36, p. 967] proves that this condition is necessary: with $\Gamma_c := (a, b) \times \{0\}$, $0 \leq a < b \leq \pi$, $m \in \mathbb{N}^*$ and $\phi_T(x, y) := \sin(mx) \sin(\pi y/L)$, we get

$$Cm \leq \|\phi_T\|_{H_0^1(\Omega)} \quad \text{and} \quad \|\partial_\nu \phi\|_{L^2((T', T'') \times \Gamma_c)} + \|\phi_T\|_{H^{-1}(\Omega)} \leq C',$$

for some constants $C, C' > 0$ independent of m .

Statement 2 is proved by Lebeau in [22] when Ω is analytic. The result is actually also true if Ω is smooth (see for instance Burq-Zworski [15]). The assumptions of contact of finite order is made in order to ensure the uniqueness of the broken geodesic flow. It is certainly also true if Ω is only C^3 using the result by Burq [13] for the wave equation and transmutation or resolvent methods (see Miller [25, 26] for instance).

Statement 3 is proved by Anatharaman, Léautaud and Macià in [1].

2.2. SIMULTANEOUS VALIDITY OF (H2) AND (H3). — The goal of this section is to prove the claim written after Theorem 1.

PROPOSITION 2. — *We assume that*

- either Ω is a disk and Γ_c is a non-empty open subset of $\partial\Omega$,
- or $\Omega = (0, \pi) \times (0, L)$ for some $L > 0$ and Γ_c contains both a horizontal and a vertical segment.

Let $T > 0$, $\psi_0 \in H^3_{(0)}(\Omega, \mathbb{C}) \cap \mathcal{S}$ and $\psi_{\text{ref}}(t) := e^{i\Delta_D t} \psi_0$. Then Assumptions **(H2)** and **(H3)** are satisfied when Γ_c is replaced by an appropriate non-empty open subset Γ'_c of Γ_c . As a consequence, the conclusion of Theorem 1 holds.

Proof of Proposition 2. — Let us assume that Ω is a disk and Γ_c is a non-empty open subset of $\partial\Omega$. By unique continuation for the Schrödinger equation, we know that the continuous function $\partial_\nu \psi_{\text{ref}}$ does not identically vanish on $(0, T) \times \Gamma_c$. This is a consequence of the Holmgren theorem (see Theorem 8.6.5 of [20]) since $\partial\Omega$ is analytic here and any hypersurface $\{\Phi = 0\}$ with $\nabla_x \Phi \neq 0$ is non-characteristic for the Schrödinger operator. We can also get this directly as a consequence of the observability estimate of [1]. Thus, there exists $0 \leq T' < T'' \leq T$ and an open subset Γ'_c of Γ_c such that $|\partial_\nu \psi_{\text{ref}}(t, x)| \geq m > 0$ for every $(t, x) \in (T', T'') \times \Gamma'_c$. Let $\tilde{T} \in (0, T'' - T')$. By the previous proposition, the Schrödinger equation on Ω is weakly observable on $(0, \tilde{T}) \times \Gamma'_c$. Therefore, both **(H2)** and **(H3)** hold with Γ_c replaced by Γ'_c . Then, Theorem 1 provides controls supported in $(0, T) \times \Gamma'_c$ that are, a fortiori, also supported in $(0, T) \times \Gamma_c$.

Now, let us assume that $\Omega = (0, \pi) \times (0, L)$ for some $L > 0$ and Γ_c contains both a horizontal segment Γ_H and a vertical segment Γ_V . By unique continuation for the Schrödinger equation (by application of the Holmgren theorem for instance), we know that $\partial_\nu \psi_{\text{ref}}$ does not identically vanish on $(0, T) \times \Gamma_H$. Thus, there exists $0 \leq T'_H < T''_H \leq T$ and an open subset Γ'_H of Γ_H such that $|\partial_\nu \psi_{\text{ref}}(t, x)| \geq m_H > 0$ for every $(t, x) \in (T'_H, T''_H) \times \Gamma'_H$. By unique continuation for the Schrödinger equation, we know that $\partial_\nu \psi_{\text{ref}}$ does not identically vanish on $(T'_H, T''_H) \times \Gamma_V$. Thus, there exists $T'_H \leq T' < T'' \leq T''_H$ and an open subset Γ'_V of Γ_V such that $|\partial_\nu \psi_{\text{ref}}(t, x)| \geq m_V > 0$ for every $(t, x) \in (T', T'') \times \Gamma'_V$. Then $|\partial_\nu \psi_{\text{ref}}(t, x)| \geq m := \min\{m_H; m_V\} > 0$ for every $(t, x) \in (T', T'') \times \Gamma'_c$ with $\Gamma'_c := \Gamma'_H \cup \Gamma'_V$. The conclusion comes as in the previous case. \square

2.3. UNIQUE CONTINUATION ASSUMPTION (H4). — The goal of this section is to prove that **(H4)** holds on rectangular domains and is related to the controllability of low frequencies.

2.3.1. The case of a rectangle

PROPOSITION 3. — Let $L > 0$, $\Omega := (0, \pi) \times (0, L)$, Γ_c be a non-empty open subset of $\partial\Omega$, $R_1, R_2 \in \mathbb{N}^*$ be such that $R_1^2 + (R_2\pi/L)^2$ is a simple eigenvalue of $(-\Delta_D)$ and $\varphi_R(x, y) := \frac{2}{\sqrt{\pi L}} \sin(R_1 x) \sin(R_2 \pi y/L)$. Then **(H4)** is satisfied.

Proof of Proposition 3. — Without loss of generality, one may assume that Γ_c contains $(a, b) \times \{0\}$ for some $0 \leq a < b \leq \pi$. Let $\lambda \in \text{Sp}(-\Delta_D)$, set

$$J := \{(p, n) \in (\mathbb{N}^*)^2; p^2 + (n\pi/L)^2 = \lambda\},$$

which is a finite set, and let

$$\Phi = \sum_{(p,n) \in J} c_{p,n} \sin(px) \sin(n\pi y/L)$$

be an eigenfunction of $(-\Delta_D)$ associated to the eigenvalue λ . From the relation $2 \sin(a) \sin(b) = \cos(a - b) - \cos(a + b)$ we deduce that

$$\varphi_R \Phi = \sum_{(p,n) \in J} \sum_{\substack{\varepsilon_1 = \pm 1, \\ \varepsilon_2 = \pm 1}} \frac{c_{p,n} \varepsilon_1 \varepsilon_2}{2\sqrt{\pi L}} \cos[(p + \varepsilon_1 R_1)x] \cos[(n + \varepsilon_2 R_2)\pi y/L]$$

thus

$$w(x, y) = \sum_{(p,n) \in J} \sum_{\substack{\varepsilon_1 = \pm 1, \\ \varepsilon_2 = \pm 1}} \frac{c_{p,n} \varepsilon_1 \varepsilon_2 \cos[(p + \varepsilon_1 R_1)x] \cos[(n + \varepsilon_2 R_2)\pi y/L]}{2\sqrt{\pi L} \left[(p + \varepsilon_1 R_1)^2 + ((n + \varepsilon_2 R_2)\pi/L)^2 + 1 \right]}$$

and in particular

$$w(x, 0) = \sum_{(p,n) \in J} \sum_{\substack{\varepsilon_1 = \pm 1, \\ \varepsilon_2 = \pm 1}} \frac{c_{p,n} \varepsilon_1 \varepsilon_2 \cos[(p + \varepsilon_1 R_1)x]}{2\sqrt{\pi L} \left[(p + \varepsilon_1 R_1)^2 + ((n + \varepsilon_2 R_2)\pi/L)^2 + 1 \right]}.$$

Let

$$P := \max\{p \in \mathbb{N}^* ; \exists n \in \mathbb{N}^* \text{ such that } (p, n) \in J \text{ and } c_{p,n} \neq 0\}.$$

Let $N \in \mathbb{N}^*$ be the unique integer such that $(P, N) \in J$. Then $x \mapsto w(x, 0)$ is a finite sum of cosine functions

$$w(x, 0) = \sum_{k=0}^{P+R_1} \alpha_k \cos(kx)$$

and its fastest oscillating term has coefficient

$$\alpha_{P+R_1} = \frac{c_{P,N}}{2\sqrt{\pi L}} \left(\frac{1}{\left[(P + R_1)^2 + ((N + R_2)\pi/L)^2 + 1 \right]} - \frac{1}{\left[(P + R_1)^2 + ((N - R_2)\pi/L)^2 + 1 \right]} \right).$$

If $w(x, 0) = 0$ for every $x \in (a, b)$, then $\alpha_k = 0$ for $k = 0, \dots, P + R_1$ and in particular $\alpha_{P+R_1} = 0$. This is impossible because $c_{P,N} \neq 0$ and $(N + R_2)^2 \neq (N - R_2)^2$ (note that $N, R_2 \in \mathbb{N}^*$). \square

2.3.2. General comments. — First, we prove the equivalence between Assumption **(H4)** and the existence of some non-vanishing integral quantities. To this end, we introduce the following notation: for $\chi \in L^2(\partial\Omega)$, V_χ denotes the unique solution in $H^{3/2}(\Omega)$ of

$$(8) \quad \begin{cases} (-\Delta + 1)V_\chi(x) = 0, & x \in \Omega, \\ \partial_\nu V_\chi(x) = \chi(x), & x \in \partial\Omega. \end{cases}$$

PROPOSITION 4. — *Let Ω be a smooth bounded open subset of \mathbb{R}^2 and $(\varphi_k)_{k \in \mathbb{N}}$ be an orthogonal basis of eigenfunctions of $(-\Delta_D)$, $R \in \mathbb{N}^*$. If the spectrum of $(-\Delta_D)$ is simple, then the following statements are equivalent.*

(1) For any eigenvector Φ of $(-\Delta_D)$, the solution w of

$$(9) \quad \begin{cases} (-\Delta + 1)w(x) = \varphi_R(x)\Phi(x), & x \in \Omega, \\ \partial_\nu w(x) = 0, & x \in \partial\Omega, \end{cases}$$

does not identically vanish on Γ_c : $w|_{\Gamma_c} \not\equiv 0$.

(2) There is an open dense set in $L^2(\Gamma_c)$ of functions χ such that $\int_\Omega V_\chi \varphi_R \varphi_k \neq 0$ for any $k \in \mathbb{N}^*$

(3) There is one $\chi \in L^2(\Gamma_c)$ such that $\int_\Omega V_\chi \varphi_R \varphi_k \neq 0$ for any $k \in \mathbb{N}^*$.

If the spectrum of $(-\Delta_D)$ is not simple, we still have (1) \Rightarrow (2) \Rightarrow (3).

Proof. — We have obviously (2) \Rightarrow (3). The main tool for the other implications will be the following formula using Equation (9) (the same as in **(H4)**) satisfied by w

$$\begin{aligned} \int_\Omega V_\chi \varphi_R \Phi &= \int_\Omega V_\chi (-\Delta + 1)w \\ &= \int_\Omega (-\Delta + 1)V_\chi w + \int_{\partial\Omega} [\partial_\nu V_\chi w - V_\chi \partial_\nu w] \\ &= \int_{\partial\Omega} \chi w. \end{aligned}$$

In the case of simple eigenvalues, we easily get (3) \Rightarrow (1) by selecting $\chi \in L^2(\Gamma_c)$ such that $\int_\Omega V_\chi \varphi_R \varphi_k = \int_{\partial\Omega} \chi w_k \neq 0$. So, the main task is to prove (1) \Rightarrow (2). For $k \in \mathbb{N}^*$, we introduce the set

$$\mathcal{G}_k := \left\{ \chi \in L^2(\Gamma_c, \mathbb{R}); \int_\Omega V_\chi \varphi_R(x) \varphi_k(x) dx \neq 0 \right\}.$$

By Assumption (1), for any k , w_k is not identically zero on Γ_c , so there exists χ such that $\int_{\partial\Omega} \chi w_k = \int_\Omega V_\chi \varphi_R \varphi_k \neq 0$. In particular, \mathcal{G}_k is not empty for every $k \in \mathbb{N}^*$. Moreover, if $\chi_0 \in L^2(\Gamma_c, \mathbb{R}) \setminus \mathcal{G}_k$ and $\chi \in \mathcal{G}_k$ then $\chi_0 + \varepsilon\chi$ belongs to \mathcal{G}_k for every $\varepsilon \in \mathbb{R}^*$ and converges to χ_0 when $\varepsilon \rightarrow 0$. Thus \mathcal{G}_k is a dense open subset of $L^2(\Gamma_c, \mathbb{R})$.

By Baire’s Lemma $\bigcap_{k \in \mathbb{N}^*} \mathcal{G}_k$ is a dense subset of $L^2(\Gamma_c, \mathbb{R})$. This proves (2). \square

With the previous proposition in hands, it should be possible to prove that **(H4)** is generic with respect to perturbations of the domain, as done in [24]. Indeed, the previous proposition proves that **(H4)** is equivalent to the existence of one χ so that some integral quantities are not zero. It is then often a good starting point to prove genericity with respect to χ and deformation of Ω . This would then directly imply that **(H4)** is generic with respect to perturbations of the domain.

Note also that with the same proof, χ can be chosen in $C_0^k(\Gamma_c)$ for some $k \in \mathbb{N}$.

2.3.3. *Control of low frequencies.* — Actually, the condition **(H4)** is a condition of controllability of the low frequencies for a control only depending on time. More precisely, we have the following result.

PROPOSITION 5. — Assume all the eigenvalues of Δ_D are simple and **(H4)** is fulfilled for one $R \in \mathbb{N}^*$. Let $\psi_{\text{ref}}(t) := \varphi_R(x)e^{-i\lambda_R t}$. Then there is one $\chi \in C^4(\Gamma_c)$ such

that $\int_{\Omega} V_{\chi} \varphi_R \Phi_k \neq 0$ for any $k \in \mathbb{N}^*$. This implies the following local controllability property for low frequencies.

For any $N \geq 0$, there exists $\delta > 0$ such that for every

$$\psi_f \in \mathcal{V} := \{\psi_f \in H_{(0)}^3(\Omega) \cap \mathcal{S}; \|\psi_f - \psi_{\text{ref}}(T)\|_{H_{(0)}^3} < \delta\},$$

there exists $u \in L^2((0, T), \mathbb{R})$ such that the solution of

$$(10) \quad \begin{cases} (i\partial_t + \Delta)\psi(t, x) = u(t)V_{\chi}(x)\psi(t, x), & (t, x) \in (0, T) \times \Omega, \\ \psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \psi(0, x) = \varphi_R(x), & x \in \Omega, \end{cases}$$

satisfies $\mathbb{P}_N^{\perp}\psi(T) = \mathbb{P}_N^{\perp}\psi_f$, where \mathbb{P}_N^{\perp} is the orthogonal projection on the N first eigenfunctions.

Proof of Proposition 5. — The map

$$\begin{aligned} \Theta : L^2((0, T), \mathbb{R}) &\longrightarrow \mathbb{P}_N^{\perp}[\mathcal{S}] \\ u &\longmapsto \mathbb{P}_N^{\perp}[\psi(T)] \end{aligned}$$

is of class C^1 and

$$\begin{aligned} d\Theta(0) : L^2((0, T), \mathbb{R}) &\longrightarrow \mathbb{P}_N^{\perp}[T_{\psi_{\text{ref}}(T)}\mathcal{S}] \\ u &\longmapsto \mathbb{P}_N^{\perp}[\Psi(T)], \end{aligned}$$

where

$$\begin{aligned} \Psi(T) &= -i \int_0^T e^{i(T-t)\Delta_D} (u(t)V_{\chi}\psi_{\text{ref}}(t)) dt \\ &= -i \sum_{k=1}^{\infty} \langle V_{\chi}\varphi_R, \varphi_k \rangle \left(\int_0^T u(t)e^{i(\lambda_k - \lambda_R)t} dt \right) e^{-i\lambda_k T} \varphi_k. \end{aligned}$$

The surjectivity of $d\Theta(0)$ can be formulated in terms of a finite trigonometric moment problem on u . Since $\langle V_{\chi}\varphi_R, \varphi_k \rangle \neq 0$ for every k and the frequencies $(\lambda_k - \lambda_R)$ are all different, this moment problem has a solution $u \in L^2((0, T), \mathbb{R})$. We can therefore apply the inverse function theorem to Θ . We skip the details since it is a simpler version of arguments used several times in the paper. \square

3. WELL-POSEDNESS OF THE SCHRÖDINGER-POISSON SYSTEM AND LINEARIZATION

The goal of this section is to prove the well-posedness of the system (1) in functional spaces that are appropriate for the controllability problem. This proof requires several preliminary results. The first one concerns Sobolev embeddings, trace theorems and elliptic estimates; they are recalled in Section 3.1. The second preliminary result concerns smoothing effects of the Schrödinger equation on a bounded domain: a smoothing effect concerning the normal derivative is recalled in Section 3.2, a smoothing effect concerning the source term is justified in Section 3.3. Finally, in Section 3.4, we prove the well-posedness of the system (1) in $H_{(0)}^3(\Omega)$ when the control g lives

in $L^2((0, T) \times \Gamma_c)$; the smoothing effects are crucial at this point. In Section 3.5, we prove the C^1 -regularity of the end-point map.

3.1. PRELIMINARY

3.1.1. *Sobolev embeddings.* — In the whole section Ω is an open subset of \mathbb{R}^n such that **(H1)** is satisfied. For $s \geq 0$ and $p \geq 1$, we use the definitions

$$\begin{aligned} W^{s,p}(\mathbb{R}^2) &:= \{ \tilde{f} \in \mathcal{S}'(\mathbb{R}^2); \mathcal{F}^{-1}[(1 + |\cdot|^s)\mathcal{F}\tilde{f}] \in L^p(\mathbb{R}^2) \}, \\ \|\tilde{f}\|_{W^{s,p}(\mathbb{R}^2)} &:= \|\mathcal{F}^{-1}[(1 + |\cdot|^s)\mathcal{F}\tilde{f}]\|_{L^p(\mathbb{R}^2)}, \\ W^{s,p}(\Omega) &:= \{ \tilde{f}|_\Omega; \tilde{f} \in W^{s,p}(\mathbb{R}^2) \}, \\ \|f\|_{W^{s,p}(\Omega)} &:= \inf \{ \|\tilde{f}\|_{W^{s,p}(\mathbb{R}^2)}; \tilde{f} \in W^{s,p}(\mathbb{R}^2), f = \tilde{f} \text{ on } \Omega \}. \end{aligned}$$

If $p = 2$, we use the notation H^s instead of $W^{s,2}$.

Remark that if $s = 1$, we also have

$$W^{1,p}(\Omega) := \{ f \in L^p(\Omega); \nabla f \in L^p(\Omega)^n \}$$

with the obvious norm equivalent to the one previously introduced.

The standard Sobolev embeddings

$$(11) \quad \begin{aligned} H^1(\Omega) &\subset L^q(\Omega), \quad \forall q \in [2, \infty), \\ W^{1,p}(\Omega) &\subset L^\infty(\Omega), \quad \forall p > 2 \end{aligned}$$

(see [12, Cor. IX.14]) and

$$W^{s,p}(\mathbb{R}^2) \subset W^{s_1,p_1}(\mathbb{R}^2) \quad \text{for } s - \frac{2}{p} = s_1 - \frac{2}{p_1}, \quad 1 < p \leq p_1 < +\infty,$$

(see [7, Th. 6.5.1]) yield to

$$(12) \quad H^{3/2}(\Omega) \subset L^\infty(\Omega),$$

$$(13) \quad H^{1/2}(\Omega) \subset L^4(\Omega),$$

$$(14) \quad \forall \varepsilon \in (0, 1/2), \exists p \in (1, 2) \text{ such that } W^{1,p}(\Omega) \subset H^{1/2+\varepsilon}(\Omega).$$

If $s \geq 0$, we denote

$$H^{-s}(\Omega) := (H_0^s(\Omega))',$$

which is a space of distributions.

3.1.2. *Regularity for elliptic boundary value problems on a smooth domain*

In this section, we recall classical results used in the article (see [23, Th. 5.4 p. 176, Th. 6.5, Th. 6.7 p. 192]).

PROPOSITION 6. — *Let Ω be a bounded open subset of \mathbb{R}^2 , of class C^∞ and locally on one side of $\partial\Omega$.*

(1) *The mapping*

$$\begin{aligned} C^\infty(\overline{\Omega}) &\longrightarrow C^\infty(\partial\Omega) \\ f &\longmapsto f|_{\partial\Omega} \end{aligned}$$

has a unique continuous extension from $D(\Delta, L^2(\Omega)) := \{f \in L^2(\Omega); \Delta f \in L^2(\Omega)\}$ into $H^{-1/2}(\partial\Omega)$.

(2) The mapping

$$\begin{aligned} H^2(\Omega) &\longrightarrow L^2(\partial\Omega) \\ f &\longmapsto \partial_\nu f \end{aligned}$$

has a unique continuous extension from $\{f \in H^{3/2}(\Omega); \Delta f \in L^2(\Omega)\}$ into $L^2(\partial\Omega)$.

(3) For every $f \in L^2(\Omega)$ and $g \in L^2(\partial\Omega)$, there exists a unique $v \in H^{3/2}(\Omega)$ such that

$$\begin{cases} (-\Delta + 1)v = f \text{ in } \Omega, \\ \partial_\nu v = g \text{ on } \partial\Omega. \end{cases}$$

Moreover, there exists $C(\Omega) > 0$ such that

$$\|v\|_{H^{3/2}(\Omega)} \leq C[\|f\|_{L^2(\Omega)} + \|g\|_{L^2(\partial\Omega)}].$$

3.1.3. Regularity and Green formula for elliptic boundary value problems on a rectangle

The goal of this section is to state the following results.

PROPOSITION 7. — Let $\Omega = (0, \pi) \times (0, L)$ for some $L > 0$, $(\Gamma_j)_{1 \leq j \leq 4}$ be its edges, $(\nu_j)_{1 \leq j \leq 4}$ be its unitary exterior normal vectors, $(S_j)_{1 \leq j \leq 4}$ be its vertices and $p \geq 1$.

(1) The mapping $u \mapsto (u|_{\Gamma_j}, \partial_{\nu_j} u)$, which is well-defined for $u \in W^{2,p}(\Omega)$ has a unique extension as an operator from

$$D(\Delta, L^p(\Omega)) := \{u \in L^p(\Omega); \Delta u \in L^p(\Omega)\}$$

into

- $W^{-1/p,p}(\Gamma_j) \times W^{-1-1/p,p}(\Gamma_j)$ when $p \neq 2$,
- $H^{-1/2-\varepsilon}(\Gamma_j) \times H^{-3/2-\varepsilon}(\Gamma_j)$, for every $\varepsilon > 0$, when $p = 2$.

(2) Moreover,

$$\int_{\Omega} ((\Delta u)v - u(\Delta v)) \, dx = \sum_{j=1}^4 \int_{\Gamma_j} ((\partial_{\nu_j} u)v - u(\partial_{\nu_j} v)) \, d\sigma_j(x)$$

for every $u \in D(\Delta, L^p(\Omega))$ and $v \in W^{2,p'}(\Omega)$ such that

- $v(S_j) = 0$ for $j = 1, \dots, 4$ when $p > 2$,
- $v(S_j) = 0$ and $\nabla v(S_j) = 0$ for $j = 1, \dots, 4$ when $p < 2$,
- $v \equiv 0$ on a neighborhood of S_j for $j = 1, \dots, 4$ when $p = 2$.

(3) For every $f \in L^2(\Omega)$ and $g \in L^2(\partial\Omega)$, there exists a unique $v \in H^{3/2}(\Omega)$ such that

$$(15) \quad \begin{cases} (-\Delta + 1)v = f \text{ in } \Omega, \\ \partial_\nu v = g \text{ on } \partial\Omega. \end{cases}$$

Moreover, there exists $C(\Omega) > 0$ such that

$$\|v\|_{H^{3/2}(\Omega)} \leq C[\|f\|_{L^2(\Omega)} + \|g\|_{L^2(\partial\Omega)}].$$

For the first two statements, see [18, Th. 1.5.3.4 & Th. 1.5.3.6]. The third one is a consequence of the same one on $(\mathbb{R}/2\pi\mathbb{Z}) \times (0, L)$ after symmetrisation and periodization. More precisely, by linearity, one may assume that $\text{Supp}(g) \subset [0, \pi] \times \{0\}$. Then we extend g as a function $\tilde{g} : (\mathbb{R}/\pi\mathbb{Z}) \times \{0\} \rightarrow \mathbb{R}$ such that $\tilde{g}(-x_1, 0) = g(x_1, 0)$ for every $x_1 \in (0, \pi)$ and $\tilde{g}(x_1, 0) = \tilde{g}(x_1 + 2\pi, 0)$. Perform similar symmetrisation and periodization for f to produce \tilde{g} well-defined on $(\mathbb{R}/2\pi\mathbb{Z}) \times (0, L)$.

Then, since $(\mathbb{R}/2\pi\mathbb{Z}) \times (0, L)$ is a smooth compact manifold with boundary, there exists a unique $\tilde{v} \in H^{3/2}((\mathbb{R}/2\pi\mathbb{Z}) \times (0, L))$ such that

$$\begin{cases} (-\Delta + 1)\tilde{v} = \tilde{f} \text{ in } \Omega, \\ \partial_\nu \tilde{v} = \tilde{g} \text{ on } (\mathbb{R}/\pi\mathbb{Z}) \times \{0, 1\}. \end{cases}$$

We easily check that the symmetry gives that if $f \in C_0^\infty(\Omega)$ and $g \in C_0^\infty(0, \pi)$, then, $\partial_{x_1} \tilde{v}(x_1, x_2) = 0$ for $x_1 \in \{0, \pi\}$ and $x_2 \in (0, L)$. Therefore, $v = \tilde{v}|_\Omega$ satisfies (15) in the weak sense. The uniqueness can be obtained by following the same process.

3.2. SMOOTHING EFFECT ON THE NORMAL DERIVATIVE. — The goal of this section is to state the following results.

PROPOSITION 8. — *Let $T > 0$ and Ω be a bounded open subset of \mathbb{R}^2 which is either C^∞ or a rectangle. There exists $C = C(T, \Omega) > 0$ such that, for every $\psi_0 \in H_0^1(\Omega)$ and $h \in L^1((0, T), H_0^1(\Omega))$, the solution ψ of*

$$(16) \quad \begin{cases} (i\partial_t + \Delta)\psi(t, x) = h(t, x), & (t, x) \in (0, T) \times \Omega, \\ \psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \psi(0, x) = \psi_0(x), & x \in \Omega, \end{cases}$$

satisfies

$$(17) \quad \int_0^T \int_{\partial\Omega} |\partial_\nu \psi(t, x)|^2 |\partial_\nu \varphi_1(x)| d\sigma(x) dt \leq C(\|\psi_0\|_{H_0^1(\Omega)} + \|h\|_{L^1((0, T), H_0^1(\Omega))}).$$

In particular, if Γ_c is an open subset of $\partial\Omega$ such that **(H1)** holds, then the following linear mapping is continuous

$$\begin{aligned} H_0^1(\Omega) \times L^1((0, T), H_0^1(\Omega)) &\longrightarrow L^2((0, T) \times \Gamma_c) \\ (\psi_0, \varphi) &\longmapsto \partial_\nu \psi. \end{aligned}$$

When Ω is smooth, Puel proves these results in [34, Lem. 3.1]. Precisely, he first proves the inequality (17) for smooth data $(\psi_0, h) \in C_c^\infty(\Omega) \times C_c^\infty((0, T) \times \Omega)$, by applying the multiplier $\nabla \varphi_1$ to Equation (16): under these assumptions ψ is regular, $\partial\psi/\partial\nu$ makes perfect sense and integrations by parts are legitimate. Then, the inequality (17) holds for $(\psi_0, h) \in H_0^1(\Omega) \times L^1((0, T), H_0^1(\Omega))$ by a density argument. Finally, Puel gets the final result because $|\partial_\nu \varphi_1(x)| \geq \beta > 0, \forall x \in \partial\Omega$

When Ω is a rectangle, Puel’s proof of the inequality (17) is still valid. Then, the conclusion of Proposition 8 follows because $|\partial_\nu \varphi_1(x)| \geq \beta > 0, \forall x \in \Gamma_c$; this is one of the reasons why we need to assume that Γ_c does not touch the vertices of Ω .

3.3. SMOOTHING EFFECT ON THE SOURCE TERM. — The goal of this section is to justify the following result.

THEOREM 5. — *Let $T > 0$, Ω be an open subset of \mathbb{R}^2 and Γ_c be an open subset of $\partial\Omega$ such that **(H1)** holds. For every $\psi_0 \in H_{(0)}^3(\Omega)$, $\mu_1 \in L^2((0, T), H^2 \cap H_0^1(\Omega))$ and $\mu_2 \in L^1((0, T), H_{(0)}^3(\Omega))$ such that*

$$(18) \quad \Delta^2 \mu_1 \equiv 0, \quad \text{Supp}(\Delta \mu_1|_{\partial\Omega}) \subset [0, T] \times \Gamma_c, \quad \Delta \mu_1|_{\Gamma_c} \in L^2((0, T) \times \Gamma_c),$$

the solution of

$$\begin{cases} (i\partial_t + \Delta)\psi(t, x) = (\mu_1 + \mu_2)(t, x), & (t, x) \in (0, T) \times \Omega, \\ \psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \psi(0, x) = \psi_0(x), & x \in \Omega, \end{cases}$$

satisfies $\psi \in C^0([0, T], H_{(0)}^3(\Omega))$. Moreover, there exists a constant $C > 0$ (independent of ψ_0, μ_1, μ_2) such that

$$(19) \quad \|\psi\|_{L^\infty([0, T], H_{(0)}^3(\Omega))} \leq C(\|\psi_0\|_{H_{(0)}^3(\Omega)} + \|\Delta \mu_1|_{\Gamma_c}\|_{L^2((0, T) \times \Gamma_c)} + \|\mu_2\|_{L^1((0, T), H_{(0)}^3(\Omega))}).$$

REMARK 1. — When Ω is smooth, the trace $\Delta \mu_1|_{\partial\Omega}$ is well-defined in the space $L^2((0, T), H^{-1/2}(\partial\Omega))$ (see Proposition 6), which gives a sense to the last two requirements in (18).

When Ω is a rectangle, then the traces along the four sides $\Delta \mu_1|_{\Gamma_j}$, $1 \leq j \leq 4$, are well-defined in $L^2((0, T), H^{-1/2-\varepsilon}(\Gamma_j))$ (see Proposition 7). The last two requirements in (18) have to be interpreted in the following sense:

$$\begin{aligned} \Gamma_j \cap \text{Supp}(\Delta \mu_1|_{\partial\Omega}) &\subset \Gamma_j \cap \Gamma_c, \quad \forall j \in \{1, \dots, 4\}, \\ \Delta \mu_1|_{\Gamma_c \cap \Gamma_j} &\in L^2((0, T) \times \Gamma_c \cap \Gamma_j), \quad \forall j \in \{1, \dots, 4\}. \end{aligned}$$

Theorem 5 emphasizes a regularizing effect, concerning the source term, because μ_1 is not assumed to belong to $L^1((0, T), H_{(0)}^3(\Omega))$. When Ω is a regular domain and $\Gamma_c = \partial\Omega$, Puel proves this result in [34, Th. 2.1]. His proof, by transposition, relies on the smoothing effect of Proposition 8.

Puel does not treat the case of a rectangle domain, but his proof would probably lead to Theorem 5 in this case too. For the sake of completeness, we propose below an alternative argument.

Proof of Theorem 5 on a rectangle. — Let $L > 0$, $\Omega := (0, \pi) \times (0, L)$, Γ_c be an open subset of $\partial\Omega$, $\psi_0 \in H_{(0)}^3(\Omega)$, $\mu_1 \in L^2((0, T), H^2 \cap H_0^1(\Omega))$ and $\mu_2 \in L^1((0, T), H_{(0)}^3(\Omega))$ be such that (18) holds. By linearity, we may assume that $\Gamma_c \subset (a, b) \times \{0\}$ with $0 < a < b < \pi$. Let $m \in \mathbb{N}^*$ be such that $(m - 1)2L^2/\pi < T \leq T_0 := 2mL^2/\pi$ and $t \in (0, T)$.

By the Duhamel formula, we have

$$\psi(t) = e^{i\Delta_D t} \psi_0 - i \int_0^t e^{i\Delta_D(t-\tau)} (\mu_1(\tau) + \mu_2(\tau)) d\tau.$$

The first and third terms in the right-hand side belong to $C^0([0, T], H^3_{(0)}(\Omega))$ because $H^3_{(0)}(\Omega)$ is stable by $e^{i\Delta_D t}$. Thus, we focus on the second one. We will make the proof in two steps: the first one when μ_1 is regular enough to perform integration by parts and the second one to get to the first case with a suitable regularization.

Step 1: μ_1 regular enough. — We assume moreover that $\Delta\mu_1 \in L^2((0, T), L^p(\Omega))$, with $p > 2$, in this step.

Using the equality $-\Delta\varphi_k = \lambda_k\varphi_k$ and integrations by part, that are licit because $\mu_1 \in L^2((0, T), H^2 \cap H^1_0(\Omega))$, we get

$$\begin{aligned} S &= \left\| \int_0^t e^{i\Delta(t-\tau)} \mu_1(\tau) d\tau \right\|_{H^3_{(0)}(\Omega)}^2 = \sum_{k=1}^{\infty} \left| \lambda_k^{3/2} \int_0^t \int_{\Omega} \mu_1(\tau, x) \varphi_k(x) dx e^{i\lambda_k \tau} d\tau \right|^2 \\ &= \sum_{k=1}^{\infty} \left| \frac{1}{\sqrt{\lambda_k}} \int_0^t \int_{\Omega} \Delta\mu_1(\tau, x) \Delta\varphi_k(x) dx e^{i\lambda_k \tau} d\tau \right|^2. \end{aligned}$$

We can apply Green’s formula (see Proposition 7) because $\Delta\mu_1(\tau, \cdot) \in D(\Delta, L^p(\Omega))$ for almost every $\tau \in (0, T)$ and $\varphi_k \in H^2(\Omega)$ vanishes on the vertices of Ω . Using $\Delta^2\mu_1 \equiv 0$, this leads to

$$\begin{aligned} S &= \sum_{k=1}^{\infty} \left| \frac{1}{\sqrt{\lambda_k}} \int_0^t \int_{\partial\Omega} \Delta\mu_1(\tau, x) \partial_{\nu} \varphi_k(x) d\sigma(x) e^{i\lambda_k \tau} d\tau \right|^2 \\ &= \sum_{k=1}^{\infty} \left| \frac{1}{\sqrt{\lambda_k}} \int_0^t \int_{\Gamma_c} \Delta\mu_1(\tau, x) \partial_y \varphi_k(x) d\sigma(x) e^{i\lambda_k \tau} d\tau \right|^2 \\ &\leq C \sum_{p, n \in \mathbb{N}^*} \left| \frac{n\pi/L}{\sqrt{p^2 + (n\pi/L)^2}} \int_0^t \int_0^{\pi} \Delta\mu_1(\tau, x_1, 0) \sin(px_1) e^{-i(p^2 + (n\pi/L)^2)\tau} dx_1 d\tau \right|^2 \\ &\leq C \sum_{p \in \mathbb{N}^*} \sum_{n \in \mathbb{N}^*} \left| \int_0^{T_0} 1_{[0, t]}(\tau) \int_0^{\pi} \Delta\mu_1(\tau, x_1, 0) \sin(px_1) dx_1 e^{-i(p^2 + (n\pi/L)^2)\tau} d\tau \right|^2 \\ &\leq C \sum_{p \in \mathbb{N}^*} \int_0^{T_0} \left| 1_{[0, t]}(\tau) \int_0^{\pi} \Delta\mu_1(\tau, x_1, 0) \sin(px_1) dx_1 e^{-ip^2\tau} \right|^2 d\tau \\ &\leq C \sum_{p \in \mathbb{N}^*} \int_0^t \left| \int_0^{\pi} \Delta\mu_1(\tau, x_1, 0) \sin(px_1) dx_1 \right|^2 d\tau \\ &\leq C \int_0^t \|\Delta\mu_1(\tau, \cdot, 0)\|_{L^2(0, \pi)}^2 d\tau. \end{aligned}$$

To go from the first line to the second one, we have used: $\text{Supp}(\Delta\mu_1|_{\partial\Omega}) \subset [0, T] \times \Gamma_c$, $\varphi_k \equiv 0$ on Γ_c . To go from the fourth line to the fifth line, we have used Bessel-Parseval’s inequality.

Note that $\Delta\mu_1$ can be computed explicitly in terms of $\Delta\mu_1|_{\Gamma_c}$, thanks to the rectangular form of the domain. This explicit expression shows the existence of a constant $C = C(\Omega) > 0$ such that

$$\|\Delta\mu_1\|_{L^2((0,t)\times\Omega)} \leq C\|\Delta\mu_1|_{\Gamma_c}\|_{L^2((0,t)\times\Gamma_c)}^2, \quad \forall t \in [0, T].$$

Thus, we get

$$\left\| \int_0^t e^{i\Delta(t-\tau)}\mu_1(\tau)d\tau \right\|_{H^3_{(0)}(\Omega)}^2 \leq C\|\Delta\mu_1|_{\Gamma_c}\|_{L^2((0,t)\times\Gamma_c)}^2.$$

This proves that the map $t \in [0, T] \mapsto \int_0^t e^{i\Delta(t-\tau)}\mu_1(\tau)d\tau$ takes values in $H^3_{(0)}(\Omega)$ and is continuous at $t = 0$. The same argument proves the continuity at any $t \in [0, T]$.

Step 2: regularization. — We have to be a bit careful with the regularization to keep the required conditions. Let $f_\varepsilon \in C^\infty_0((0, T) \times \Gamma_c)$ converging strongly to $\Delta\mu_1|_{\Gamma_c}$ in $L^2((0, T) \times \Gamma_c)$. Denote

$$\begin{cases} \Delta g_\varepsilon(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ g_\varepsilon(t, x) = f_\varepsilon(t, x), & (t, x) \in (0, T) \times \partial\Omega, \end{cases}$$

By elliptic regularity (similar to Proposition 7), g_ε converges strongly to $\Delta\mu_1$ in $L^2((0, T), H^{1/2}(\Omega))$.

Now, denote $\mu_{1,\varepsilon}$ the solution of

$$\begin{cases} \Delta\mu_{1,\varepsilon}(t, x) = g_\varepsilon(t, x), & (t, x) \in (0, T) \times \Omega, \\ \mu_{1,\varepsilon}(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega. \end{cases}$$

By elliptic regularity, $\mu_{1,\varepsilon}$ converges to μ_1 in $L^2((0, T), H^2 \cap H^1_0(\Omega))$. The estimate (19) (that is proven for $\mu_{1,\varepsilon}$ regular enough thanks to Step 1) allows us to prove that the related ψ_ε make a Cauchy sequence in $L^\infty([0, T], H^3_{(0)}(\Omega))$. The limit can only be the expected ψ by uniqueness of the solution and convergence of $\mu_{1,\varepsilon}$ to μ_1 . The estimate (19) follows in this case. \square

3.4. WELL-POSEDNESS OF THE LINEAR SCHRÖDINGER-POISSON SYSTEM. — The goal of this section is the proof of the following result, thanks to Theorem 5.

THEOREM 6. — *Let $T > 0$, Ω be an open subset of \mathbb{R}^2 and Γ_c be an open subset of $\partial\Omega$ such that **(H1)** holds. For every $\psi_0 \in H^3_{(0)}(\Omega)$ and $g \in L^2((0, T) \times \partial\Omega, \mathbb{R})$, there exists a unique solution $\psi \in C^0([0, T], H^3_{(0)}(\Omega))$ of the system (1).*

To this end, we will need the following preliminary result.

PROPOSITION 9. — *Let $T > 0$ and $\psi \in C^0([0, T], H^3_{(0)}(\Omega))$. For every function $g \in L^2((0, T) \times \partial\Omega, \mathbb{R})$, the solution of*

$$(20) \quad \begin{cases} (-\Delta + 1)v(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu v(t, x) = g(t, x)1_{\Gamma_c}(x), & (t, x) \in (0, T) \times \partial\Omega, \end{cases}$$

satisfies $v \in L^2((0, T), H^{3/2}(\Omega))$, $v\psi \in L^2((0, T), H^2 \cap H_0^1(\Omega))$ and $\Delta^2[v\psi]$ belongs to $L^2((0, T), H^{-1}(\Omega))$. Moreover, the operator

$$\begin{aligned} L^2((0, T) \times \partial\Omega, \mathbb{R}) &\longrightarrow L^2((0, T), H^{-1}(\Omega)) \\ g &\longmapsto \Delta^2(v\psi). \end{aligned}$$

is continuous.

Proof of Proposition 9

Step 1. — We prove that $v\psi \in L^2((0, T), H^2 \cap H_0^1(\Omega))$. By Proposition 6 and 7, we know that $v \in L^2((0, T), H^{3/2}(\Omega))$. Thus $v\psi \in L^2((0, T), H^{3/2}(\Omega) \cap H_0^1(\Omega))$ because $H^{3/2}(\Omega) \subset L^\infty$ is an algebra. To end Step 1, it is sufficient to prove that $\Delta(v\psi) \in L^2((0, T), L^2(\Omega))$. Using $\Delta v = v$ in $(0, T) \times \Omega$, we get

$$\Delta(v\psi) = v(\psi + \Delta\psi) + 2\nabla v \cdot \nabla\psi.$$

– The Sobolev embedding (12) justifies that

$$H^{3/2}(\Omega) * H_0^1(\Omega) \subset L^\infty(\Omega) * L^2(\Omega) = L^2(\Omega)$$

(where $*$ stands for the multiplication of scalar valued functions). Thus $v(\psi + \Delta\psi) \in L^2((0, T), L^2(\Omega))$.

– The Sobolev embedding (12) justifies that

$$H^{1/2}(\Omega) * H^2(\Omega) \subset L^2(\Omega) * L^\infty(\Omega) = L^2(\Omega).$$

Thus $\nabla v \cdot \nabla\psi \in L^2((0, T), L^2(\Omega))$.

Step 2. — We prove that $\Delta^2[v\psi] \in L^2((0, T), H^{-1}(\Omega))$. Using $\Delta v = v$, we get

$$(21) \quad \Delta^2[v\psi] = v(\psi + 2\Delta\psi + \Delta^2\psi) + 4\nabla v \cdot \nabla(\psi + \Delta\psi) + 4\text{Tr}[D^2v \cdot D^2\psi].$$

– The Sobolev embeddings (12) and (13) justify that $H^{3/2}(\Omega) * H_0^1(\Omega) \subset H_0^1(\Omega)$. Thus, by duality, $H^{3/2}(\Omega) * H^{-1}(\Omega) \subset H^{-1}(\Omega)$. This proves that $v(\psi + 2\Delta\psi + \Delta^2\psi) \in L^2((0, T), H^{-1}(\Omega))$.

– The Sobolev embedding (13) justifies that

$$H^{1/2}(\Omega) * H_0^1(\Omega) \subset L^4(\Omega) * L^4(\Omega) \subset L^2(\Omega).$$

Thus, by duality, $H^{1/2}(\Omega) * L^2(\Omega) \subset H^{-1}(\Omega)$. This proves that $\nabla v \cdot \nabla(\psi + \Delta\psi) \in L^2((0, T), H^{-1}(\Omega))$.

– The Sobolev embedding (11) and then (14) justify that

$$H^1(\Omega) * H_0^1(\Omega) \subset H^{1/2+\varepsilon}(\Omega).$$

Thus, by duality, $H^1(\Omega) * H^{-1/2-\varepsilon}(\Omega) \subset H^{-1}(\Omega)$. This proves that $\text{Tr}[D^2v \cdot D^2\psi] \in L^2((0, T), H^{-1}(\Omega))$. □

Proof of Theorem 6. — Let $T > 0$, $\psi_0 \in H_{(0)}^3(\Omega)$, $g \in L^2((0, T) \times \partial\Omega, \mathbb{R})$ and v be the solution of (20) We apply the fixed point theorem to the map

$$\begin{aligned} F : C^0([0, T], H_{(0)}^3(\Omega)) &\longrightarrow C^0([0, T], H_{(0)}^3(\Omega)) \\ \psi &\longmapsto \xi, \end{aligned}$$

where $\xi = F(\psi)$ is the solution of

$$\begin{cases} (i\partial_t + \Delta)\xi(t, x) = v(t, x)\psi(t, x), & (t, x) \in (0, T) \times \Omega, \\ \xi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \xi(0, x) = \psi_0(x), & x \in \Omega. \end{cases}$$

Step 1. — We prove that F takes values in $C^0([0, T], H^3_{(0)}(\Omega))$.

Let $\psi \in C^0([0, T], H^3_{(0)}(\Omega))$. We introduce the solution μ_2 of

$$(22) \quad \begin{cases} \Delta^2 \mu_2(t, x) = \Delta^2[v\psi](t, x), & (t, x) \in (0, T) \times \Omega, \\ \mu_2(t, x) = \Delta \mu_2(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \end{cases}$$

and the function $\mu_1 := v\psi - \mu_2$. To prove that $\xi = F(\psi)$ belongs to $C^0([0, T], H^3_{(0)}(\Omega))$, it suffices to prove that μ_1 and μ_2 satisfy the assumptions of Theorem 5.

By Proposition 9, $\Delta^2[v\psi]$ belongs to $L^2((0, T), H^{-1}(\Omega))$ thus, by elliptic regularity, $\mu_2 \in L^2((0, T), H^3_{(0)}(\Omega))$. Then, μ_1 belongs to $L^2((0, T), H^2 \cap H^1_0(\Omega))$, by Proposition 9. Moreover, $\Delta^2 \mu_1 \equiv 0$ by (22) and $\Delta \mu_1|_{\partial\Omega} = 2g \partial_\nu \psi 1_{\Gamma_c}$.

Step 2. — We prove that F is a contraction when g is small enough. Let $\psi, \tilde{\psi} \in C^0([0, T], H^3_{(0)}(\Omega))$. From (19) and elliptic regularity, we get

$$\begin{aligned} \|F(\psi) - F(\tilde{\psi})\|_{C^0([0, T], H^3_{(0)}(\Omega))} &\leq C(\|\Delta(\mu_1 - \tilde{\mu}_1)\|_{L^2((0, T) \times \Gamma_c)} + \|\mu_2 - \tilde{\mu}_2\|_{L^1((0, T), H^3_{(0)}(\Omega))}) \\ &\leq C' \|g\|_{L^2((0, T) \times \Gamma_c)} \|\psi - \tilde{\psi}\|_{C^0([0, T], H^3_{(0)}(\Omega))}. \end{aligned}$$

for some constant C' independent of ψ and $\tilde{\psi}$. Thus F is a contraction when $C' \|g\|_{L^2((0, T) \times \Gamma_c)} < 1$. Otherwise, one may subdivide the interval $(0, T)$ into a finite number of intervals on which this assumption is satisfied and iterate the previous result. □

3.5. C^1 -REGULARITY OF THE END-POINT MAP

PROPOSITION 10. — *Let $T > 0$, Ω be an open subset of \mathbb{R}^2 and Γ_c be an open subset of $\partial\Omega$ such that **(H1)** holds. Consider the end-point map*

$$\begin{aligned} \Theta : [H^3_{(0)}(\Omega) \cap \mathcal{S}] \times L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow H^3_{(0)}(\Omega) \cap \mathcal{S} \\ (\psi_0, g) &\longmapsto \psi(T), \end{aligned}$$

where ψ is the solution of (1). Then, Θ is C^1 and for every $\psi_0 \in [H^3_{(0)}(\Omega) \cap \mathcal{S}]$,

$$\begin{aligned} d\Theta(\psi_0, 0) : [H^3_{(0)}(\Omega) \cap T_{\mathcal{S}}\psi_0] \times L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow H^3_{(0)}(\Omega) \cap T_{\mathcal{S}}\psi_{\text{ref}}(T) \\ (\Psi_0, G) &\longmapsto \Psi(T), \end{aligned}$$

where $\psi_{\text{ref}}(t) := e^{i\Delta_D t}\psi_0$ and Ψ is the solution of the linearized system

$$(23) \quad \begin{cases} (i\partial_t + \Delta)\Psi(t, x) = V(t, x)\psi_{\text{ref}}(t, x), & (t, x) \in (0, T) \times \Omega, \\ \Psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \Psi(0, x) = \Psi_0(x), & x \in \Omega, \\ (-\Delta + 1)V(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu V(t, x) = G(t, x)1_{\Gamma_c}(x), & (t, x) \in (0, T) \times \partial\Omega. \end{cases}$$

This result is a consequence of estimate (19). Its proof is classical and follows the same steps as in [5, Proof of Prop. 3, pp. 531–532].

4. LOCAL EXACT CONTROL OF HIGH FREQUENCIES

The goal of this section is the proof of Theorem 1, via a perturbation argument. In Section 4.1, we prove the controllability of the linearized system at high frequency, thanks to the weak observability inequality of the adjoint system and the Hilbert Uniqueness Method. In Section 4.2, we prove Theorem 1 by applying the inverse mapping theorem.

4.1. CONTROL OF HIGH FREQUENCIES FOR THE LINEARIZED SYSTEM. — The goal of this section is to prove the following result.

PROPOSITION 11. — *Let $T > 0$, $\psi_0 \in H^3_{(0)}(\Omega) \cap \mathcal{S}$ and $\psi_{\text{ref}}(t) := e^{i\Delta_D t}\psi_0$. We assume that **(H1)**, **(H2)** and **(H3)** hold. Then, there exists $K \in \mathbb{N}^*$ and a continuous linear map*

$$L : \mathbb{P}_K[H^3_{(0)}(\Omega)] \longrightarrow L^2((0, T) \times \Gamma_c, \mathbb{R})$$

such that, for every $\Psi_f \in \mathbb{P}_K[H^3_{(0)}(\Omega)]$, the solution of (23) with $\Psi_0 = 0$ and control $G = L(\Psi_f)$ satisfies $\mathbb{P}_K[\Psi(T)] = \Psi_f$.

The following proposition will allow us to work on $r := \Delta^2\Psi$ instead of Ψ .

PROPOSITION 12. — *Let $T > 0$, $\psi_0 \in H^3_{(0)}(\Omega) \cap \mathcal{S}$, $\psi_{\text{ref}}(t) := e^{i\Delta_D t}\psi_0$, $G \in L^2((0, T) \times \Gamma_c, \mathbb{R})$ and $\Psi \in C^0([0, T], H^3_{(0)}(\Omega))$ be the solution of the system (23) with $\Psi_0 = 0$. Then, the function $r := \Delta^2(\Psi)$ belongs to $C^0([0, T], H^{-1}(\Omega))$ and is the solution by transposition of*

$$(24) \quad \begin{cases} (i\partial_t + \Delta)r(t, x) = \mathcal{K}(G)(t, x), & (t, x) \in (0, T) \times \Omega, \\ r(t, x) = 2G\partial_\nu\psi_{\text{ref}}1_{\Gamma_c}, & (t, x) \in (0, T) \times \partial\Omega, \\ r(0, x) = 0, & x \in \Omega, \end{cases}$$

where

$$(25) \quad \begin{aligned} \mathcal{K} : L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow L^2((0, T), H^{-1}(\Omega)) \\ G &\longmapsto \Delta^2(V\psi_{\text{ref}}), \text{ where } \begin{cases} (-\Delta + 1)V = 0 \text{ in } (0, T) \times \Omega, \\ \partial_\nu V = G1_{\Gamma_c} \text{ on } (0, T) \times \partial\Omega. \end{cases} \end{aligned}$$

This means that, for every $\phi_T \in H_0^1(\Omega)$ and $\varphi \in L^1((0, T), H_0^1(\Omega))$ the following equality holds

$$(26) \quad \int_0^T \langle r(t), \overline{\varphi(t)} \rangle_{H^{-1}, H_0^1} dt = \int_0^T \langle \mathcal{K}(G)(t), \overline{\phi(t)} \rangle_{H^{-1}, H_0^1} dt - i \langle r(T), \overline{\phi_T} \rangle_{H^{-1}, H_0^1} + 2 \int_0^T \int_{\Gamma_c} G(t, x) \partial_\nu \psi_{\text{ref}}(t, x) \overline{\partial_\nu \phi(t, x)} d\sigma(x) dt,$$

where $\phi \in C^0([0, T], H_0^1(\Omega))$ is the solution of

$$\begin{cases} (i\partial_t + \Delta)\phi(t, x) = \varphi(t, x), & (t, x) \in (0, T) \times \Omega, \\ \phi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega \\ \phi(T, x) = \phi_T(x), & x \in \Omega. \end{cases}$$

Note that there is no ambiguity in this definition of a solution by transposition of (24). Indeed, for every $\phi_T \in H_0^1(\Omega)$ and $\varphi \in L^1((0, T), H_0^1(\Omega))$, then $\partial_\nu \phi$ belongs to $L^2((0, T) \times \Gamma_c)$ by Proposition 8, thus, the last integral in (26) is well-defined. Moreover, $\Delta^2(V\psi_{\text{ref}}) \in L^2((0, T), H^{-1}(\Omega))$ thanks to Proposition 9.

Proof of Proposition 12. — When $G \in C_c^\infty((0, T) \times \Gamma_c)$, $\varphi \in C_c^\infty((0, T) \times \Omega)$ and $\phi_T \in C_c^\infty(\Omega)$, then, formula (26) can be proved with integrations by part. The conclusion follows thanks to the continuity of the maps \mathcal{K} (see Proposition 9) and the continuity of the following linear mappings

$$\begin{aligned} L^2((0, T) \times \Gamma_c) &\longrightarrow C^0([0, T], H_{(0)}^3(\Omega)) \longrightarrow C^0([0, T], H^{-1}(\Omega)) \\ G &\longmapsto \Psi \longmapsto r = \Delta^2(\Psi) \\ L^1((0, T), H_0^1(\Omega)) \times H_0^1(\Omega) &\longrightarrow L^2((0, T) \times \Gamma_c) \\ (\varphi, \phi_T) &\longmapsto \partial_\nu \phi \end{aligned}$$

stated in Theorem 5 and Proposition 8. □

From now on, we denote by \mathcal{K}^* the adjoint operator of \mathcal{K} ,

$$\mathcal{K}^* : L^2((0, T), H_0^1(\Omega)) \longrightarrow L^2((0, T) \times \Gamma_c, \mathbb{R}),$$

i.e.,

$$(27) \quad \Re \left(\int_0^T \langle \mathcal{K}(G), \bar{\xi} \rangle_{H^{-1}, H_0^1} dt \right) = \int_0^T \int_{\Gamma_c} G \mathcal{K}^*(\xi) d\sigma(x) dt$$

for every $\xi \in L^2((0, T), H_0^1(\Omega))$ and $G \in L^2((0, T) \times \Gamma_c, \mathbb{R})$.

Thus Proposition 11 is a consequence of the following result.

PROPOSITION 13. — *Let $T > 0$, $\psi_0 \in H_{(0)}^3(\Omega) \cap \mathcal{S}$ and $\psi_{\text{ref}}(t) := e^{i\Delta_D t} \psi_0$. We assume that **(H1)**, **(H2)** and **(H3)** hold. Then, there exists $K \in \mathbb{N}^*$ and a continuous linear map*

$$\tilde{L} : \mathbb{P}_K[H^{-1}(\Omega)] \longrightarrow L^2((0, T) \times \Gamma_c, \mathbb{R})$$

such that, for every $r_f \in \mathbb{P}_K[H^{-1}(\Omega)]$, the solution of (24) with control $G = \tilde{L}(r_f)$ satisfies $\mathbb{P}_K[r(T)] = r_f$.

The controllability result of Proposition 13 is a consequence of the following weak observability result.

PROPOSITION 14. — *Let $T > 0$, $\psi_0 \in H_{(0)}^3(\Omega) \cap \mathcal{S}$ and $\psi_{\text{ref}}(t) := e^{i\Delta_D t} \psi_0$. We assume that **(H1)**, **(H2)** and **(H3)** hold. There exists $\mathcal{C}_1 > 0$ such that, for every $\phi_T \in H_0^1(\Omega, \mathbb{C})$ the solution of*

$$(28) \quad \begin{cases} (i\partial_t + \Delta)\phi(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \phi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \phi(T, x) = \phi_T, & x \in \Omega, \end{cases}$$

satisfies

$$(29) \quad \|\phi_T\|_{H_0^1(\Omega)} \leq \mathcal{C}_1 (\|2\Im(\partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi}) + \mathcal{K}^*(i\phi)\|_{L^2((0, T) \times \Gamma_c)} + \|\phi_T\|_{H^{-1}(\Omega)}).$$

Proof of Proposition 13, assuming Proposition 14. — By [12, Th. II.10 & II.19], the existence of a continuous right inverse to the continuous operator

$$\begin{aligned} F_K : L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow \mathbb{P}_K[H^{-1}(\Omega)] \\ G &\longmapsto \mathbb{P}_K[r(T)] \end{aligned}$$

is equivalent to the existence of $\mathcal{C}_2 > 0$ such that

$$(30) \quad \|\phi_T\|_{H_0^1(\Omega)} \leq \mathcal{C}_2 \|F_K^*(\phi_T)\|_{L^2((0, T) \times \Gamma_c)}, \quad \forall \phi_T \in \mathbb{P}_K[H_0^1(\Omega)],$$

where $F_K^* : \mathbb{P}_K[H_0^1(\Omega)] \rightarrow L^2((0, T) \times \Gamma_c, \mathbb{R})$ is the adjoint operator associated to F_K .

For $G \in L^2((0, T) \times \Gamma_c, \mathbb{R})$ and $\phi_T \in \mathbb{P}_K[H_0^1(\Omega)]$, we have (note that \mathcal{K} is \mathbb{R} -linear)

$$\begin{aligned} \Re(\langle F_K(G), \overline{\phi_T} \rangle_{H^{-1}, H_0^1}) &= \Re(\langle r(T), \overline{\phi_T} \rangle_{H^{-1}, H_0^1}) \quad \text{because } \phi_T \in \mathbb{P}_K[H_0^1(\Omega)] \\ &= \Re\left(-i \int_0^T \int_{\Gamma_c} 2G \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi} d\sigma(x) dt \right. \\ &\quad \left. - i \int_0^T \int_\Omega \langle \mathcal{K}(G), \overline{\phi} \rangle_{H^{-1}, H_0^1} dx dt \right) \quad \text{by (26)} \\ &= \int_0^T \int_{\Gamma_c} G (2\Im(\partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi}) + \mathcal{K}^*(i\phi)) d\sigma(x) dt \quad \text{by (27)}. \end{aligned}$$

Thus F_K^* has the following explicit expression

$$\begin{aligned} F_K^* : \mathbb{P}_K[H_0^1(\Omega)] &\longrightarrow L^2((0, T) \times \Gamma_c, \mathbb{R}) \\ \phi_T &\longmapsto 2\Im(\partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi}) + \mathcal{K}^*(i\phi). \end{aligned}$$

Let \mathcal{C}_1 be as in Proposition 14. There exists $K \in \mathbb{N}^*$ such that

$$\mathcal{C}_1 \|\phi_T\|_{H^{-1}(\Omega)} \leq \frac{1}{2} \|\phi_T\|_{H_0^1(\Omega)}, \quad \forall \phi_T \in \mathbb{P}_K[H_0^1(\Omega)]$$

and then (30) follows from (29) with $\mathcal{C}_2 := 2\mathcal{C}_1$. □

Proof of Proposition 14. — Let \tilde{T} be as in **(H2)** and T', T'' be as in **(H3)**. Let $\varepsilon := (T'' - T' - \tilde{T})/2$ and $\rho \in C_c^\infty(\mathbb{R}, \mathbb{R})$ be such that $\rho \equiv 1$ on $(T' + \varepsilon, T' + \varepsilon + \tilde{T})$ and $\text{Supp}(\rho) \subset (T', T'')$.

Step 1. — We prove the existence of $C_1 > 0$ such that, for every $\phi_T \in H_0^1(\Omega)$, the solution of (28) satisfies

$$(31) \quad \|\phi_T\|_{H_0^1(\Omega)} \leq C_1 \left(\|2\Im(\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi})\|_{L^2(\mathbb{R} \times \Gamma_c)} + \|\phi_T\|_{H^{-1}(\Omega)} \right).$$

Let $\phi_T \in H_0^1(\Omega)$. By **(H2)**, we have

$$\begin{aligned} \|\phi_T\|_{H_0^1(\Omega)}^2 &\leq 2\mathcal{C}_0^2 \left[\int_{T'+\varepsilon}^{T'+\varepsilon+\tilde{T}} \int_{\Gamma_c} |\partial_\nu \phi(t, x)|^2 d\sigma(x) dt + \|\phi_T\|_{H^{-1}}^2 \right] \\ &\leq 2\mathcal{C}_0^2 \left[\int_{\mathbb{R}} \int_{\Gamma_c} |\rho(t) \overline{\partial_\nu \phi(t, x)}|^2 d\sigma(x) dt + \|\phi_T\|_{H^{-1}}^2 \right] \end{aligned}$$

because $\rho \equiv 1$ on $(T'+\varepsilon, T'+\varepsilon+\tilde{T})$. Then, taking into account that $\text{Supp}(\rho) \subset (T', T'')$ and $|\partial_\nu \psi_{\text{ref}}(t, x)| \geq m > 0$ for every $(t, x) \in (T', T'') \times \Gamma_c$, i.e **(H3)**, we obtain

$$\begin{aligned} \|\phi_T\|_{H_0^1(\Omega)}^2 &\leq \frac{3\mathcal{C}_0^2}{m^2} \int_{\mathbb{R}} \int_{\Gamma_c} |\rho(t) \partial_\nu \psi_{\text{ref}}(t, x) \overline{\partial_\nu \phi(t, x)}|^2 d\sigma(x) dt \\ &\quad - \mathcal{C}_0^2 \int_{\mathbb{R}} \int_{\Gamma_c} |\rho(t) \partial_\nu \phi(t, x)|^2 d\sigma(x) dt + 2\mathcal{C}_0^2 \|\phi_T\|_{H^{-1}}^2. \end{aligned}$$

Moreover,

$$\begin{aligned} 2 \int_{\mathbb{R}} \int_{\Gamma_c} |\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi}|^2 d\sigma dt &= \int_{\mathbb{R}} \int_{\Gamma_c} \left(|\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi}|^2 + |\rho \partial_\nu \overline{\psi_{\text{ref}}} \partial_\nu \phi|^2 \right) d\sigma dt \\ &= \int_{\mathbb{R}} \int_{\Gamma_c} \left(|2\Im(\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi})|^2 + 2\Re((\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi})^2) \right) d\sigma dt. \end{aligned}$$

Note here that in the end of the proof, we will prove that the second term is compact. It will be crucial for that to notice that it is $\Re[(\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi})^2]$ and not $|\Re[\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi}]|^2$.

Thus

$$\begin{aligned} \|\phi_T\|_{H_0^1(\Omega)}^2 &\leq \frac{3\mathcal{C}_0^2}{2m^2} \int_{\mathbb{R}} \int_{\Gamma_c} \left(|2\Im(\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi})|^2 + 2\Re[(\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi})^2] \right) d\sigma dt \\ &\quad - \mathcal{C}_0^2 \int_{\mathbb{R}} \int_{\Gamma_c} |\rho \partial_\nu \phi|^2 d\sigma dt + 2\mathcal{C}_0^2 \|\phi_T\|_{H^{-1}}^2. \end{aligned}$$

In order to get (31), it suffices to prove the existence of $C > 0$ such that

$$(32) \quad \int_{\mathbb{R}} \int_{\Gamma_c} \Re[(\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi})^2] d\sigma dt \leq \frac{m^2}{3} \int_{\mathbb{R}} \int_{\Gamma_c} |\rho \partial_\nu \phi|^2 d\sigma dt + C \|\phi_T\|_{H^{-1}(\Omega)}^2.$$

Let $R \in \mathbb{N}^*$ that will be chosen later on,

$$(33) \quad \psi_{\text{ref}}^H(t) := \mathbb{P}_R[\psi_{\text{ref}}(t)] \quad \text{and} \quad \psi_{\text{ref}}^L := \psi_{\text{ref}} - \psi_{\text{ref}}^H.$$

Then

$$\begin{aligned} \int_{\mathbb{R}} \int_{\Gamma_c} \Re[(\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi})^2] d\sigma dt \\ = \int_{\mathbb{R}} \int_{\Gamma_c} \Re \left((\rho \partial_\nu \psi_{\text{ref}}^L \overline{\partial_\nu \phi})^2 + (\rho \overline{\partial_\nu \phi})^2 \partial_\nu \psi_{\text{ref}}^H (2 \partial_\nu \psi_{\text{ref}}^L + \partial_\nu \psi_{\text{ref}}^H) \right) d\sigma dt. \end{aligned}$$

In order to get (32), it is sufficient to prove that, for R large enough

$$(34) \quad \int_{\mathbb{R}} \int_{\Gamma_c} \Re[(\rho \partial_\nu \psi_{\text{ref}}^L \overline{\partial_\nu \phi})^2] d\sigma dt \leq C \|\phi_T\|_{H^{-1}},$$

$$(35) \quad \int_{\mathbb{R}} \int_{\Gamma_c} |(\rho \overline{\partial_\nu \phi})^2 \partial_\nu \psi_{\text{ref}}^H (2 \partial_\nu \psi_{\text{ref}}^L + \partial_\nu \psi_{\text{ref}}^H)| d\sigma dt \leq \frac{m^2}{2} \int_{\mathbb{R}} \int_{\Gamma_c} |\rho \partial_\nu \phi|^2 d\sigma dt.$$

Note that $\psi_{\text{ref}}^H \rightarrow 0$ in $C^0([0, T], H_{(0)}^3(\Omega))$ when $R \rightarrow +\infty$ because $\psi_0 \in H_{(0)}^3(\Omega)$. Thus, $\partial_\nu \psi_{\text{ref}}^H|_{\Gamma_c} \rightarrow 0$ in $L^\infty((0, T) \times \Gamma_c)$ when $R \rightarrow +\infty$, because of the Sobolev embeddings $\partial_\nu [H_{(0)}^3(\Omega)] \subset H^{3/2}(\Gamma_c) \subset L^\infty(\Gamma_c)$. As a consequence, there exists R_0 such that for any $R \geq R_0$, we have

$$\|\partial_\nu \psi_{\text{ref}}^H (2 \partial_\nu \psi_{\text{ref}}^L + \partial_\nu \psi_{\text{ref}}^H)\|_{L^\infty((0, T) \times \Gamma_c)} \leq \frac{m^2}{2}.$$

This implies (35). Until the end of Step 2, R is fixed. The function ρ is $C_c^\infty(\mathbb{R})$ thus, for every $N \in \mathbb{N}^*$, there exists $C_N > 0$ such that

$$\left| \int_{\mathbb{R}} \rho(t)^2 e^{i[\lambda_j + \lambda_J - \lambda_k - \lambda_K]t} dt \right| \leq \frac{C_N}{(\lambda_j + \lambda_J)^N}, \quad \forall j, J \in \mathbb{N}^*, k, K \in \{1, \dots, R-1\}.$$

We have

$$\begin{aligned} & \left| \int_{\mathbb{R}} \int_{\Gamma_c} \Re[(\rho \partial_\nu \psi_{\text{ref}}^L \overline{\partial_\nu \phi})^2] d\sigma dt \right| \\ &= \left| \sum_{k, K=1}^{R-1} \sum_{j, J=1}^{\infty} \psi_k^0 \psi_K^0 \overline{\phi_j^0 \phi_J^0} \left(\int_{\mathbb{R}} \rho(t)^2 e^{i[\lambda_j + \lambda_J - \lambda_k - \lambda_K]t} dt \right) \cdot \left(\int_{\Gamma_c} \partial_\nu \varphi_k \partial_\nu \varphi_K \partial_\nu \varphi_j \partial_\nu \varphi_J d\sigma \right) \right| \\ &\leq \sum_{k, K=1}^{R-1} \sum_{j, J=1}^{\infty} |\psi_k^0 \psi_K^0 \phi_j^0 \phi_J^0| \frac{C_N}{(\lambda_j + \lambda_J)^N} \lambda_k^\alpha \lambda_K^\alpha \lambda_j^\alpha \lambda_J^\alpha \quad \text{for some } \alpha > 0 \\ &\leq C(\psi_0, R, N) \left(\sum_j \lambda_j^{\alpha-N} |\phi_j^0| \right)^2 \\ &\leq C'(\psi_0, R, N) \|\phi_T\|_{H^{-1}(\Omega)}^2 \text{ for } N \text{ large enough.} \end{aligned}$$

Here, we have used Weyl law to get that for another N large enough, $\sum_j^\infty \lambda_j^{-N} < +\infty$. Inequality (34) is proved, which ends Step 1.

Step 2. — We prove the existence of $C_2 > 0$ such that, for every $\phi_T \in H_0^1(\Omega)$, the solution of (28) satisfies

$$(36) \quad \|\phi_T\|_{H_0^1(\Omega)} \leq C_2 \left(\|2\Im(\partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi}) + \mathcal{K}^*(i\phi)\|_{L^2((0, T) \times \Gamma_c)} + \|\phi_T\|_{H^{-1}(\Omega)} \right).$$

From (31), we deduce that

$$\begin{aligned} & \|\phi_T\|_{H_0^1(\Omega)} \\ & \leq C_1 \left(\|2\Im(\rho \partial_\nu \psi_{\text{ref}} \overline{\partial_\nu \phi}) + \mathcal{K}^*(i\phi)\|_{L^2(\mathbb{R} \times \Gamma_c)} + \|\mathcal{K}^*(i\phi)\|_{L^2(\mathbb{R} \times \Gamma_c)} + \|\phi_T\|_{H^{-1}(\Omega)} \right). \end{aligned}$$

In order to prove (36), it suffices to prove the existence of $C > 0$ such that,

$$(37) \quad \|\mathcal{K}^*(i\phi)\|_{L^2((0,T)\times\Gamma_c)} \leq \frac{1}{2C_1} \|\phi_T\|_{H_0^1(\Omega)} + C \|\phi_T\|_{H^{-1}}, \quad \forall \phi_T \in H_0^1(\Omega).$$

Let $\tilde{R} \in \mathbb{N}^*$ that will be chosen later on and ψ_{ref}^H be defined as in (33) but with R replaced by \tilde{R} . From (25) and the expansion (21), we see that $\mathcal{K} = \mathcal{K}_1 + \mathcal{K}_2$, where

$$\begin{aligned} \mathcal{K}_1 : L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow L^2((0, T), H^{-1}(\Omega)) \\ G &\longmapsto \Delta^2(V\psi_{\text{ref}}^H) \end{aligned}$$

and

$$\begin{aligned} \mathcal{K}_2 : L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow L^2((0, T), H^{-1/2-\varepsilon}(\Omega)) \\ G &\longmapsto \Delta^2(V\psi_{\text{ref}}^L) \end{aligned}$$

is continuous. We denote by $\|\mathcal{K}_1\|$ and $\|\mathcal{K}_2\|$ their operator norm in this functional frame.

Note that $\psi_{\text{ref}}^H \rightarrow 0$ in $C^0([0, T], H_0^3(\Omega))$ when $\tilde{R} \rightarrow +\infty$ because $\psi_0 \in H_0^3(\Omega)$. Thus, there exists $\tilde{R} \in \mathbb{N}^*$ such that

$$\|\mathcal{K}_1\| \leq \frac{1}{2C_1\sqrt{T}}.$$

From now on \tilde{R} is fixed. Then, for every $\phi_T \in H_0^1(\Omega)$,

$$\begin{aligned} \|\mathcal{K}^*(i\phi)\|_{L^2((0,T)\times\Gamma_c)} &\leq \|\mathcal{K}_1^*(i\phi)\|_{L^2((0,T)\times\Gamma_c)} + \|\mathcal{K}_2^*(i\phi)\|_{L^2((0,T)\times\Gamma_c)} \\ &\leq \frac{1}{2C_1\sqrt{T}} \|\phi\|_{L^2((0,T), H_0^1(\Omega))} + \|\mathcal{K}_2\| \|\phi\|_{L^2((0,T), H^{1/2+\varepsilon}(\Omega))} \\ &\leq \frac{1}{2C_1} \|\phi_T\|_{H_0^1(\Omega)} + \sqrt{T} \|\mathcal{K}_2\| \|\phi_T\|_{H^{1/2+\varepsilon}(\Omega)}. \end{aligned}$$

This implies (37) after interpolation, which ends Step 2. \square

4.2. CONTROL OF HIGH FREQUENCIES FOR THE NONLINEAR SYSTEM. — The goal of this section is the proof of Theorem 1. Thus, in the whole section, $T > 0$, $\psi_0 \in H_0^3(\Omega) \cap \mathcal{S}$, $\psi_{\text{ref}}(t) := e^{i\Delta_D t} \psi_0$ are fixed and **(H1)**, **(H2)**, **(H3)** are assumed to hold. We consider the end-point map

$$\begin{aligned} \tilde{\Theta}_K : L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow \mathbb{P}_K[H_0^3(\Omega)] \\ g &\longmapsto \mathbb{P}_K[\psi(T)], \end{aligned}$$

where ψ solves (1). By Proposition 10, $\tilde{\Theta}_K$ is of class C^1 and

$$\begin{aligned} d\tilde{\Theta}_K(0) : L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow \mathbb{P}_K[H_0^3(\Omega)] \\ G &\longmapsto \mathbb{P}_K[\Psi(T)], \end{aligned}$$

where Ψ solves (23) with $\Psi_0 = 0$. By Proposition 11, there exists $K \in \mathbb{N}^*$ such that $d\tilde{\Theta}_K(0)$ has a continuous right inverse. By the inverse mapping theorem, $\tilde{\Theta}_K$ is a local C^1 -diffeomorphism on a neighborhood of 0. The first statement of Theorem 1 holds with $\Upsilon := \tilde{\Theta}_K^{-1}$ which is locally well-defined.

There exists $K' \geq K$ such that

$$\|\psi_{\text{ref}}(T) - \mathbb{P}_{K'}[\psi_{\text{ref}}(T)]\|_{H^3_{(0)}} < \delta.$$

Then, $\psi_f := (\mathbb{P}_K - \mathbb{P}_{K'})[\psi_{\text{ref}}(T)]$ belongs to \mathcal{V} thus $g := \Upsilon(\psi_f)$ is well-defined in $L^2((0, T) \times \Gamma_c, \mathbb{R})$ and the associated solution of (1) satisfies $\mathbb{P}_K[\psi(T)] = \psi_f$ thus $\mathbb{P}_{K'}[\psi(T)] = 0$. As a consequence $\psi(T)$ is a finite sum of eigenfunctions of $(-\Delta_D)$ so it is a smooth function. \square

5. LOCAL EXACT CONTROL AROUND EIGENFUNCTIONS

The goal of this section is to prove Theorem 2, by following essentially the same strategy as in the previous section. In Section 5.1, we prove the observability of the adjoint of the linearized system. In Section 5.2, we prove Theorem 2.

5.1. OBSERVABILITY RESULTS. — The goal of this section is to prove the following observability inequality.

PROPOSITION 15. — *Let $T > 0$, $R \in \mathbb{N}^*$. We assume that **(H1)**, **(H2)**, **(H3')** and **(H4)** hold. There exists $\mathcal{C}_2 > 0$ such that, for every $\phi_T \in H^1_0(\Omega) \cap T_S\varphi_R$ the solution of*

$$(38) \quad \begin{cases} (i\partial_t + \Delta + \lambda_R)\phi(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \phi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \phi(T, x) = \phi_T, & x \in \Omega, \end{cases}$$

satisfies

$$(39) \quad \|\phi_T\|_{H^1_0(\Omega)} \leq \mathcal{C}_2 \|2\Im(\partial_\nu \varphi_R \overline{\partial_\nu \phi}) + \tilde{\mathcal{K}}^*(i\phi)\|_{L^2((0, T) \times \Gamma_c)},$$

where

$$(40) \quad \begin{aligned} \tilde{\mathcal{K}} : L^2(\Gamma_c, \mathbb{R}) &\longrightarrow L^2(\Omega, \mathbb{C}) \\ G &\longmapsto \Delta^2(V\varphi_R), \quad \text{where } \begin{cases} (-\Delta + 1)V = 0, & \text{in } \Omega, \\ \partial_\nu V = G1_{\Gamma_c} & \text{on } \partial\Omega. \end{cases} \end{aligned}$$

The proof of Proposition 15 relies on two key ingredients: the weak observability of the Schrödinger equation **(H2)** and a unique continuation result given by the following statement.

PROPOSITION 16. — *Let $T > 0$, $R \in \mathbb{N}^*$. We assume that **(H1)**, **(H2)**, **(H3')** and **(H4)** hold. Then for every non-zero $\phi_T \in H^1_0(\Omega) \cap T_S\varphi_R$, the function $2\Im(\partial_\nu \varphi_R \overline{\partial_\nu \phi}) + \tilde{\mathcal{K}}^*(i\phi)$ is not identically zero on $(0, T) \times \Gamma_c$, where $\phi(t) := e^{i\Delta_D(t-T)}\phi_T$.*

Proof of Proposition 16. — Let $T' \in (\tilde{T}, T)$, where \tilde{T} is as in **(H2)**. By Proposition 14 (with T replaced by T') and the change of phase $\phi(t, x) \leftarrow \phi(t, x)e^{i\lambda_R t}$, we know that there exists $\mathcal{C}_1 > 0$ such that, for every $\phi_T \in H^1_0(\Omega)$, the solution of (38) satisfies

$$(41) \quad \|\phi_T\|_{H^1_0(\Omega)} \leq \mathcal{C}_1 (\|2\Im(\partial_\nu \varphi_R \overline{\partial_\nu \phi}) + \tilde{\mathcal{K}}^*(i\phi)\|_{L^2((0, T') \times \Gamma_c)} + \|\phi_T\|_{H^{-1}(\Omega)}).$$

We introduce

$$N_T := \left\{ \phi_0 \in H^1_0(\Omega) \cap T_S\varphi_R ; 2\Im(\partial_\nu \varphi_R \overline{\partial_\nu \phi}) + \tilde{\mathcal{K}}^*(i\phi) = 0 \text{ in } L^2((0, T) \times \Gamma_c) \right\},$$

where $\phi(t) := e^{i(\Delta_D + \lambda_R)t} \phi_0 := S(t)\phi_0$. N_T is a \mathbb{R} -vector subspace of $H_0^1(\Omega)$. We want to prove that $N_T = \{0\}$.

Step 1. — We prove that N_T contains eigenfunctions of $-\Delta_D$. Let $\phi_0 \in N_T$. Since $\phi_0 \in H_0^1(\Omega)$, then $\phi_\varepsilon := (S(\varepsilon)\phi_0 - \phi_0)/\varepsilon$ is bounded in $H^{-1}(\Omega)$ uniformly when $\varepsilon \rightarrow 0$. Moreover, for $\varepsilon < T - T'$, ϕ_ε belongs to $N_{T'}$. Applying (41) to ϕ_ε , with $\varepsilon < T - T'$, we get $\|\phi_\varepsilon\|_{H_0^1(\Omega)} \leq C_1 \|\phi_\varepsilon\|_{H^{-1}}$. Thus ϕ_ε is bounded in $H_0^1(\Omega)$ when $\varepsilon \rightarrow 0$. Therefore $\phi_0 \in H_{(0)}^3(\Omega)$. Indeed, by Fatou's lemma,

$$\begin{aligned} \|\phi_0\|_{H_{(0)}^3}^2 &= \sum_{k=1}^{\infty} |\lambda_k^{3/2} \langle \phi_0, \varphi_k \rangle|^2 = \sum_{k=1}^{\infty} \liminf_{\varepsilon \rightarrow 0} \left| \sqrt{\lambda_k} \frac{e^{i\lambda_k \varepsilon} - 1}{\varepsilon} \langle \phi_0, \varphi_k \rangle \right|^2 \\ &\leq \liminf_{\varepsilon \rightarrow 0} \sum_{k=1}^{\infty} \left| \sqrt{\lambda_k} \frac{e^{i\lambda_k \varepsilon} - 1}{\varepsilon} \langle \phi_0, \varphi_k \rangle \right|^2 \\ &\leq \liminf_{\varepsilon \rightarrow 0} \|\phi_\varepsilon\|_{H_0^1}^2 < \infty. \end{aligned}$$

For $T'' < T$, the map

$$\begin{aligned} \mathcal{J}_{T''} : H_0^1(\Omega) &\longrightarrow L^2((0, T'') \times \Gamma_c) \\ \phi_0 &\longmapsto 2\Im[\partial_\nu \varphi_R \overline{\partial_\nu \phi}] + \mathcal{K}^*(i\phi) \end{aligned}$$

is continuous (see Propositions 8 and 9, which imply the continuity of $\mathcal{K} : L^2(\Gamma_c, \mathbb{R}) \rightarrow L^2(\Omega)$ and thus of $\mathcal{K}^* : L^2(\Omega) \mapsto L^2(\Gamma_c, \mathbb{R})$). Moreover, $\mathcal{J}_{T''}(\phi_\varepsilon) = 0$ for every $\varepsilon < T - T''$, and $\phi_\varepsilon \rightarrow i(\Delta_D + \lambda_R)\phi_0$ in $H_0^1(\Omega)$ thus $\mathcal{J}_{T''}[i(\Delta_D + \lambda_R)\phi_0] = 0$. This holds for every $T'' \in (0, T)$, thus $i(\Delta_D + \lambda_R)\phi_0 \in N_T$.

We have proved that N_T is stable by $i(\Delta_D + \lambda_R)$ and only contains smooth functions in $D[(-\Delta_D)^s]$, $\forall s \in \mathbb{N}^*$. Applying again the estimate (41) to $i(\Delta_D + \lambda_R)\phi_0$, we get

$$\|(\Delta_D + \lambda_R)\phi_0\|_{H_0^1(\Omega)} \leq C_1 \|(\Delta_D + \lambda_R)\phi_0\|_{H^{-1}(\Omega)}.$$

This shows that the unit ball of N_T is compact (for the $H_0^1(\Omega)$ -topology), thus N_T has finite dimension (Rellich's theorem).

Then the finite dimensional space N_T is stable by the real symmetric operator $-(\Delta_D + \lambda_R)^2 = [i(\Delta_D + \lambda_R)]^2$, thus $-(\Delta_D + \lambda_R)^2$ has an eigenfunction in N_T . This candidate is also an eigenfunction of $(-\Delta_D)$.

Step 2. — We prove that

$$(42) \quad \tilde{\mathcal{K}}^*(z\varphi_k) = \Re(z) (\lambda_k^2 w_k - 2 \partial_\nu \varphi_R \partial_\nu \varphi_k)_{|\Gamma_c}, \quad \forall z \in \mathbb{C}, k \in \mathbb{N}^*,$$

where w_k is defined in (H4). Let $z \in \mathbb{C}$ and $k \in \mathbb{N}^*$. For every $G \in L^2(\Gamma_c, \mathbb{R})$, we have

$$\int_{\Gamma_c} G \tilde{\mathcal{K}}^*(z\varphi_k) d\sigma(x) = \Re \left(\int_{\Omega} \tilde{\mathcal{K}}(G) \overline{z\varphi_k} dx \right) = \Re(z) \int_{\Omega} \tilde{\mathcal{K}}(G) \varphi_k dx,$$

thus $\tilde{\mathcal{K}}^*(z\varphi_k) = \Re(z) \tilde{\mathcal{K}}^*(\varphi_k)$.

If $G \in C_c^\infty((0, T) \times \Gamma_c, \mathbb{R})$, then the potential V defined in (40) belongs to $C^\infty((0, T) \times \Omega, \mathbb{R})$ thus the following integrations by part are legitimate:

$$\begin{aligned} \int_{\Omega} \tilde{\mathcal{K}}(G)\varphi_k dx &= \int_{\Omega} \Delta^2(V\varphi_R)\varphi_k dx \quad \text{with } V \text{ as in (40)} \\ &= \int_{\partial\Omega} (\partial_\nu[\Delta(V\varphi_R)]\varphi_k - \Delta(V\varphi_R)\partial_\nu\varphi_k) d\sigma(x) + \int_{\Omega} \Delta(V\varphi_R)\Delta\varphi_k dx \\ &= -2 \int_{\Gamma_c} G\partial_\nu\varphi_R\partial_\nu\varphi_k d\sigma(x) - \lambda_k \int_{\Omega} \Delta(V\varphi_R)\varphi_k dx, \\ \int_{\Omega} \Delta(V\varphi_R)\varphi_k dx &= -\lambda_k \int_{\Omega} V\varphi_R\varphi_k dx \\ &= -\lambda_k \int_{\Omega} V(-\Delta + 1)w_k dx \\ &= -\lambda_k \int_{\Gamma_c} Gw_k d\sigma(x). \end{aligned}$$

This proves that

$$\int_{\Omega} \tilde{\mathcal{K}}(G)\varphi_k dx = \int_{\Gamma_c} G(\lambda_k^2 w_k - 2\partial_\nu\varphi_R\partial_\nu\varphi_k) d\sigma$$

for every $G \in C_c^\infty((0, T) \times \Gamma_c, \mathbb{R})$. The same equality holds by density for every $G \in L^2((0, T) \times \Gamma_c, \mathbb{R})$. Therefore, $\tilde{\mathcal{K}}^*(\varphi_k) = \lambda_k^2 w_k - 2\partial_\nu\varphi_R\partial_\nu\varphi_k$.

Step 3. — We prove that $N_T = \{0\}$. Working by contradiction, we assume that $N_T \neq \{0\}$. By Step 1, there exists $z \in \mathbb{C} \setminus \{0\}$ and $k \in \mathbb{N}^*$ such that $z\varphi_k \in N_T$. Moreover, by Step 2,

$$2\Im(\partial_\nu\varphi_R\overline{\partial_\nu[z\varphi_k e^{i(\lambda_R - \lambda_k)t}]} + \tilde{\mathcal{K}}^*(iz\varphi_k e^{i(\lambda_R - \lambda_k)t}) = \Re(iz e^{i(\lambda_R - \lambda_k)t})\lambda_k^2 w_k.$$

Thus $\Re(iz e^{i(\lambda_R - \lambda_k)t})w_k$ vanishes on $(0, T) \times \Gamma_c$. By Assumption **(H4)**, we deduce that

$$(43) \quad \Re(iz e^{i(\lambda_R - \lambda_k)t}) \equiv 0 \quad \text{on } (0, T).$$

First case: $k = R$. Then $z \in i\mathbb{R}$ because $z\varphi_k \in T_S\varphi_R$. From (43), we deduce that $0 = \Re(iz) = iz$, which is impossible because $z \neq 0$.

Second case: $k \neq R$. Then $\lambda_k \neq \lambda_R$ because λ_R is a simple eigenvalue (see **(H4)**). From (43), we deduce that $z = 0$, which is a contradiction. \square

REMARK 2. — Note that Assumption **(H4)** is necessary for the controllability of the linearized system 23 (with $\Psi_0 = 0$) around $(\psi_{\text{ref}}(t) = \varphi_R e^{-i\lambda_R t}, g_{\text{ref}} \equiv 0)$. Indeed, for any eigenfunction Φ of $(-\Delta_D)$ associated with the eigenvalue λ_Φ , we have

$$\begin{aligned} \langle \Psi(T), \Phi \rangle &= \left\langle i \int_0^T e^{-i\Delta_D(T-t)} (V(t)\psi_{\text{ref}}(t)) dt, \Phi \right\rangle \\ &= ie^{-i\lambda_\Phi T} \int_0^T e^{-i(\lambda_R - \lambda_\Phi)t} \langle V(t), \varphi_R \Phi \rangle dt \\ &= ie^{-i\lambda_\Phi T} \int_0^T e^{-i(\lambda_R - \lambda_\Phi)t} \langle G(t), w \rangle_{L^2(\Gamma_c)} dt, \end{aligned}$$

where w is as in **(H4)** (apply the Green formula, as in the proof of Proposition 2). In particular, if **(H4)** is not fulfilled, we are in one of the following situation

– if λ_R is a multiple eigenvalue, then, for any eigenfunction Φ associated to the eigenvalue $\lambda_\Phi = \lambda_R$ and linearly independent of φ_R , one cannot control the complex component $i\langle \Psi(T), \Phi \rangle e^{i\lambda_\Phi T}$ because it only takes real values

$$i\langle \Psi(T), \Phi \rangle e^{i\lambda_\Phi T} = \int_0^T \langle G(t), w \rangle dt \in \mathbb{R},$$

– if there exists an eigenfunction Φ for which $w \equiv 0$ on Γ_c , then, one cannot control the component $\langle \Psi(T), \Phi \rangle$. Indeed, whatever the control G is, the final state satisfies

$$\langle \Psi(T), \Phi \rangle = ie^{-i\lambda_\Phi T} \int_0^T e^{-i(\lambda_R - \lambda_\Phi)t} \langle G(t), w \rangle_{L^2(\Gamma_c)} dt = 0.$$

Proof of Proposition 15. – Working by contradiction, we assume that, for every $n \in \mathbb{N}^*$, there exists $\phi_T^n \in H_0^1(\Omega)$ such that

$$(44) \quad 1 = \|\phi_T^n\|_{H_0^1(\Omega)} > n \|2\Im(\partial_\nu \varphi_R \overline{\partial_\nu \phi^n}) + \tilde{\mathcal{K}}^*(i\phi^n)\|_{L^2((0,T) \times \Gamma_c)}.$$

Up to a subsequence, one may assume that $\phi_T^n \rightharpoonup \phi_T^\infty$ weakly in $H_0^1(\Omega)$. The strong continuity of the operator

$$\begin{aligned} H_0^1(\Omega) &\longrightarrow L^2((0, T) \times \Gamma_c) \\ \phi_T &\longmapsto 2\Im(\partial_\nu \varphi_R \overline{\partial_\nu \phi}) + \tilde{\mathcal{K}}^*(i\phi) \end{aligned}$$

implies its continuity for the weak topology [12, Th. III.9]. Thus

$$2\Im(\partial_\nu \varphi_R \overline{\partial_\nu \phi^n}) + \tilde{\mathcal{K}}^*(i\phi^n) \xrightarrow{n \rightarrow \infty} 2\Im(\partial_\nu \varphi_R \overline{\partial_\nu \phi^\infty}) + \tilde{\mathcal{K}}^*(i\phi^\infty) \text{ in } L^2((0, T) \times \Gamma_c).$$

From (44), we deduce that

$$2\Im(\partial_\nu \varphi_R \overline{\partial_\nu \phi^\infty}) + \tilde{\mathcal{K}}^*(i\phi^\infty) = 0 \text{ on } (0, T) \times \Gamma_c.$$

By Proposition 16, $\phi_T^\infty = 0$. Thus, up to a subsequence, $\|\phi_T^n\|_{H^{-1}(\Omega)} \rightarrow 0$ when $n \rightarrow \infty$. We deduce from (41) that

$$1 = \|\phi_T^n\|_{H_0^1(\Omega)} \leq \mathcal{C}_1 \left(\frac{1}{n} + \|\phi_T^n\|_{H^{-1}(\Omega)} \right) \xrightarrow{n \rightarrow \infty} 0,$$

which is a contradiction. □

5.2. PROOF OF THEOREM 2. – Let $T > 0$, $R \in \mathbb{N}^*$ and $\psi_{\text{ref}}(t) := \varphi_R(x)e^{-i\lambda_R t}$. We assume that **(H1)**, **(H2)**, **(H3')** and **(H4)** hold. By Proposition 10, the end-point map

$$\begin{aligned} \tilde{\Theta} : [H_{(0)}^3(\Omega) \cap \mathcal{S}] \times L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow [H_{(0)}^3(\Omega) \cap \mathcal{S}]^2 \\ (\psi_0, g) &\longmapsto (\psi_0, \psi(T)e^{i\lambda_R T}) \end{aligned}$$

is of class C^1 and

$$\begin{aligned} d\tilde{\Theta}(\varphi_R, 0) : [H_{(0)}^3(\Omega) \cap T_S \varphi_R] \times L^2((0, T) \times \Gamma_c, \mathbb{R}) &\longrightarrow [H_{(0)}^3(\Omega) \cap T_S \varphi_R]^2 \\ (\Psi_0, G) &\longmapsto (\Psi_0, \Psi(T)), \end{aligned}$$

where

$$(45) \quad \begin{cases} (i\partial_t + \Delta + \lambda_R)\Psi(t, x) = V(t, x)\varphi_R, & (t, x) \in (0, T) \times \Omega, \\ \Psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \Psi(0, x) = \Psi_0(x), & x \in \Omega, \\ (-\Delta + 1)V(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu V(t, x) = G(t, x)1_{\Gamma_c}(x), & (t, x) \in (0, T) \times \partial\Omega. \end{cases}$$

The same arguments as in the previous section prove that $d\tilde{\Theta}(\varphi_R, 0)$ has a continuous right inverse thanks to Proposition 15. The inverse mapping theorem proves that $\tilde{\Theta}$ is a local C^1 -diffeomorphism. \square

5.3. SOME COMMENTS ABOUT CONTROL ONLY DEPENDING ON TIME. — In this subsection, we comment on the information given by our previous study for a control that would be of the form

$$(46) \quad \begin{cases} (i\partial_t + \Delta)\psi(t, x) = v(t, x)\psi(t, x), & (t, x) \in (0, T) \times \Omega, \\ \psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \psi(0, x) = \psi_0(x) & x \in \Omega, \\ (-\Delta + 1)v(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu v(t, x) = g(t)\mu(x)1_{\Gamma_c}(x) & (t, x) \in (0, T) \times \partial\Omega, \end{cases}$$

where the control $g \in L^2(0, T)$ only depends on time and $\mu \in L^\infty(\partial\Omega)$ is fixed.

This is very close to the configuration studied in several papers described in the introduction (except that we impose that the potential $v = g(t)V_\mu$ is harmonic).

If we followed the framework described in the previous section, we would obtain that the observability estimate necessary to obtain the controllability of the linearized system, would be of the form

$$(47) \quad \|\phi_T\|_{H_0^1(\Omega)}^2 \leq C_2 \int_0^T \left| \int_{\partial\Omega} 2\Im(\mu(x)\partial_\nu\varphi_R\overline{\partial_\nu\phi}) + \mathcal{K}^*(i\phi)d\sigma(x) \right|^2 dt,$$

instead of the classical observability estimate (39), where \mathcal{K} would be another (but similar) compact operator.

Then, it becomes quite clear that (47) is really not likely to be true in general. Indeed, to contradict (47), it suffices to find a sequence of solutions with H_0^1 norm equals to 1 but whose Neumann trace on the boundary weakly converges to zero. This can be easily done for instance

- on the square $(0, \pi)^2$: with the normalized initial data $\phi_n(x, y) = c_n \sin(nx)\sin(ny)$
- on the disk: with some initial data $\phi_n(r, \theta) = e^{in\theta}g_n(r)$, where g_n are appropriate normalized Bessel functions.

Also, this analysis can also help to identify some class of initial data that could be observable. It is necessary that there is no oscillation of the trace on the boundary. This could be true for instance

- on the square: with some initial data that would depend only on x or y .
- on the disk: with some radial initial data.

Also, from a microlocal point of view, we can see that (47) can only be true for sequences of solutions whose trace do not weakly converge to zero. That means some data whose Neumann trace on the boundary is concentrated in the frequency $\xi' = 0$. Yet, the theorems of propagation of microlocal defect measure (described for example for the wave equation in [14]) could suggest that it implies that the data are concentrated close to some rays that intersect the boundary orthogonally.

6. SCHRÖDINGER EQUATION WITH REAL-VALUED BOUNDARY CONTROLS

The goal of this section is to prove Theorem 3. The arguments have already been developed in the Step 1 of the proof of Proposition 14 but here, we give a simpler proof in this simpler case.

PROPOSITION 17. — *Let Ω be an open subset of \mathbb{R}^2 , Γ be an open subset of $\partial\Omega$ and $0 < \tilde{T} < T < \infty$. If the Schrödinger equation on Ω is weakly observable on $(0, \tilde{T}) \times \Gamma$ then, there exists $\mathcal{C}_1 > 0$ such that*

$$(48) \quad \|\phi_0\|_{H_0^1(\Omega)} \leq \mathcal{C}_1 (\|\Im(\partial_\nu \phi)\|_{L^2((0, T) \times \Gamma)} + \|\phi_0\|_{H^{-1}(\Omega)}), \quad \forall \phi_0 \in H_0^1(\Omega, \mathbb{C}),$$

where $\phi(t) := e^{it\Delta_D} \phi_0$.

Proof. — Let $T', T'' \in (0, T)$ and $\rho \in C_c^\infty(\mathbb{R}, \mathbb{R}^+)$ be such that $T'' - T' > \tilde{T}$, $\rho \equiv 1$ on (T', T'') and $\text{Supp}(\rho) \subset (0, T)$. The assumption and conservation of the norm give

$$(49) \quad \|\phi_0\|_{H_0^1(\Omega)}^2 \leq \mathcal{C}_0 (\|\rho \partial_\nu \phi\|_{L^2(\mathbb{R}_t \times \Gamma)}^2 + \|\phi_0\|_{H^{-1}(\Omega)}^2), \quad \forall \phi_0 \in H_0^1(\Omega, \mathbb{C})$$

Since $|\Im(\partial_\nu \phi)|^2 = |\partial_\nu \phi|^2/2 + \Re((\partial_\nu \phi)^2)/2$, it is enough to prove

$$\left| \int_{\mathbb{R}_t} \int_{\Gamma} \rho^2 (\partial_\nu \phi)^2 \right| \leq C \|\phi_0\|_{H^{-1}(\Omega)}^2.$$

This inequality is obviously false if $(\partial_\nu \phi)^2$ is replaced by $|\partial_\nu \phi|^2$. It is actually true because $(\partial_\nu \phi)^2$ is the product of two terms that oscillate in the same direction, which is false for $|\partial_\nu \phi|^2$.

More precisely, write $\phi_0 = \sum_k \varphi_k \phi_{0,k}$, where $(\varphi_k)_{k \in \mathbb{N}^*}$ is an orthonormal basis of eigenfunctions for Δ_D with eigenvalues $-\lambda_k$. We have $\phi = \sum_k e^{-i\lambda_k t} \varphi_k \phi_{0,k}$

$$\int_{\mathbb{R}_t} \int_{\Gamma} \rho^2 (\partial_\nu \phi)^2 = \sum_{k,l} \phi_{0,l} \phi_{0,k} \left(\int_{\mathbb{R}_t} e^{-i(\lambda_k + \lambda_l)t} \rho^2 \right) \left(\int_{\Gamma} (\partial_\nu \varphi_k)(\partial_\nu \varphi_l) \right).$$

The boundary term can be bounded for instance by trace estimates (actually, finer estimates would give a better exponent $\lambda_k^{1/2}$)

$$\begin{aligned} \int_{\Gamma} |(\partial_\nu \varphi_k)(\partial_\nu \varphi_l)| &\leq \|\partial_\nu \varphi_k\|_{L^2(\Gamma)} \|\partial_\nu \varphi_l\|_{L^2(\Gamma)} \leq C \|\varphi_k\|_{H^2(\Omega)} \|\varphi_l\|_{H^2(\Omega)} \\ &\leq C \lambda_k \lambda_l \leq C(\lambda_k + \lambda_l)^2. \end{aligned}$$

The function ρ is $C_c^\infty(\mathbb{R})$ thus, for every $N \in \mathbb{N}^*$, there exists $C_N > 0$ (whose value may change) such that

$$\left| \int_{\mathbb{R}} \rho(t)^2 e^{-i[\lambda_k + \lambda_l]t} dt \right| \leq \frac{C_N}{(\lambda_k + \lambda_l)^N} \leq \frac{C_N}{(\lambda_k \lambda_l)^{N/2}}, \quad \forall k, l \in \mathbb{N}^*$$

So, choosing N large enough (and replacing $N/2 - 2$ by N), we obtain the bound

$$\left| \int_{\mathbb{R}_t} \int_{\Gamma} \rho^2 (\partial_\nu \phi)^2 \right| \leq \sum_{k,l} |\phi_{0,l} \phi_{0,k}| \frac{C_N}{(\lambda_k \lambda_l)^N} \leq C_N \left(\sum_k \frac{|\phi_{0,k}|}{\lambda_k^N} \right)^2 \leq C \|\phi_0\|_{H^{-1}(\Omega)}^2$$

when N is chosen large enough so that $\lambda_k^{1/2-N}$ is summable, which is always possible (for instance using Weyl law). □

The following result can be proved thanks to the arguments developed in the proof of Proposition 16 (in a simpler way, because \mathcal{K}^* does not appear anymore), together with Proposition 17 and the time-oscillation of $\partial_\nu \phi$.

PROPOSITION 18. — *Let Ω be an open subset of \mathbb{R}^2 , Γ be an open subset of $\partial\Omega$ and $0 < \tilde{T} < T < \infty$. If the Schrödinger equation on Ω is weakly observable on $(0, \tilde{T}) \times \Gamma$ then, for every non-zero $\phi_0 \in H_0^1(\Omega) \cap T_S \varphi_R$,*

$$\Im(\partial_\nu \phi) \neq 0 \quad \text{on } (0, T) \times \Gamma_c,$$

where $\phi(t) := e^{it\Delta_D} \phi_0$.

Proof. — Since the method is classical and was already performed in a more complicated case, we only detail the difference with Proposition 16. We define similarly

$$N_T := \{ \phi_0 \in H_0^1(\Omega) ; \Im(\partial_\nu \phi) = 0 \text{ in } L^2((0, T) \times \Gamma_c) \},$$

where $\phi(t) := e^{it\Delta_D} \phi_0$.

Following Step 1, we obtain that if $N_T \neq \{0\}$, N_T only contains one eigenfunction. Assume $\varphi \in N_T \setminus \{0\}$ is an eigenfunction of Δ_D . Then, $e^{it\Delta_D} \varphi = e^{-it\lambda} \varphi$. Note that the Dirichlet boundary condition implies $\lambda \neq 0$ and therefore $\varphi \in N_T$ implies $\partial_\nu \varphi = 0$ on Γ_c . By unique continuation for eigenfunctions, we get $\varphi = 0$, a contradiction. □

REMARK 3. — This result would be false if we considered Neumann boundary conditions. The real constants solutions $\Phi(t) = c$ with $c \in \mathbb{R}$ create a zero observation. The system is therefore not controllable with a real control. Yet, the space of non-controllable data is only one dimensional.

Then, arguing by contradiction as in the proof of Proposition 15, we obtain the following observability Proposition which directly implies Theorem 3 by the HUM method (a variant with \mathbb{R} -linear vector space).

PROPOSITION 19. — *Let Ω be an open subset of \mathbb{R}^2 , Γ be an open subset of $\partial\Omega$ and $0 < \tilde{T} < T < \infty$. If the Schrödinger equation on Ω is weakly observable on $(0, \tilde{T}) \times \Gamma$ then, there exists $\mathcal{C}_1 > 0$ such that*

$$(50) \quad \|\phi_0\|_{H_0^1(\Omega)} \leq \mathcal{C}_1 \|\Im(\partial_\nu \phi)\|_{L^2((0, T) \times \Gamma)}, \quad \forall \phi_0 \in H_0^1(\Omega, \mathbb{C}),$$

where $\phi(t) := e^{it\Delta_D} \phi_0$.

7. NONLINEAR EQUATION ON A RECTANGLE

The goal of this section is to prove Theorem 4. Note that the system (4) may be written

$$(51) \quad \begin{cases} i\partial_t\psi = -\Delta\psi + (-\Delta_N + 1)^{-1}(\varepsilon|\psi|^2)\psi + v\psi, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu\psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \psi(0, x) = \psi_0(x), & x \in \Omega, \\ (-\Delta + 1)v = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu v(t, x) = g(t, x)1_{\Gamma_c}(x), & (t, x) \in (0, T) \times \partial\Omega. \end{cases}$$

In Section 7.1, we prove a smoothing effect. In Section 7.2, we prove the well-posedness of the system (51) in functional spaces appropriate for the controllability problem. In Section 7.3, we prove the C^1 -regularity of the end-point map. In Section 7.4, we prove the controllability of the linearized system. In Section 7.4.5, we prove Theorem 4 by applying the inverse mapping theorem.

7.1. SMOOTHING EFFECT ON THE SOURCE TERM. — We introduce the operator \mathcal{A} defined by

$$D(\mathcal{A}) := H_N^2(\Omega, \mathbb{C}), \quad \mathcal{A}\phi := -\Delta\phi + \frac{2\varepsilon}{\pi L}(-\Delta_N + 1)^{-1}(\Re(\phi)).$$

On $H_N^2(\Omega, \mathbb{C})$, we use the norm

$$\|\varphi\|_{H_N^2} := \|(-\Delta + 1)\varphi\|_{L^2(\Omega)} = \left(\sum_{p,n \in \mathbb{N}} |(p^2 + (n\pi/L)^2 + 1)\langle \varphi, \xi_{p,n} \rangle|^2 \right)^{1/2},$$

where $\langle \cdot, \cdot \rangle$ is the usual $L^2(\Omega)$ -scalar product and

$$(52) \quad \begin{aligned} \xi_{p,n}(x_1, x_2) &= \zeta_p(x_1)\phi_n(x_2) \\ \zeta_p(x_1) &= \begin{cases} 1/\sqrt{\pi} & \text{if } p = 0, \\ \sqrt{2/\pi} \cos(px_1) & \text{if } p \in \mathbb{N}^*, \end{cases} & \phi_n(x_2) &= \begin{cases} 1/\sqrt{L} & \text{if } n = 0, \\ \sqrt{2/L} \cos(n\pi x_2/L) & \text{if } n \neq 0. \end{cases} \end{aligned}$$

The goal of this section is the proof of the following result.

THEOREM 7. — *Let $T > 0$, $L > 0$, $\Omega = (0, \pi) \times (0, L)$ and Γ_c be an open subset of $\partial\Omega$ such that $\bar{\Gamma}_c$ does not contain any vertex of Ω . For every $\psi_0 \in H_N^2(\Omega)$, $\mu_1 \in L^2((0, T), H^{3/2}(\Omega))$ and $\mu_2 \in L^1((0, T), H_N^2(\Omega))$ such that*

$$(53) \quad (-\Delta + 1)\mu_1 \equiv 0, \quad \partial_\nu\mu_1 \in L^2((0, T) \times \partial\Omega), \quad \text{Supp}(\partial_\nu\mu_1) \subset \Gamma_c,$$

the solution of

$$\begin{cases} (i\partial_t - \mathcal{A})\psi(t, x) = (\mu_1 + \mu_2)(t, x), & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu\psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \psi(0, x) = \psi_0(x), & x \in \Omega, \end{cases}$$

satisfies $\psi \in C^0([0, T], H_N^2(\Omega))$. Moreover, there exists a constant $C > 0$ (independent of ψ_0, μ_1, μ_2) such that

$$(54) \quad \|\psi\|_{L^0([0, T], H_N^2(\Omega))} \leq C(\|\psi_0\|_{H_N^2(\Omega)} + \|\partial_\nu \mu_1\|_{L^2((0, T) \times \Gamma_c)} + \|\mu_2\|_{L^1((0, T), H_N^2(\Omega))}).$$

REMARK 4. — By Proposition 7, $\partial_\nu \mu_1|_{\Gamma_j}$ is well-defined in $L^2((0, T), H^{-3/2-\varepsilon}(\Gamma_j))$, for $j = 1, \dots, 4$. This gives a sense to the second assumption in (53) i.e., $\partial_\nu \mu_1|_{\Gamma_j} \in L^2((0, T) \times \Gamma_j)$ for $j = 1, \dots, 4$.

To prove Theorem 7, we need the following preliminary result.

PROPOSITION 20. — Under the assumptions of the previous statement, the map

$$\mathcal{H} : t \longmapsto \int_0^t e^{-i\Delta_N s} \mu_1(s) ds$$

belongs to $C^0([0, T], H_N^2(\Omega))$ and

$$(55) \quad \|\mathcal{H}\|_{L^\infty((0, T), H_N^2(\Omega))} \leq C\|\partial_\nu \mu_1\|_{L^2((0, T) \times \Gamma_c)},$$

for some constant $C = C(T) > 0$ independent of μ_1 .

Proof of Proposition 20. — The proof is quite close to that of Theorem 5 on the rectangle. We only perform Step 1, since the step 2 of regularization is very similar. Actually, following similar arguments as Step 2, we can first assume $\partial_\nu \mu_1|_{\partial\Omega} \in C_0^\infty((0, T) \times \Gamma_c)$ which gives by elliptic regularity $\mu_1 \in C^\infty((0, T) \times \Omega)$. The inequality (55) that we obtain for smooth functions will then allow us to get the same result under the above regularity assumptions.

So, up to now, we assume that μ_1 is regular enough. By linearity, we may assume that $\Gamma_c \subset (a, b) \times \{0\}$ with $0 < a < b < \pi$. We have

$$\begin{aligned} S = \|\mathcal{H}(t)\|_{H_N^2(\Omega)}^2 &= \sum_{p, n \in \mathbb{N}} \left| (p^2 + (n\pi/L)^2 + 1) \int_0^t \langle \mu_1(s), \xi_{p, n} \rangle e^{-i(p^2 + (n\pi/L)^2)s} ds \right|^2 \\ &= \sum_{p, n \in \mathbb{N}} \left| \int_0^t \langle \mu_1(s), (-\Delta + 1)\xi_{p, n} \rangle e^{-i(p^2 + (n\pi/L)^2)s} ds \right|^2. \end{aligned}$$

We can apply the Green formula because μ_1 and $\xi_{p, n}$ are smooth. This leads to

$$\begin{aligned} S &= \sum_{p, n=0}^\infty \left| \int_0^t e^{-i(p^2 + (n\pi/L)^2)s} \int_{\Gamma_c} \partial_\nu \mu_1(s, x) \xi_{p, n}(x) d\sigma(x) ds \right|^2 \\ &\leq C \sum_{p=0}^\infty \sum_{n=0}^\infty \left| \int_0^{T_0} 1_{(0, t)}(s) e^{-ip^2 s} \int_0^\pi \partial_{x_2} \mu_1(s, x_1, 0) \cos(px_1) dx_1 e^{-i(n\pi/L)^2 s} ds \right|^2 \\ &\leq C \sum_{p=0}^\infty \int_0^{T_0} \left| 1_{(0, t)}(s) e^{-ip^2 s} \int_0^\pi \partial_{x_2} \mu_1(s, x_1, 0) \cos(px_1) dx_1 \right|^2 ds \end{aligned}$$

thanks to the Bessel-Parseval inequality in $L^2(0, T_0)$ with $T_0 \geq T$ of the form $2mL^2/\pi$ with $m \in \mathbb{N}^*$. Thus, we have

$$\begin{aligned} S &\leq C \sum_{p=0}^{\infty} \int_0^t \left| \int_0^\pi \partial_{x_2} \mu_1(s, x_1, 0) \cos(px_1) dx_1 \right|^2 ds \\ &\leq C \int_0^t \sum_{p=0}^{\infty} \left| \int_0^\pi \partial_{x_2} \mu_1(s, x_1, 0) \cos(px_1) dx_1 \right|^2 ds \\ &\leq C \int_0^t \|\partial_{x_2} \mu_1(s, \cdot, 0)\|_{L^2(0, \pi)}^2 ds \end{aligned}$$

thanks to the orthogonality of the functions $(\cos(px_1))_{p \in \mathbb{N}}$ in $L^2(0, \pi)$.

Note that μ_1 can be computed explicitly in terms of $\partial_\nu \mu_1|_{\Gamma_c}$ thanks to the rectangular form of the domain Ω . This explicit expression shows the existence of a constant $C = C(\Omega) > 0$ such that

$$\|\mu_1\|_{L^2((0,t), H^1(\Omega))}^2 \leq C \|\partial_\nu \mu_1\|_{L^2((0,t) \times \partial\Gamma_c)}, \quad \forall t \in (0, T).$$

Therefore,

$$\|\mathcal{H}(t)\|_{H_N^2(\Omega)}^2 \leq C(T) \|\partial_\nu \mu_1\|_{L^2((0,t) \times \partial\Gamma_c)}, \quad \forall t \in (0, T).$$

This inequality proves that the map \mathcal{H} takes values in $H_N^2(\Omega)$ on $[0, T]$ and that $\mathcal{H} : [0, T] \rightarrow H_N^2(\Omega)$ is continuous at $t = 0$. The same proof shows that \mathcal{H} is continuous at any $t \in (0, T]$. □

Proof of Theorem 7. — By the Duhamel formula, we have

$$(56) \quad \psi(t) = e^{i\Delta_N t} \psi_0 - i \int_0^t e^{i\Delta_N(t-s)} (\tilde{\mu}_2(s) + \mu_1(s)) ds,$$

where

$$\tilde{\mu}_2(t) := \mu_2(t) - \frac{2\varepsilon}{\pi L} (-\Delta_N + 1)^{-1} (\Re \psi).$$

In particular, $\tilde{\mu}_2 \in L^1((0, T), H_N^2(\Omega))$ and

$$(57) \quad \|\tilde{\mu}_2\|_{L^1((0,T), H_N^2(\Omega))} \leq \|\mu_2\|_{L^1((0,T), H_N^2(\Omega))} + \frac{2\varepsilon T}{\pi L} \|\psi\|_{L^\infty((0,T), L^2(\Omega))}.$$

For every $\tau \in \mathbb{R}$, the operator $e^{i\Delta_N \tau}$ is an isometry of $H_N^2(\Omega)$ thus, the first two terms in the right-hand side of (56) belong to $C^0([0, T], H_N^2(\Omega))$. The third term also does, by Proposition 20, thus $\psi \in C^0([0, T], H_N^2(\Omega))$.

Step 1. — Proof of (54) when T is small enough so that

$$(58) \quad \frac{2\varepsilon T}{\pi L} \leq \frac{1}{2}.$$

We deduce from (56) that, for every $t \in (0, T)$

$$\|\psi(t)\|_{H_N^2(\Omega)} \leq \|\psi_0\|_{H_N^2(\Omega)} + \int_0^t \|\tilde{\mu}_2(s)\|_{H_N^2(\Omega)} ds + \left\| \int_0^t e^{-i\Delta_N s} \mu_1(s) ds \right\|_{H_N^2(\Omega)}.$$

Using (57), Proposition 20 and (58) we obtain

$$\|\psi\|_{L^\infty((0,T), H_N^2(\Omega))} \leq 2(\|\psi_0\|_{H_N^2(\Omega)} + \|\mu_2\|_{L^1((0,T), H_N^2(\Omega))} + C \|\partial_\nu \mu_1\|_{L^2((0,T) \times \Gamma_c)}).$$

Step 2. — Proof of (54) when T is arbitrary. Let $0 = T_1 < T_2 < \dots < T_m = T$ be a subdivision of $[0, T]$ such that $2\varepsilon(T_{k+1} - T_k)/\pi L \leq 1/2$ for $k = 1, \dots, m-1$. Applying Step 1 on each subinterval (T_k, T_{k+1}) we obtain (54). \square

7.2. WELL-POSEDNESS OF THE NONLINEAR SCHRÖDINGER-POISSON SYSTEM. — The goal of this section is the proof of the following result.

PROPOSITION 21. — *Let $T > 0$. There exists $\rho, \delta > 0$ such that, for every $\psi_0 \in H_N^2(\Omega)$, $g \in L^2((0, T) \times \partial\Omega)$ with $\|\psi_0 - \psi_{\text{ref}}(0)\|_{H_N^2} < \rho$ and $\|g\|_{L^2((0, T) \times \partial\Omega)} < \delta$, there exists a unique solution $\psi \in C^0([0, T], H_N^2(\Omega))$ of (51). Moreover $\|\psi(t)\|_{L^2(\Omega)} = \|\psi_0\|_{L^2(\Omega)}$ for every $t \in [0, T]$.*

We search ψ in the form

$$\psi(t, x, y) = \frac{e^{-i\varepsilon t/\pi L}}{\sqrt{\pi L}} (1 + \zeta(t, x, y)),$$

where ζ is the solution of

$$(59) \quad \begin{cases} (i\partial_t - \mathcal{A})\zeta = \Phi(\zeta) + v(1 + \zeta), & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu \zeta(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \zeta(0, x) = \zeta_0(x), & x \in \Omega, \\ (-\Delta + 1)v(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu v(t, x) = g(t, x)1_\Gamma(x), & (t, x) \in (0, T) \times \partial\Omega, \end{cases}$$

and

$$(60) \quad \Phi(\zeta) := \frac{2\varepsilon}{\pi L} (-\Delta_N + 1)^{-1} (\Re(\zeta))\zeta + \frac{\varepsilon}{\pi L} (-\Delta_N + 1)^{-1} (|\zeta|^2)(1 + \zeta).$$

Proposition 21 is a consequence of the following result.

PROPOSITION 22. — *Let $T > 0$. There exists $\rho, \delta > 0$ such that, for every $\zeta_0 \in H_N^2(\Omega)$, $g \in L^2((0, T) \times \partial\Omega, \mathbb{R})$ with $\|\zeta_0\|_{H_N^2} < \rho$ and $\|g\|_{L^2((0, T) \times \partial\Omega)} < \delta$, there exists a unique weak solution $\zeta \in C^0([0, T], H_N^2(\Omega))$ of (59).*

Proof of Proposition 22. — Let $T > 0$.

Step 1. — We prove the existence of $C_0 > 0$ such that, for every $\zeta \in H_N^2(\Omega)$ and $v \in H^{3/2}(\Omega)$ satisfying $(-\Delta + 1)v = 0$ in Ω , then $(-\Delta + v)(v\zeta) \in L^2(\Omega)$ and

$$\|(-\Delta + v)(v\zeta)\|_{L^2(\Omega)} \leq C_0 \|v\|_{H^{3/2}(\Omega)} \|\zeta\|_{H_N^2(\Omega)}.$$

We have

$$(-\Delta + 1)[v\zeta] = -2\nabla v \cdot \nabla \zeta - v\Delta \zeta.$$

– We have $\Delta \zeta \in L^2(\Omega)$ and $v \in L^\infty(\Omega)$, by the Sobolev embedding (12), thus $v\Delta \zeta \in L^2(\Omega)$.

– We have $\nabla v \in H^{1/2}(\Omega)$ and $\nabla \zeta \in H^1(\Omega)$. From the Sobolev embedding (13), we deduce that ∇v and $\nabla \zeta$ belong to $L^4(\Omega)$. Thus $\nabla v \cdot \nabla \zeta \in L^2(\Omega)$.

Step 2. — Choice of R, δ, ρ . Let $C > 0$ be as in Proposition 20 and $C' > 0$ be the constant of the third statement of Proposition 7. Let $C_1, C_2, C_3 > 0$ be such that

$$\begin{aligned} \|\Phi(\zeta)\|_{H_N^2(\Omega)} &\leq C_1 \|\zeta\|_{H_N^2(\Omega)}^2, \quad \forall \zeta \in H_N^2(\Omega) \text{ such that } \|\zeta\|_{H_N^2(\Omega)} \leq 1, \\ \|\zeta\|_{L^\infty(\bar{\Omega})} &\leq C_2 \|\zeta\|_{H_N^2(\Omega)}, \quad \forall \zeta \in H_N^2(\Omega), \\ \|\Phi(\zeta) - \Phi(\tilde{\zeta})\|_{H_N^2(\Omega)} &\leq C_3 \|\zeta - \tilde{\zeta}\|_{H_N^2(\Omega)} \max \{ \|\zeta\|_{H_N^2(\Omega)}; \|\tilde{\zeta}\|_{H_N^2(\Omega)} \}, \\ &\quad \forall \zeta, \tilde{\zeta} \in H_N^2(\Omega) \text{ such that } \|\zeta\|_{H_N^2(\Omega)}, \|\tilde{\zeta}\|_{H_N^2(\Omega)} \leq 1, \end{aligned}$$

where $\Phi(\zeta)$ is defined in (60). Let

$$(61) \quad \begin{aligned} R &:= \min \left\{ 1; \frac{1}{4C_1T}; \frac{1}{4C_3T} \right\}, \quad \rho := \frac{R}{2} \\ \delta &:= \min \left\{ \frac{R}{4[C(1+C_2)+C_0C'\sqrt{T}]}; \frac{1}{4(CC_2+C_0C'T)} \right\}, \end{aligned}$$

and $\zeta_0 \in H_N^2(\Omega)$ be such that $\|\zeta_0\|_{H_N^2(\Omega)} < \rho$.

We consider the map

$$\begin{aligned} F : \overline{B_R}[C^0([0, T], H_N^2(\Omega))] &\longrightarrow \overline{B_R}[C^0([0, T], H_N^2(\Omega))] \\ \zeta &\longmapsto \xi, \end{aligned}$$

where $\xi := F(\zeta)$ is the solution of

$$\begin{cases} (i\partial_t - \mathcal{A})\xi = \Phi(\zeta) + v(1 + \zeta), & (t, x, y) \in (0, T) \times \Omega, \\ \partial_\nu \xi(t, x, y) = 0, & (t, x, y) \in (0, T) \times \partial\Omega, \\ \xi(0, x, y) = \zeta_0(x, y), & (x, y) \in \Omega. \end{cases}$$

The end of the proof consists in applying the Banach fixed point theorem to the map F .

Step 3. — We prove that F takes values in $\overline{B_R}[C^0([0, T], H_N^2)]$.

Let $\zeta \in \overline{B_R}[C^0([0, T], H_N^2)]$. We introduce the solution μ_3 of

$$\begin{cases} (-\Delta + 1)\mu_3(t, x_1, x_2) = (-\Delta + 1)(v\zeta)(t, x_1, x_2), & (t, x_1, x_2) \in (0, T) \times \Omega, \\ \partial_\nu \mu_3(t, x_1, x_2) = 0, & (t, x_1, x_2) \in (0, T) \times \partial\Omega \end{cases}$$

and the functions

$$\mu_1 := v + v\zeta - \mu_3, \quad \mu_2 := \mu_3 + \Phi(\zeta).$$

By Step 1, $(-\Delta + 1)(v\zeta) \in L^2((0, T) \times \Omega)$ and thus $\mu_3 \in L^2((0, T), H_N^2(\Omega))$. Therefore μ_1 and μ_2 satisfy the assumptions of Theorem 7 and $\xi := F(\zeta) \in C^0([0, T], H_N^2(\Omega))$.

By Proposition 20 and (61)

$$\begin{aligned} \|\xi\|_{L^\infty((0,T),H_N^2)} &\leq \|\zeta_0\|_{H_N^2} + C\|\partial_\nu\mu_1\|_{L^2((0,T)\times\Gamma_c)} \\ &\quad + \int_0^T (\|\mu_3(s)\|_{H_N^2} + \|\Phi[\zeta(s)]\|_{H_N^2}) ds \\ &\leq \frac{R}{2} + C\|g(1+\zeta)\|_{L^2((0,T)\times\Gamma_c)} \\ &\quad + \sqrt{T}\|(-\Delta+1)(v\zeta)\|_{L^2((0,T)\times\Omega)} + TC_1R^2 \\ &\leq \frac{3R}{4} + C\delta(1+C_2) + \sqrt{T}C_0C'\delta R \\ &\leq R. \end{aligned}$$

Step 4. — We prove that F is a contraction. Let $\zeta, \tilde{\zeta} \in \overline{B_R}[C^0([0, T], H_N^2)]$. By Proposition 20 and (61),

$$\begin{aligned} \|\xi - \tilde{\xi}\|_{L^\infty((0,T),H_N^2)} &\leq C\|\partial_\nu(\mu_1 - \tilde{\mu}_1)\|_{L^2((0,T)\times\Gamma_c)} \\ &\quad + \int_0^T (\|(\mu_3 - \tilde{\mu}_3)(s)\|_{H_N^2} + \|\Phi(\zeta(s)) - \Phi(\tilde{\zeta}(s))\|_{H_N^2}) ds \\ &\leq C\|g(\zeta - \tilde{\zeta})\|_{L^2((0,T)\times\Gamma_c)} + \sqrt{T}\|(-\Delta+1)[v(\zeta - \tilde{\zeta})]\|_{L^2((0,T)\times\Omega)} \\ &\quad + TC_3R\|\zeta - \tilde{\zeta}\|_{L^\infty((0,T),H_N^2)} \\ &\leq (C\delta C_2 + C_0C'\delta + TC_3R)\|\zeta - \tilde{\zeta}\|_{L^\infty((0,T),H_N^2)} \\ &\leq \frac{1}{2}\|\zeta - \tilde{\zeta}\|_{L^\infty((0,T),H_N^2)}. \quad \square \end{aligned}$$

7.3. C^1 -REGULARITY OF THE END-POINT MAP. — The end-point map is defined by

$$(62) \quad \begin{aligned} \Theta_T : [\mathcal{S} \cap H_N^2(\Omega, \mathbb{C})] \times L^2((0, T) \times \partial\Omega, \mathbb{R}) &\longrightarrow [\mathcal{S} \cap H_N^2(\Omega, \mathbb{C})]^2 \\ (\psi_0, g) &\longmapsto (\psi(0), \psi(T)), \end{aligned}$$

where ψ is the solution of (51). The goal of this section is to state the following result, which is a consequence of the estimate (54).

PROPOSITION 23. — *Let $T > 0$ and $\rho, \delta > 0$ be as in Proposition 21. The end-point map Θ_T defined by (62) is C^1 on*

$$\{(\psi_0, g) \in H_N^2 \times L^2((0, T) \times \partial\Omega); \|\psi_0 - \psi_{\text{ref}}(0)\|_{H_N^2} < \rho \text{ and } \|g\|_{L^2((0,T)\times\partial\Omega)} < \delta\}$$

and

$$d\Theta_T(\psi_{\text{ref}}(0), 0) \cdot (\Psi_0, G) = \left(\Psi_0, \frac{e^{-i\varepsilon T/\sqrt{\pi L}}}{\sqrt{\pi L}} \Psi(T) \right),$$

where Ψ is the solution of

$$(63) \quad \begin{cases} (i\partial_t - \mathcal{A})\Psi = V, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu \Psi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \Psi(0, x, y) = \Psi_0(x), & x \in \Omega, \\ (-\Delta + 1)V = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu V(t, x) = G(t, x)1_{\Gamma_c}(x), & (t, x) \in (0, T) \times \partial\Omega. \end{cases}$$

7.4. CONTROLLABILITY OF THE LINEARIZED SYSTEM. — Let

$$T_S := \left\{ \varphi \in L^2(\Omega, \mathbb{C}); \Re \left(\int_{\Omega} \varphi(x, y) \, dx dy \right) = 0 \right\}.$$

The goal of this section is the proof of the following result.

PROPOSITION 24. — *Let $T > 0$. There exists a continuous map*

$$L : (T_S \cap H_N^2(\Omega, \mathbb{C}))^2 \longrightarrow L^2((0, T) \times \partial\Omega) \\ (\Psi_0, \Psi_f) \longmapsto G$$

such that, for every $(\Psi_0, \Psi_f) \in (T_S \cap H_N^2(\Omega, \mathbb{C}))^2$ the solution of (63) satisfies $\Psi(T) = \Psi_f$.

7.4.1. *Hilbert Uniqueness Method.* — The proof of Proposition 24 requires the following observability result, where \mathcal{B} is defined by

$$D(\mathcal{B}) := H_N^2(\Omega, \mathbb{C}), \quad \mathcal{B}\phi := -\Delta\phi + i\frac{2\varepsilon}{\pi L}(-\Delta_N + 1)^{-1}(\mathfrak{I}(\phi)).$$

PROPOSITION 25. — *Let $T > 0$. There exists $\mathcal{C} > 0$ such that, for every $\phi_T \in L^2(\Omega)$, the solution of*

$$(64) \quad \begin{cases} (i\partial_t - \mathcal{B})\phi = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu\phi(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \phi(T, x) = \phi_T(x), & x \in \Omega, \end{cases}$$

satisfies

$$(65) \quad \|\phi_T\|_{L^2(\Omega)} \leq \mathcal{C} \|\mathfrak{I}(\phi)\|_{L^2((0, T) \times \Gamma_c)}.$$

Note that the solution of (64) can be computed explicitly (as in the case $\varepsilon = 0$) because \mathcal{B} preserves $\mathcal{C}\xi_{p,n}$. The boundary condition in (64) has to be understood in the sense of the semi-group. The proof of Proposition 25 relies on two intermediate results:

- the weak observability of (64), proved in Section 7.4.2 below,
- a unique continuation property, proved in Section 7.4.3.

Note that the trace $\mathfrak{I}(\phi)|_{\partial\Omega}$ is well-defined in $L^2((0, T) \times \partial\Omega)$ (see Lemma 1 below).

Proof of Proposition 24, assuming Proposition 25. — We want to prove the existence of a continuous right inverse for the continuous operator

$$F : L^2((0, T) \times \Gamma_c, \mathbb{R}) \longrightarrow T_S \\ G \longmapsto (-\Delta + 1)\Psi(T),$$

where Ψ is the solution of (63) with $\Psi_0 = 0$. By [12, Th. II.10 & II.19], it is equivalent to prove the existence of a constant $\mathcal{C} > 0$ such that

$$\|\phi_T\|_{L^2(\Omega)} \leq \mathcal{C} \|F^*(\phi_T)\|_{L^2((0, T) \times \Gamma_c)}, \quad \forall \phi_T \in T_S.$$

Thus, to deduce Proposition 24 from Proposition 25, it is sufficient to prove that

$$F^*(\phi_T) = -\mathfrak{S}(\phi)|_{\Gamma_c}, \quad \forall \phi_T \in T_S,$$

where ϕ is associated to ϕ_T by (64). The trace $\mathfrak{S}(\phi)|_{\Gamma_c}$ is defined as an extension, thus, it is sufficient to prove that

$$(66) \quad F^*(\phi_T) = -\mathfrak{S}(\phi)|_{\Gamma_c}, \quad \forall \phi_T \in T_S \cap C_c^\infty(\Omega).$$

Let $\phi_T \in T_S \cap C_c^\infty(\Omega)$. Then $\phi \in C^\infty([0, T], H_N^s(\Omega))$ for every $s > 0$, thus $\tilde{\phi} := (-\Delta + 1)\phi \in C^\infty([0, T], H_N^s(\Omega))$ for every $s > 0$ and

$$\begin{cases} (i\partial_t + \mathcal{B})\tilde{\phi} = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu \tilde{\phi}(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \tilde{\phi}(T, x) = (-\Delta + 1)\phi_T(x), & x \in \Omega, \end{cases}$$

because \mathcal{B} and $(-\Delta + 1)$ commute. The functions Ψ and $\tilde{\phi}$ are solutions in the sense of the semi-group, thus

$$\begin{aligned} \Re\langle \Psi(T), \tilde{\phi}(T) \rangle &= \mathfrak{S}\left(\int_0^T \int_\Omega V(t, x) \overline{\tilde{\phi}(t, x)} dx dt\right) \\ &= \mathfrak{S}\left(\int_0^T \int_\Omega V(t, x) \overline{(-\Delta + 1)\phi(t, x)} dx dt\right). \end{aligned}$$

Therefore,

$$\begin{aligned} \Re\langle F(G), \phi_T \rangle &= \Re\langle (-\Delta + 1)\Psi(T), \phi_T \rangle \\ &= \Re\langle \Psi(T), (-\Delta + 1)\phi_T \rangle \quad \text{because } \Psi(T), \phi_T \in H_N^2(\Omega) \\ &= \mathfrak{S}\left(\int_0^T \int_\Omega V(t, x) \overline{(-\Delta + 1)\phi(t, x)} dx dt\right) \\ &= -\int_0^T \int_{\Gamma_c} G(t, x) \mathfrak{S}(\phi(t, x)) d\sigma(x) dt. \end{aligned}$$

This proves (66). □

LEMMA 1. — *Let $T > 0$. There exists $C = C(T) > 0$ such that, for every $\phi_0 \in L^2(\Omega)$, the solution $\phi \in C^0([0, T], L^2(\Omega))$ of (64) satisfies*

$$\int_0^T \int_{\partial\Omega} |\phi(t, x)|^2 d\sigma(x) dt \leq C \|\phi_0\|_{L^2(\Omega)}^2.$$

Proof of Lemma 1

Step 1. — Proof when $\varepsilon = 0$. The following equality holds in $C^0([0, T], L^2(\Omega))$

$$\phi(t, x_1, x_2) = \sum_{p, n=0}^{\infty} \langle \phi_0, \xi_{p, n} \rangle e^{-i[p^2 + (n\pi/L)^2]t} \xi_{p, n}(x_1, x_2).$$

Using (52) and the orthogonality of $(\zeta_p)_{p \in \mathbb{N}}$ in $L^2(0, \pi)$, we get

$$\begin{aligned} \int_0^T \int_0^\pi |\phi(t, x_1, 0)|^2 dx_1 dt &= \int_0^T \int_0^\pi \left| \sum_{p=0}^\infty \left(\sum_{n=0}^\infty \langle \phi_0, \xi_{p,n} \rangle e^{-i[p^2 + (n\pi/L)^2]t} \right) \zeta_p(x_1) \right|^2 dx_1 dt \\ &= \int_0^T \sum_{p=0}^\infty \left| \sum_{n=0}^\infty \langle \phi_0, \xi_{p,n} \rangle e^{-i[p^2 + (n\pi/L)^2]t} \right|^2 dt \\ &= \sum_{p=0}^\infty \int_0^T \left| \sum_{n=0}^\infty \langle \phi_0, \xi_{p,n} \rangle e^{-i(n\pi/L)^2 t} \right|^2 dt \\ &\leq \sum_{p=0}^\infty C_{\text{Ing}} \sum_{n=0}^\infty |\langle \phi_0, \xi_{p,n} \rangle|^2 = C_{\text{Ing}} \|\phi_0\|_{L^2(\Omega)}^2, \end{aligned}$$

where $C_{\text{Ing}}(T) > 0$ is the Ingham constant associated to the family $(e^{-i(n\pi/L)^2 t})_{n \in \mathbb{N}}$ in $L^2(0, T)$ (see [19]).

Step 2. — Proof when $\varepsilon \neq 0$. The following equality holds in $L^2(\Omega)$.

$$\phi(t) = e^{-i\Delta_N t} \phi_0 + \frac{2\varepsilon}{\pi L} \int_0^t e^{-i\Delta_N(t-s)} (-\Delta_N + 1)^{-1} (\Im(\phi(s))) ds, \quad \forall t \in [0, T].$$

By Step 1, the trace on $\partial\Omega$ of $e^{-i\Delta_N t} \phi_0$ has an $L^2((0, T) \times \partial\Omega)$ -norm bounded by $\sqrt{C_{\text{Ing}}} \|\phi_0\|_{L^2(\Omega)}$. The second term of the right-hand side belongs to $C^0([0, T], H_N^2(\Omega))$ and its $L^\infty((0, T), H_N^2(\Omega))$ -norm is bounded by

$$\frac{2\varepsilon}{\pi L} \int_0^T \|\phi(s)\|_{L^2(\Omega)} ds = \frac{2\varepsilon T}{\pi L} \|\phi_0\|_{L^2(\Omega)}.$$

Thus, this term as a trace on $\partial\Omega$ with an $L^2((0, T) \times \partial\Omega)$ -norm that satisfies a similar estimate. \square

7.4.2. Weak observability. — The goal of this section is the proof of the following result.

PROPOSITION 26. — *Let $T > 0$. There exists $\mathcal{C}' = \mathcal{C}'(T) > 0$ such that, for every $\phi_T \in L^2(\Omega)$, the solution of (64) satisfies*

$$(67) \quad \|\phi_T\|_{L^2(\Omega)} \leq \mathcal{C}' (\|\Im(\phi)\|_{L^2((0, T) \times \Gamma_c)} + \|\phi_T\|_{H^{-2}(\Omega)}).$$

Proof of Proposition 26. — We recall Tenenbaum and Tucsnak's result [36, Th. 1.1]: for every $T > 0$ and every non-empty open subset Γ of $\partial\Omega$, there exists $\mathcal{C}_0 = \mathcal{C}_0(T, \Gamma) > 0$ such that, for every $\phi_T \in L^2(\Omega)$, the solution $\tilde{\phi}$ of

$$(68) \quad \begin{cases} (i\partial_t + \Delta)\tilde{\phi}(t, x) = 0, & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu \tilde{\phi}(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \tilde{\phi}(T, x) = \phi_T(x), & x \in \Omega, \end{cases}$$

satisfies

$$(69) \quad \|\phi_T\|_{L^2(\Omega)} \leq \mathcal{C}_0 \|\tilde{\phi}\|_{L^2((0, T) \times \Gamma)}.$$

Step 1. — We prove that, for every $T > 0$, there exists $\mathcal{C}_1 = \mathcal{C}_1(T) > 0$ such that for every $\phi_T \in L^2(\Omega)$, the solution $\tilde{\phi}$ of (68) satisfies

$$(70) \quad \|\phi_T\|_{L^2(\Omega)} \leq \mathcal{C}_1 (\|\Im(\tilde{\phi})\|_{L^2((0,T)\times\Gamma_c)} + \|\phi_T\|_{H^{-1}(\Omega)}).$$

Let $T > 0$ and $\rho \in C_c^\infty(\mathbb{R}, \mathbb{R}^+)$ be such that $\rho \equiv 1$ on $(T/3, 2T/3)$ and $\text{Supp}(\rho) \subset (0, T)$. Let $\phi_T \in L^2(\Omega)$. We have

$$\begin{aligned} \|\phi_T\|_{L^2(\Omega)}^2 &\leq \mathcal{C}_0^2 \int_{T/3}^{2T/3} \int_{\Gamma_c} |\tilde{\phi}(t, x)|^2 d\sigma(x) dt \quad \text{with } \mathcal{C}_0 = \mathcal{C}_0(T/3) \text{ by (69)} \\ &\leq \mathcal{C}_0^2 \int_{\mathbb{R}} \int_{\Gamma_c} |\rho(t)\tilde{\phi}(t, x)|^2 d\sigma(x) dt \\ &\leq \frac{\mathcal{C}_0^2}{2} \int_{\mathbb{R}} \int_{\Gamma_c} (|\rho(t)\tilde{\phi}(t, x)|^2 + |\rho(t)\overline{\tilde{\phi}(t, x)}|^2) d\sigma(x) dt \\ &\leq \frac{\mathcal{C}_0^2}{2} \int_{\mathbb{R}} \int_{\Gamma_c} (|\rho(t)\Im(\tilde{\phi}(t, x))|^2 + 2\Re(\rho(t)^2\tilde{\phi}(t, x)^2)) d\sigma(x) dt. \end{aligned}$$

In order to get (70), it suffices to prove the existence of $C > 0$ such that

$$\int_{\mathbb{R}} \int_{\Gamma_c} \Re[\rho(t)^2\tilde{\phi}(t, x)^2] d\sigma(x) dt \leq C \|\phi_T\|_{H^{-1}(\Omega)}^2.$$

The function ρ^2 belongs to $C_c^\infty(\mathbb{R}, \mathbb{R})$, thus, for every $N \in \mathbb{N}^*$, there exists $C_N > 0$ such that

$$\left| \int_{\mathbb{R}} \rho(t)^2 e^{-i\omega t} dt \right| \leq \frac{C_N}{(1 + \omega)^N}, \quad \forall \omega > 0.$$

We have

$$\begin{aligned} &\left| \int_{\mathbb{R}} \int_{\Gamma_c} \rho(t)^2 \tilde{\phi}(t, x)^2 d\sigma(x) dt \right| \\ &= \left| \sum_{j, J \in \mathbb{N}^*} \phi_{T,j} \phi_{T,J} \left(\int_{\mathbb{R}} \rho(t)^2 e^{-(\lambda_j + \lambda_J)t} dt \right) \left(\int_{\Gamma_c} \varphi_j(x) \varphi_J(x) d\sigma(x) \right) \right| \\ &\leq \sum_{j, J \in \mathbb{N}^*} |\phi_{T,j} \phi_{T,J}| \frac{C_N}{(1 + \lambda_j + \lambda_J)^N} C \lambda_j^\alpha \lambda_J^\alpha \quad \text{for some } C, \alpha > 0 \\ &\leq CC_N \left(\sum_{j \in \mathbb{N}^*} |\phi_{T,j}| \frac{\lambda_j^\alpha}{(1 + \lambda_j)^N} \right)^2 \\ &\leq CC_N \left(\sum_{j \in \mathbb{N}^*} \frac{|\phi_{T,j}|^2}{1 + \lambda_j} \right)^2 \left(\sum_{j \in \mathbb{N}^*} \frac{\lambda_j^{2\alpha} (1 + \lambda_j)}{(1 + \lambda_j)^{2N}} \right)^2 \\ &\leq C \|\phi_T\|_{H^{-1}(\Omega)}^2 \quad \text{for } N \text{ large enough.} \end{aligned}$$

Step 2. — **Conclusion.** Working by contradiction, we assume the existence of a sequence $(\phi_T^n)_{n \in \mathbb{N}^*}$ such that the associated solutions ϕ^n of (64) satisfy

$$(71) \quad 1 = \|\phi_T^n\|_{L^2(\Omega)} > n (\|\Im(\phi^n)\|_{L^2((0,T)\times\Gamma_c)} + \|\phi_T^n\|_{H^{-2}(\Omega)}).$$

We introduce the solutions $\tilde{\phi}^n$ of (68) associated to final condition $\tilde{\phi}^n(T) = \phi_T^n$. Then $\xi^n := \phi^n - \tilde{\phi}^n$ solves

$$\begin{cases} (i\partial_t + \Delta)\xi^n = i\frac{2\varepsilon}{\pi L}(-\Delta_N + 1)^{-1}\mathfrak{I}(\phi^n), & (t, x) \in (0, T) \times \Omega, \\ \partial_\nu \xi^n(t, x) = 0, & (t, x) \in (0, T) \times \partial\Omega, \\ \xi^n(T, x) = 0. & x \in \Omega, \end{cases}$$

From (71), we deduce that

$$\phi_T^n \xrightarrow[n \rightarrow \infty]{} 0 \quad \text{in } H^{-2}(\Omega).$$

Moreover this sequence is bounded in $L^2(\Omega)$, thus, by interpolation

$$\phi_T^n \xrightarrow[n \rightarrow \infty]{} 0 \quad \text{in } H^{-1}(\Omega).$$

Therefore,

$$(-\Delta_N + 1)^{-1}\mathfrak{I}(\phi^n) \xrightarrow[n \rightarrow \infty]{} 0 \quad \text{in } L^\infty((0, T), H^1(\Omega))$$

and

$$\xi^n \xrightarrow[n \rightarrow \infty]{} 0 \quad \text{in } L^\infty((0, T), H^1(\Omega)).$$

As a consequence $\mathfrak{I}(\xi^n)|_{\Gamma_c} \xrightarrow[n \rightarrow \infty]{} 0$ in $L^\infty((0, T) \times \Gamma_c)$. Moreover, $\mathfrak{I}(\phi^n)|_{\Gamma_c} \xrightarrow[n \rightarrow \infty]{} 0$ in $L^2((0, T) \times \Gamma_c)$ by (71), thus $\mathfrak{I}(\tilde{\phi}^n)|_{\Gamma_c} \xrightarrow[n \rightarrow \infty]{} 0$ in $L^2((0, T) \times \Gamma_c)$. From Step 1, we get

$$1 = \|\phi_T^n\|_{L^2(\Omega)} \leq \mathcal{C}_1 (\|\mathfrak{I}(\tilde{\phi}^n)\|_{L^2((0, T) \times \Gamma_c)} + \|\phi_T^n\|_{H^{-1}(\Omega)}) \xrightarrow[n \rightarrow \infty]{} 0,$$

which is a contradiction. \square

7.4.3. Unique continuation. — The goal of this section is to prove the following result.

PROPOSITION 27. — *We assume that ε satisfies (6). Let $T > 0$, Γ_c be an open subset of $\partial\Omega$ and $\phi_T \in \mathcal{T}_S \setminus \{0\}$ and ϕ be the solution of (64). Then $\mathfrak{I}(\phi)|_{\Gamma_c}$ does not identically vanish on $(0, T) \times \Gamma_c$.*

Proof of Proposition 27. — To simplify, we assume that $\Gamma_c = (a, b) \times \{0\}$, where $0 < a < b < 1$. We introduce

$$N_T := \{\phi_0 \in \mathcal{T}_S; \mathfrak{I}(\phi) = 0 \text{ on } (0, T) \times \Gamma_c\},$$

where $\phi(t) := S(t)\phi_0$ solves (64) with initial condition at $t = 0$: $\phi(0) = \phi_0$. N_T is an \mathbb{R} -vector subspace of $L^2(\Omega)$. Working by contradiction, we assume that $N_T \neq \{0\}$.

Step 1. — We prove that N_T is stable by $(-i\mathcal{B})$ and has finite dimension. Let $\phi_0 \in N_T$. Since $\phi_0 \in L^2(\Omega)$, then $\phi_\varepsilon := (S(\varepsilon)\phi_0 - \phi_0)/\varepsilon$ is bounded in $H_N^{-2}(\Omega)$ when $[\varepsilon \rightarrow 0]$. Moreover, for $\varepsilon < T/2$, $\phi_\varepsilon \in N_{T/2}$. Applying (67) to ϕ_ε , we get

$$\|\phi_\varepsilon\|_{L^2(\Omega)} \leq \mathcal{C}(T/2) \|\phi_\varepsilon\|_{H_N^{-2}(\Omega)}, \quad \forall \varepsilon \in (0, T/2).$$

Thus ϕ_ε is bounded in $L^2(\Omega)$ when $[\varepsilon \rightarrow 0]$. Therefore $\phi_0 \in H^2_N(\Omega)$. Indeed, by Fatou's lemma,

$$\begin{aligned} \|\phi_0\|_{H^2_N(\Omega)}^2 &= \sum_{k=1}^\infty |\lambda_k \langle \phi_0, \varphi_k \rangle|^2 = \sum_{k=1}^\infty \liminf_{\varepsilon \rightarrow 0} \left| \frac{e^{-i\lambda_k \varepsilon} - 1}{\varepsilon} \langle \phi_0, \varphi_k \rangle \right|^2 \\ &\leq \liminf_{\varepsilon \rightarrow 0} \sum_{k=1}^\infty \left| \frac{e^{-i\lambda_k \varepsilon} - 1}{\varepsilon} \langle \phi_0, \varphi_k \rangle \right|^2 \\ &\leq \liminf_{\varepsilon \rightarrow 0} \|\phi_\varepsilon\|_{L^2(\Omega)}^2 < \infty. \end{aligned}$$

Let $T' \in (0, T)$. The map

$$\begin{aligned} \mathcal{J}_{T'} : L^2(\Omega) &\longrightarrow L^2((0, T') \times \Gamma_c) \\ \phi_0 &\longmapsto \mathfrak{I}(\phi)|_{\Gamma_c} \end{aligned}$$

is continuous (see Lemma 1). Moreover, $\mathcal{J}_{T'}(\phi_\varepsilon) = 0$ for every $\varepsilon \in (0, T - T')$ and $\phi_\varepsilon \rightarrow -i\mathcal{B}\phi_0$ in $L^2(\Omega)$ when $[\varepsilon \rightarrow 0]$ thus $\mathcal{J}_{T'}(-i\mathcal{B}\phi_0) = 0$. This holds for every $T' \in (0, T)$, therefore $-i\mathcal{B}\phi_0 \in N_T$. We have proved that N_T is stable by $-i\mathcal{B}$ and only contains smooth functions in $D[\mathcal{B}^s]$, $\forall s \in \mathbb{N}^*$. Applying estimate (67) to $(-i\mathcal{B}\phi_0)$, we get

$$\|\mathcal{B}\phi_0\|_{L^2(\Omega)} \leq C' \|\mathcal{B}\phi_0\|_{H^{-2}(\Omega)}.$$

This shows that the unit ball of N_T is compact (for the $L^2(\Omega)$ -topology), thus N_T has finite dimension, by the Riesz theorem.

Step 2. — We prove that N_T contains a finite sum of functions $\xi_{p,n}$. The finite dimensional \mathbb{R} -vector space N_T is stable by the operator $-i\mathcal{B}$, thus there exists $\alpha, \beta \in \mathbb{R}$ such that

$$\text{Ker}[P(-i\mathcal{B})] \cap N_T \neq \{0\}, \quad \text{where } P(X) = X^2 + \alpha X + \beta.$$

Let $\phi_0 \in \text{Ker}[P(-i\mathcal{B})] \cap N_T$. Let $a_{p,n}, b_{p,n} \in \mathbb{R}$ be such that

$$\langle \phi_0, \xi_{p,n} \rangle = a_{p,n} + ib_{p,n}, \quad \forall p, n \in \mathbb{N}.$$

Note that $a_{0,0} = 0$ because $\phi_0 \in T_S$. Explicit computations show that the relation $P(-i\mathcal{B})\phi_0 = 0$ reads

$$\sum_{p,n=0}^\infty \left(P[-i\lambda_{p,n}]a_{p,n} + iP \left[-i \left(\lambda_{p,n} + \frac{2\varepsilon}{\pi L(\lambda_{p,n} + 1)} \right) \right] b_{p,n} \right) \xi_{p,n} = 0,$$

where $\lambda_{p,n} := p^2 + (n\pi/L)^2$. Thus

$$P[-i\lambda_{p,n}]a_{p,n} = P \left[-i \left(\lambda_{p,n} + \frac{2\varepsilon}{\pi L(\lambda_{p,n} + 1)} \right) \right] b_{p,n} = 0, \quad \forall p, n \in \mathbb{N}.$$

The polynomial P has at most two roots and the two quantities

$$\lambda_{p,n} \quad \text{and} \quad \lambda_{p,n} + 2\varepsilon/\pi L(\lambda_{p,n} + 1)$$

diverge when $\|(n, p)\| \rightarrow \infty$, thus only a finite number of coefficients $\langle \phi_0, \xi_{p,n} \rangle$ can be different from zero.

Step 3. — Conclusion. Let $N \in \mathbb{N}$ be such that $\phi_0 = \sum_{n,p=0}^N \langle \phi_0, \xi_{p,n} \rangle \xi_{n,p}$.

Step 3.1. — Assumption (6) implies that $\lambda_{p,n} + 2\varepsilon/\pi L(\lambda_{p,n} + 1) > 0$ for every $n, p \in \mathbb{N}$. Explicit computations show that

$$\Im[\phi(t, x_1, 0)] = \sum_{p,n=0}^N (b_{p,n} \cos(\omega_{p,n}t) - \gamma_{p,n} a_{p,n} \sin(\omega_{p,n}t)) \zeta_p(x_1),$$

where

$$\omega_{p,n} := \sqrt{\lambda_{p,n} \left(\lambda_{p,n} + \frac{2\varepsilon}{\pi L(\lambda_{p,n} + 1)} \right)}, \quad \forall p, n \in \mathbb{N},$$

$$\gamma_{p,n} := \sqrt{\frac{\lambda_{n,p}}{\lambda_{p,n} + 2\varepsilon/\pi L(\lambda_{p,n} + 1)}}.$$

The continuous function $(t, x) \mapsto \Im[\phi(t, x_1, 0)]$ vanishes on $(0, T) \times (a, b)$ because $\phi_0 \in N_T$. In particular, for every $t \in [0, T]$, the function $x_1 \mapsto \Im[\phi(t, x_1, 0)]$ vanishes on (a, b) . But the functions $(x_1 \mapsto \cos(px_1))_{0 \leq p \leq N}$ are linearly independent on (a, b) , thus

$$\sum_{n=0}^N (b_{p,n} \cos(\omega_{p,n}t) - \gamma_{p,n} a_{p,n} \sin(\omega_{p,n}t)) = 0, \quad \forall t \in (0, T), 1 \leq p \leq N.$$

We notice that Assumption (6) on ε imply

$$\omega_{p,n}^2 \neq \omega_{p,m}^2, \quad \forall p, n, m \in \mathbb{N} \text{ such that } n \neq m \text{ and } (p, n), (p, m) \neq (0, 0),$$

where

$$\omega_{p,n}^2 := (p^2 + (n\pi/L)^2) \left(p^2 + (n\pi/L)^2 + \frac{2\varepsilon}{\pi L(p^2 + n\pi/L + 1)} \right), \quad \forall p, n, m \in \mathbb{N}.$$

Under Assumption (6), the map $f(s) := s(s + 2\varepsilon/\pi L(s + 1))$ is strictly increasing on $[m, \infty)$ and thus the above property holds.

So, the frequencies $\{\omega_{p,n}; 0 \leq n \leq N\}$ are all different for every $p \in \mathbb{N}$. Thus, the previous relations imply that $b_{p,n} = a_{p,n} = 0$ for every $0 \leq p, n \leq N$, i.e., $\phi_0 = 0$, which is a contradiction. \square

7.4.4. Observability. — Proposition 25 follows from Propositions 26 and 27, by working as in the proof of Proposition 15.

7.4.5. Controllability of the nonlinear PDE. — The proof of Theorem 4 consists in applying the inverse mapping theorem to the end-point map Θ defined in (62), as in Section 4.2.

REFERENCES

- [1] N. ANANTHARAMAN, M. LÉAUTAUD & F. MACIÀ — “Wigner measures and observability for the Schrödinger equation on the disk”, *Invent. Math.* **206** (2016), no. 2, p. 485–599.
- [2] F. D. ARARUNA, E. CERPA, A. MERCADO & M. SANTOS — “Internal null controllability of a linear Schrödinger-KdV system on a bounded interval”, *J. Differential Equations* **260** (2016), no. 1, p. 653–687.
- [3] J. M. BALL, J. E. MARSDEN & M. SLEMROD — “Controllability for distributed bilinear systems”, *SIAM J. Control Optim.* **20** (1982), no. 4, p. 575–597.

- [4] K. BEAUCHARD – “Local controllability of a 1-D Schrödinger equation”, *J. Math. Pures Appl. (9)* **84** (2005), no. 7, p. 851–956.
- [5] K. BEAUCHARD & C. LAURENT – “Local controllability of 1D linear and nonlinear Schrödinger equations”, *J. Math. Pures Appl. (9)* **94** (2010), no. 5, p. 520–554.
- [6] ———, “Bilinear control of high frequencies for a 1D Schrödinger equation”, *Math. Control Signals Systems* (to appear), [hal-01333625](#).
- [7] J. BERGH & J. LÖFSTRÖM – *Interpolation spaces. An introduction*, Grundlehren Math. Wiss., vol. 223, Springer-Verlag, Berlin-New York, 1976.
- [8] U. BOSCAIN, M. CAPONIGRO, T. CHAMBRION & M. SIGALOTTI – “A weak spectral condition for the controllability of the bilinear Schrödinger equation with application to the control of a rotating planar molecule”, *Comm. Math. Phys.* **311** (2012), no. 2, p. 423–455.
- [9] U. BOSCAIN, M. CAPONIGRO & M. SIGALOTTI – “Multi-input Schrödinger equation: controllability, tracking and application to the quantum angular momentum”, *J. Differential Equations* **256** (2014), no. 11, p. 3524–3551.
- [10] U. BOSCAIN, T. CHAMBRION & M. SIGALOTTI – “On some open questions in bilinear quantum control”, in *Proceeding ECC (Zürich, 2013)*, IEEE, 2013, [hal-00818216](#).
- [11] N. BOUSSAÏD, M. CAPONIGRO & T. CHAMBRION – “Regular propagators of bilinear quantum systems”, [hal-01016299](#), 2014.
- [12] H. BREZIS – *Analyse fonctionnelle. Théorie et applications*, Collection Mathématiques Appliquées pour la Maîtrise, Masson, Paris, 1983.
- [13] N. BURQ – “Contrôle de l’équation des ondes dans des ouverts peu réguliers”, *Asymptot. Anal.* **14** (1997), p. 157–191.
- [14] ———, “Mesures semi-classiques et mesures de défaut”, in *Séminaire Bourbaki, Vol. 1996/97*, Astérisque, vol. 245, Société Mathématique de France, Paris, 1997, Exp. No. 826, p. 167–195.
- [15] N. BURQ & M. ZWORSKI – “Geometric control in the presence of a black box”, *J. Amer. Math. Soc.* **17** (2004), no. 2, p. 443–471.
- [16] T. CHAMBRION, P. MASON, M. SIGALOTTI & U. BOSCAIN – “Controllability of the discrete-spectrum Schrödinger equation driven by an external field”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **26** (2009), no. 1, p. 329–349.
- [17] J.-M. CORON – *Control and nonlinearity*, Mathematical Surveys and Monographs, vol. 136, American Mathematical Society, Providence, RI, 2007.
- [18] P. GRISVARD – *Elliptic problems in nonsmooth domains*, Monographs and Studies in Mathematics, vol. 24, Pitman, Boston, MA, 1985.
- [19] A. HARAUX – “Séries lacunaires et contrôle semi-interne des vibrations d’une plaque rectangulaire”, *J. Math. Pures Appl. (9)* **68** (1989), no. 4, p. 457–465.
- [20] L. HÖRMANDER – *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis*, Grundlehren Math. Wiss., vol. 256, Springer-Verlag, Berlin, 1983.
- [21] I. LASIECKA & R. TRIGGIANI – “Sharp regularity theory for second order hyperbolic equations of Neumann type. I. L_2 nonhomogeneous data”, *Ann. Mat. Pura Appl. (4)* **157** (1990), p. 285–367.
- [22] G. LEBEAU – “Contrôle de l’équation de Schrödinger”, *J. Math. Pures Appl. (9)* **71** (1992), no. 3, p. 267–291.
- [23] J.-L. LIONS & E. MAGENES – *Non-homogeneous boundary value problems and applications. Vol. I*, Grundlehren Math. Wiss., vol. 181, Springer-Verlag, New York-Heidelberg, 1972.
- [24] F. MÉHATS, Y. PRIVAT & M. SIGALOTTI – “On the controllability of quantum transport in an electronic nanostructure”, *SIAM J. Appl. Math.* **74** (2014), no. 6, p. 1870–1894.
- [25] L. MILLER – “How violent are fast controls for Schrödinger and plate vibrations?”, *Arch. Rational Mech. Anal.* **172** (2004), no. 3, p. 429–456.
- [26] ———, “Controllability cost of conservative systems: resolvent condition and transmutation”, *J. Funct. Anal.* **218** (2005), no. 2, p. 425–444.
- [27] M. MORANCEY – “Simultaneous local exact controllability of 1D bilinear Schrödinger equations Morgan Morancey”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **31** (2014), no. 3, p. 501–529.
- [28] M. MORANCEY & V. NERSESYAN – “Global exact controllability of 1d Schrödinger equations with a polarizability term”, *Comptes Rendus Mathématique* **352** (2014), no. 5, p. 425–429.
- [29] ———, “Simultaneous global exact controllability of an arbitrary number of 1d bilinear Schrödinger equations”, *J. Math. Pures Appl. (9)* **103** (2015), no. 1, p. 228–254.

- [30] V. NERSESYAN – “Growth of Sobolev norms and controllability of the Schrödinger equation”, *Comm. Math. Phys.* **290** (2009), no. 1, p. 371–387.
- [31] ———, “Global approximate controllability for Schrödinger equation in higher Sobolev norms and applications”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **27** (2010), no. 3, p. 901–915.
- [32] V. NERSESYAN & H. NERSISYAN – “Global exact controllability in infinite time of Schrödinger equation: multidimensional case”, *J. Math. Pures Appl. (9)* **97** (2012), no. 4, p. 295–317.
- [33] Y. PRIVAT & M. SIGALOTTI – “The squares of the Laplacian-Dirichlet eigenfunctions are generically linearly independent”, *ESAIM Control Optim. Calc. Var.* **16** (2010), no. 3, p. 794–805, Erratum: *Ibid.*, p. 806–807.
- [34] J.-P. PUEL – “A regularity property for Schrödinger equations on bounded domains”, *Rev. Mat. Univ. Complut. Madrid* **26** (2013), no. 1, p. 183–192.
- [35] D. TATARU – “On the regularity of boundary traces for the wave equation”, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **26** (1998), no. 1, p. 185–206.
- [36] G. TENENBAUM & M. TUCSNAK – “Fast and strongly localized observation for the Schrödinger equation”, *Trans. Amer. Math. Soc.* **361** (2009), p. 951–977.
- [37] G. TURINICI – “On the controllability of bilinear quantum systems”, in *Mathematical Models and Methods for Ab Initio Quantum Chemistry* (C. Le Bris & M. Defranceschi, eds.), Lect. Notes in Chemistry, vol. 74, Springer, 2000.

Manuscript received June 18, 2016

accepted January 26, 2017

KARINE BEAUCHARD, IRMAR, École Normale Supérieure de Rennes, UBL
Avenue Robert Schumann, 35170 Bruz, France
E-mail : Karine.Beauchard@ens-rennes.fr
Url : <http://w3.bretagne.ens-cachan.fr/math/people/karine.beauchard/>

CAMILLE LAURENT, Sorbonne Universités, CNRS UMR 7598 and UPMC Univ. Paris 06, Laboratoire
Jacques-Louis Lions
F-75005, Paris, France
E-mail : camille.laurent@upmc.fr
Url : <https://www.ljll.math.upmc.fr/~laurent/>