

JOURNAL de Théorie des Nombres de BORDEAUX

anciennement Séminaire de Théorie des Nombres de Bordeaux

Carsten ELSNER, Shun SHIMOMURA et Iekata SHIOKAWA

Asymptotic representations for Fibonacci reciprocal sums and Euler's formulas for zeta values

Tome 21, n° 1 (2009), p. 145-157.

<http://jtnb.cedram.org/item?id=JTnb_2009__21_1_145_0>

© Université Bordeaux 1, 2009, tous droits réservés.

L'accès aux articles de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>), implique l'accord avec les conditions générales d'utilisation (<http://jtnb.cedram.org/legal/>). Toute reproduction en tout ou partie cet article sous quelque forme que ce soit pour tout usage autre que l'utilisation à fin strictement personnelle du copiste est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

Asymptotic representations for Fibonacci reciprocal sums and Euler's formulas for zeta values

par CARSTEN ELSNER, SHUN SHIMOMURA et IEKATA SHIOKAWA

RÉSUMÉ. Nous présentons les représentations asymptotiques pour certaines sommes des réciproques des nombres de Fibonacci et des nombres de Lucas quand un paramètre tend vers une valeur critique. Comme cas limite de nos résultats, nous obtenons les formules d'Euler pour les valeurs des fonctions de zeta.

ABSTRACT. We present asymptotic representations for certain reciprocal sums of Fibonacci numbers and of Lucas numbers as a parameter tends to a critical value. As limiting cases of our results, we obtain Euler's formulas for values of zeta functions.

1. Introduction

Let $\{F_n\}_{n \geq 0}$ and $\{L_n\}_{n \geq 0}$ be the Fibonacci numbers and the Lucas numbers, defined by

$$\begin{aligned} F_0 &= 0, & F_1 &= 1, & F_{n+2} &= F_{n+1} + F_n & (n \geq 0), \\ L_0 &= 2, & L_1 &= 1, & L_{n+2} &= L_{n+1} + L_n & (n \geq 0). \end{aligned}$$

For various reciprocal sums of these numbers, transcendence and algebraic relations were studied. Duverney, Ke. Nishioka, Ku. Nishioka, and the last named author [3] (see also [2]) proved the transcendence of the numbers $\sum_{n=1}^{\infty} F_n^{-2s}$, $\sum_{n=1}^{\infty} L_n^{-2s}$, $\sum_{n=1}^{\infty} F_{2n-1}^{-s}$, $\sum_{n=1}^{\infty} L_{2n}^{-s}$ ($s \in \mathbb{N}$) by using Nesterenko's theorem ([10]) on Ramanujan functions. In [4], for $\zeta_F(s) := \sum_{n=1}^{\infty} F_n^{-s}$, we proved that the values $\zeta_F(2)$, $\zeta_F(4)$, $\zeta_F(6)$ are algebraically independent, and that for any integer $s \geq 4$

$$\zeta_F(2s) - r_s \zeta_F(4) \in \mathbb{Q}(\zeta_F(2), \zeta_F(6))$$

with some $r_s \in \mathbb{Q}$; for example

$$\begin{aligned} \zeta_F(8) - \frac{15}{14} \zeta_F(4) &= \frac{1}{378(4u+5)^2} \left(256u^6 - 3456u^5 + 2880u^4 + 1792u^3v \right. \\ &\quad \left. - 11100u^3 + 20160u^2v - 10125u^2 + 7560uv + 3136v^2 - 1050v \right) \end{aligned}$$

with $u := \zeta_F(2)$, and $v := \zeta_F(6)$.

Assume that $\alpha, \beta \in \mathbb{C}$ satisfy

$$(1) \quad \alpha\beta = -1, \quad |\beta| < 1.$$

Let $\{U_n\}_{n \geq 0}$ and $\{V_n\}_{n \geq 0}$ be sequences defined by

$$(2) \quad U_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad V_n = \alpha^n + \beta^n \quad (n \geq 0),$$

which are generalized Fibonacci and Lucas numbers, respectively. Indeed, if $\beta = (1 - \sqrt{5})/2$, then $U_n = F_n$, $V_n = L_n$. For $s \in \mathbb{N}$ consider the reciprocal sums

$$(3) \quad h_{2s} := (\alpha - \beta)^{-2s} \sum_{n=1}^{\infty} \frac{1}{U_{2n}^{2s}},$$

$$(4) \quad g_{2s-1}^* := \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{V_{2n-1}^{2s-1}},$$

which are holomorphic for $|\beta| < 1$. For $\{h_{2s}\}_{s \in \mathbb{N}}$ (respectively, $\{g_{2s-1}^*\}_{s \in \mathbb{N}}$) algebraic relations were discussed in [6] (respectively, [5]). These sums may also be regarded as functions of the modulus k of Jacobian elliptic functions (see Sections 4 and 5, also [4], [5], [6]).

In this paper we present asymptotic representations for these sums as $\beta \rightarrow -1+0$ (or $k \rightarrow 1-0$). Degenerate cases of our expressions coincide with Euler's formulas for $\zeta(2s) = \sum_{n=1}^{\infty} n^{-2s}$ and $L(2s-1) = \sum_{n=1}^{\infty} (-1)^{n+1}(2n-1)^{-(2s-1)}$, respectively (see Section 3).

2. Statement of results

Let $B_{2\nu}$ and $E_{2\nu}$ ($\nu = 0, 1, 2, \dots$) denote, respectively, the Bernoulli and the Euler numbers:

$$\begin{aligned} B_0 &= 1, & B_2 &= 1/6, & B_4 &= -1/30, & B_6 &= 1/42, \dots; \\ E_0 &= 1, & E_2 &= -1, & E_4 &= 5, & E_6 &= -61, \dots, \end{aligned}$$

and let s be a positive integer. For $s \geq 2$, let $\sigma_1(s), \dots, \sigma_{s-1}(s)$ be the elementary symmetric functions of the $s-1$ numbers $-1^2, -2^2, \dots, -(s-1)^2$ defined by

$$\sigma_i(s) = (-1)^i \sum_{1 \leq r_1 < \dots < r_i \leq s-1} r_1^2 \cdots r_i^2 \quad (1 \leq i \leq s-1),$$

and for $s \geq 0$, set $\sigma_0(s) = 1$. For $s \geq 1$ let $\tau_1(s), \dots, \tau_s(s)$ be the elementary symmetric functions of the s numbers $-1^2, -3^2, \dots, -(2s-1)^2$ given by

$$\tau_i(s) = (-1)^i \sum_{1 \leq r_1 < \dots < r_i \leq s} (2r_1 - 1)^2 \cdots (2r_i - 1)^2 \quad (1 \leq i \leq s),$$

and for $s \geq 0$ set $\tau_0(s) = 1$. Our results are given as follows:

Theorem 1. For $-1 < \beta < -1 + \delta_0$, we have

$$(\alpha^2 - \beta^2)^{2s} h_{2s} = \Phi_s(\eta)(1 + O(e^{-\pi^2/(2\eta)})),$$

$$\eta := -\log(-\beta) = (1 + \beta)(1 + O(1 + \beta)).$$

Here δ_0 is a sufficiently small positive number, and $\Phi_s(X)$ is an entire function written in the form

$$\Phi_s(X) := \frac{2^s}{(2s-1)!} \left(X^{-2} (\cosh(4X) - 1) \right)^s \sum_{p=1}^s \Lambda_p(X)$$

with

$$\Lambda_p(X) := \begin{cases} \frac{\sigma_{s-1}(s)}{96} X^{2s-2} (\pi^2 - 12X + 4X^2), & \text{if } p = 1; \\ \frac{(-1)^{p-1}}{2^{2p+2} p} \sigma_{s-p}(s) B_{2p} X^{2s-2p} (\pi^{2p} - (-1)^p 2^{2p} X^{2p}), & \text{if } p \geq 2. \end{cases}$$

Several coefficients of the series expansion

$$\Phi_s(X) = \sum_{j=0}^{\infty} \lambda_j^{(s)} X^j, \quad \lambda_j^{(s)} \in \mathbb{Q}[\pi]$$

are given by

$$\lambda_0^{(s)} = \frac{2^{2s-1} (-1)^{s-1} B_{2s}}{(2s)!} \pi^{2s},$$

and (i) for $s = 1, 2$,

$$\lambda_1^{(1)} = -2, \quad \lambda_2^{(1)} = \frac{2}{9} \pi^2 + \frac{2}{3}, \quad \lambda_3^{(1)} = -\frac{8}{3}, \quad \lambda_4^{(1)} = \frac{16}{135} \pi^2 + \frac{8}{9},$$

$$\lambda_1^{(2)} = 0, \quad \lambda_2^{(2)} = \frac{4}{135} \pi^4 - \frac{4}{9} \pi^2, \quad \lambda_3^{(2)} = \frac{16}{3},$$

(ii) for $s \geq 3$,

$$\lambda_1^{(s)} = 0, \quad \lambda_2^{(s)} = \frac{2^{2s-1} (-1)^{s-1}}{3 \cdot (2s-1)!} \pi^{2s-2} (2B_{2s} \pi^2 + s(2s-1)B_{2s-2}), \quad \lambda_3^{(s)} = 0.$$

Theorem 2. For $-1 < \beta < -1 + \delta_0$, we have

$$(\alpha + \beta)^{2s-1} g_{2s-1}^* = \Psi_s(\eta)(1 + O(e^{-\pi^2/(2\eta)})).$$

Here δ_0 and η are as in Theorem 1, and $\Psi_s(X)$ is an entire function written in the form

$$\Psi_s(X) := \frac{(X^{-1} \sinh X)^{2s-1}}{(2s-2)!}$$

$$\times \sum_{p=0}^{s-1} (-1)^p 2^{-2p-2} \pi^{2p+1} \tau_{s-1-p}(s-1) E_{2p} X^{2s-2p-2}.$$

Several coefficients of the series expansion

$$\Psi_s(X) = \sum_{j=0}^{\infty} \mu_j^{(s)} X^{2j}, \quad \mu_j^{(s)} \in \mathbb{Q}[\pi]$$

are given by

$$\mu_0^{(s)} = \frac{(-1)^{s-1} E_{2s-2}}{2^{2s}(2s-2)!} \pi^{2s-1},$$

and (i) for $s = 1$,

$$\mu_m^{(1)} = \frac{\pi}{4 \cdot (2m+1)!} \quad (m \geq 1),$$

(ii) for $s \geq 2$,

$$\mu_1^{(s)} = \frac{(-1)^{s-1} (2s-1)}{2^{2s+1} \cdot 3 \cdot (2s-2)!} \pi^{2s-3} (E_{2s-2} \pi^2 + 8(s-1)(2s-3)E_{2s-4}).$$

Remark 1. The asymptotic formulas above are also valid as $\beta \rightarrow -1$ through the sector $|\arg(\beta+1)| < \pi/2 - \theta_0$, where $\theta_0 > 0$ is an arbitrary small number (then $\eta \rightarrow 0$, $|\arg \eta| < \pi/2 - \theta'_0$ for some $\theta'_0 > 0$) (see Remark 4 and Section 5).

Remark 2. The quantity β may be regarded as a function of the modulus k of Jacobian elliptic functions, which satisfies $\beta \rightarrow -1 + 0$ as $k \rightarrow 1 - 0$ (see Section 4). Moreover, $\beta \rightarrow i$ and $\alpha \rightarrow i$ as $k \rightarrow \infty$ through a suitable sector. In this case, for

$$(\alpha - \beta)^{2s} h_{2s} = \sum_{n=1}^{\infty} \frac{1}{U_{2n}^{2s}} \quad \text{and} \quad (\alpha - \beta)^{2s-1} f_{2s-1} = \sum_{n=1}^{\infty} \frac{1}{U_{2n-1}^{2s-1}},$$

we obtain similar asymptotic formulas as $\beta \rightarrow i$.

3. Euler's formulas for zeta functions

For $-1 < \beta < -1 + \delta_0$ and for $n \geq 1$, observing that

$$\begin{aligned} \left| \frac{\alpha^{2n} - \beta^{2n}}{\alpha^2 - \beta^2} \right| &= \left| \sum_{\nu=0}^{n-1} \alpha^{2n-2-2\nu} \beta^{2\nu} \right| \\ &= \left| \sum_{\nu=0}^{n-1} \beta^{4\nu-2n+2} \right| = \left| \frac{1}{2} \sum_{\nu=0}^{n-1} (\beta^{4\nu-2n+2} + \beta^{-(4\nu-2n+2)}) \right| \geq n, \end{aligned}$$

we have for $s \in \mathbb{N}$

$$\lim_{\beta \rightarrow -1+0} (\alpha^2 - \beta^2)^{2s} h_{2s} = \sum_{n=1}^{\infty} \frac{1}{n^{2s}}.$$

Therefore, letting β tend to $-1 + 0$ in Theorem 1, we obtain

$$\zeta(2s) = \frac{2^{2s-1}(-1)^{s-1}B_{2s}}{(2s)!}\pi^{2s} \quad (s \in \mathbb{N}).$$

For each $s \in \mathbb{N}$ a similar argument concerning Theorem 2 leads us

$$\lim_{\beta \rightarrow -1+0} (\alpha + \beta)^{2s-1} g_{2s-1}^* = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^{2s-1}} = \frac{(-1)^{s-1} E_{2s-2}}{2^{2s}(2s-2)!} \pi^{2s-1}.$$

4. Preliminaries

Consider the complete elliptic integrals of the first and the second kinds with the modulus k

$$(5) \quad K = K(k) := \int_0^1 \frac{dt}{\sqrt{(1-t^2)(1-k^2t^2)}}, \quad E = E(k) := \int_0^1 \sqrt{\frac{1-k^2t^2}{1-t^2}} dt$$

for $k^2 \in \mathbb{C} \setminus (\{0\} \cup [1, +\infty))$, where the branch of each integrand is chosen so that it tends to 1 as $t \rightarrow 0$. Put

$$(6) \quad \beta = \beta(k) = -e^{-\pi K'/(2K)}, \quad \alpha = \alpha(k) = -1/\beta(k),$$

where

$$(7) \quad K' := K(k'), \quad k' := \sqrt{1-k^2},$$

the branch of k' being chosen so that $k' > 0$ for $0 < k < 1$. These integrals are expressible as follows (cf. [1, 17.3.9 and 17.3.10]):

$$(8) \quad K = \frac{\pi}{2} F(1/2, 1/2, 1, k^2),$$

$$(9) \quad K' = \frac{\pi}{2} F(1/2, 1/2, 1, 1-k^2),$$

$$(10) \quad E = \frac{\pi}{2} F(-1/2, 1/2, 1, k^2),$$

with the hypergeometric function

$$F(a, b, c, x) = \sum_{n=0}^{\infty} \frac{(a)_n(b)_n}{(c)_n n!} x^n, \quad (a)_n := \frac{\Gamma(a+n)}{\Gamma(a)} = a(a+1) \cdots (a+n-1).$$

Lemma 3. *As $k \rightarrow 1 - 0$, we have*

$$(11) \quad K = l(k)(1 + 2 \log 2 \cdot l(k)^{-1})(1 + O(1 - k^2)),$$

$$(12) \quad K' = \frac{\pi}{2}(1 + O(1 - k^2)),$$

$$(13) \quad E = 1 + O((1 - k^2)l(k)),$$

with

$$l(k) := \log(1/k') = \log(1/\sqrt{1-k^2}).$$

Furthermore $l(k) \rightarrow +\infty$ and $K \rightarrow +\infty$ as $k \rightarrow 1 - 0$.

Proof. Formula (12) follows immediately from (9). Recall the connection formula for $F(1/2, 1/2, 1, z)$ around $z = 0$ and $z = 1$:

$$\begin{aligned} F(1/2, 1/2, 1, z) &= \\ \frac{\Gamma(1)}{\Gamma(1/2)^2} \sum_{n=0}^{\infty} \frac{(1/2)_n^2}{(n!)^2} &\left(2\psi(n+1) - 2\psi(n+1/2) - \log(1-z) \right) (1-z)^n, \\ \psi(t) &= \frac{\Gamma'(t)}{\Gamma(t)} = -\gamma + \sum_{m=0}^{\infty} \left(\frac{1}{m+1} - \frac{1}{m+t} \right) \end{aligned}$$

for $|1-z| < 1$, $|\arg(1-z)| < \pi$ (cf. [1, 15.3.10]). Putting $z = k^2$, we derive for $|1-k^2| < 1$, $|\arg(1-k^2)| < \pi$,

$$\begin{aligned} K &= \frac{1}{2} \sum_{n=0}^{\infty} \frac{(1/2)_n^2}{(n!)^2} \\ &\times \left(2 \sum_{m=0}^{\infty} \left(\frac{1}{m+n+1/2} - \frac{1}{m+n+1} \right) - \log(1-k^2) \right) (1-k^2)^n \\ &= 2 \log 2 - \frac{1}{2} \log(1-k^2) + O((1-k^2) \log(1-k^2)) \\ &= l(k) + 2 \log 2 + O((1-k^2)l(k)), \end{aligned}$$

which implies (11), where the branch of $\log(1-k^2)$ is chosen so that $\operatorname{Im} \log(1-k^2) = 0$ for $0 < k < 1$, namely for $\arg k = \arg(1-k) = 0$. Using the formula

$$\begin{aligned} F(-1/2, 1/2, 1, z) &= \\ \frac{\Gamma(1)^2}{\Gamma(1/2)\Gamma(3/2)} + \frac{\Gamma(1)}{\Gamma(-1/2)\Gamma(1/2)} \sum_{n=0}^{\infty} \frac{(1/2)_n(3/2)_n}{n!(n+1)!} &(1-z)^{n+1} \\ &\times \left(\log(1-z) - \psi(n+1) - \psi(n+2) + \psi(n+1/2) + \psi(n+3/2) \right) \end{aligned}$$

for $|1-z| < 1$, $|\arg(1-z)| < \pi$ (cf. [1, 15.3.11]), we obtain (13). \square

From (11) it follows that

$$(14) \quad 1 - k^2 = e^{-2l(k)} = O(e^{-2K})$$

as $k \rightarrow 1 - 0$. Hence, by (12), for $1 - \delta_1 < k < 1$,

$$(15) \quad \beta(k) = -\exp\left(-\frac{\pi^2}{4K}(1 + O(e^{-2K}))\right),$$

and $|\beta(k)| < 1$, provided that $\delta_1 > 0$ is sufficiently small. This implies that $\beta(k) \rightarrow -1 + 0$ as $k \rightarrow 1 - 0$.

Remark 3. The connection formulas in the proof of Lemma 3 are also valid, as long as we consider the analytic continuation of $\log(1 - k^2)$ on the universal covering of the punctured disk $0 < |k - 1| < 1$. Hence as $k \rightarrow 1$ (without any condition on $\arg(1 - k^2)$) we have $\operatorname{Re} K(k) \rightarrow +\infty$ and $\beta(k) \rightarrow -1$, $|\arg(\beta(k) + 1)| < \pi/2 - \theta_0$, where θ_0 is as in Remark 1. Conversely, for every path C_β tending to $\beta = -1$ through the sector $|\arg(\beta + 1)| < \pi/2$, there exists a curve C_k tending to $k = 1$ such that $\beta(C_k) = C_\beta$.

Lemma 4. *There exists a small positive number δ_1 such that, for $1 - \delta_1 < k < 1$, the quantities $\beta = \beta(k)$ and $\alpha = \alpha(k) = -1/\beta(k)$ satisfy*

$$(16) \quad \alpha(k)^2 - \beta(k)^2 = (e^{\pi^2/(2K)} - e^{-\pi^2/(2K)})(1 + O(e^{-2K})),$$

$$(17) \quad \alpha(k) + \beta(k) = (e^{\pi^2/(4K)} - e^{-\pi^2/(4K)})(1 + O(e^{-2K})).$$

Proof. Using (6), (12) and (14), we have

$$\begin{aligned} \alpha(k)^2 - \beta(k)^2 &= e^{\pi K'/K} - e^{-\pi K'/K} \\ &= e^{-\pi K'/K}(e^{2\pi K'/K} - 1) \\ &= \exp\left(-\frac{\pi^2}{2K}(1 + O(e^{-2K}))\right)\left(\exp\left(\frac{\pi^2}{K}(1 + O(e^{-2K}))\right) - 1\right) \\ &= e^{\frac{-\pi^2}{2K}}(1 + O(K^{-1}e^{-2K}))\left(e^{\frac{\pi^2}{K}}(1 + O(K^{-1}e^{-2K})) - 1\right) \\ &= e^{-\pi^2/(2K)}(1 + O(e^{-2K}))\left((e^{\pi^2/K} - 1) + O(K^{-1}e^{-2K})\right) \\ &= e^{-\pi^2/(2K)}(e^{\pi^2/K} - 1)(1 + O(e^{-2K})) \end{aligned}$$

as $k \rightarrow 1 - 0$ (and $K \rightarrow +\infty$), which implies (16). Estimate (17) is obtained by a similar argument. \square

We note the following lemmas concerning the Jacobian elliptic functions

$$\operatorname{ns}(k, z) = 1/\operatorname{sn}(k, z), \quad \operatorname{cn}(k, z) = \sqrt{1 - \operatorname{sn}^2(k, z)}$$

with $w = \operatorname{sn}(k, z)$ defined by

$$z = \int_0^w \frac{dv}{\sqrt{(1 - v^2)(1 - k^2v^2)}}.$$

Lemma 5. *Around $z = 0$,*

$$\operatorname{ns}^2(k, z) = z^{-2} + \sum_{j=0}^{\infty} c_j(k)z^{2j},$$

where $c_j(k) \in \mathbb{Q}[k^2]$. In particular,

$$\operatorname{ns}^2(1, z) = \coth^2 z = z^{-2} + \sum_{j=0}^{\infty} c_j(1)z^{2j},$$

where

$$c_0(1) = \frac{2}{3}, \quad c_j(1) = -\frac{2^{2j+2}(2j+1)B_{2j+2}}{(2j+2)!} \quad (j \geq 1).$$

Proof. The function $u = \text{ns}^2(k, z)$ satisfies the differential equations

$$(u')^2 = 4u(u-1)(u-k^2) \quad \text{and} \quad u'' = 6u^2 - 4(k^2 + 1)u + 2k^2.$$

Substituting the expansion $z^{-2} + \sum_{j=0}^{\infty} c_j(k)z^{2j}$ of $\text{ns}^2(k, z)$ into these equations, we can determine the coefficients $c_j(k)$ (see [4, Lemma 2] and the recursive relation in it). The relation $\text{ns}^2(1, z) = \coth^2 z$ immediately follows from the fact that $u = \text{ns}^2(1, z)$ satisfies the equation $(u')^2 = 4u(u-1)^2$ admitting the general solution $\coth^2(z - z_0)$. \square

The function $u = k \operatorname{cn}(k, z)$ satisfies

$$(u')^2 = (k^2 - u^2)(u^2 + 1 - k^2), \quad u(0) = k.$$

If $k = 1$, then $(u')^2 = u^2(1 - u^2)$ admits the general solution $\operatorname{sech}(z - z_0)$. From this fact we obtain the following (see also [6, Lemma 2]):

Lemma 6. *Around $z = 0$,*

$$k \operatorname{cn}(k, z) = k + \sum_{j=1}^{\infty} c_j^*(k)z^{2j},$$

where $c_j^*(k) \in \mathbb{Q}[k]$. In particular,

$$\operatorname{cn}(1, z) = \operatorname{sech} z = 1 + \sum_{j=1}^{\infty} c_j^*(1)z^{2j}, \quad c_j^*(1) = \frac{E_{2j}}{(2j)!}.$$

5. Proofs of results

5.1. Proof of Theorem 1.

Recall that

$$\begin{aligned} (18) \quad (\alpha^2 - \beta^2)^{2s} h_{2s} &= \left(\frac{\alpha^2 - \beta^2}{2}\right)^{2s} \sum_{\nu=1}^{\infty} \operatorname{cosech}^{2s}(\nu\pi K'/K) \\ &= \frac{(\alpha^2 - \beta^2)^{2s}}{(2s-1)!} \sum_{j=0}^{s-1} \sigma_{s-j-1}(s) A_{2j+1}(\beta^2), \end{aligned}$$

where

$$A_{2j+1}(q) := \sum_{n=1}^{\infty} \frac{n^{2j+1} q^{2n}}{1 - q^{2n}}$$

(cf. [12, Table 1(i)], [4]). The series $A_{2j+1}(\beta(k)^2)$ are generated by

$$\left(\frac{2K}{\pi}\right)^2 \text{ns}^2\left(\frac{2Kx}{\pi}\right) = \frac{4K(K-E)}{\pi^2} + \operatorname{cosec}^2 x - 8 \sum_{j=0}^{\infty} (-1)^j A_{2j+1} \frac{(2x)^{2j}}{(2j)!}$$

(cf. [12, Tables 1(i)], [9], [11, p. 535]), namely

$$(19) \quad A_1 = \frac{1}{24} \left(1 - \left(\frac{2K}{\pi} \right)^2 \left(\frac{3E}{K} - 2 + k^2 \right) \right),$$

$$(20) \quad \left(\frac{2K}{\pi} \right)^{2j+2} c_j(k) = a_j - (-1)^j \frac{2^{2j+3}}{(2j)!} A_{2j+1} \quad (j \geq 1).$$

Here $c_j(k)$ are the coefficients given in Lemma 5, and a_j are the coefficients of the expansion

$$\operatorname{cosec}^2 z = z^{-2} + \sum_{j=0}^{\infty} a_j z^{2j},$$

namely, $a_0 = 1/3$, $a_j = (-1)^{j-1} c_j(1)$ ($j \geq 1$).

Recall that $K \rightarrow +\infty$ and that $\beta(k) \rightarrow -1 + 0$ as $k \rightarrow 1 - 0$ (cf. (15) and Lemma 3). By (16) we have

$$(21) \quad \begin{aligned} (\alpha^2 - \beta^2)^{2s} &= (e^{\pi^2/K} + e^{-\pi^2/K} - 2)^s (1 + O(e^{-2K})) \\ &= \pi^{4s} K^{-2s} \left(2\pi^{-4} K^2 (\cosh(\pi^2/K) - 1) \right)^s (1 + O(e^{-2K})) \end{aligned}$$

as $k \rightarrow 1 - 0$. Note that

$$2\pi^{-4} K^2 (\cosh(\pi^2/K) - 1) = \sum_{m=1}^{\infty} \frac{2\pi^{4(m-1)}}{(2m)!} K^{-2(m-1)} = 1 + O(K^{-2}).$$

By (19), (14) and Lemma 3, the quantity $A_1(\beta^2)$ can be written as

$$\begin{aligned} A_1 &= \frac{1}{24} \left(1 - \frac{4K^2}{\pi^2} \left(\frac{3}{K} (1 + O(l(k)e^{-2K})) - 1 + O(e^{-2K}) \right) \right) \\ &= \frac{1}{24} \left(1 - \frac{4K^2}{\pi^2} (-1 + 3K^{-1} + O(e^{-2K})) \right) \\ &= \frac{K^2}{6\pi^2} \left(1 - 3K^{-1} + \frac{\pi^2}{4} K^{-2} \right) (1 + O(e^{-2K})). \end{aligned}$$

Combining this with (21), we have for $s \geq 1$

$$(22) \quad \begin{aligned} (\alpha^2 - \beta^2)^{2s} A_1 &= \frac{\pi^{4s-2}}{6} K^{-2(s-1)} \left(2\pi^{-4} K^2 (\cosh(\pi^2/K) - 1) \right)^s \\ &\quad \times \left(1 - 3K^{-1} + \frac{\pi^2}{4} K^{-2} \right) (1 + O(e^{-2K})) \end{aligned}$$

as $k \rightarrow 1 - 0$. For $s \geq 2$ and for $2 \leq p \leq s$, using (20), (21), and Lemma 5 with the facts $a_{p-1} = (-1)^p c_{p-1}(1)$, $c_{p-1}(k) = c_{p-1}(1) + O(e^{-2K})$, we have

$$\begin{aligned}
(\alpha^2 - \beta^2)^{2s} A_{2p-1} &= \frac{(-1)^p (2p-2)!}{2} \pi^{4s-2p} c_{p-1}(1) K^{-2(s-p)} \\
&\quad \times \left(2\pi^{-4} K^2 (\cosh(\pi^2/K) - 1) \right)^s \\
&\quad \times (1 - (-1)^p 2^{-2p} \pi^{2p} K^{-2p}) (1 + O(e^{-2K})) \\
(23) \quad &= \frac{(-1)^{p-1}}{p} 2^{2p-2} \pi^{4s-2p} B_{2p} K^{-2(s-p)} \\
&\quad \times \left(2\pi^{-4} K^2 (\cosh(\pi^2/K) - 1) \right)^s \\
&\quad \times (1 - (-1)^p 2^{-2p} \pi^{2p} K^{-2p}) (1 + O(e^{-2K}))
\end{aligned}$$

as $k \rightarrow 1 - 0$. By (15),

$$\begin{aligned}
K^{-1} &= -\frac{4}{\pi^2} \log(-\beta) (1 + O(e^{-2K})), \\
K &= \frac{\pi^2}{4\eta} (1 + O(e^{-2K})) = \frac{\pi^2}{4\eta} + O(K e^{-2K}) = \frac{\pi^2}{4\eta} + O(1),
\end{aligned}$$

and hence

$$(24) \quad K^{-1} = \frac{4}{\pi^2} \eta (1 + O(e^{-\pi^2/(2\eta)})), \quad e^{-2K} = O(e^{-\pi^2/(2\eta)}).$$

For $s \geq 1$ we derive from (18), (22) and (23) that

$$\begin{aligned}
(\alpha^2 - \beta^2)^{2s} h_{2s} &= \frac{(\alpha^2 - \beta^2)^{2s}}{(2s-1)!} \sum_{p=1}^s \sigma_{s-p}(s) A_{2p-1} \\
(25) \quad &= \frac{\left(2\pi^{-4} K^2 (\cosh(\pi^2/K) - 1) \right)^s}{(2s-1)!} \sum_{p=1}^s \sigma_{s-p}(s) F_{s,p}(K)
\end{aligned}$$

with

$$F_{s,1}(K) := \frac{\pi^{4s-2}}{6} K^{-2(s-1)} \left(1 - 3K^{-1} + \frac{\pi^2}{4} K^{-2} \right) (1 + O(e^{-2K})),$$

and

$$\begin{aligned}
F_{s,p}(K) &:= \frac{(-1)^{p-1}}{p} 2^{2p-2} \pi^{4s-2p} B_{2p} K^{-2(s-p)} \\
&\quad \times (1 - (-1)^p 2^{-2p} \pi^{2p} K^{-2p}) (1 + O(e^{-2K}))
\end{aligned}$$

for $p \geq 2$. Now we note the following: for a function $\phi(z)$ holomorphic for $|z| < r_0$ satisfying $\phi(0) \neq 0$,

$$\phi(z + \Delta z) = \phi(z) + \int_0^{\Delta z} \phi'(z+t) dt = \phi(z) + O(\Delta z) = \phi(z)(1 + O(\Delta z))$$

as $\Delta z \rightarrow 0$, uniformly for $|z| < r'_0$, provided that r'_0 is sufficiently small. Substituting (24) into (25), and using this fact with $z = 4\pi^{-2}\eta$ and $\Delta z = O(\eta e^{-\pi^2/(2\eta)})$, we have the required formula. Using

$$2X^{-2}(\cosh(4X) - 1) = \sum_{m=0}^{\infty} \frac{2^{4m+5}}{(2m+2)!} X^{2m}$$

and

$$\sigma_1(s) = -\sum_{r=1}^{s-1} r^2 = -\frac{1}{6}s(s-1)(2s-1),$$

we obtain the coefficients mentioned in the theorem.

5.2. Proof of Theorem 2.

Note that, for $s \in \mathbb{N}$,

$$\begin{aligned} (\alpha + \beta)^{2s-1} g_{2s-1}^* &= \left(\frac{\alpha + \beta}{2}\right)^{2s-1} \sum_{\nu=1}^{\infty} (-1)^{\nu+1} \operatorname{cosech}^{2s-1}((\nu - 1/2)\pi K'/K) \\ (26) \quad &= \frac{(\alpha + \beta)^{2s-1}}{2^{2s-2}(2s-2)!} \sum_{j=0}^{s-1} \tau_{s-1-j}(s-1) T_{2j}(\beta^2), \end{aligned}$$

where

$$T_{2j}(q) := \sum_{n=1}^{\infty} \frac{(2n-1)^{2j} q^{n-1/2}}{1 + q^{2n-1}}$$

(cf. [12, Table 1(xv)]). The series $T_{2j}(\beta(k)^2)$ are generated by

$$\left(\frac{2K}{\pi}\right) k \operatorname{cn}\left(\frac{2Kx}{\pi}\right) = 4 \sum_{j=0}^{\infty} (-1)^j T_{2j} \frac{x^{2j}}{(2j)!},$$

namely

$$(27) \quad \left(\frac{2K}{\pi}\right)^{2j+1} c_j^*(k) = \frac{(-1)^j 4}{(2j)!} T_{2j} \quad (j \geq 0),$$

where $c_j^*(k)$ are coefficients given in Lemma 6. By (17)

$$\begin{aligned} (28) \quad \alpha + \beta &= (e^{\pi^2/(4K)} - e^{-\pi^2/(4K)})(1 + O(e^{-2K})) \\ &= \frac{\pi^2}{2} K^{-1} \left(4\pi^{-2} K \sinh(\pi^2/(4K))\right) (1 + O(e^{-2K})), \end{aligned}$$

where

$$4\pi^{-2} K \sinh(\pi^2/(4K)) = \sum_{m=1}^{\infty} \frac{\pi^{4(m-1)}}{4^{2(m-1)} (2m-1)!} K^{-2(m-1)} = 1 + O(K^{-2}).$$

Note that $c_{j-1}^*(k) = c_{j-1}^*(1) + O(1-k) = c_{j-1}^*(1) + O(e^{-2K})$ as $k \rightarrow 1-0$, because $c_{j-1}^*(k) \in \mathbb{Q}[k]$. By (27), (28) and Lemma 6, we have, for $s \geq 1$

and for $0 \leq p \leq s - 1$,

$$(29) \quad (\alpha + \beta)^{2s-1} T_{2p} = (-1)^p 2^{2p-2s} \pi^{4s-2p-3} E_{2p} K^{-2(s-p-1)} \\ \times \left(4\pi^{-2} K \sinh(\pi^2/(4K)) \right)^{2s-1} (1 + O(e^{-2K})).$$

Substituting these quantities into (26), we obtain

$$(\alpha + \beta)^{2s-1} g_{2s-1}^* = \frac{(\alpha + \beta)^{2s-1}}{2^{2s-2} (2s-2)!} \sum_{p=0}^{s-1} \tau_{s-1-p}(s-1) T_{2p} \\ = \frac{1}{(2s-2)!} \left(4\pi^{-2} K \sinh(\pi^2/(4K)) \right)^{2s-1} \\ \times \sum_{p=0}^{s-1} \left((-1)^p 2^{2p-4s+2} \pi^{4s-2p-3} \tau_{s-1-p}(s-1) \right. \\ \left. \times E_{2p} K^{-2(s-p-1)} (1 + O(e^{-2K})) \right).$$

This combined with (24) yields the required formula. Using

$$2X^{-1} \sinh X = 2 \sum_{m=0}^{\infty} \frac{X^{2m}}{(2m+1)!}$$

and

$$\tau_1(s-1) = - \sum_{r=1}^{s-1} (2r-1)^2 = - \frac{1}{3} (s-1)(2s-1)(2s-3),$$

we can check the coefficients given in the theorem.

Acknowledgements

The authors are grateful to Professor Yu. V. Nesterenko for stimulating discussions on the degeneration of reciprocal sums of Fibonacci numbers, and also to the referee for his/her very helpful comments.

References

- [1] M. ABRAMOWITZ AND I. A. STEGUN, *Handbook of Mathematical Functions*. Dover, New York, 1965.
- [2] D. DUVERNEY, KE. NISHIOKA, KU. NISHIOKA, AND I. SHIOKAWA, *Transcendence of Jacobi's theta series*. Proc. Japan Acad. Ser. A Math. Sci. **72** (1996), 202–203.
- [3] D. DUVERNEY, KE. NISHIOKA, KU. NISHIOKA, AND I. SHIOKAWA, *Transcendence of Rogers-Ramanujan continued fraction and reciprocal sums of Fibonacci numbers*. Proc. Japan Acad. Ser. A Math. Sci. **73** (1997), 140–142.
- [4] C. ELSNER, S. SHIMOMURA, AND I. SHIOKAWA, *Algebraic relations for reciprocal sums of Fibonacci numbers*. Acta Arith. **130** (2007), 37–60.
- [5] C. ELSNER, S. SHIMOMURA, AND I. SHIOKAWA, *Algebraic relations for reciprocal sums of odd terms in Fibonacci numbers*. Ramanujan J. **17** (2008), 429–446.
- [6] C. ELSNER, S. SHIMOMURA, AND I. SHIOKAWA, *Algebraic relations for reciprocal sums of even terms in Fibonacci numbers*. To appear in St. Petersburg Math. J.
- [7] H. HANCOCK, *Theory of Elliptic Functions*. Dover, New York, 1958.

- [8] A. HURWITZ AND R. COURANT, *Vorlesungen über allgemeine Funktionentheorie und elliptische Funktionen*. Springer, Berlin, 1925.
- [9] C. G. J. JACOBI, *Fundamenta Nova Theoriae Functionum Ellipticarum*. Königsberg, 1829.
- [10] YU. V. NESTERENKO, *Modular functions and transcendence questions*. Mat. Sb. **187** (1996), 65–96; English transl. Sb. Math. **187** (1996), 1319–1348.
- [11] E. T. WHITTAKER AND G. N. WATSON, *Modern Analysis*, 4th ed. Cambridge Univ. Press, Cambridge, 1927.
- [12] I. J. ZUCKER, *The summation of series of hyperbolic functions*. SIAM J. Math. Anal. **10** (1979), 192–206.

Carsten ELSNER

Fachhochschule für die Wirtschaft
University of Applied Sciences
Freundallee 15
30173 Hannover, Germany
E-mail: carsten.elsner@fhwd.de

Shun SHIMOMURA

Department of Mathematics
Keio University
3-14-1 Hiyoshi, Kohoku-ku
Yokohama 223-8522 Japan
E-mail: shimomur@math.keio.ac.jp

Iekata SHIOKAWA

Department of Mathematics
Keio University
3-14-1 Hiyoshi, Kohoku-ku
Yokohama 223-8522 Japan
E-mail: shiokawa@beige.ocn.ne.jp