

ANNALES DE L'I. H. P., SECTION A

A. H. TAUB

Space-times admitting a covariantly constant spinor field

Annales de l'I. H. P., section A, tome 41, n° 2 (1984), p. 227-236

http://www.numdam.org/item?id=AIHPA_1984__41_2_227_0

© Gauthier-Villars, 1984, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Space-times admitting a covariantly constant spinor field

by

A. H. TAUB

Department of Mathematics, University of California,
Berkeley, CA 94720

ABSTRACT. — It is shown that if a space-time V admits a covariantly constant spinor field ψ_A , and hence a covariantly constant null vector field l_μ determined by ψ_A , its Ricci tensor is proportional to the tensor product of l_μ by itself. Further, the conformal tensor of V is shown to be of Petrov-Penrose type N. That is the four index symmetric spinor determined by the conformal tensor of V is proportional to the spinor product of ψ_A with itself. The Bianchi identities are used to show that empty asymptotically flat space-times with infinitely extendible null geodesics tangent to l_μ are flat.

RÉSUMÉ. — On montre que, si un espace temps V admet un champ spinoriel constant par covariance ψ_A , et par suite un champ de vecteurs isotrope constant par covariance l_μ déterminé par ψ_A , son tenseur de Ricci est proportionnel au tenseur produit de l_μ par lui-même. En outre, on montre que le tenseur conforme de V est du type N de Petrov-Penrose, c'est-à-dire que le spineur symétrique à quatre indices déterminé par le tenseur conforme de V est proportionnel au produit spinoriel de ψ_A par lui-même. On utilise les identités de Bianchi pour montrer que des espaces temps vides et asymptotiquement plats, dont les géodésiques isotropes tangentées à l_μ sont prolongeables indéfiniment, sont plats.

1. INTRODUCTION

It is the purpose of this paper to characterize those space-times that admit a covariantly constant two component spinor field. In addition, we shall prove that asymptotically flat empty space-times that contain such a spinor field are flat.

We shall use the notation and results of reference [1] where spinors ϕ^A and ψ^A satisfying

$$\phi^A \psi_A = 1 ; \quad (1.1)$$

are introduced. Spinor indices are manipulated with the asymmetric Levi-Civita tensor density ε_{AB} and ε^{AB} by the rules

$$\phi_A = \varepsilon_{AB} \phi^B$$

and

$$\phi^A = \phi_B \varepsilon^{BA}$$

Equation (1.1) is equivalent to

$$\phi^A \psi_B - \psi^A \phi_B = \delta_B^A \quad (1.2)$$

In terms of these the Newman-Penrose spin coefficients are defined as

$$\begin{aligned} A_{0v} &= \phi^A \phi_{A;v} \\ A_{1v} &= \phi^A \psi_{A;v} = \psi^A \phi_{A;v} \\ A_{2v} &= \psi^A \psi_{A;v} \end{aligned} \quad (1.3)$$

where the semi-colon denotes the spinor covariant derivative with respect to the metric of the space-time V .

The spin-coefficients define three two forms by the equation

$$R_{i\mu\nu} = A_{i\mu;v} - A_{i\mu;v} - \sqrt{2} E_{ijk} A_\mu^j A_\nu^k \quad i, j, k = (1, 2, 3) \quad (1.4)$$

where

$$E_{ijk} = \sqrt{a} \varepsilon_{ijk}$$

with ε_{ijk} the Levi-Civita alternating tensor density and

$$\begin{aligned} a &= \det \| a_{ij} \| \\ \| a_{ij} \| &= \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1/2 & 0 \\ 1 & 0 & 0 \end{pmatrix}; \quad \| a^{ij} \| = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -2 & 0 \\ 1 & 0 & 0 \end{pmatrix} \\ a_{ij} a^{jk} &= \delta_i^k \end{aligned}$$

Lower case latin indices are manipulated with the a_{ij} and a^{ij} .

The quantities X_{iAB} are defined by

$$\begin{aligned} X_{0AB} &= \phi_A \phi_B &= X_{AB}^2 \\ X_{1AB} &= \frac{1}{2}(\phi_A \phi_B + \psi_A \psi_B) &= -\frac{1}{2} X_{AB}^1 \\ X_{2AB} &= \psi_A \psi_B &= X_{AB}^0 \end{aligned}$$

and satisfy

$$X_{iAB} X^{iCD} = \frac{1}{2}(\delta_A^C \delta_B^D + \delta_B^C \delta_A^D)$$

The $R_{i\mu\nu}$ are related to the spinor curvature two-forms $R_{AB\mu\nu}$ by the equations

$$R_{i\mu\nu} = X_i^{AB} R_{AB\mu\nu} = \frac{1}{4} X_i^{AB} p_{AB}^{\rho\sigma} R_{\rho\sigma\mu\nu}$$

where $R_{\rho\sigma\mu\nu}$ are the components of the Riemann curvature tensor of V whose metric is $g_{\mu\nu}$. This metric determines 2×2 matrices

$$q_\mu = \|\gamma_\mu \dot{A}_B\|$$

such that

$$\bar{q}_\mu q_\nu + \bar{q}_\nu q_\mu = 2g_{\mu\nu} 1_2$$

when the bar over a quantity denotes its complex conjugate. Matrices $\|p_{\mu\nu} \dot{A}_B\|$ are defined by the equations

$$2p_{\mu\nu} = \bar{q}_\mu q_\nu - \bar{q}_\nu q_\mu.$$

Greek indices are manipulated by $g_{\mu\nu}$ and $g^{\mu\nu}$. Various algebraic properties of the matrices $p_{\mu\nu}$ are given in the appendix.

We may also write

$$R_{i\mu\nu} = \frac{1}{8} \left((C_{ij} - \frac{R}{3} a_{ij}) p_{\mu\nu}^j + S_{i\bar{j}} \bar{P}_{\mu\nu}^{\bar{j}} \right)$$

where R is the scalar curvature of the space-time,

$$C_{ij} = \frac{1}{4} p_{AB}^{\rho\sigma} C_{\rho\sigma\mu\nu} p_{CD}^{\mu\nu} X_i^{AB} X_j^{CD} = \dot{R}_{i\mu\nu} p_j^{\mu\nu}$$

where $C_{\rho\sigma\mu\nu}$ is the conformal tensor of V , and

$$S_{i\bar{j}} = \frac{1}{4} p_{AB}^{\rho\sigma} R_{\rho\sigma\mu\nu} \bar{p}_{CD}^{\mu\nu} X_i^{AB} \bar{X}_j^{\bar{C}\bar{D}}.$$

When we write

$$C_{ABCD} = X_{AB}^i C_{ij} X_{CD}^j$$

we have

$$\begin{aligned} C_{ABCD} &= C_0 \phi_A \phi_B \phi_C \phi_D + 4C_1 \phi_{(A} \phi_B \phi_C \psi_{D)} \\ &\quad + 6C_2 \phi_{(A} \phi_B \psi_C \psi_{D)} + 4C_3 \psi_{(A} \psi_B \psi_C \phi_{D)} + C_4 \psi_A \psi_B \psi_C \psi_D \end{aligned}$$

where the parentheses denote the symmetric sum and

$$\|C_{ij}\| = \left\| \begin{array}{ccc} C_4 & -C_3 & C_2 \\ -C_3 & C_2 & -C_1 \\ C_2 & -C_1 & C_0 \end{array} \right\|$$

2. THE COVARIANTLY CONSTANT CONDITION

This condition may be expressed as the equations

$$\phi_{A;\mu} = 0 \quad (2.1)$$

The integrability conditions of these equations are

$$\phi_{A;\mu\nu} - \phi_{A;\nu\mu} = \phi_C R_{A\mu\nu}^C = 0 \quad (2.2)$$

In this section we shall discuss consequences of equations (2.1). It follows from these equations and equations (1.3) that

$$A_{0\mu} = A_{1\mu} = 0 \quad (2.3)$$

Equations (1.4) then imply that

$$R_{i\mu\nu} = \delta_i^2 (A_{2\nu;\mu} - A_{2\mu;\nu}) \quad (2.4)$$

Since the quantities $p_{\mu\nu}^i$ (and $\bar{p}_{\mu\nu}^i$) provide a basis for self-dual (and anti-self-dual) two forms in V , we may write

$$A_{2\nu;\mu} - A_{2\mu;\nu} = \frac{1}{8} (Z^i p_{i\mu\nu} + \zeta^i \bar{p}_{i\mu\nu})$$

where

$$\begin{aligned} Z^i &= p^{i\mu\nu} (A_{2\nu;\mu} - A_{2\mu;\nu}) \\ \zeta^i &= \bar{p}^{i\mu\nu} (A_{2\nu;\mu} - A_{2\mu;\nu}) \end{aligned}$$

Hence

$$R_{i\mu\nu} = \frac{1}{8} (\delta_i^2 Z_j p_{\mu\nu}^j + \delta_i^2 \zeta_j \bar{p}_{\mu\nu}^j)$$

Then

$$C_{ij} - \frac{R}{3} A_{ij} = \delta_i^2 Z_j$$

Since the left hand side of this equation is symmetric in i and j we must have

$$Z_j = \delta_j^2 Z$$

Similarly we have

$$S_{i\bar{j}} = \delta_i^2 \zeta_j = \bar{S}_{j\bar{2}}$$

Hence

$$\zeta_i = \zeta \delta_i^2$$

with

$$\zeta = \bar{\zeta}.$$

Thus

$$C_{ij} - \frac{1}{3} R a_{ij} = Z \delta_i^2 \delta_j^2.$$

Since

$$a^{ij} C_{ij} = 0$$

we have

$$-R = Z a^{22} = 0$$

and

$$C_{ij} = Z \delta_i^2 \delta_j^2.$$

Hence

$$\begin{aligned} C_0 &= Z \\ C_1 &= C_2 = C_3 = C_4 = 0 \end{aligned}$$

and

$$C_{ABCD} = Z \phi_A \phi_B \phi_C \phi_D \quad (2.5)$$

Since

$$R_{AB\dot{C}\dot{D}} = S_{ij} X_{AB}^i \bar{X}_{CD}^j = - \left(R_{\mu\rho} - \frac{R}{4} g_{\mu\rho} \right) \gamma_{CA}^\rho \gamma_{DB}^\mu$$

it follows from the expression for S_{ij} that

$$\zeta \phi_A \phi_B \bar{\phi}_C \bar{\phi}_D = - \left(R_{\mu\rho} - \frac{R}{4} g_{\mu\rho} \right) \gamma_{CA}^\rho \gamma_{DB}^\mu$$

The spinors ϕ_A and ψ_A define a null tetrad of vectors by the equations

$$l_\mu = \frac{1}{\sqrt{2}} \gamma_{\mu\dot{A}\dot{B}} \bar{\phi}_A \phi_B$$

$$n_\mu = \frac{1}{\sqrt{2}} \gamma_{\mu\dot{A}\dot{B}} \bar{\psi}_A \psi_B$$

$$m_\mu = \frac{1}{\sqrt{2}} \gamma_{\mu\dot{A}\dot{B}} \bar{\psi}_A \psi_B$$

$$\bar{m}_\mu = \frac{1}{\sqrt{2}} \gamma_{\mu\dot{A}\dot{B}} \bar{\phi}_A \psi_B$$

On using the fact that

$$\gamma_{CA}^\rho \gamma_\sigma^{\dot{C}A} = -2\delta_\sigma^\rho$$

it follows from the above expression that

$$R_{\sigma\tau} = -\frac{1}{2} \zeta l_\sigma l_\tau. \quad (2.6)$$

Where l_σ is the covariantly constant null vector associated with the covariantly constant spinor ϕ_A .

Note that since

$$R_{AB\mu\nu} = X_{AB}^i R_{i\mu\nu} = 1/8 \phi_A \phi_B (Z p_{\mu\nu}^2 + \zeta \bar{p}_{\mu\nu}^2).$$

Hence equations (2.2), the integrability equations of equations (2.1) are satisfied.

3. THE BIANCHI IDENTITIES

These identities imply that

$$C_{\lambda\mu\nu;\kappa}^{\kappa} = -\frac{1}{2} \left[\left(R_{\lambda\nu} - \frac{R}{6} g_{\lambda\nu} \right)_{;\mu} - \left(R_{\lambda\mu} - \frac{R}{6} g_{\lambda\mu} \right)_{;\nu} \right]. \tag{3.1}$$

The spinor

$$C_{ABCD} = \frac{1}{4} p_{AB}^{\rho\sigma} C_{\rho\sigma\mu\nu} p_{CD}^{\mu\nu}.$$

Hence

$$\begin{aligned} \gamma^{\tau\dot{E}A} C_{ABCD;\tau} &= -\frac{1}{4} \gamma^{\tau\dot{E}}_A p^{\rho\sigma A}_B C_{\rho\sigma\mu\nu;\tau} p_{CD}^{\mu\nu} \\ &= -C^{\tau}_{\alpha\mu\nu;\tau} \gamma^{\alpha\dot{E}}_B p_{CD}^{\mu\nu} \end{aligned} \tag{3.2}$$

as follows from the fact that

$$\gamma^{\tau\dot{E}A} p^{\sigma\rho A}_B = - [E^{\alpha\tau\sigma\rho} + g^{\alpha\sigma} g^{\tau\rho} - g^{\alpha\rho} g^{\tau\sigma}] \gamma^{\dot{E}}_{\alpha B}$$

and the fact that the conformal tensor is equal to its double dual.

Substituting from equations (2.5), (2.6) and (3.1) into equations (3.2) we obtain

$$\gamma^{\tau\dot{E}A} Z_{,\tau} \phi_A \phi_B \phi_C \phi_D = -\frac{1}{2} l_{\alpha} \gamma^{\alpha\dot{E}}_B \zeta_{,\mu} l_{\nu} p_{CD}^{\mu\nu}. \tag{3.3}$$

On multiplying these equations by $\psi^B \psi^C \psi^D$ and summing one finds that

$$\gamma^{\tau\dot{E}A} Z_{,\tau} \phi_A = l_{\alpha} \gamma^{\alpha\dot{E}A} \psi_A \zeta_{,\mu} \bar{m}^{\mu}$$

on using the fact that

$$p_{CD}^{\mu\nu} \psi^C \psi^D = 2(n^{\mu} \bar{m}^{\nu} - n^{\nu} \bar{m}^{\mu})$$

Hence

$$l^{\tau} Z_{,\tau} = 0 \tag{3.4}$$

$$m^{\tau} Z_{,\tau} = -\zeta_{,\tau} m^{\tau} \tag{3.5}$$

since

$$l_{\alpha} m^{\alpha} = l_{\alpha} n^{\alpha} + 1 = 0.$$

Equations (3.4) and (3.5) are equivalent to equations (3.3).

It follows from equation (3.3) that Z is a constant along the null geodesic with tangent l_{μ} . If the space-time is asymptotically flat and such a null geodesic reaches null infinity Z must vanish there. Therefore it must vanish all along the null geodesic. Hence Z must vanish at all events

of space-time which lie on the infinitely extendible null geodesics with covariantly constant tangent vectors l_μ . At these events of V the conformal tensor of V vanishes. The Ricci tensor of V is given by equation (2.6) with ζ satisfying

$$\bar{m}^r \zeta_{,r} = 0$$

at these events of V . The space time V is flat if all events of V are of the type described above and $\zeta = 0$, that is the space time is empty.

APPENDIX

Algebraic Properties of the Spinors $\gamma^{\mu\dot{A}B}$ and $p^{\mu\nu C D}$.

The former spinors are defined as follows:

$$\|\gamma^{\mu\dot{A}B}\| = q^\mu$$

where

$$\bar{q}^\mu q^\nu + \bar{q}^\nu q^\mu = 2g^{\mu\nu}1_2,$$

the bar over a quantity denoting its complex conjugate. That is,

$$\bar{\gamma}^{\mu\dot{A}B}\gamma^{\nu\dot{B}C} + \bar{\gamma}^{\nu\dot{A}B}\gamma^{\mu\dot{B}C} = 2g^{\mu\nu}\delta^{\dot{A}C} \quad (\text{A.1})$$

Hence

$$\gamma^{\mu\dot{A}B}\gamma^{\nu\dot{B}A} = 2g^{\mu\nu} \quad (\text{A.2})$$

The spinors

$$\gamma^{\mu\dot{A}B} = \bar{e}_{AC}\gamma^{\mu\dot{C}B} \quad (\text{A.3})$$

satisfy

$$\bar{\gamma}^{\mu\dot{A}B} = \gamma^{\mu\dot{B}A} \quad (\text{A.4})$$

that is form Hermitian matrices. Equations (A.2) are thus equivalent to

$$\gamma^{\mu\dot{A}B}\gamma_{\nu\dot{A}B} = -2\delta_\nu^\mu \quad (\text{A.5})$$

with

$$\begin{aligned} \gamma_{\nu\dot{A}B} &= g_{\nu\rho}\gamma_{\dot{A}B}^\rho \\ \gamma_{\nu\dot{A}B} &= \gamma_{\nu\dot{C}B}^{\dot{A}CB} \end{aligned}$$

Since the four matrices $\gamma^{\mu\dot{A}B}$ form a basis for 2×2 Hermitian matrices $H_{\dot{A}B}$ we may write

$$H_{\dot{A}B} = h_\sigma \gamma_{\dot{A}B}^\sigma$$

with

$$h_\sigma = -\frac{1}{2} H_{\dot{A}B} \gamma_{\dot{A}B}^{\dot{A}B\sigma} = \bar{h}_\sigma.$$

Since the mapping from $H_{\dot{A}B}$ to h^σ is one to one, we must have

$$\gamma_{\dot{A}B}^{\dot{A}B\sigma} \gamma_{\dot{C}D}^{\dot{A}B\sigma} = -2\delta_C^{\dot{A}} \delta_D^{\dot{B}} \quad (\text{A.6})$$

The spinor $p_{\mu\nu}^{\dot{A}B}$ defined by the equations

$$\|p_{\mu\nu}^{\dot{A}B}\| = p_{\mu\nu} = \frac{1}{2}(\bar{q}_\mu q_\nu - \bar{q}_\nu q_\mu) = -p_{\nu\mu} \quad (\text{A.7})$$

satisfies

$$\text{trace } p_{\mu\nu} = p_{\mu\nu}^{\dot{A}A} = 0 \quad (\text{A.8})$$

$$p^{\mu\nu} = \frac{1}{2} E^{\mu\nu\sigma\tau} p_{\sigma\tau} = g^{\mu\sigma} g^{\nu\tau} p_{\sigma\tau} \quad (\text{A.9})$$

$$\bar{p}^{\mu\nu} = \frac{1}{2} E^{\mu\nu\sigma\tau} \bar{p}_{\sigma\tau} = -g^{\mu\sigma} g^{\nu\tau} p_{\sigma\tau} \quad (\text{A.10})$$

where

$$E^{\sigma\tau\mu\nu} = \frac{1}{\sqrt{g}} e^{\sigma\tau\mu\nu}; \quad E_{\sigma\tau\mu\nu} = \sqrt{g} \varepsilon_{\sigma\tau\mu\nu}$$

$\varepsilon^{\sigma\tau\mu\nu}$ is the Levi-Civita tensor density, g is the determinant of the metric tensor $g_{\mu\nu}$ and hence is a pure imaginary tensor. The $p_{\mu\nu}$ form a (redundant) basis for all traceless 2×2 matrices.

It follows from equations (A.1) and (A.7) that

$$\bar{\gamma}^{\mu\dot{A}}_B \gamma^{\nu\dot{B}}_C = g^{\mu\nu} \delta^A_C + p^{\mu\nu A}_C \quad (\text{A.11})$$

It may be verified by choosing a particular coordinate system in V and in the spin space that

$$p_{\lambda\mu} p_{\sigma\tau} = - (E_{\lambda\mu\sigma\tau} - g_{\lambda\tau} g_{\mu\sigma} + g_{\lambda\sigma} g_{\mu\tau}) 1_2 \quad (\text{A.12})$$

Hence

$$\| \gamma^{\dot{A}}_{\mu B} p^B_{\sigma\tau C} \| = q_{\mu} p_{\sigma\tau} = - (E_{\nu\mu\sigma\tau} - g_{\nu\tau} g_{\mu\sigma} + g_{\nu\sigma} g_{\mu\tau}) q^{\nu} \quad (\text{A.13})$$

$$\| p^A_{\sigma\tau B} \gamma^{\dot{B}}_C \| = p_{\sigma\tau} \bar{q}_\mu = (E_{\nu\mu\sigma\tau} - g_{\nu\tau} g_{\mu\sigma} + g_{\nu\sigma} g_{\mu\tau}) \bar{q}^{\nu} \quad (\text{A.14})$$

It follows from (A.12) that

$$p_{\lambda\mu AB} p^{\dot{A}\dot{B}}_{\sigma\tau} = - p^{\dot{A}}_{\lambda\mu B} p^{\dot{B}}_{\sigma\tau A} = 2(E_{\lambda\mu\sigma\tau} + g_{\lambda\sigma} g_{\mu\tau} - g_{\lambda\tau} g_{\mu\sigma}) \quad (\text{A.15})$$

It may be verified from the definition of $p_{\mu\nu AB}$ and equation (A.6) that

$$p^{\sigma\tau}_{CD} p^{\dot{A}\dot{B}}_{\sigma\tau} = 4(\delta^A_C \delta^B_D + \delta^B_C \delta^A_D) \quad (\text{A.16})$$

$$\gamma^{\sigma\dot{C}\dot{D}} p_{\sigma\tau AB} = - (\gamma^{\dot{C}}_{\tau A} \delta^D_B + \gamma^{\dot{D}}_{\tau B} \delta^C_A) \quad (\text{A.17})$$

$$p_{\mu\nu}{}^{AB} \bar{p}^{\nu}_{\sigma}{}^{CD} = - 2\gamma_{\mu}{}^{\dot{C}\dot{A}} \gamma_{\sigma}{}^{\dot{D}\dot{B}} - g_{\mu\sigma} \varepsilon^{AB} \varepsilon^{\dot{C}\dot{D}} - p_{\mu\sigma}{}^{AB} \varepsilon^{\dot{C}\dot{D}} - \bar{p}_{\mu\sigma}{}^{CD} \varepsilon^{\dot{A}\dot{B}} \quad (\text{A.18})$$

If ϕ^A and ψ^A are a pair of spinors satisfying

$$\phi^A \psi_A = 1 \quad (\text{A.19})$$

then

$$\phi^A \psi_B - \psi^A \phi_B = \delta^A_B. \quad (\text{A.20})$$

The vectors

$$\begin{aligned} l_{\mu} &= \frac{1}{\sqrt{2}} \gamma_{\mu\dot{A}\dot{B}} \bar{\phi}^A \phi^B \\ n_{\mu} &= \frac{1}{\sqrt{2}} \gamma_{\mu\dot{A}\dot{B}} \bar{\psi}^A \psi^B \\ m_{\mu} &= \frac{1}{\sqrt{2}} \gamma_{\mu\dot{A}\dot{B}} \bar{\psi}^A \phi^B \\ \bar{m}_{\mu} &= \frac{1}{\sqrt{2}} \gamma_{\mu\dot{A}\dot{B}} \bar{\phi}^A \psi^B \quad p^{\nu}_{\sigma} \end{aligned} \quad (\text{A.21})$$

form a null tetrad such that

$$g_{\mu\nu} = - l_{\mu} n_{\nu} - n_{\mu} l_{\nu} + m_{\mu} \bar{m}_{\nu} + \bar{m}_{\mu} m_{\nu} \quad (\text{A.22})$$

The quantities

$$p_0{}^{\mu\nu} = p^{\mu\nu}_{AB} \phi^A \phi^B = 2(m^{\mu} l^{\nu} - m^{\nu} l^{\mu}) = \dot{p}_0{}^{\mu\nu}$$

$$p_1{}^{\mu\nu} = p^{\mu\nu}_{AB} \phi^A \phi^B = (m^{\mu} \bar{m}^{\nu} - \bar{m}^{\mu} m^{\nu} + n^{\mu} l^{\nu} - n^{\nu} l^{\mu}) = \dot{p}_1{}^{\mu\nu}$$

$$p_2{}^{\mu\nu} = p^{\mu\nu}_{AB} \psi^A \psi^B = 2(n^{\mu} \bar{m}^{\nu} - n^{\nu} \bar{m}^{\mu}) = p_2{}^{\nu\mu}$$

where

$$\dot{p}_i{}^{\mu\nu} = \frac{1}{2} E^{\mu\nu\sigma\tau} g_{\sigma\alpha} g_{\tau\beta} p_i{}^{\alpha\beta}.$$

That is the $p_i^{\mu\nu}$ are three linearly independent self-dual anti-symmetric tensors. Their complex conjugates are anti-self-dual anti-symmetric tensors.

It may be verified that

$$\begin{aligned} p_i^{\mu\nu} p_{j\mu\nu} &= 8a_{ij} \\ p_i^{\mu\nu} \bar{p}_{j\mu\nu} &= 0 \end{aligned}$$

where

$$\| a_{ij} \| = \begin{vmatrix} 0 & 0 & 1 \\ 0 & -\frac{1}{2} & 0 \\ 1 & 0 & 0 \end{vmatrix}.$$

REFERENCE

- [J] A. H. TAUB, « Curvature Invariants, Characteristic Classes and Petrov Classification of Space-Times », in *Differential Geometry and Relativity* Cahen & Flato (eds.), D. Reidel Publishing Co., Dordrecht-Holland, 1976.

(Manuscrit reçu le 12 Janvier 1984)

Reproduction in whole or in part without the permission of the author or his representative is prohibited (law of March 11, 1957, Article 40, line 1). Such reproduction by whatever means, constitutes an infringement forbidden by Article 425 and those following it, of the Penal Code. The law of March 11, 1957, lines 2 and 3 of Article 41, authorizes only those copies or reproductions made for the exclusive use of the copyst, and not intended for collective use and such analyses and short quotations as are made for the purposes of an example or illustration.

© GAUTHIER-VILLARS 1984

Toute représentation ou reproduction intégrale ou partielle, faite sans le consentement de l'auteur ou de ses ayants droit ou ayants cause, est illicite (loi du 11 mars 1957, alinéa 1^{er} de l'article 40). Cette représentation ou reproduction, par quelque procédé que ce soit, constituerait une contrefaçon sanctionnée par les articles 425 et suivants du Code pénal. La loi du 11 mars 1957 n'autorise, aux termes des alinéas 2 et 3 de l'article 41, que les copies ou reproductions strictement réservées à l'usage privé du copiste et non destinées à une utilisation collective d'une part et d'autre part que les analyses et les courtes citations dans un but d'exemple et d'illustration.

© GAUTHIER-VILLARS 1984

COPYRIGHT

The appearance of the code at the bottom of the first page of an article in this journal indicates the copyright owner's consent that copies of the article may be made for personal or internal use, or for the personal or internal use of specific clients. This consent is given on the condition, however, that the copier pay the stated per-copy fee through the Copyright Clearance Center, Inc., Operations Center, 21 Congress St., Salem, Mass. 01970, U.S.A. for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purpose, for creating new collective works, or for resale.

© GAUTHIER-VILLARS 1984