

# ANNALES DE L'I. H. P., SECTION B

KAI LAI CHUNG

## **The lifetime of conditional brownian motion in the plane**

*Annales de l'I. H. P., section B*, tome 20, n° 4 (1984), p. 349-351

[http://www.numdam.org/item?id=AIHPB\\_1984\\_\\_20\\_4\\_349\\_0](http://www.numdam.org/item?id=AIHPB_1984__20_4_349_0)

© Gauthier-Villars, 1984, tous droits réservés.

L'accès aux archives de la revue « *Annales de l'I. H. P., section B* » (<http://www.elsevier.com/locate/anihpb>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

## The lifetime of conditional Brownian motion in the plane

by

Kai Lai CHUNG (\*)

Department of Mathematics, Stanford University, CA 94305, USA

SUMMARY. — In this note I give a short and perspicacious proof of a recent remarkable result due to Cranston and McConnell [3].

RÉSUMÉ. — Cette note est consacrée à une démonstration courte d'un résultat remarquable et récent de Cranston et McConnell [3].

Let  $D$  be a bounded domain in  $\mathbb{R}^d$ ,  $d \geq 1$ ;  $H(D)$  the class of strictly positive harmonic functions in  $D$ ;  $X = \{X_t, t \geq 0\}$  the standard Brownian motion in  $\mathbb{R}^d$ ;  $\tau_B = \inf \{t > 0: X_t \notin B\}$  for any Borel set  $B$ ;  $m$  the Lebesgue measure in  $\mathbb{R}^d$ ;  $E_h^x$  the expectation associated with the  $h$ -conditioned Brownian motion starting at  $x \in D$ .

THEOREM. — Let  $d = 2$ . There exists a constant  $C$  depending only on  $D$  such that

$$(1) \quad \sup_{\substack{x \in D \\ h \in H(D)}} E_h^x \{ \tau_D \} \leq Cm(D).$$

We begin by stating explicitly the case where  $h \equiv 1$ , namely unconditioned Brownian motion, for a general Borel set  $B$  in  $\mathbb{R}^d$ ,  $d \geq 1$ .

(\*) Research supported in part by NSF grant MCS83-01072 at Stanford University.

LEMMA. — We have

$$\sup_{x \in D} E^x \{ \tau_B \} \leq A_d m(D)^{2/d}$$

where

$$A_d = \frac{1}{2\pi d^2} (d + 1)^{\frac{2(d+1)}{d}}.$$

This lemma can be proved by an elementary method using only the strong Markov property of  $X$  and the form of its transition density. It is generalizable and adaptable to similar estimates; see [1], p. 148 ff.

As the first simplification in the proof of the theorem, we deal directly with a general  $h$  in  $H(D)$ . This spares us some unnecessary « hard theory », such as the famous Martin representation, and the behavior of a minimal harmonic function at the boundary. Cf. Lemma 2.2 in [3], which is actually a result due to Doob. Thus for any  $h \in H(D)$ , we put for clarity:

$$(2) \quad Y(t) = \begin{cases} \left(\frac{1}{h}\right)(X_t) & \text{for } 0 \leq t < \tau_D, \\ 0 & \text{for } t \geq \tau_D. \end{cases}$$

It is a basic idea in  $h$ -conditioning that  $\{Y_t, \mathcal{F}_t, t \geq 0\}$  is a super-martingale, where  $\{\mathcal{F}_t\}$  is the natural filtration of  $\{X_t\}$ . Let  $0 < a < b < \infty$ ; let  $D'[a, b]$  and  $U'[a, b]$  denote respectively the number of downcrossings and upcrossings of  $[a, b]$  by  $\{Y_t, t \geq 0\}$ . Then we have for any  $x \in D$ :

$$(3) \quad E^x \{ D'[a, b] \} \leq \frac{b}{b-a}; \quad E^x \{ U'[a, b] \} \leq \frac{a}{b-a}.$$

For the first inequality (due to G. A. Hunt), see e. g. [2], p. 341; the second does not follow trivially from the first, but both follow from Dubins's inequalities (*loc. cit.*). Taking reciprocals, we deduce that if  $D[a, b]$  and  $U[a, b]$  denote the corresponding numbers for  $\{h(X_t), t \geq 0\}$ , then

$$(4) \quad E^x \{ U[a, b] \} \leq \frac{a}{b-a}; \quad E^x \{ D[a, b] \} \leq \frac{b}{b-a}.$$

We now define for any  $x_0 \in D$ :

$$C_n = \{ x \in D : h(x) = 2^n h(x_0) \},$$

$$D_n = \{ x \in D : 2^{n-1} h(x_0) < h(x) < 2^{n+1} h(x_0) \},$$

where  $n$  is an integer. Furthermore, we denote by  $N_n$  the total number of times a path moves from inside  $D_n$  to outside  $D_n$ . If it starts from  $C_n$ , this

can be done either by a downcrossing of  $[2^n h(x_0), 2^{n-1} h(x_0)]$ , or an upcrossing of  $[2^n h(x_0), 2^{n+1} h(x_0)]$ . Hence we have by (5):

$$(6) \quad \sup_{x \in C_n} E_h^x \{ N_n \} \leq \frac{2^{n-1}}{2^n - 2^{n-1}} + \frac{2^{n+1}}{2^{n+1} - 2^n} = 3.$$

Next, it is plain that for any  $x \in C_n$  :

$$(7) \quad E_h^x \{ \tau_{D_n} \} = \frac{1}{h(x)} E^x \left\{ \int_0^{\tau_{D_n}} h(X_t) dt \right\} \leq 2 E^x \{ \tau_{D_n} \}.$$

It remains to add up all the crossings, *without ordering*. But we must not forget that a path may leave  $D$  before completing the last crossing. In this case 1 must be added to  $N_n$  in the counting. Therefore our final estimate is as follows:

$$(8) \quad E_h \{ \tau_D \} \leq (3 + 1) \sum_{n=-\infty}^{\infty} 2 \sup_{x \in C_n} E^x \{ \tau_{D_n} \} \leq 8 A_d \sum_{n=-\infty}^{\infty} m(D_n)^{\frac{2}{d}}$$

by (6), (7) and the lemma. For  $d = 2$ , this yields (1) with  $C = 8A_d$ .

REFERENCES

[1] K. L. CHUNG, *Lectures from Markov Processes to Brownian motion, Grundlehren der mathematischen Wissenschaft*, t. 249, Springer-Verlag, 1982.  
 [2] K. L. CHUNG, *A course in Probability Theory*, 2nd edition, Academic Press, 1974.  
 [3] M. CRANSTON and T. R. MCCONNELL, The lifetime of conditioned Brownian motion, *Z. Wahrsch. Verw. Gebiete*, t. 65, 1983, p. 1-11.

(Manuscrit reçu le 16 mars 1984)