

Astérisque

UDO LEISERING

On the p -part of character degrees of solvable groups

Astérisque, tome 181-182 (1990), p. 217-220

http://www.numdam.org/item?id=AST_1990__181-182__217_0

© Société mathématique de France, 1990, tous droits réservés.

L'accès aux archives de la collection « Astérisque » (<http://smf4.emath.fr/Publications/Asterisque/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ON THE p -PART OF CHARACTER DEGREES OF SOLVABLE GROUPS ¹

by
Udo Leisering

1 Introduction

The Fong-Swan Theorem shows a relation between irreducible Brauer characters and ordinary irreducible characters by the following: Let φ be an irreducible Brauer character of a p -solvable group G . There exists a p -rational irreducible character χ of G , such that $\chi = \varphi$ as a Brauer character. Especially: Every condition on ordinary characters is valid for Brauer characters (in a p -solvable group). We now ask for a kind of inversion of this relation and consider the character degrees. Let q be a prime, such that $q^2 \nmid \beta(1)$ for all $\beta \in IBr_p(G)$. Do we get a bound $n \in \mathbb{N}$, such that $q^n \nmid \chi(1)$ for all $\chi \in Irr(G)$? In general this is impossible.

2 Example

Let $\Gamma(8)$ be the group of all semilinear maps on $GF(8)$. It is easy to prove that the set of degrees of all irreducible Brauer characters in characteristic 7 is $cd_7(\Gamma(8)) := \{1, 7\}$. The set of ordinary character degrees is $cd(\Gamma(8)) := \{1, 7, 3\}$. We put

$$H_n := \Gamma(8) \times \dots \times \Gamma(8).$$

Clearly: $3 \nmid \beta(1)$ for all $\beta \in IBr_7(H_n)$, but there exists a $\chi \in Irr(H_n)$, such that $\chi(1) = 3^n$. (There exists many other examples for primes $p \neq 7$ too.)

For the prime p it is possible to show the following:

3 Theorem

Let G be solvable with $O_p(G) = E$ and $p^2 \nmid \beta(1)$ for all $\beta \in IBr_p(G)$. It follows:

- a) G has elementary abelian Sylow- p -subgroups.
- b) $p^2 \nmid \chi(1)$ for all $\chi \in Irr(G)$.

Before we start to prove this theorem, we need some short lemma:

¹This is a small part of a Dissertation at Mainz.

4 Lemma

Let $N \trianglelefteq G$, V an irreducible KN -modul and K a field. Further let $T := T_G(V)$ be the inertia group of V and W an irreducible KT -modul, such that $W_N = eV$ with $e \in IN$. Then W^G is an irreducible KG -modul and

$$|G : T_G(V)| \cdot \dim V \mid \dim W.$$

Proof: Manz[4]; Lemma 1 .

5 Lemma

Let N be an abelian normal subgroup of G , such that $(|G/C_G(N)|, |N|) = 1$ and $G/C_G(N)$ is abelian.

- a) $G/C_G(N)$ has a regular orbit on N , $Irr(N)$ and $IBr_p(N)$ if $p \nmid |N|$.
- b) There exists a $\chi \in Irr(G)$ resp. $\beta \in IBr_p(G)$ if $p \nmid |N|$, such that

$$|G/C_G(N)| \mid \chi(1) \quad \text{resp.} \quad |G/C_G(N)| \mid \beta(1).$$

Proof:

- a) Isaacs[3], 13.24 yields that $Irr(N)$ and N are isomorphic as permutation modules. Then the result follows by Passman[5], lemma 2.2 .
- b) This follows by a) and Lemma 4.

6 Proof of theorem 3

We denote by $F(G)$ the Fitting subgroup of G and define $F_j/F_{j-1} := F(G/F_{j-1})$, ($F_0 := E$).

- a) Let G be a minimal counterexample to statement a).

- (i) $\Phi(G) = E$, where $\Phi(G)$ is the Frattini subgroup of G :

Proof: Clear, since $p \nmid |F_1|$ and $F(G/\Phi(G)) = F_1/\Phi(G)$.

- (ii) $p \mid |F_2/F_1|$, but $p^2 \nmid |F_2/F_1|$:

Proof: Assume that $p \nmid |F_2/F_1|$, hence $O_p(G/F_1) = E$. Since G is a minimal counterexample, G/F_1 has elementary abelian Sylow- p -subgroups and out of $p \nmid |F_1|$ it follows, that G has elementary abelian Sylow- p -subgroups too; a contradiction. Therefore $p \mid |F_2/F_1|$.

Let $P_0/F_1 \in Syl_p(F_2/F_1)$ and $A \leq G$, such that A/F_1 is a maximal abelian normal subgroup of P_0/F_1 , esp. $A \trianglelefteq G$. Hence $p^2 \nmid \alpha(1)$ for all $\alpha \in IBr_p(A)$ (Clifford theory). Since we have $C_A(F_1) = F_1$, Lemma 5 yields an $\alpha \in IBr_p(A)$, such that $|A/F_1| = \alpha(1)$. Hence $|A/F_1| \mid p$ and therefore $A = P_0$.

- (iii) Conclusion: Let $P_0 \trianglelefteq G$ be as defined in (ii). Lemma 5a) yields a $\lambda_0 \in IBr_p(F_1)$, such that

$$T_G(\lambda_0) \cap P_0 = F_1.$$

In particular $p \mid |G : T_G(\lambda_0)|$. But by Lemma 4 we now have, that $p^2 \nmid |G : T_G(\lambda_0)|$ since $p^2 \nmid \beta(1)$ for all $\beta \in IBr_p(G)$. Hence

$$|T_G(\lambda_0)|_p = \frac{1}{p} |G|_p.$$

CHARACTER DEGREES

Now let $P_1/F_1 \in \text{Syl}_p(T_G(\lambda_0)/F_1)$. Obviously

$$P_0/F_1 \cap P_1/F_1 = E$$

and because of the order of $T_G(\lambda_0)$ we get

$$(P_0P_1)/F_1 \in \text{Syl}_p(G/F_1).$$

We claim, that P_1/F_1 is elementary abelian:

Since $P_0/F_1 = O_p(G/F_1)$ it is obviously, that $O_p(G/P_0) = E$. Furthermore G is a minimal counterexample and therefore G/P_0 has elementary abelian Sylow- p -subgroups. Hence $P_1/F_1 \cong (P_1P_0)/P_0$ is elementary abelian.

Step (ii) yields $|P_0/F_1| = p$ and therefore P_1/F_1 is operating trivial on P_0/F_1 . So $(P_0P_1)/F_1$ is elementary abelian and for any $P \in \text{Syl}_p(G)$ we have:

$$P = P/(P \cap F_1) \cong (PF_1)/F_1 \cong (P_0P_1)/F_1 \text{ is elementary abelian.}$$

- b) Assume $p^2 \mid \chi(1)$ for a $\chi \in \text{Irr}(G)$. Let D be the defect group of the block of χ . Then it follows by a) that D is elementary abelian. If the p -part of the order of G is $|G|_p = p^a$ and $|D| = p^d$, Brauer's Theorem about the height of characters yields, that $p^{(a-d)} \mid \chi(1)$ and $p^{a-d+1} \nmid \chi(1)$ (Fong[2];Thm.3C). Since we have the assumption $p^2 \mid \chi(1)$ it follows, that $a - d \geq 2$. Let now $\beta \in \text{IBr}_p(G)$ belong to the same block as χ . Then $p^{a-d} \mid \beta(1)$ (Feit[1];IV,4.5) and therefore $p^2 \mid \beta(1)$; a contradiction.

7 Remark

Tsushima proved in [6]; Theorem 5 the following:

If G is solvable and $p^2 \nmid \chi(1)$ for all $\chi \in \text{Irr}(G)$, then $G/O_p(G)$ has elementary abelian Sylow- p -subgroups.

With the theorem of Fong-Swan (Feit[1];X,2.1) this is a corollary of theorem 1. (Or use the proof of theorem 1 directly).

References

[1] W.Feit - "The Representation Theory of Finite Groups", North-Holland, Amsterdam, 1982.
 [2] P.Fong - On the Characters of p -solvable groups, Trans. Amer. Math. Soc. 98 (1961), 263-284.
 [3] I.M.Isaacs - "Character Theory of Finite Groups", Academic Press, New York, San Francisco, London 1976.
 [4] O.Manz - On the modular version of Ito's Theorem on character degrees for groups of odd order, Nagoya Math. J. 105 (1987), 121-128.

[5] D.S.Passman - Groups with normal, solvable Hall p' -subgroups,
Trans. Amer. Math. Soc. 123 (1966), 99-111.

[6] Y.Tsushima - On the weakly regular p -blocks with respect to $O_{p'}(G)$,
Osaka J. Math. 14 (1977), 465-470.

Udo Leisering
Fachbereich 17, Mathematik
Johannes Gutenberg-Universität
Saarstr.21
6500 Mainz