

COMPOSITIO MATHEMATICA

JAMES B. CARRELL

Orbits of the Weyl group and a theorem of DeConcini and Procesi

Compositio Mathematica, tome 60, n° 1 (1986), p. 45-52

http://www.numdam.org/item?id=CM_1986__60_1_45_0

© Foundation Compositio Mathematica, 1986, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

ORBITS OF THE WEYL GROUP AND A THEOREM OF DECONCINI AND PROCESI

James B. Carrell *

§1. Introduction

In [DP], C. DeConcini and C. Procesi proved a remarkable connection between dual SL_n conjugacy classes of nilpotent complex matrices. The goal of this paper is to show that weaker form of this connection holds for certain pairs of nilpotent conjugacy classes for an arbitrary semi-simple algebraic group G over \mathbb{C} . We also obtain a short intrinsic proof of the formula of [DP]. The main techniques in this paper are analytic, namely applications of the ideas on torus actions developed in [ACL].

First, recall the SL_n case. For an $n \times n$ nilpotent complex matrix σ , let $\pi(\sigma) = \sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_k$ be the partition of n determined by the sizes of the Jordan blocks of σ . Two nilpotent conjugacy classes O_σ and O_τ are called dual if $\pi(\sigma)$ and $\pi(\tau)$ are dual partitions of n . Let $A_\sigma = A(\overline{O_\sigma} \cap \mathcal{D}_n)$ be the coordinate ring of the scheme theoretic intersection of $\overline{O_\sigma}$ and the diagonal $n \times n$ matrices \mathcal{D}_n over \mathbb{C} with trace zero. Also, let X^τ be the variety of Borel subalgebras of sl_n containing τ . In [DP], it was shown that there exists an S_n -equivariant \mathbb{C} -algebra isomorphism

$$A_\sigma \cong H^*(X^\tau)(\mathbb{C}\text{-coefficients}), \quad (1)$$

provided σ and τ are dual. Moreover, the S_n -module structure is the induced representation $\text{Ind}_{S_{\sigma_1} \times \dots \times S_{\sigma_m}}^{S_n}(\mathbb{C})$ which verifies a conjecture of H. Kraft [Kr] (inspired by some questions of Kostant). In [Ta], Tanisaki simplified the proof of (1) and extended the result on induced representations to certain symplectic conjugacy classes. Kraft explicitly pointed out, however, that $A(\overline{O} \cap \mathcal{D}_6)$ is not an induced representation for the class of type (3, 3) in $sl(6)$, [Kr].

The present treatment proceeds by showing that (1) factors into two homomorphisms, either of which may fail to be an isomorphism even when defined. One of these morphisms arises from the theory of sheets

* Supported in part by a grant from the Natural Sciences and Engineering Research Council of Canada

in a Lie algebra [BK]. The other relates the cohomology of a variety of fixed flags to a graded algebra associated to a certain Weyl group orbit.

Let $G \supset B \supset H$ be respectively a semi-simple algebraic group over \mathbb{C} , a Borel subgroup, and a maximal torus. By convention, Lie algebras will be denoted by the corresponding lower case script letter. Let W be the Weyl group of $(\mathfrak{g}, \mathfrak{h})$. For $s \in \mathfrak{h}$, consider the orbit $W \cdot s \subset \mathfrak{h}$ as a finite reduced variety with ring $A(W \cdot s)$ of regular functions, i.e. $A(W \cdot s) = A(\mathfrak{h})I(W \cdot s)$. $A(W \cdot s)$ is a W -module with a canonical W -invariant filtration $D_0 \subset D_1 \subset D_2 \subset \dots$, where D_i consists of restrictions of polynomials on \mathfrak{h} of degree $\leq i$. $D_i D_j \subset D_{i+j}$, so the graded ring $\text{Gr } A(W \cdot s) = \bigoplus D_i / D_{i-1}$ is canonically defined.

THEOREM 1: *Given $s \in \mathfrak{h}$, let τ be a regular nilpotent in the Levi subalgebra $\mathfrak{e} = \mathfrak{z}_{\mathfrak{g}}(s)$, and let X^τ denote the variety of Borel subalgebras of X containing τ . Let $i_\tau: X^\tau \rightarrow X$ be the inclusion into the flag variety of all Borel subalgebras of \mathfrak{g} . Then there exists a W -equivariant \mathbb{C} -algebra homomorphism $\psi_s: \text{Gr } A(W \cdot s) \rightarrow H^*(X^\tau)$. ψ_s doubles degree. The image of ψ_s is $i_\tau^* H^*(X)$. Moreover, ψ_s is an isomorphism if and only if i_τ^* is surjective. In fact, the kernel and cokernel of ψ have the same dimension.*

Note that W acts on $H^*(X^\tau)$ via the Springer representation ([Spr]), while the W -module $\text{Gr } A(W \cdot s)$ is $\text{Ind}_{W_L}^W(\mathbb{C})$, where W_L is the Weyl group of $(\mathfrak{e}, \mathfrak{h})$.

The second step of the factorization is the definition of a map $A(\overline{O}_\sigma \cap \mathfrak{h}) \rightarrow \text{Gr } A(W \cdot s)$ for a suitable nilpotent $\sigma \in \mathfrak{g}$. This step appears implicitly in [Kr, Prop. 4]. Begin by fixing a parabolic subalgebra \mathfrak{p} containing \mathfrak{e} as a Levi subalgebra, and let σ be a Richardson element of the nilradical $\text{nil } \mathfrak{p}$ (i.e. the P conjugacy class of σ is Zariski dense in $\text{nil } \mathfrak{p}$). One imposes two conditions on σ :

- (N) the closure \overline{O}_σ of the nilpotent G orbit is normal in \mathfrak{g} , and
- (S) $Z_P(\sigma) = Z_G(\sigma)$, i.e. the P and G stabilizers of σ coincide.

PROPOSITION 1: *Assuming (N) and (S), there exists a W -equivariant, surjective, graded \mathbb{C} -algebra homomorphism $\phi_\sigma: A_\sigma = A(\overline{O}_\sigma \cap \mathfrak{h}) \rightarrow \text{Gr } A(W \cdot s)$.*

Combining this with Theorem 1 we obtain our “generalization” of (1).

COROLLARY 1: *Let P be a parabolic in G whose nilradical contains a Richardson element σ satisfying (N) and (S). Then there exists a surjective degree doubling W -equivariant \mathbb{C} -algebra homomorphism*

$$\mu: A(\overline{O}_\sigma \cap \mathfrak{h}) \rightarrow i_\tau^* H^*(X)$$

where τ is a regular nilpotent in a Levi subalgebra of \mathfrak{g} .

The homomorphism μ admits a definition which is independent of s . Let \mathcal{N} be the nilpotent cone in \mathfrak{g} , and let $\beta: A(\mathcal{N} \cap \mathfrak{h}) \rightarrow H^*(X)$ be the

W -equivariant isomorphism obtained as the composition of isomorphisms

$$A(\mathcal{N} \cap \mathfrak{h}) \xrightarrow{\kappa} A(\mathfrak{h})/I_W \xrightarrow{\beta'} H^*(X).$$

Here I_W is the ideal generated by $\{f \in A(\mathfrak{h})^W \mid f(0) = 0\}$, κ is the natural map [Ko], and β' is the isomorphism of Borel which assigns to any element χ of the group $X(H)$ of characters on H the first Chern class $c_1(L_\chi)$ of the holomorphic line bundle L_χ on X . It will be clear from the definition of μ and the proof of Theorem 1 that we obtain a commutative diagram

$$\begin{array}{ccc} A(\mathcal{N} \cap \mathfrak{h}) & \xrightarrow{\beta} & H^*(X) \\ \downarrow & & \downarrow \\ A(\overline{O}_\sigma \cap \mathfrak{h}) & \xrightarrow{\mu_\sigma} & H^*(X^\tau) \end{array} \quad (2)$$

As an application, we now prove (1). Let σ and τ lie on dual nilpotent orbits in \mathfrak{sl}_n . Then τ may be chosen as a regular nilpotent in $\mathfrak{l} = \mathfrak{sl}_{\tau_1} \times \cdots \times \mathfrak{sl}_{\tau_m}$ and σ as a Richardson in $\mathfrak{nil} \mathfrak{p}$, where \mathfrak{p} is the parabolic corresponding to \mathfrak{l} . In \mathfrak{sl}_n condition (N) holds by [KP] and (S) is well known. Thus we obtain a surjection $A_\sigma \rightarrow i_\tau^* H^*(X)$, where here X is the flag variety of SL_n . By a result of Spaltenstein i_τ^* is always surjective [Sp], hence it will suffice to show that $\dim A_\sigma \leq \dim H^*(X^\tau)$. By [Ta], $\dim A_\sigma \leq \frac{n!}{\tau_1! \cdots \tau_m!}$ which is precisely $\dim H^*(X^\tau)$.

Acknowledgement: I would like to thank T.A. Springer for a remark that led to Proposition 1 and Roger Richardson for several interesting discussions about the results herein and for catching some errors in the original version. I also thank W. Borho and P. Slodowy for useful comments.

§2. Torus actions on varieties of fixed flags

The point of this section will be to establish the existence of a torus with finitely many fixed points on X^σ , for any regular nilpotent σ in a Levi subalgebra \mathfrak{l} , and to determine the combinatorial properties of the fixed point set. It is convenient, for this purpose, to consider X as the variety of a Borel subgroups of G with G acting on X by conjugation. For $M \subset G$, let X^M denote the set of all Borels containing M . Fix a Cartan subalgebra \mathfrak{h} in \mathfrak{g} with group H , and let $\mathfrak{l} \supset \mathfrak{h}$ be a Levi subalgebra with group L . Let W be the Weyl group of $(\mathfrak{g}, \mathfrak{h})$ and W_L the Weyl group of $(\mathfrak{l}, \mathfrak{h})$.

THEOREM 2: *Let u be a regular unipotent in L . Then the torus $S = Z(L)$ acts on X^u with exactly $[W : W_L]$ fixed points. In fact, if we fix a Borel B*

containing u and S , then there exists a one to one correspondence $\phi_u: X^u \cap X^S \rightarrow W_L \backslash W$ defined as follows: if $B' \in X^u \cap X^S$ and $w = w(B')$ is the unique element of W so that $wBw^{-1} = B'$, then $\phi_u(B') = \bar{w}$ (the right coset of w modulo W_L). The set $W_u = \{w \in W \mid wBw^{-1} \in X^u \cap X^S\}$ gives a complete set of representatives for $W_L \backslash W$.

PROOF: First, it is well known that X^S has $[W: W_L]$ components, each being a flag variety $L/L \cap B$. L being connected, u acts on each component of X^S and has exactly one fixed point by regularity. Hence $X^u \cap X^S$ has exactly $[W: W_L]$ points.

Next, Let H be the maximal torus corresponding to ℓ . Also suppose $B \supset H$. There exists a surjection $\phi_u: X^u \cap X^H \rightarrow W_L \backslash W$ defined exactly as above. To show surjectivity, it is enough to show that for any $w \in W$ there is a $\tilde{w} \in W_L$ so that $B' = (\tilde{w}w)B(\tilde{w}w)^{-1}$ contains u and H . It suffices to show $u \in B'$. Note that $wBw^{-1} \cap L$ is a Borel in L containing H , and hence there exists a $\tilde{w} \in W_L$ so that $\tilde{w}(wBw^{-1} \cap L)\tilde{w}^{-1} = B \cap L$. But $u \in B \cap L$ and $\tilde{w}(wBw^{-1} \cap L)\tilde{w}^{-1} = (\tilde{w}w)B(\tilde{w}w)^{-1} \cap L$, so $u \in B' = (\tilde{w}w)B(\tilde{w}w)^{-1}$. Hence ϕ_u is surjective. But clearly, $X^u \cap X^H \subset X^u \cap X^S$, so by $\#X^u \cap X^H \geq [W: W_L] = \#X^u \cap X^S$ we conclude $X^u \cap X^S = X^u \cap X^H$, and ϕ_u as defined above is a bijection, which completes the proof.

COROLLARY 2: Let $P \supset B$ be a parabolic with Levi subgroup L . Then

$$\dim H^*(X^u) = \dim H^0(X^S) = \dim H^*(G/P) = [W: W_L].$$

PROOF: It suffices to verify the first equality. Use the fact that for an algebraic action of a torus S on a projective variety V , the Euler characteristic $\chi(V)$ equals $\chi(V^S)$. In particular, $\chi(X^u) = [W: W_L]$. But, every X_u has vanishing odd rational homology, so $\chi(X^u) = \dim H^*(X^u)$.

§3. Torus actions and cohomology

In this section we will explain the main result of [ACL] which will be the important tool in the proof of Theorem 1. Let X be a smooth projective variety over \mathbb{C} of dimension n on which an algebraic torus S acts algebraically. The fixed point set $Z = X^S$ is automatically nontrivial, but not necessarily finite. Further, let Y denote a closed S -invariant subvariety of X so that $Y \cap Z$ is finite (it is automatically nonempty). For simplicity, assume also that $H^q(X; \Omega_X^p)$ vanishes if $p \neq q$, Ω_X^p being the sheaf on X of germs of holomorphic p -forms. As a consequence of this, $H^{\text{odd}}(X)$ vanishes and $H^{2p}(X) = H^p(X; \Omega_X^p)$ for all $p \geq 0$. Note that all cohomology has complex coefficients.

THEOREM 3: ([ACL]). *Associated to any regular element s of \mathfrak{o} , there exists a filtration of $H^*(Z)$*

$$H^*(Z) = F_n \supset F_{n-1} \supset \cdots \supset F_1 \supset F_0 \supset \cdots \quad (n = \dim X)$$

so that

- (a) $F_i F_j \subset F_{i+j}$
- (b) $F_i/F_{i-1} \cong H^{2i}(X)$, and
- (c) $H^{2^*}(X) \cong \text{Gr } H^*(Z) = \sum_{i \geq 0} F_i/F_{i-1}$ (as graded rings).

Let $j: Y \cap Z \rightarrow Z$ be the inclusion, $G_k = j^* F_k \subset H^0(Y \cap Z)$, and assume j^* is surjective. Then there exists a surjective homomorphism of graded rings

$$\psi_Y: \text{Gr } H^0(Y \cap Z) = \sum_{i \geq 0} G_i/G_{i-1} \rightarrow I(X, Y)$$

where $I(X, Y)$ denotes the graded ring $\sum_{k \geq 0} I(X, Y)_k$ with

$$I(X, Y)_k = \text{Im}[H^{2k}(X) \rightarrow H^{2k}(Y)].$$

Finally, if all odd Betti numbers of Y vanish, then ψ_Y is an isomorphism if and only if $I(X, Y) = H^*(Y)$.

To see how to calculate $H^*(X)$ in $\text{Gr } H^*(Z)$, chose a Leray cover \mathcal{U} of X and form the complex $K_X^* = \sum C^q(\mathcal{U}, \Omega_X^p)$ with differential $D: K_X^j \rightarrow K_X^{j+1}$ given on $C^q(\mathcal{U}, \Omega_X^p)$ by $\delta + (-1)^p i(V)$. By [CL₂], $\underline{H}_X^k = H^k(K_X^*, D)$ vanishes if $k \neq 0$, and \underline{H}_X^0 is a ring with filtration

$$\underline{H}_X^0 = \bar{F}_N \supset \bar{F}_{n-1} \supset \cdots \supset \bar{F}_0 \supset \cdots$$

so that $\bar{F}_i \bar{F}_j \subset \bar{F}_{i+j}$, and there exists a graded ring isomorphism

$$\text{Gr } \underline{H}_X^0 = \oplus \bar{F}_i/\bar{F}_{i-1} \cong \oplus H^i(X; \Omega_X^i).$$

The inclusion map $i_Z: Z \rightarrow X$ induces a quasi-isomorphism $i_Z^*: K_X^* \rightarrow K_Z^*$, hence $\underline{H}_X^0 \cong \underline{H}_Z^0 = \sum_{p \geq 0} H^p(Z; \Omega_Z^p)$. But by various vanishing theorems, in particular [CS], the right hand side is $H^*(Z)$. In particular, if $F_j = i_Z^*(\bar{F}_j)$, then $\text{Gr } \underline{H}_X^0 \cong \text{Gr } \underline{H}_Z^0$ which finishes the calculation of $H^*(X)$.

The next step is to use this calculation to compute $c_1(L)$ in $\text{Gr } H^*(Z)$, where L is any holomorphic line bundle on X . Let V be the holomorphic vector field on X determined by a regular element s of \mathfrak{o} . The fundamental fact is that since V has zeros, there exists a lifting of the derivation $V: \mathcal{O}_X \rightarrow \mathcal{O}_X$ on the sheaf of germs of holomorphic functions

O_X on X to a $\tilde{V}: O_X(L) \rightarrow O_X(L)$ such that if $f \in O_{X,x}$ and $\sigma \in O_X(L)_x$, then

$$\tilde{V}(f\sigma) = V(f)\sigma + f\tilde{V}(\sigma) \quad (\text{see [CL}_2]).$$

To calculate the class $\tilde{c}(L) \in \bar{F}_1 \subset \underline{H}_X^0$ which defines $c_1(L)$ in the associated graded, one chooses a local \bar{h} olomorphic connection D_α for $L|U_\alpha$ for any $U_\alpha \in \mathcal{U}$. Thus $D_\alpha: O_{U_\alpha}(L) \rightarrow O_{U_\alpha}(L) \otimes \Omega_{U_\alpha}^1$ satisfies $D_\alpha(f\sigma) = \sigma \otimes df + fD_\alpha(\sigma)$. Then a Cech cocycle representing $\tilde{c}(L)$ in K_X^0 is $\{\theta_{\alpha,\beta}\} + \{L_\alpha\}$, where $\theta_{\alpha,\beta} = D_\alpha - D_\beta$ and $L_\alpha = \tilde{V} - i(V)D_\alpha$. In K_Z^0 , $\tilde{c}(L)$ is represented by $\{i_Z^*\theta_{\alpha,\beta}\} + \tilde{V}|Z$, where $i_Z: Z \rightarrow X$ is the inclusion. In the associated graded, both classes give $c_1(L) \in H^2(X)$.

Suppose now that Y is an S -invariant subvariety of X so that $Y \cap Z$ is finite, and let $j: Y \cap Z \rightarrow Z$ be the inclusion. Note first that $\tilde{V}|Z$ is a holomorphic function on Z . For $\tilde{V}|Z \in H^0(Z, \text{Hom}(O_X(L), O_X(L))) = H^0(Z, O_Z)$. Consequently, since Z is compact, $\tilde{V}|Z \in H^0(Z)$. Thus $j^*(\{i_Z^*\theta_{\alpha,\beta}\} + \tilde{V}|Z) = j^*(\tilde{V}|Z) \in G_1 \subset H^0(Y \cap Z)$, and $j^*(V|Z)$ determines $i_Y^*c_1(L) \in H^2(Y)$ (modulo the kernel of the mapping $G_1/G_0 \rightarrow \text{Im } i_Y^* \subset H^2(Y)$).

We now return to the case where X is the flag variety of an arbitrary semio-simple group G . Let $s \in \mathfrak{h}$, and suppose σ is a regular nilpotent in $\mathfrak{e} = \mathfrak{z}_\mathfrak{g}(s)$. Denote by S the torus $Z(L)$. We will apply the above construction to the bundle L_χ on X associated to a character $\chi \in X(H)$. Let V be the vector field associated to s , and denote the lift of V to $O_X(L_\chi)$ by V_χ . Recall that in Theorem 2 we exhibited a bijection $\phi_\sigma^{*-1}: H^0(X^\sigma \cap X^S) \cong H^0(X^S) \rightarrow \mathbb{C}^{W_L \setminus W}$, the ring of complex functions on $W_L \setminus W$.

LEMMA: *Let \mathfrak{e}' be a Borel subalgebra of \mathfrak{g} in $X^\sigma \cap X^S$ and suppose $\psi_\sigma(\mathfrak{e}') = \bar{w}$. Then*

$$V_\chi(\mathfrak{e}') = -d\chi(w^{-1} \cdot s)$$

Consequently, $j_\sigma^*V_\chi(\mathfrak{e}') = -d\chi(w^{-1} \cdot s)$ where $j_\sigma: X^\sigma \cap X^S \rightarrow X^S$ is the inclusion.

The proof is identical with lemma 1 of [C], so we will omit it.

§4. Proof of Theorem 1

Using Theorem 2, we have an identification $\phi_\sigma^*: \mathbb{C}^{W_L \setminus W} \rightarrow H^0(X^\sigma \cap X^S)$, where the notation is as in §3. Recall (§1) that $A(W \cdot s)$ has a canonical filtration $D_0 \subset D_1 \subset \dots$. Let $G_0 \subset G_1 \subset \dots$ be the filtration of $H^0(X^\sigma \cap X^S)$ defined in Theorem 3. We will now define an isomorphism

$$\tilde{\psi}_s: A(W \cdot s) \rightarrow \mathbb{C}^{W_L \setminus W}$$

and show that $\tilde{\psi}_s(D_i) = \phi_\sigma^*{}^{-1}(G_i)$ for all i . The hypotheses of Theorem 3 hold, so we obtain the desired map ψ_s as the composition

$$\mathrm{Gr} A(W \cdot s) \rightarrow \mathrm{Gr} H^0(X^\sigma \cap X^S) \rightarrow \mathrm{Im} i_\sigma^* = I(X, X^\sigma)$$

To define ψ_s , it suffices to define $\psi_s(\omega)$ for $\omega \in \mathfrak{h}^*$. We set $\psi_s(\omega)(\bar{w}) = -\omega(w^{-1} \cdot s)$.

Now W acts on $\mathbb{C}^{W_L \setminus W}$ on the right: $(w \cdot f)(\bar{v}) = f(\overline{vw})$ if $v, w \in W$ and $f \in \mathbb{C}^{W_L \setminus W}$.

LEMMA: $\tilde{\psi}_s$ is a W -equivariant isomorphism.

PROOF: A straight forward calculation.

Since $A(W \cdot s)$ is generated by D_1 and $H^1(X)$ is generated by $H^2(X)$, the surjectivity assertion of Theorem 1 will be proved if we show $\tilde{\psi}_s(D_1) = \phi_\sigma^*{}^{-1}(G_1)$. But the Lemma of §3 says that if $\chi \in X(H)$, then $\tilde{\psi}_s(d\chi) = j_\sigma^*(\tilde{c}(L_\chi))$, where $d\chi \in \mathfrak{h}^*$ is the differential of χ , so ψ_s is surjective. To finish the proof, we must show W -equivariance and that ψ_s is an isomorphism if and only if i_σ^* is surjective. The proof of equivariance will be left to the reader. The second statement follows immediately from Theorem 3 due to the fact that $H^{\mathrm{odd}}(X^\sigma)$ vanishes for all σ [BS].

§5. Proof of Proposition 1

Let $A(\mathcal{J})$ denote the ring of polynomials on \mathcal{J} and Let $I = I(O_s) \subset A(\mathcal{J})$ be the ideal of O_s . Recall that $\mathrm{gr} I$ is the ideal generated by leading terms in I . By [BK, Satz. 1.8 and pp. 80–82], the hypotheses imply that $I(\overline{O}_\sigma) = \mathrm{gr} I$. Thus, $I(\overline{O}_\sigma \cap \mathfrak{h}) = I(\overline{O}_\sigma) + I(\mathfrak{h}) = \mathrm{gr} I + I(\mathfrak{h})$. Set theoretically, $O_s \cap \mathfrak{h} = W \cdot s$. However, $O_s \cap \mathfrak{h}$ is also smooth (see e.g. [Hu, pp. 116–117]), so $I(O_s \cap \mathfrak{h}) = I(W \cdot s)$. In particular,

$$I(W \cdot s) = I + I(\mathfrak{h}).$$

For an arbitrary pair of ideals I_1 and I_2 in $A(\mathcal{J})$, $\mathrm{gr} I_1 + \mathrm{gr} I_2 \subset \mathrm{gr}(I_1 + I_2)$. Thus, since $I(\mathfrak{h})$ is homogeneous and consequently $\mathrm{gr}(I(\mathfrak{h})) = I(\mathfrak{h})$,

$$\mathrm{gr}(I(W \cdot s)) \subset \mathrm{gr} I + I(\mathfrak{h}).$$

We immediately obtain a W -equivariant surjective algebra homomorphism

$$A(\overline{O}_\sigma \cap \mathfrak{h}) = A(\mathfrak{h})/\mathrm{gr} I + I(\mathfrak{h}) \rightarrow A(\mathcal{J})/\mathrm{gr}(I(W \cdot s)).$$

Furthermore, since $A(\mathcal{G})$ is graded,

$$A(\mathcal{G})/\text{gr}(I(W \cdot s)) \cong \text{Gr } A(W \cdot s). \quad (3)$$

Moreover, $W \cdot s \subset \mathcal{H}$ and W acts homogeneously on $A(\mathcal{H})$, so (3) is W -equivariant. Combining these maps yields the desired morphism.

References

- [ACL] E. AKYILDIZ, J.B. CARRELL and D.I. LIEBERMAN: Zeros of holomorphic vector fields on singular varieties and intersection rings of Schubert varieties. *Compositio Math.* 57 (1986) 237–248.
- [BK] W. BORHO and H. KRAFT: Über Bahnen und deren Deformation bei linearen Aktionen reductiver Gruppen. *Comment. Math. Helvetici* 54 (1979) 62–104.
- [BS] W.M. BEYNON and N. SPALTENSTEIN: Green functions of finite Chevalley groups of type E ($n = 6, 7, 8$), preprint.
- [C] J.B. CARRELL: Vector fields and the cohomology of G/B , Manifolds and Lie groups. Papers in honor of Y. Matsushima. *Progress in Mathematics*, Vol. 14 Birkhauser, Boston (1981).
- [C-L₁] J.B. CARRELL and D.I. LIEBERMAN: Holomorphic vector fields and compact Kaehler manifolds. *Invent. Math.* 21 (1973) 303–309.
- [C-L] J.B. CARRELL and D.I. LIEBERMAN: Vector fields and Chern numbers. *Math. Ann.* 225 (1977) 263–273.
- [CS] J.B. CARRELL and A.J. SOMMESE: Some topological aspects of \mathbb{C}^* -actions on compact Kaehler manifolds. *Comment. Math. Helv.* 54 (1979) 567–587.
- [DP] C. DECONCINI and C. PROCESI: Symmetric functions, conjugacy classes, and the flag variety. *Invent. Math.*, 64 (1981) 203–219.
- [Hu] J. HUMPHREYS: *Linear algebraic groups*. Springer Verlag, Berlin and New York (1975).
- [Ko] B. KOSTANT: Lie group representations on polynomial rings. *Amer. J. Math.* 85 (1963) 327–404.
- [Kr] H. KRAFT: Conjugacy classes and Weyl group representations; Tableaux de Young et foncteurs de Schur en algebre et geometrie (Conference international, Torun Pologne 1980). *Asterisque* 87–88 (1981) 195–205.
- [KP] H. KRAFT and C. PROCESI: Closures of conjugacy classes of matrices are normal. *Invent. Math.* 53 (1979) 227–247.
- [Sp] N. SPALTENSTEIN: The fixed point set of a unipotent transformation on the flag manifold. *Nederl. Akad. Wetensch. Proc. Ser. A.* 79 (1976) 452–456.
- [Spr] T.A. SPRINGER: A construction of representations of Weyl groups. *Invent. Math.* 44 (1978) 279–293.
- [Ta] T. TANISAKI: Defining ideals of the closures of the conjugacy classes and representations of the Weyl groups. *Tohoku Math. J.* 34 (1982) 575–585.

(Oblatum 10-IV-1985 & 9-VII-1985)

J.B. Carrell
 Mathematics Department
 The University of British Columbia
 Vancouver
 Canada V6T 1Y4