

COMPOSITIO MATHEMATICA

MARTIN LÜBKE

CHRISTIAN OKONEK

**Differentiable structures of elliptic surfaces
with cyclic fundamental group**

Compositio Mathematica, tome 63, n° 2 (1987), p. 217-222

http://www.numdam.org/item?id=CM_1987__63_2_217_0

© Foundation Compositio Mathematica, 1987, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Differentiable structures of elliptic surfaces with cyclic fundamental group

MARTIN LÜBKE¹ & CHRISTIAN OKONEK²

¹*Mathematisch Instituut, Niels Bohrweg 1, Postbus 9512, NL-2300 RA Leiden, The Netherlands;* ²*Mathematisches Institut, Universität Göttingen, Bunsenstrasse 3–5, D-3400 Göttingen, Federal Republic of Germany*

Received 6 February 1987; accepted 2 March 1987

1. Introduction

Recently two very different results on differentiable structures of elliptic surfaces have been proved. On one hand, there is the following theorem of Ue [U]:

THEOREM *Let X, X' be relatively minimal elliptic surfaces over smooth curves S, S' such that the Euler number $e(X)$ is positive and $\pi_1(X)$ is not cyclic. Then X and X' are oriented diffeomorphic if and only if $e(X) = e(X')$ and $\pi_1(X) \cong \pi_1(X')$.*

In particular, the diffeomorphism type of such a surface is already determined by its homeomorphism type.

The elliptic surfaces not covered by Ue's result are elliptic surfaces over \mathbb{P}^1 with at most two multiple fibres F_p, F_q of multiplicities p and q ; their fundamental group is isomorphic to \mathbb{Z}/k where $k = \text{g.c.d.}(p, q)$ ([Dv], [U]). If $k = 1$ and $p_g = 0$ these are the so-called Dolgachev surfaces $X_{p,q}$, which are all homeomorphic to \mathbb{P}^2 blown up in nine points [F]. For these surfaces Friedman and Morgan [FM] resp. Okonek and Van de Ven [OV] proved the following theorem which is in sharp contrast to Ue's result.

THEOREM *The Dolgachev surfaces $X_{2,q}$ with $q \equiv 1 \pmod{2}$ are pairwise differentially inequivalent.*

In [FM] it is furthermore proved that the mapping $(p, q) \mapsto X_{p,q}$ which associates to a pair (p, q) of relatively prime integers the diffeomorphism type of a Dolgachev surface $X_{p,q}$ is finite-to-one.

In this paper we tackle the case $p_g = 0$ and $\text{g.c.d.}(p, q) = k \geq 1$. We will show, that for every fixed k there are infinitely many differentially inequivalent surfaces $X_{p,q}$. For odd k all $X_{p,q}$ are homeomorphic by a result of Hambleton and Kreck [HK], whereas for even k the topological classification is still incomplete [HK];* the case $k = 2$ has been treated in [O].

The main tool in the proof of this result is Donaldson's invariant introduced in [D1], [D2]. Most of the techniques which we will use have already been developed in [OV] and [LO], so we refer to these papers for some details.

2. Precise statement of results

To describe the surfaces we are dealing with, let Y_0, Y_1, Y_2 resp. x_0, x_1 be homogeneous coordinates in \mathbb{P}^2 resp. \mathbb{P}^1 and Q_0, Q_1 two irreducible homogeneous cubic polynomials in Y_0, Y_1, Y_2 . Let $X \subset \mathbb{P}^1 \times \mathbb{P}^2$ be the zero-set of the polynomial $x_0 Q_0 + x_1 Q_1$. For generic Q_0, Q_1 the surface X is smooth. The induced projection to \mathbb{P}^1 defines an elliptic fibration with irreducible fibres and without multiple fibres. Applying logarithmic transformations of multiplicities p and q along two smooth fibres, we obtain the surface $X_{p,q}$ with multiple fibres F_p and F_q . The fundamental group of this surface is $\pi_1(X_{p,q}) \cong \mathbb{Z}/k$ where $k = \text{g.c.d.}(p, q)$.

If we write $p = kp', q = kq'$, then there are uniquely determined integers β, γ with $\gamma p' + \beta q' = 1$ and $0 \leq \gamma \leq q' - 1$; we define two divisors as follows:

$$\begin{aligned} C_{p,q} &= \beta F_p + \gamma F_q \\ T_{p,q} &= p' F_p - q' F_q. \end{aligned}$$

Clearly $T_{p,q}$ is a torsion element in H^2 is a torsion element in $H^2(X_{p,q}, \mathbb{Z})$ because

$$kT_{p,q} = pF_p - qF_q \sim 0$$

(here \sim denotes linear equivalence). Every vertical divisor D on $X_{p,q}$ can be written in the form

$$D \sim aF + bF_p + cF_q$$

where $a, b, c \in \mathbb{Z}$ and F is a generic fibre. We define

$$N(D) = ap'q'k + bq' + cp'.$$

* See: Note added in proof.

In particular, since the canonical divisor of $X_{p,q}$ is

$$K_{p,q} \sim -F + (p - 1)F_p + (q - 1)F_q \sim F - F_p - F_q,$$

we have $N(K_{p,q}) = p'q'k - p' - q'$. An easy calculation shows:

LEMMA For every vertical divisor $D \sim aF + bF_p + cF_q$ we have

$$D \sim N(D)C_{p,q} + (b\gamma - c\beta)T_{p,q}.$$

In other words, the group of vertical divisors modulo torsion is isomorphic to \mathbb{Z} with generator $C_{p,q}$.

If L is any ample divisor on $X_{p,q}$, then

$$N(D) = \text{deg}_L(D)/\text{deg}_L(C_{p,q}),$$

so every vertical divisor of degree 0 is torsion.

What we need to know about the topology of $X_{p,q}$ is the following. First of all, $c_1^2(X_{p,q}) = 0$ since $K_{p,q}$ is vertical. Then, since the geometric genus and the topological Euler characteristic are invariant under logarithmic transformations, we have $p_g(X_{p,q}) = 0$, $e(X_{p,q}) = 12$. Therefore the signature of $X_{p,q}$ is $\sigma(X_{p,q}) = -8$. The intersection form

$$S_{X_{p,q}} : H^2(X_{p,q}, \mathbb{Z})/\text{torsion} \times H^2(X_{p,q}, \mathbb{Z})/\text{torsion} \rightarrow \mathbb{Z}$$

is even if and only if $k \equiv 0 \pmod 2$ and $p' + q' \equiv 0 \pmod 2$ [O]. Thus

$$S_{X_{p,q}} \triangleq \begin{cases} -E_8 \oplus H & \text{if } k \equiv 0 \pmod 2, p' + q' \equiv 0 \pmod 2, \\ \langle 1 \rangle \oplus 9\langle -1 \rangle & \text{otherwise.} \end{cases}$$

We will use the following result of Hambleton and Kreck [HK]:

THEOREM Let M be a closed, oriented, differentiable manifold of real dimension 4 with $\pi_1(M) \cong \mathbb{Z}/k$. If k is odd, then the oriented homeomorphism type of M is determined by the intersection form on $H^2(M, \mathbb{Z})/\text{torsion}$.

COROLLARY For fixed odd* k , all surfaces $X_{p,q}$ with $\text{g.c.d.}(p, q) = k$ are homeomorphic.

Since the surfaces $X_{p,q}$ are algebraic with $p_g(X_{p,q}) = 0$, we have $b_+(X_{p,q}) = 1$ and the Donaldson invariant Γ is defined for every $X_{p,q}$. For the definition

* See: Note added in proof.

and the properties of Γ see [D1], [D2], [FM], [OV]. Now we state our main result.

THEOREM: *For every pair of integers $p, q \geq 1$ there exists an ample divisor $L_{p,q}$ on $X_{p,q}$ and an integer $n_{p,q} \geq N(K_{p,q})$ such that*

$$\Gamma(L_{p,q}) \equiv n_{p,q} C_{p,q}$$

in $H^2(X_{p,q}, \mathbb{Z})/\text{torsion}$. If the surfaces $X_{p,q}$ and $X_{r,s}$ are diffeomorphic, then $n_{p,q} = n_{r,s}$.

COROLLARY *Given $p_0, q_0 \geq 1$ there exist only finitely many pairs p, q such $X_{p,q}$ is diffeomorphic to X_{p_0,q_0} .*

3. Sketch of proofs

Choose an ample divisor $L_{p,q}^0$ on $X_{p,q}$ and let $L_{p,q} = L_{p,q}^0 + nK_{p,q}, n \geq 0$. The main ingredient for calculating the Donaldson invariant is the moduli space of $L_{p,q}$ -stable 2-bundles E with Chern classes $c_1(E) = 0, c_2(E) = 1$ on $X_{p,q}$. We will denote this space by $M_{p,q}$. It can be determined by the same methods as in [LO] and [OV]: Each stable bundle E is given by an extension

$$0 \rightarrow \mathcal{O}(D - K_{p,q}) \rightarrow E \rightarrow \mathcal{I}_z \otimes \mathcal{O}(K_{p,q} - D) \rightarrow 0 \tag{*}$$

where $D = bF_p + cF_q$ is a vertical divisor with $0 \leq b \leq p - 1, 0 \leq c \leq q - 1$, and z is a simple point in $F_p \cup F_q$. Since there may be torsion in $H^2(X_{p,q}, \mathbb{Z})$, a vertical curve D is not necessarily determined by its degree, but still there is a *unique* pair (D, z) defining E by (*) if we require D to have maximal degree *and* maximal b . It is not hard to check that given a pair (D, z_0) maximal (in the above sense) for a stable bundle E_0 with $z_0 \in F_p$ (or F_q), then also for every other point $z \in F_p$ (or F_q) the bundle E defined by (*) is stable and D is maximal for E . Hence $M_{p,q}$ is as a set the disjoint union of a finite number of copies of F_p and F_q , but the analytic structure of $M_{p,q}$ is in general non-reduced. If $\mathbb{E}_{p,q}$ is the universal bundle over $M_{p,q} \times X_{p,q}$, which can be constructed as in [OV], [LO], then from [D2] we get

$$\Gamma(L_{p,q}) \equiv 2a_1[M_{p,q}^{\text{red}}]c_2(\mathbb{E}_{p,q}) + a_2K_{p,q}$$

in $H^2(X_{p,q}, \mathbb{Z})/\text{torsion}$, where a_1, a_2 are suitable positive integers (this is the only difference from the formula for Γ used in [FM], [OV]). The coefficient

a_1 comes from the multiplicities of $M_{p,q}$ ([D2], Prop. 3.13), and a_2 depends on the torsion group $H_1(X_{p,q}, \mathbb{Z})$ ([D2], Appendix). As in [OV] the first term on the right hand side of the equation above consists of a certain number of copies of F_p and F_q , so

$$\Gamma(L_{p,q}) \equiv a_2 K_{p,q} + C$$

where C is a vertical divisor with $N(C) \geq 0$, or

$$\Gamma(L_{p,q}) \equiv n_{p,q} C_{p,q}$$

with $n_{p,q} \geq N(K_{p,q})$.

Now the same arguments as in [OV] show that for large n the chamber in the positive cone in $H^2(X_{p,q}, \mathbb{R})$ containing $L_{p,q} = L_{p,q}^0 + nK_{p,q}$ is independent of n ; also if $f: X_{p,q} \rightarrow X_{r,s}$ is an orientation preserving diffeomorphism and $L_{r,s}$ is a suitable ample divisor on $X_{r,s}$, then $L_{p,q}$ and $f^*(L_{r,s})$ are up to sign in the same chamber. From the constance of Γ on the chambers and naturality we get

$$n_{p,q} C_{p,q} \equiv \Gamma(L_{p,q}) \equiv \pm \Gamma(f^*(L_{r,s})) \equiv \pm f^*(\Gamma(L_{r,s})) \equiv \pm n_{r,s} f^*(C_{r,s}).$$

Now let D be a divisor on $X_{p,q}$ representing the class $f^*(C_{r,s}) \in H^2(X_{p,q}, \mathbb{Z})$. Then D is vertical, and modulo torsion we get from our lemma

$$n_{p,q} C_{p,q} \equiv \pm n_{r,s} N(D) C_{p,q}$$

implying $n_{r,s} | n_{p,q}$. Since the same argument works also the other way, we conclude $n_{p,q} = n_{r,s}$; this proves the theorem.

Finally for a given $N \in \mathbb{N}$ there are only finitely many pairs (p, q) with $n_{p,q} \leq N$, thus the corollary follows immediately.

Acknowledgements

The second author was supported by the Heisenberg program of the DFG.

Note added in proof: Hambleton and Kreck have meanwhile extended their topological classification to smooth, oriented 4-manifolds with fundamental group $\pi_1 = \mathbb{Z}/k$, $k \equiv 0 \pmod 2$ (I. Hambleton, M. Kreck: Smooth structures on algebraic surfaces with cyclic fundamental group, preprint 1987). Their result implies in particular that the oriented homeomorphism type of the surfaces $X_{p,q}$ with $\text{g.c.d.}(p, q) = k \equiv 0 \pmod 2$ is also determined by their intersection form. From the corollary to our main theorem it follows that all these surfaces have infinitely many smooth structures too.

References

- [D1] S.K. Donaldson: *La topologie différentielle des surfaces complexes*. C.R. Acad. Sc. Paris t.301, Série 1 No. 6, (1985) 317–320.
- [D2] S.K. Donaldson: Irrationality and the h-cobordism conjecture. Preprint (1986).
- [Dv] I. Dolgachev: Algebraic surfaces with $q = p_g = 0$. In: *Algebraic surfaces, proceedings of 1977 C.I.M.E.*. Cortona, Liguori, Napoli (1981) 97–215.
- [F] M.H. Freedman: The topology of four dimensional manifolds. *J. Diff. Geom.* 17 (1982) 357–453.
- [FM] R. Friedman and J. Morgan: On the diffeomorphism type of certain elliptic surfaces. Preprint (1985).
- [HK] I. Hambleton and M. Kreck: On the classification of topological 4-manifolds with finite fundamental group. Preprint, Mainz (1986).
- [LO] M. Lübke and C. Okonek: Stable bundles on regular elliptic surfaces. To appear in *J. reine angew. Math.*
- [O] C. Okonek: Fake Enriques surfaces. Preprint, Göttingen (1987).
- [OV] C. Okonek and A. Van de Ven: Stable bundles and differentiable structures on certain elliptic surfaces. *Invent. Math.* 86 (1986) 357–370.
- [U] M. Ue: On the diffeomorphism types of elliptic surfaces with multiple fibres. *Invent. Math.* 84 (1986) 633–643.