

COMPOSITIO MATHEMATICA

V. B. MEHTA

A. RAMANATHAN

Schubert varieties in $G/B \times G/B$

Compositio Mathematica, tome 67, n° 3 (1988), p. 355-358

<http://www.numdam.org/item?id=CM_1988__67_3_355_0>

© Foundation Compositio Mathematica, 1988, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Schubert varieties in $G/B \times G/B$

V.B. MEHTA & A. RAMANATHAN

*School of Mathematics, Tata Institute of Fundamental Research, Homi Bhabha Road,
Colaba, Bombay 400 005, India*

Received 1 December 1987; accepted 11 February 1988

Introduction

Let G be a semi-simple, simply connected algebraic group defined over an algebraically closed field of characteristic $p > 0$. Let $T \subset G$ be a maximal torus, $B \supset T$ a Borel subgroup and $W = N(T)/T$ the Weyl group. G acts on the homogeneous space G/B and also on $G/B \times G/B$ by the diagonal action: for $g, x_1, x_2, \in G, g(x_1B, x_2B) = (gx_1B, gx_2B)$. By *Schubert Varieties* in $G/B \times G/B$ we mean the closures of the G -orbits in $G/B \times G/B$. It is known ([11, 12]) that these orbit closures are in 1–1 correspondence with the elements of W , the element $w \in W$ corresponding to the closure of the orbit of (eB, wB) , where $e \in G$ is the identity element. In particular, taking $w = e$, G/B gets imbedded diagonally in $G/B \times G/B$.

In this paper we prove that these Schubert Varieties are Frobenius-split in the sense of [4, Def. 2]. Our method is as follows: fix $w \in W$ with $l(w) = i$ and denote the Schubert variety in G/B corresponding to w by X_i . Then B acts on X_i on the left and one may form the associated fibre-space $\tilde{X}_i = G \times^B X_i$. The map $f: \tilde{X}_i \rightarrow G/B \times G/B$ given by $f(g, x) = (gB, gxB)$ is an isomorphism onto the G -orbit closure of (eB, wB) (cf. [11]). Hence we may work with \tilde{X}_i instead. Express w as a product of reflections associated to the simple roots, $w = s_{\alpha_1} s_{\alpha_2} \dots s_{\alpha_i}$ and $Z_i \rightarrow X_i$ be the corresponding Demazure desingularization of X_i (cf. [2, 3]) and let $\psi_i: Z_i \rightarrow X_i$ be the birational map. B acts on Z_i on the left and we may construct the associated fibre-space $\tilde{Z}_i = G \times^B Z_i$. The map ψ_i is B -equivariant and descends to a birational map $\tilde{\psi}_i: \tilde{Z}_i \rightarrow \tilde{X}_i$. Since X_i is normal [1, 5, 7, 10] and $\tilde{X}_i \rightarrow G/B$ is a fibre-space with fibre X_i it follows that \tilde{X}_i is also normal and that $\tilde{\psi}_i(\mathcal{O}_{\tilde{Z}_i}) = \mathcal{O}_{\tilde{X}_i}$. So to prove that \tilde{X}_i is Frobenius-split, it is sufficient to prove that \tilde{Z}_i is Frobenius-split. We calculate the canonical bundle $K_{\tilde{Z}_i}$ of \tilde{Z}_i (this has been done, without detail, in [11]). From this description of $K_{\tilde{Z}_i}$ it follows from [4 prop. 8] that \tilde{Z}_i is Frobenius-split. It also follows from [6, 8] that \tilde{X}_i is Cohen-Macaulay and has rational singularities. We first recall the basic

facts about the standard resolutions of Schubert Varieties in G/B and Frobenius-splitting from [4, 8] and then we prove the main result. Our result should prove useful in the study of the decomposition of the G -module $H^0(G/B, L) \times H^0(G/B, M)$, where L and M are line bundles on G/B , see [11].

Section I

Let G, B and W be as in the introduction, let $w \in W$ with $l(w) = i$ and denote by X_i the Schubert variety in G/B corresponding to w . Then according to [2, 3, 8] there exists a smooth projective variety Z_i , and a map $\psi_i: Z_i \rightarrow X_i$ with the following properties:

- (1) ψ_i is birational.
- (2) There exists i smooth subvarieties of codim 1 in Z_i , denoted by $Z_{i,1} \dots Z_{i,i}$ intersecting transversally. Further if we denote $\bigcup_{j=1}^i Z_{i,j}$ by ∂Z_i , then $\psi_i^{-1}(\partial X_i) = \partial Z_i$, where ∂X_i is the union of the codim 1 Schubert varieties in X_i .
- (3) Put $v = w s_{\alpha_i}$ and $X_{i-1} = \overline{BvB/B}$. Then there exists a map $f_i: Z_i \rightarrow Z_{i-1}$ such that f_i is a locally trivial \mathbb{P}^1 fibration with a section $\sigma_i: Z_{i-1} \rightarrow Z_i$. Further, $\partial Z_i = f_i^{-1}(\partial Z_{i-1}) \cup \sigma_i(Z_{i-1})$.
- (4) The canonical bundle K_{Z_i} is given by the formula $K_{Z_i} = \mathcal{O}_{Z_i}(-\partial Z_i) \times \psi_i^* L_e^{-1}$ is the line bundle on $X_i \subset G/B$ associated to half sum of the positive roots.

The varieties Z_i and the morphisms ψ_i are constructed by induction on $l(w)$, see [3, 8] for more details. We recall one proposition from [8].

PROPOSITION 1. *Z_i is Frobenius-split and any sub-intersection of the divisors in ∂Z_i is compatibly split in Z_i .*

Proof. This is [8, Remark 2.5].

Now consider the varieties $\tilde{Z}_i = G \times^B Z_i$ as in the introduction. The maps $f_i: Z_i \rightarrow Z_{i-1}$ and $\sigma_i: Z_{i-1} \rightarrow Z_i$ are B -equivariant, hence we get maps $\tilde{f}_i: \tilde{Z}_i \rightarrow \tilde{Z}_{i-1}$ and $\tilde{\sigma}_i: \tilde{Z}_{i-1} \rightarrow \tilde{Z}_i$. It follows that there exist i smooth subvarieties of \tilde{Z}_i denoted by $\tilde{Z}_{i,1} \dots \tilde{Z}_{i,i}$, intersecting transversally, whose union we denote by $\partial \tilde{Z}_i$. Let p_1 and p_2 denote the two projections of $G/B \times G/B$ and for any pair of line bundles L, M on G/B , denote $(p_1^* L \times p_2^* M)$ by (L, M) .

PROPOSITION 2. *The canonical bundle $K_{\tilde{Z}_i}$ is given by*

$$K_{\tilde{Z}_i} = \mathcal{O}_{\tilde{Z}_i}(-\partial \tilde{Z}_i) \times \tilde{\psi}_i^*(L_e^{-1}, L_e^{-1}).$$

Proof. (See also [11]). We prove this by induction on $l(w)$. If $l(w) = 0$ then $\tilde{Z}_0 = G/B$ and $\partial\tilde{Z}_0 = \emptyset$. So $\mathcal{O}_{\tilde{Z}_0}(-\partial\tilde{Z}_0) \times \tilde{\psi}_0^*(L_\varrho^{-1}, L_\varrho^{-1})$ is the line-bundle L_ϱ^{-2} on G/B , as $\tilde{\psi}_0: G/B \rightarrow G/B \times G/B$ is the diagonal imbedding. Assume the result for $l(w) = i - 1$. Now it follows from [8, Lemma 3] that $K_{\tilde{Z}_i/\tilde{Z}_{i-1}} = \mathcal{O}_{\tilde{Z}_i}(-\sigma_i(\tilde{Z}_{i-1})) \times \tilde{\psi}_i^*(1, L_\varrho^{-1}) \times \tilde{f}_i^* \tilde{\sigma}_i^* \tilde{\psi}_i^*(1, L_\varrho)$. Denote this line bundle on \tilde{Z}_i by A . Then $K_{\tilde{Z}_i} = A \times \tilde{f}_i^*(K_{\tilde{Z}_{i-1}}) = A \times \tilde{f}_i^*[\mathcal{O}_{\tilde{Z}_{i-1}}(-\partial\tilde{Z}_{i-1}) \times \tilde{\psi}_{i-1}^*(L_\varrho^{-1}, L_\varrho^{-1})] = \mathcal{O}_{\tilde{Z}_i}(-\partial\tilde{Z}_i) \times \tilde{\psi}_i^*(1, L_\varrho^{-1}) \times \tilde{f}_i^* \tilde{\sigma}_i^* \tilde{\psi}_i^*(1, L_\varrho) \times \tilde{f}_i^* \tilde{\psi}_{i-1}^*(L_\varrho^{-1}, L_\varrho^{-1})$. But $\tilde{\psi}_{i-1}: \tilde{Z}_{i-1} \rightarrow G/B \times G/B = \tilde{\psi}_i \tilde{\sigma}_i: \tilde{Z}_{i-1} \rightarrow G/B \times G/B$. Hence we get $K_{\tilde{Z}_i} = \mathcal{O}_{\tilde{Z}_i}(-\partial\tilde{Z}_i) \times \tilde{\psi}_i^*(1, L_\varrho^{-1}) \times \tilde{f}_i^* \tilde{\psi}_{i-1}^*(L_\varrho^{-1}, 1)$. But $\tilde{f}_i^* \tilde{\psi}_{i-1}^*(L_\varrho^{-1}, 1) = \tilde{\psi}_i^*(L_\varrho^{-1}, 1)$ as both of them are isomorphic to $q^*(L_\varrho^{-1}, 1)$ where q is the projection $\tilde{Z}_i \rightarrow G/B$. Hence we get $K_{\tilde{Z}_i} = \mathcal{O}_{\tilde{Z}_i}(-\partial\tilde{Z}_i) \times \tilde{\psi}_i^*(L_\varrho^{-1}, L_\varrho^{-1})$.

THEOREM 1. \tilde{Z}_i is Frobenius-split and any sub-intersection of the divisors in $\partial\tilde{Z}_i$ is compatibly split in \tilde{Z}_i .

Proof. From Prop. 2, we know that $K_{\tilde{Z}_i}^{-1} = \mathcal{O}_{\tilde{Z}_i}(\partial\tilde{Z}_i) \times \tilde{\psi}_i^*(L_\varrho, L_\varrho)$. From [8, Remark 2], we know that $\sigma = D + \tilde{D}$ is an element of $H^0(G/B, L_\varrho^2)$ such that σ^{p-1} splits G/B . Consider the section $t = \partial\tilde{Z}_i + \tilde{\psi}_i^*(D, \tilde{D})$ of $K_{\tilde{Z}_i}$. It follows from [4, Prop. 8] that t^{p-1} splits \tilde{Z}_i and that any sub-intersection of the divisors in $\partial\tilde{Z}_i$ is compatibly split in \tilde{Z}_i by t^{p-1} .

COROLLARY 1. Let N be the length of the maximal element $w_0 \in W$. Then by the above, \tilde{Z}_0 is compatibly-split in \tilde{Z}_N . So it follows from [4, Prop. 4] that $\tilde{\psi}_0(\tilde{Z}_0) = G/B$ is compatibly-split in $\tilde{\psi}_N(\tilde{Z}_N) = G/B \times G/B$, where G/B is imbedded diagonally in $G/B \times G/B$.

This was first proved by the second author by other methods (cf. [9]).

COROLLARY 2. From Corollary 1 and [9, Cor. 2.3] it follows that any imbedding of G/B by a complete linear system is projectively normal.

This was first proved in [7] (see also [9]).

COROLLARY 3. It follows from [6, 8] that the Schubert varieties \tilde{X}_i in $G/B \times G/B$ are Cohen–Macaulay and have rational singularities.

Remark. It can be proved, using the methods of [9], that these Schubert varieties in $G/B \times G/B$ are scheme-theoretically defined by quadrics. This will be taken up in a later paper. Analogues follow for $G/P_1 \times G/P_2$.

References

1. H.H. Andersen, Schubert Varieties and Demazure’s character formula. *Invent. Math.* 79 (1985) 611–618.

2. Demazure, M., Desingularization des Varietes de Schubert generalisees. *Ann. Sc. Ec. Norm. Super.* 7 (1974) 53–88.
3. G. Kempf, Linear systems on homogeneous spaces. *Ann. Math.* 103 (1976) 557–591.
4. V.B. Mehta and A. Ramanathan, Frobenius splitting and cohomology vanishing for Schubert varieties. *Ann. Math.* 122 (1985) 27–40.
5. V.B. Mehta and V. Srinivas, Normality of Schubert varieties. *Amer. Jour. Math.* 109 (1987) 987–989.
6. V.B. Mehta and V. Srinivas, A note on Schubert varieties in G/B . Preprint.
7. S. Ramanan and A. Ramanathan, Projective normality of flag varieties and Schubert varieties. *Invent. Math.* 79 (1985) 217–224.
8. A. Ramanathan, Schubert Varieties are arithmetically Cohen–Macaulay. *Invent. Math.* 80 (1985) 283–294.
9. A. Ramanathan, Equations defining Schubert Varieties and Frobenius-splitting of the diagonal. *Publ. I.H.E.S.* 65 (1987) 61–90.
10. C.S. Seshadri, Line bundles on Schubert varieties Vector bundles on Algebraic Varieties, *Proceedings of the Bombay Colloquium* 1984.
11. Kumar, Shrawan, Proof of the Parthasarathy-Ranga Rao-Varadarajan conjecture. Preprint.
12. T. Springer, Quelques applications de la cohomologie d’intersection. *Seminaire Bourbaki Expose* 589, Vol. 1981/1982.