

COMPOSITIO MATHEMATICA

G. A. CAMERA

J. GIMENEZ

**The nonlinear superposition operator acting
on Bergman spaces**

Compositio Mathematica, tome 93, n° 1 (1994), p. 23-35

http://www.numdam.org/item?id=CM_1994__93_1_23_0

© Foundation Compositio Mathematica, 1994, tous droits réservés.

L'accès aux archives de la revue « Compositio Mathematica » (<http://www.compositio.nl/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

The nonlinear superposition operator acting on Bergman spaces

G. A. CÁMERA & J. GIMÉNEZ

Department of Mathematics, I.V.I.C., Caracas, Venezuela

Received 30 August 1992; accepted in revised form 20 June 1993

1. Introduction

Let Δ denote the unit disc $\{z: |z| < 1\}$ in the complex plane and let $H(\Delta)$ denote the space of analytic functions in Δ with the topology of the uniform convergence in compact subsets of Δ . Given a function $f: \mathbb{C} \rightarrow \mathbb{C}$ we associate to it the operator F_f defined by

$$F_f(u)(z) = f(u(z)), \quad u \in H(\Delta).$$

This operator is known as the autonomous nonlinear superposition (or composition) operator [1]. If A and B are linear subspaces of $H(\Delta)$ and $F_f(u) \in B$ whenever $u \in A$ we shall say that F_f acts from A to B . It is easy to see that if F_f acts from $H(\Delta)$ to $H(\Delta)$, then f must be an entire function and conversely. In this case mere action implies the continuity and the boundedness of the operator [2]. That mere action implies continuity has already been proved for various spaces of real functions, for instance L^p spaces [6] and Sobolev spaces [7]. Necessary and sufficient conditions have been given in [2] in order that F_f acts from H^p to H^q , $0 < p, q \leq +\infty$, where H^p denotes the classical Hardy space in the unit disc. It is also true in this case that mere action implies continuity [2]. If N denotes the Nevanlinna space of functions in $H(\Delta)$ of bounded characteristic then the actions from $\bigcup_{p < q} H^q$ to N and from N to N have been studied in [3].

In this note we shall consider the problem of action and continuity between the Bergman space B_p defined by

$$B_p = \{u \in H(\Delta) : u \in L^p(dx dy)\}, \quad 0 < p < \infty.$$

The space B_∞ is the usual one of bounded analytic functions in Δ . The topology in these spaces is given by the metric induced (when $p \geq 1$) by

$$\|u\|_{B_p} = \left(\frac{1}{\pi} \iint_{\Delta} |u(z)|^p dx dy \right)^{1/p}.$$

If $p < 1$ the topology induced by the metric $\|u\|_{B_p}^p$ is used. We also consider the action between B_p and the Hardy space H^q and vice versa. For functions in H^p we use the standard notation

$$M_p(r, u) = \left(\frac{1}{2\pi} \int_0^{2\pi} |u(re^{i\theta})|^p d\theta \right)^{1/p} \quad \text{and} \quad \|u\|_p = \lim_{r \rightarrow 1} M_p(r, u).$$

The symbol BN (which stands for Bergman-Nevanlinna) shall denote the set of functions u in $H(\Delta)$ such that

$$\iint_{\Delta} \log^+ |u(z)| dx dy < \infty.$$

Clearly $H^p \subset B_p$ and $B_p \subset BN$ for all p . Finally, we study the action between Hardy functions and Bergman-Nevanlinna functions.

We would like to thank the referee for his helpful comments.

2. The action in B_p

We shall need the following lemma.

LEMMA 1. *Let $0 < p < \infty$. If $u \in B_p$ then*

$$|u(z)| \leq \frac{\|u\|_{B_p}}{(1 - |z|)^{2/p}}, \quad z \in \Delta.$$

Proof. This is an easy consequence of the subharmonicity of $|u|^p$.

Next we are ready to prove the following result. In what follows the symbol “[s]” denotes the integer part of s .

THEOREM 1. *Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be an entire function. Then F_f acts from B_p to B_q , $0 < p, q \leq \infty$ if and only if f is a polynomial of degree less than or equal to $\left[\frac{p}{q} \right]$.*

Proof. If f is a polynomial of degree $n \leq \left[\frac{p}{q} \right]$ then $f \circ u \in B_q$, $\forall u \in B_p$. In fact, it is enough to see that if $k \leq \left[\frac{p}{q} \right]$, $k \in \mathbb{N}$, then $u^k \in B_q$. This is true since

$$\|u^k\|_{B_q} = \left(\frac{1}{\pi} \iint_{\Delta} |u(z)|^{kq} dx dy \right)^{1/q} = \left(\left(\frac{1}{\pi} \iint_{\Delta} (|u(z)|)^{kq} dx dy \right)^{1/kq} \right)^k$$

$$\leq \left(\frac{1}{\pi} \iint_{\Delta} (|u(z)|)^p dx dy \right)^{k/p} = \|u\|_{B_p}^k.$$

Next, we assume that F_f acts from B_p to B_q , $0 < p, q < \infty$. Let $\varepsilon > 0$ and define $u_\varepsilon^{(1)}(z) = \left(\frac{1}{1-z} - \frac{1}{2} \right)^{2/p+\varepsilon}$. Clearly $u_\varepsilon^{(1)}$ belongs to B_p . Therefore $f \circ u_\varepsilon^{(1)} \in B_q$ and by Lemma 1 one can write

$$|f(u_\varepsilon^{(1)}(z))| \leq \frac{\|f \circ u_\varepsilon^{(1)}\|_{B_q}}{(1-|z|)^{2/q}}, \quad z \in \Delta. \tag{2.1}$$

Set $w_1 = u_\varepsilon^{(1)}(z)$. Assume first that $p < 1$ and take $\varepsilon < 1 - p$. Then the range set of $u_\varepsilon^{(1)}$ is $\mathbb{C} \setminus 0$. Given $w_1 \in \mathbb{C} \setminus 0$ let $z \in \Delta$ such that $w_1 = u_\varepsilon^{(1)}(z)$ and

$$|z| = \frac{|w_1^{p+\varepsilon/2} - \frac{1}{2}|}{|w_1^{p+\varepsilon/2} + \frac{1}{2}|}.$$

Thus, from (2.1) we get

$$\begin{aligned} |f(w_1)| &\leq \frac{\|f \circ u_\varepsilon^{(1)}\|_{B_q}}{\left(1 - \frac{|w_1^{p+\varepsilon/2} - \frac{1}{2}|}{|w_1^{p+\varepsilon/2} + \frac{1}{2}|}\right)^{2/q}} \\ &= \frac{\|f \circ u_\varepsilon^{(1)}\|_{B_q} |w_1^{p+\varepsilon/2} + \frac{1}{2}|^{2/q}}{(|w_1^{p+\varepsilon/2} + \frac{1}{2}| - |w_1^{p+\varepsilon/2} - \frac{1}{2}|)^{2/q}}. \end{aligned}$$

If w_1 is such that $|w_1| > e^{1/2(p+\varepsilon)}$ then $z_1 = w_1^{p+\varepsilon/2} (-\pi \leq \text{Arg } w_1 < \pi)$ satisfies $\text{Re } z_1 > e^{1/4} \cos(-\pi/2(p+\varepsilon)) > 0$. In fact, since $\text{Arg } w_1 \geq -\pi$ then $((p+\varepsilon)/2) \text{Arg } w_1 \geq -\pi/2(p+\varepsilon)$. On the other hand

$$\text{Re } z_1 = |w_1|^{p+\varepsilon/2} \cos\left(\frac{p+\varepsilon}{2} \text{Arg } w_1\right) > |w_1|^{p+\varepsilon/2} \cos\left(-\frac{\pi}{2}(p+\varepsilon)\right).$$

Hence, if $|w_1| > e^{1/2(p+\varepsilon)}$ then $\text{Re } z_1 > e^{1/4} \cos(\pi/2(p+\varepsilon)) > 0$. Therefore, one can find a positive constant c (depending on ε and p) such that

$$|w_1^{p+\varepsilon/2} + \frac{1}{2}| - |w_1^{p+\varepsilon/2} - \frac{1}{2}| > c, \quad |w_1| > e^{1/2(p+\varepsilon)}.$$

If we use this inequality in (2.2) we obtain

$$|f(w_1)| \leq C(p, q, \varepsilon) (|w_1|^{p+\varepsilon/q} + 2^{-2/q})$$

for all w_1 such that $|w_1| > e^{1/2(p+\varepsilon)}$. Thus f is a polynomial of degree less than or equal to $p + \varepsilon/q$. By letting $\varepsilon \rightarrow 0$ we obtain the desired result when $p < 1$.

Now, we assume that $p \geq 1$. Let

$$S_1 = \left\{ w_1 : \frac{-\pi}{2(p+\varepsilon)} \leq \text{Arg } w_1 < \frac{\pi}{2(p+\varepsilon)}, |w_1| > 1 \right\}.$$

If $w_1 \in S_1$ we choose z_1 such that $w_1 = z_1^{2/p+\varepsilon}$ with $-\pi/4 < \text{Arg } z_1 < \pi/4$. Thus $(|w_1|^{p+\varepsilon/2} + \frac{1}{2}) - |w_1|^{p+\varepsilon/2} - \frac{1}{2})^{2/q} \geq c > 0$, for some constant c . Combining this inequality with (2.2) we obtain

$$\begin{aligned} |f(w_1)| &\leq \frac{\|f \circ u_\varepsilon^{(1)}\|_{B_q}}{c} (|w_1|^{p+\varepsilon/2} + \frac{1}{2})^{2/q} \\ &\leq c(\varepsilon, q) \left(|w_1|^{p+\varepsilon/q} + \frac{1}{2^{2/q}} \right). \end{aligned} \tag{2.3}$$

Let

$$w_2 \in S_2 = \left\{ w_2 : \frac{\pi}{2(p+\varepsilon)} \leq \text{Arg } w_2 < \frac{3\pi}{2(p+\varepsilon)}, |w_2| > 1 \right\}$$

and

$$u_\varepsilon^{(2)}(z) = \left(\frac{1}{1-z} - \frac{1}{2} \right)^{2/p+\varepsilon} e^{i\pi/p+\varepsilon}.$$

Clearly $u_\varepsilon^{(2)} \in B_p$. By hypothesis $f \circ u_\varepsilon^{(2)} \in B_q$ and by Lemma 1

$$|f(u_\varepsilon^{(2)})(z)| \leq \frac{\|f \circ u_\varepsilon^{(2)}\|_{B_q}}{(1-|z|)^{2/q}}, \quad z \in \Delta. \tag{2.4}$$

Given $w_2 \in S_2$ we choose z_2 such that $|z_2| > 1$ and $w_2 = z_2^{2/p+\varepsilon} e^{i\pi/p+\varepsilon}$ with $-\pi/4 \leq \text{Arg } z_2 \leq \pi/4$. From (2.4) we obtain

$$\begin{aligned} |f(w_2)| &\leq \frac{\|f \circ u_\varepsilon^{(2)}\|_{B_q}}{\left(1 - \frac{|w_2|^{p+\varepsilon/2} e^{i\pi/2} - \frac{1}{2}}{|w_2|^{p+\varepsilon/2} e^{i\pi/2} + \frac{1}{2}} \right)^{2/q}} \\ &= \frac{\|f \circ u_\varepsilon^{(2)}\|_{B_q} |w_2|^{p+\varepsilon/2} e^{i\pi/2} + \frac{1}{2}|^{2/q}}{\left(|w_2|^{p+\varepsilon/2} e^{i\pi/2} + \frac{1}{2} - |w_2|^{p+\varepsilon/2} e^{i\pi/2} - \frac{1}{2} \right)^{2/q}}. \end{aligned} \tag{2.5}$$

Since $-\pi/4 \leq \text{Arg } z_2 \leq \pi/4$ then

$$|w_2^{p+\varepsilon/2} e^{-i\pi/2} + \frac{1}{2}| - |w_2^{p+\varepsilon/2} e^{-i\pi/2} - \frac{1}{2}| \geq c^{q/2} > 0.$$

Hence, from (2.5) we obtain

$$|f(w_2)| \leq c^{-1} \|f \circ u_\varepsilon^{(2)}\|_{B_q} |w_2^{p+\varepsilon/2} e^{-i\pi/2} + \frac{1}{2}|^{2/q}.$$

By repeating the same argument n times, where n is such that $\bar{\Delta} = \mathbb{C} \setminus \bigcup_{i=1}^n S_i$, we obtain

$$|f(w)| \leq c_1 (|w|^{p+\varepsilon/2} + \frac{1}{2})^{2/q}, \quad w \in \mathbb{C} \setminus \bar{\Delta},$$

where c_1 depends on f , ε and q . This proves that f is an entire function of order at most $p + \varepsilon/q$. Letting ε tend to zero gives the desired result for $p \geq 1$.

3. Continuity of F_f

We shall prove in this section that if F_f acts from B_p to B_q then it is necessarily continuous. We also prove local Lipschitzness.

First of all let us prove the following lemma.

LEMMA 2. If $u_k \rightarrow u$ in B_p , $n \in \mathbb{N}$ and $n \leq \left\lfloor \frac{p}{q} \right\rfloor$, then $u_k^n \rightarrow u^n$ in B_q .

Proof. The proof is similar to the analogue lemma given in [2]. We give it here somewhat simplified. The case $n = 1$ is obvious. Let us assume that $n > 1$. The functions u_k^n and u^n belong to B_q . In fact, since $nq \leq p$ then for every $u \in B_p$ we have

$$\|u^n\|_{B_q} \leq \|u\|_{B_p}^n.$$

On the other hand,

$$\begin{aligned} \|u_k^n - u^n\|_{B_q} &= \left(\frac{1}{\pi} \iint_{\Delta} |u_k^n - u^n|^q dx dy \right)^{1/q} \\ &= \left(\left(\frac{1}{\pi} \iint_{\Delta} (|u_k^n - u^n|^{1/n})^{qn} dx dy \right)^{1/nq} \right)^n \\ &\leq \left(\frac{1}{\pi} \iint_{\Delta} |u_k^n - u^n|^{p/n} dx dy \right)^{n/p} \\ &= \left(\frac{1}{\pi} \iint_{\Delta} |u_k - u|^{p/n} |u_k^{n-1} + \dots + u^{n-1}|^{p/n} dx dy \right)^{n/p}. \end{aligned}$$

Now we use Hölder inequality

$$\left| \int fg \right| \leq \left(\int f^r \right)^{1/r} \left(\int g^s \right)^{1/s},$$

with $f = |u_k - u|^{p/n}$, $g = |u_k^{n-1} + \dots + u^{n-1}|^{p/n}$, $r = n > 1$ and

$$\frac{1}{n} + \frac{1}{s} = 1 \left(s = \frac{n}{n-1} \right),$$

and obtain

$$\begin{aligned} \|u_k^n - u^n\|_{B_q} &\leq \left(\frac{1}{\pi} \iint_{\Delta} |u_k - u|^p dx dy \right)^{1/p} \\ &\quad \left(\frac{1}{\pi} \iint_{\Delta} |u_k^{n-1} + \dots + u^{n-1}|^{p/n-1} dx dy \right)^{n-1/p} \\ &\leq c \|u_k - u\|_{B_p} \sum_{l=0}^{n-1} \|u_k^{n-1-l} u^l\|_{B_{p/n-1}}, \end{aligned} \quad (3.1)$$

where c is a constant. Again by a refined version of Hölder inequality we get that $u_k^{n-1-l} u^l \in B_{p/n-1}$, $l = 0, 1, \dots, n-1$ and

$$\|u_k^{n-1-l} u^l\|_{B_{p/n-1}} \leq \|u_k^{n-1-l}\|_{B_{p/n-1-l}} \|u^l\|_{B_{p/l}} = u_k \|u\|_{B_p}^{n-1-l} \|u\|_{B_p}^l \quad (3.2)$$

This inequality implies that all summands on the right-hand side of (3.1) are bounded (for all k). Hence $u_k^n \rightarrow u^n$ in B_q as required.

THEOREM 2. *If F_f acts from B_p to B_q then it is necessarily continuous, bounded and locally Lipschitz.*

Proof. Since F_f acts from B_p to B_q then, by Theorem 1, f is a polynomial of degree $n \leq \left\lfloor \frac{p}{q} \right\rfloor$. Set $f(z) = a_n z^n + \dots + a_0$. Let $u_k(z) \rightarrow u(z)$, as $k \rightarrow \infty$, in B_p .

Then

$$F_f(u_k)(z) - F_f(u)(z) = a_n(u_k^n(z) - u^n(z)) + \dots + a_1(u_k(z) - u(z)).$$

Thus

$$\|F_f(u_k) - F_f(u)\|_{B_q} \leq C(\|u_k^n - u^n\|_{B_q} + \dots + \|u_k - u\|_{B_q}) \rightarrow 0 \quad \text{as } k \rightarrow \infty$$

by Lemma 2. The boundedness of F_f comes from the inequality

$$\|F_f(u)\|_{B_q} \leq C(|a_n| \|u\|_{B_p}^n + \dots + |a_1| \|u\|_{B_p} + |a_0|), \quad u \in B_p$$

which can be deduced from $\|u^n\|_{B_q} \leq \|u\|_{B_p}^n$, for all $u \in B_p$ and all $n \leq \left\lfloor \frac{p}{q} \right\rfloor$.

In order to prove that F_f is locally lipschitz we must see that if $u, v \in B(0, R) \subset B_p$ then there exists a constant $C = C(p, q, R, f)$ such that $\|F_f(u) - F_f(v)\|_{B_q} \leq C\|u - v\|_{B_p}$.

On the one hand,

$$\|F_f(u) - F_f(v)\|_{B_q} \leq C(\|u^n - v^n\|_{B_q} + \dots + \|u - v\|_{B_q}).$$

On the other hand, in the same way that we deduced (3.1) and (3.2) we obtain

$$\begin{aligned} \|u^n - v^n\|_{B_q} &\leq C_1 \|u - v\|_{B_p} \sum_{l=0}^{n-1} \|u^{n-1-l} - v^l\|_{B_{p/n-1}} \\ &\leq C_1 \|u - v\|_{B_p} \left(\sum_{l=0}^{n-1} \|u\|_{B_p}^{n-1-l} \|v\|_{B_p}^l \right) \leq C(p, q, R, f) \|u - v\|_{B_p}, \end{aligned}$$

for all $n \leq \left\lfloor \frac{p}{q} \right\rfloor$.

4. The action from B_p to the Bergman-Nevanlinna space

The Bergman-Nevanlinna space is defined by

$$BN = \left\{ u \in H(\Delta) : \|u\|_{BN} = \frac{1}{\pi} \iint_{\Delta} \log^+ |u(z)| \, dx \, dy < \infty \right\}.$$

It is easy to see that $B_p \subset BN, \forall p > 0$. The following result is an easy consequence of Theorem 1.

COROLLARY 1. *Let f be an entire function such that F_f acts from BN to $B_q, 0 < q \leq \infty$. Then f is constant.*

Proof. In particular F_f acts from B_p to B_q , for all $p > 0$. Then from Theorem 1 we conclude that f is a polynomial of degree at most $\left\lfloor \frac{p}{q} \right\rfloor$.

Taking p less than q one obtains the desired conclusion.

LEMMA 3. *If $u \in BN$ then*

$$\log^+ |u(z)| \leq \frac{\|u\|_{BN}}{(1 - |z|)^2}, \quad z \in \Delta.$$

This lemma is a consequence of the subharmonicity of $\log^+ |u|$.

THEOREM 3. *Let f be an entire function. Then F_f acts from $\bigcup_{p < q} B_q$ to BN ($0 < p < \infty$) if and only if f has order at most p .*

Proof. Assume that F_f acts from $\bigcup_{p < q} B_q$ to BN . Let $\varepsilon > 0$. The functions

$$w_1 = u_\varepsilon(z) = \left\{ \frac{1}{1-z} - \frac{1}{2} \right\}^{2/p+\varepsilon}$$

belong to $\bigcup_{p < q} B_q$. Hence, $f \circ u_\varepsilon \in BN$ and so, by Lemma 3

$$|f(w_1)| \leq e^{C/(1-|z|)^2}. \quad (4.1)$$

If $p < 1$ we take ε so that $p + \varepsilon < 1$. Then the range of u_ε is $\mathbb{C} \setminus 0$. Thus, given $w_1 \in \mathbb{C} \setminus 0$ we take $z \in \Delta$ such that

$$|z| = \frac{|w_1^{p+\varepsilon/2} - \frac{1}{2}|}{|w_1^{p+\varepsilon/2} + \frac{1}{2}|}.$$

From (4.1) we get

$$\begin{aligned} \log |f(w_1)| &\leq \frac{C}{(1-|z|)^2} \\ &= \frac{C}{\left(1 - \frac{|w_1^{p+\varepsilon/2} - \frac{1}{2}|}{|w_1^{p+\varepsilon/2} + \frac{1}{2}|}\right)^2} = \frac{C|w_1^{p+\varepsilon/2} + \frac{1}{2}|^2}{(|w_1^{p+\varepsilon/2} + \frac{1}{2}| - |w_1^{p+\varepsilon/2} - \frac{1}{2}|)^2}. \end{aligned} \quad (4.2)$$

If $|w_1| > e^{1/2(p+\varepsilon)}$ then $z_1 = w_1^{p+\varepsilon/2} (-\pi \leq \text{Arg } w < \pi)$ satisfies

$$\text{Re } z_1 > e^{1/4} \cos \left(-\frac{\pi}{2} (p + \varepsilon) \right) > 0.$$

Therefore, there is a positive constant C such that

$$|w_1^{p+\varepsilon/2} + \frac{1}{2}| - |w_1^{p+\varepsilon/2} - \frac{1}{2}| > C, \quad |w_1| > e^{1/2(p+\varepsilon)}. \quad (4.3)$$

Combining (4.2) and (4.3) we obtain

$$\log |f(w_1)| \leq C_1 |w_1^{p+\varepsilon/2} + \frac{1}{2}|^2,$$

for a suitable positive constant C_1 and all w_1 such that $|w_1| > e^{1/2(p+\varepsilon)}$. This

shows that f is of order at most $p + \varepsilon$. Letting ε tend to zero permits us to conclude that f is of order at most p .

Next, we assume that $p \geq 1$. In this case we argue as in Theorem 1 and obtain from (4.2) that

$$\log |f(w)| = O(|w|^{p+\varepsilon})$$

for all w outside a ball. Hence f is an entire function of order at most p .

Let us suppose now that f is an entire function of order less than or equal to p and $u \in \bigcup_{p < q} B_q$. Take $\varepsilon > 0$ such that $u \in B_{p+\varepsilon}$. There is a constant C so that

$$\log^+ M(r, f) \leq r^{p+\varepsilon} + C, \quad \forall r \geq 0.$$

To prove that $f \circ u \in BN$ we write

$$\begin{aligned} \iint_{\Delta} \log^+ |f(u(z))| \, dx \, dy &= \int_0^1 r \, dr \int_0^{2\pi} \log^+ |f(u(r e^{i\theta}))| \, d\theta \\ &\leq \int_0^1 r \, dr \int_0^{2\pi} \log^+ M(|u(r e^{i\theta})|, f) \, d\theta \leq \int_0^1 r \, dr \int_0^{2\pi} |u(r e^{i\theta})|^{p+\varepsilon} \, d\theta + \pi C \\ &= \iint_{\Delta} |u(z)|^{p+\varepsilon} \, dx \, dy + \pi C < \infty, \end{aligned}$$

since $u \in B_{p+\varepsilon}$.

THEOREM 4. *Let f be an entire function of order less than p or of order p and finite type ($0 < p < \infty$). Then F_f acts from B_p to BN .*

Proof. We may assume that f is of order p and finite type $\sigma - \delta > 0$. Hence, there is a constant C such that

$$\log^+ M(r, f) \leq \sigma r^p + C, \quad r \geq 0.$$

If $u \in B_p$ then

$$\begin{aligned} \iint_{\Delta} \log^+ |f(u(z))| \, dx \, dy &= \int_0^1 r \, dr \int_0^{2\pi} \log^+ |f(u(r e^{i\theta}))| \, d\theta \\ &\leq \int_0^1 r \, dr \int_0^{2\pi} \log^+ M(|u(r e^{i\theta})|, f) \, d\theta \\ &\leq \sigma \iint_{\Delta} |u(z)|^p \, dx \, dy + \pi C < \infty, \end{aligned}$$

as required.

5. Transforming Hardy functions into Bergman functions and vice versa

We shall begin by stating the following classical results by Hardy and Littlewood [5].

LEMMA 4. *Let u be analytic in Δ and*

$$M_p(r, u) \leq \frac{C}{(1-r)^\beta}, \quad 0 < p < \infty, \quad \beta \geq 0.$$

Then there is a constant $K = K(p, \beta)$ such that

$$M_{q_1}(r, u) \leq \frac{KC}{(1-r)^{\beta+1/p-1/q_1}}, \quad p < q_1 \leq \infty.$$

A proof of this result can be found in [4, p. 84].

LEMMA 5. *If $0 < p < q_1 \leq \infty$, $u \in H^p$, $\lambda \geq p$, and $\alpha = 1/p - 1/q_1$ then*

$$\int_0^1 (1-r)^{\lambda\alpha-1} M_{q_1}(r, f)^\lambda dr < \infty.$$

The reader can find a proof of this result in [4, p. 87].

The next result shows that one cannot transform Bergman functions into Hardy functions by means of nonlinear superposition. In case $p = \infty$ it is trivial that F_f acts for any f .

THEOREM 5. *Let f be an entire function. If $p \neq \infty$ then F_f acts from B_p to H^q if and only if f is constant.*

Proof. If F_f acts from B_p to H^q then, by Theorem 1, f is a polynomial. Now we get the desired conclusion by noting that for a non-constant polynomial f it is not true that $f \circ u \in H^q$, $\forall u \in B_p$. If this were true then the zeros of all Bergman functions would have to satisfy the Blaschke condition, and this is false.

THEOREM 6. *Let f be an entire function. Then F_f acts from H^p to B_q if and only if f is a polynomial of degree at most $\left\lfloor \frac{2p}{q} \right\rfloor$.*

If $p = \infty$ then F_f acts from H^p to B_q for any f . In the proof of this theorem we shall rule out this case.

COROLLARY 2. *The operator F_f acts from H^p to B_p if and only if f is a polynomial of degree one or two.*

Proof of Theorem 6. The proof that f must be a polynomial of degree less than or equal to $\left[\frac{2p}{q}\right]$ can be done as in Theorem 1. Let us assume now that f is a polynomial of degree $n \leq \left[\frac{2p}{q}\right]$. We shall prove that if $u \in H^p$ then $u^n \in B_q$. Let us suppose, first of all, that $n < \frac{2p}{q}$. Then

$$\begin{aligned} \iint_{\Delta} |u^n(z)|^q \, dx \, dy &= \int_0^1 r \, dr \int_0^{2\pi} |u(r e^{i\theta})|^{nq} \, d\theta \\ &\leq 2\pi \int_0^1 r \, dr \left(\frac{1}{2\pi} \int_0^{2\pi} |u(r e^{i\theta})|^{2p} \, d\theta \right)^{nq/2p} \\ &= 2\pi \int_0^1 r M_{2p}(r, u)^{nq} \, dr. \end{aligned} \tag{5.1}$$

Now, using Lemma 4 with $\beta = 0$, $q_1 = 2p$ we obtain

$$M_{2p}(r, u) \leq \frac{C}{(1-r)^{1/p-1/2p}} = \frac{C}{(1-r)^{1/2p}},$$

for some constant C . Combining this inequality with (5.1) one gets

$$\iint_{\Delta} |u^n(z)|^q \, dx \, dy \leq 2\pi C \int_0^1 \frac{r \, dr}{(1-r)^{nq/2p}} < \infty.$$

Thus $u^n \in B_q$, as required.

Next we assume that $n = 2p/q$. In this case we use Lemma 5 with $q_1 = 2p$, $\alpha = 1/2p$, and $\lambda = 2p$ to conclude that

$$\int_0^1 M_{2p}(r, u)^{2p} \, dr < \infty$$

as required.

6. The action from H^p to BN

If $p = \infty$ and f is any entire function then F_f acts from H^∞ to H^∞ and consequently it acts from H^∞ to BN . When $p < \infty$ we have the following result.

THEOREM 7. *Let $f: \mathbb{C} \rightarrow \mathbb{C}$. Then F_f acts from $\bigcup_{p < q} H^q$ to BN if and only if f is an entire function of order at most $2p$.*

Proof. Let us assume that F_f acts from $\bigcup_{p < q} H^q$ to BN . Clearly f must be an entire function. On the other hand, given $\varepsilon > 0$, the function

$$u_\varepsilon(z) = \left\{ \frac{1}{1-z} - \frac{1}{2} \right\}^{1/p+\varepsilon}$$

belongs to $\bigcup_{p < q} H^q$. Using these functions and Lemma 3 and arguing as in Theorem 1 we deduce that f has order at most $2p$. Conversely, let us suppose that f has order at most $2p$. Let $u \in \bigcup_{p < q} H^q$ and $\varepsilon > 0$ such that $u \in H^{p+\varepsilon}$. Next we take a constant C such that

$$\log^+ M(r, f) \leq r^{2(p+\varepsilon)} + C, \quad \forall r \geq 0,$$

and use this inequality to get

$$\iint_{\Delta} \log^+ |f(u(z))| \, dx \, dy \leq \iint_{\Delta} |u(z)|^{2(p+\varepsilon)} \, dx \, dy + \pi C < \infty,$$

since $u^2 \in B_{p+\varepsilon}$ in view of Corollary 2.

As a corollary we have

COROLLARY 3. *If $u \in N$, $u(z) \neq 0$ in Δ , and $\log u(z) \neq 0$ in Δ , then $e^{(\log u)^{2-\varepsilon}} \in BN$, $\forall \varepsilon, 0 < \varepsilon \leq 2$.*

Proof. For $\varepsilon = 0$ the result breaks down as it is shown by the example

$$u(z) = \exp \left\{ \frac{1+z}{1-z} \right\}.$$

To prove the corollary we proceed as follows. Since u does not vanish in Δ then $\log u$ is analytic there. Moreover $\log |u| \in h^1$, where h^1 is the space of harmonic functions in Δ which satisfy the Riesz-Herglotz representation. This can be seen from the following relations

$$\begin{aligned} \int_0^{2\pi} |\log |u(re^{i\theta})|| \, d\theta &= \int_0^{2\pi} \log^+ |u(re^{i\theta})| \, d\theta + \int_0^{2\pi} \log^- |u(re^{i\theta})| \, d\theta \\ &= 2 \int_0^{2\pi} \log^+ |u(re^{i\theta})| \, d\theta - 2\pi \log |u(0)|, \end{aligned}$$

and the fact that $u \in N$. Then $\log u \in H^p, \forall p < 1$. Thus $(\log u)^{2-\varepsilon} \in \bigcup_{1/2 < q} H^q$,

$\forall \varepsilon, 0 < \varepsilon \leq 2$. Since $f(z) = e^z$ is an entire function of order 1 then, by Theorem 7, $e^{(\log u)^{2-\varepsilon}} \in BN$, as required.

Finally, we have

THEOREM 8. *Let f be an entire function of order less than $2p$ or of order $2p$ and finite type. Then F_f acts from H^p to BN .*

Proof. We may assume that f is of order $2p$ and finite type $\sigma - \delta > 0$. There is a constant C such that

$$\log M(r, f) \leq \sigma r^{2p} + C, \quad r \geq 0.$$

If $u \in H^p$ we get from the last inequality

$$\iint_{\Delta} \log^+ |f(u(z))| \, dx \, dy \leq \sigma \iint_{\Delta} |u(z)|^{2p} \, dx \, dy + \pi C < \infty,$$

since $u^2 \in B_p$ in view of Corollary 2.

7. Some open questions

We finish this article by posing some questions. (1) If F_f acts from B_p to BN then, in particular, it acts from $\bigcup_{p < q} B_q$ to BN and, by Theorem 3, it has order at most p . In case that f has order p is it true that f has finite type? (2) One may ask the corresponding question for the action between H^p and BN .

References

1. Appel, J., Zabrejko, P. P.: *Nonlinear Superposition Operators*. Cambridge University Press, 1990.
2. Cámara, G. A.: The autonomous nonlinear superposition operator in H^p (to appear).
3. Cámara, G. A.: On entire functions acting via nonlinear superposition on Nevanlinna functions (to appear).
4. Duren, P.: *Theory of H^p Spaces*. Academic Press, New York, 1970.
5. Hardy, G. H. and Littlewood, J. E.: Some properties of fractional integrals. II. *Math. Z.* 34 (1932) 403–439.
6. Krasnosel'skij, M. A.: On the continuity of the operator $Fu(x) = f(x, u(x))$ (Russian). *Doklady Akad. Nauk SSSR* 77(2) (1951) 185–188.
7. Marcus, M. and Mizel, V. J.: Every superposition operator mapping one Sobolev space into another is continuous. *J. Funct. Anal.* 33 (1979) 217–229.