

CAHIERS DE TOPOLOGIE ET GÉOMÉTRIE DIFFÉRENTIELLE CATÉGORIQUES

SYMEON BOZAPALIDES

ANESTIS FIRARIDES

Automata and categories

Cahiers de topologie et géométrie différentielle catégoriques, tome
23, n° 2 (1982), p. 149-156

http://www.numdam.org/item?id=CTGDC_1982__23_2_149_0

© Andrée C. Ehresmann et les auteurs, 1982, tous droits réservés.

L'accès aux archives de la revue « Cahiers de topologie et géométrie différentielle catégoriques » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

AUTOMATA AND CATEGORIES

by *Symeon BOZAPALIDES and Anestis FIRARIDES*

Throughout this paper \mathcal{U} denotes a monoidal closed category with tensor product \otimes and internal-hom $[,]$. We assume that \mathcal{U} has countable colimits and kernel pairs.

I. Given a sequence $\Omega = (\Omega_n)$ of objects of \mathcal{U} , an Ω -algebra is simply a pair (A, a_n) with $A \in \mathcal{U}$ and $a_n : \Omega_n \otimes A^n \rightarrow A$. (Ω_n is the object of symbols of n -ary operations of the algebra.) Homomorphisms are defined classically : they are morphisms which respect the Ω -algebra structure :

$$\begin{array}{ccc}
 \Omega_n \otimes A^n & \xrightarrow{a_n} & A \\
 \Omega_n \otimes f^n \downarrow & & \downarrow f \\
 \Omega_n \otimes B^n & \xrightarrow{b_n} & B
 \end{array}$$

We can now construct the free Ω -algebra generated by an object X ; its carrier object is the coproduct $\Omega(X) = \coprod_{n \geq 1} W_n$, where the objects W_n (trees of length n) are inductively defined as follows :

$$\begin{aligned}
 W_1 &= X \amalg \Omega_0 \\
 &\dots\dots\dots \\
 W_{n+1} &= \coprod_{\lambda_1 + \dots + \lambda_k = n} \Omega_k \otimes W_{\lambda_1} \otimes \dots \otimes W_{\lambda_k} .
 \end{aligned}$$

II. One thing of central importance in this paper is the «free approximation» of the translation-monoid of an Ω -algebra (A, a_n) .

Consider the object

$$A^* = \coprod_{1 \leq i \leq n < \omega} \Omega_n \otimes A^{i-1} \otimes \hat{A} \otimes A^{n-i}$$

where « \wedge » means omission of the underlying factor, and ω denotes the first infinite ordinal.

We have a morphism $a^*: A^* \rightarrow [A, A]$ such that the adjoint transform of $a^* \circ in_{i,n}$ is a_n followed by a symmetry morphism.

Call tA the free monoid generated by A^* , i. e., $tA = \coprod_{n \geq 0} (A^*)^n$.

Then a^* is uniquely extended to a monoid homomorphism

$$a^+ : tA \rightarrow [A, A].$$

This says that tA acts on A by $s_A : tA \otimes A \rightarrow A$ obtained from a^+ via $\otimes \dashv [,]$.

III. In the framework described above we examine the realization problem for tree or algebra automata; precisely an Ω -algebra automaton in \mathcal{U} is a 5-tuple $\mathcal{A} = (Q, X, \tau, Y, \beta)$, where $Q = (Q, \delta_n)$ is an Ω -algebra (Q the state object, δ_n the transition arrows) and $\tau : X \rightarrow Q$, $\beta : Q \rightarrow Y$ are morphisms of \mathcal{U} called initial state and output.

The reachability map of \mathcal{A} is the Ω -algebra homomorphism induced by τ , $\bar{\tau} : \Omega(X) \rightarrow Q$

$$\begin{array}{ccc} X & \xrightarrow{j_X} & \Omega(X) \\ & \searrow \tau & \downarrow \bar{\tau} \\ & & Q \end{array}$$

where j_X is the morphism «insertion of generators».

Call \mathcal{A} \mathcal{E} -reachable if $\bar{\tau}$ belongs to a certain class \mathcal{E} of \mathcal{U} -epimorphisms.


Finally the arrow

$$\Omega(X) \xrightarrow{\bar{\tau}} Q \xrightarrow{\beta} Y$$

is the response of \mathcal{A} . Now given a morphism $f : \Omega(X) \rightarrow Y$, we say that \mathcal{A} realizes f iff its response is just f .

Our scope is to construct an \mathcal{E} -reachable automaton (\mathcal{E} = coequalizers of reflexive pairs) $\mathcal{A}_f = (Q_f, X, \tau_f, Y, \beta_f)$ which realizes f in the most economical way, i. e. for every \mathcal{E} -reachable automaton \mathcal{A} which also

realizes f , there exists a unique algebra-homomorphism $\lambda: Q \rightarrow Q_f$ making



commutative.

THEOREM. Any morphism $f: \Omega(X) \rightarrow Y$ has a reflexive coequalizer-reachable minimal realization.

PROOF. Let

$$\tilde{s}_{\Omega(X)}: \Omega(X) \rightarrow [t\Omega(X), \Omega(X)]$$

be the adjoint transform of $s_{Q(X)}$,

$$E_f \begin{array}{c} \xrightarrow{\alpha} \\ \xrightarrow{\gamma} \end{array} \Omega(X)$$

the kernel pair of the morphism

$$\Omega(X) \xrightarrow{\tilde{s}_{\Omega(X)}} [t\Omega(X), \Omega(X)] \xrightarrow{[I, f]} [t\Omega(X), Y]$$

and $r_f: \Omega(X) \rightarrow Q_f$ the coequalizer of (α, γ) .

We use Theorem 2.6 [A. A. M.]. Postulates i and iv of this theorem are evident. To prove Postulate iii, we need the following lemma:

LEMMA. The monoid $t\Omega(X)$ acts on the object Q_f in such a way making r_f an action homomorphism.

PROOF. We have

$$\begin{aligned} f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes s_{\Omega(X)}) \circ (t\Omega(X) \otimes t\Omega(X) \otimes \alpha) &= \\ = f \circ s_{\Omega(X)} \circ (m \otimes \Omega(X)) \circ (t\Omega(X) \otimes t\Omega(X) \otimes \alpha) &= \\ = f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes \alpha) \circ (m \otimes E_f), & \end{aligned}$$

m the multiplication of the monoid $t\Omega(X)$. Similarly

$$\begin{aligned} f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes s_{\Omega(X)}) \circ (t\Omega(X) \otimes t\Omega(X) \otimes \gamma) &= \\ = f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes \gamma) \circ (m \otimes E_f). & \end{aligned}$$

Using the identity

$$f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes a) = f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes \gamma)$$

and passing to adjoint transforms we obtain

$$\begin{aligned} [l, f] \circ \tilde{s}_{\Omega(X)} \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes a) &= \\ &= [l, f] \circ \tilde{s}_{\Omega(X)} \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes \gamma). \end{aligned}$$

Because (a, γ) is the kernel pair of $[l, f] \circ \tilde{s}_{\Omega(X)}$ the above equality implies the existence of a morphism $\psi : t\Omega(X) \otimes E_f \rightarrow E_f$ such that

$$a \circ \psi = s_{\Omega(X)} \circ (t\Omega(X) \otimes a), \quad \gamma \circ \psi = s_{\Omega(X)} \circ (t\Omega(X) \otimes \gamma),$$

and consequently

$$\begin{aligned} r_f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes a) &= r_f \circ a \circ \psi = r_f \circ \gamma \circ \psi = \\ &= r_f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes \gamma). \end{aligned}$$

But then the fact that

$$t\Omega(X) \otimes r_f = \text{coeq}(t\Omega(X) \otimes a, t\Omega(X) \otimes \gamma)$$

gives us the desired action $s : t\Omega(X) \otimes Q_f \rightarrow Q_f$. ■

Return to Postulate iii. We must prove that Q_f admits an Ω -algebra structure making r_f an algebra homomorphism. If we define the morphism

$$h_n^i : \Omega_n \otimes \Omega(X)^{i-1} \otimes Q_f \otimes \Omega(X)^{n-i} \rightarrow Q_f$$

to be the composite

$$\begin{aligned} \Omega_n \otimes \Omega(X)^{i-1} \otimes Q_f \otimes \Omega(X)^{n-i} &\simeq \Omega_n \otimes \Omega(X)^{n-1} \otimes Q_f \xrightarrow{\text{---}} \\ \xrightarrow{\text{---}} \xrightarrow{in_n^i} \Omega(X)^* \otimes Q_f &\xrightarrow{\text{---}} \xrightarrow{in_1} \Omega(X) \otimes Q_f \xrightarrow{\text{---}} \xrightarrow{s} Q_f, \end{aligned}$$

then from the previous lemma we get

$$r_f \circ \mu_n = h_n^i \circ (\Omega_n \otimes \Omega(X)^{i-1} \otimes r_f \otimes \Omega(X)^{n-i})$$

(where $\mu_n : \Omega_n \otimes \Omega(X)^n \rightarrow \Omega(X)$ are the structural arrows of the free Ω -algebra $\Omega(X)$); therefore

$$\left. \begin{aligned} h_n^i \circ (\Omega_n \otimes \Omega(X)^{i-1} \otimes r_f \otimes \Omega(X)^{n-i}) &= \\ = h_n^j \circ (\Omega_n \otimes \Omega(X)^{j-1} \otimes r_f \otimes \Omega(X)^{n-j}) &= r_f \circ \mu_n \end{aligned} \right\} (\Delta_n^i, j)$$

for any $n \in \mathbb{N}$ and $1 \leq i, j \leq n$.

$$k_\lambda \circ (\Omega_n \otimes \Omega(X)^{n-\lambda} \otimes r_f \otimes Q_f^{\lambda-1}) = k_{\lambda-1}.$$

We take $k_1 = h_n^n$. k_1 coequalizes

$$(\Omega_n \otimes \Omega(X)^{n-2} \otimes \alpha \otimes Q_f, \Omega_n \otimes \Omega(X)^{n-2} \otimes \gamma \otimes Q_f).$$

(This comes if we put (1) together with


$$\Omega_n \otimes E_f^{n-1} \otimes r_f = \text{coeq}(\Omega_n \otimes E_f^{n-1} \otimes \alpha, \Omega_n \otimes E_f^{n-1} \otimes \gamma)$$

and the fact that

$$(\Omega_n \otimes \alpha^{n-2} \otimes E_f \otimes Q_f, \Omega_n \otimes \gamma^{n-2} \otimes E_f \otimes Q_f)$$

is reflexive.) Consequently there exists a unique

$$k_2: \Omega_n \otimes \Omega(X)^{n-2} \otimes Q_f^2 \rightarrow Q_f \quad \text{with} \quad k_2 \circ (\Omega_n \otimes \Omega(X)^{n-2} \otimes r_f \otimes Q_f) = k_1.$$



k_3 results from k_2 by a similar as above argument, and so on. Finally, we determine $k_n: \Omega_n \otimes Q_f^n \rightarrow Q_f$ which is the n -th structural arrow of Q_f .

It remains to show that r_f is an algebra homomorphism. We have

$$\begin{aligned} r_f \circ \mu_n &= k_1 \circ (\Omega_n \otimes \Omega(X)^{n-1} \otimes r_f) \\ &= k_2 \circ (\Omega_n \otimes \Omega(X)^{n-2} \otimes r_f \otimes Q_f) \circ (\Omega_n \otimes \Omega(X)^{n-1} \otimes r_f) \\ &\dots \dots \dots \\ &= k_{n-1} \circ (\Omega_n \otimes \Omega(X) \otimes r_f \otimes Q_f^{n-2}) \circ \dots \circ (\Omega_n \otimes \Omega(X)^{n-1} \otimes r_f) \end{aligned}$$

$$\begin{aligned}
 &= k_n \circ (\Omega_n \otimes r_f \otimes Q_f^{n-1}) \circ \dots \circ (\Omega_n \otimes \Omega(X)^{n-1} \otimes r_f) \\
 &= k_n \circ (\Omega_n \otimes r_f^n).
 \end{aligned}$$

We now establish the postulate i. It suffices to prove that f coequalizes the pair $(\bar{a}, \bar{\gamma})$ in a universal way, i. e. wherever $f \circ \bar{p} = f \circ \bar{q}$ also holds for a reflexive pair

$$R \begin{array}{c} \xrightarrow{p} \\ \xrightarrow{q} \end{array} \Omega(X),$$

there exists a unique arrow $h: R \rightarrow E_f$ such that $a \circ h = p$, $\gamma \circ h = q$. (\bar{a} , $\bar{\gamma}$, \bar{p} , \bar{q} are the unique algebra homomorphisms deduced by a, γ, p, q respectively.) The equality $f \circ \bar{a} = f \circ \bar{\gamma}$ comes easily if we prove that $f \circ a = f \circ \gamma$. Let $\phi: Q_f \rightarrow [t\Omega(X), Y]$ be the unique arrow such that

$$\phi \circ r_f = [l, f] \circ \bar{s}_{\Omega(X)}.$$

If $u: l \rightarrow t\Omega(X)$ is the unit of the monoid $t\Omega(X)$, then we have

$$\psi \circ (u \otimes E_f) = Id_{E_f},$$

because ψ is an action. Therefore

$$\begin{aligned}
 f \circ a &= f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes a) \circ (u \otimes E_f) \\
 &= \bar{\phi} \circ (t\Omega(X) \otimes r_f) \circ (t\Omega(X) \otimes a) \circ (u \otimes E_f) \\
 &= \bar{\phi} \circ (t\Omega(X) \otimes r_f) \circ (t\Omega(X) \otimes \gamma) \circ (u \otimes E_f) \\
 &= f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes \gamma) \circ (u \otimes E_f) = f \circ \gamma,
 \end{aligned}$$

where $\bar{\phi}$ is the transform of ϕ via $\otimes \dashv [,]$.

Now, let $\epsilon: \Omega(X) \rightarrow \Omega(R)$ be the common right inverse of \bar{p} and \bar{q} : $\bar{p} \circ \epsilon = Id = \bar{q} \circ \epsilon$. We have

$$\bar{q} \circ s_{\Omega(R)} = s_{\Omega(X)} \circ (t\bar{q} \otimes \bar{q}), \quad \bar{p} \circ s_{\Omega(R)} = s_{\Omega(X)} \circ (t\bar{p} \otimes \bar{p}),$$

where $t\bar{p}$, $t\bar{q}$ are the free monoid homomorphisms induced by \bar{p} , \bar{q} respectively. Hence

$$\begin{aligned}
 f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes \bar{p}) &= f \circ s_{\Omega(X)} \circ (t\bar{p} \otimes \bar{p}) \circ (t\epsilon \otimes \Omega(R)) = \\
 &= f \circ \bar{p} \circ s_{\Omega(R)} \circ (t\epsilon \otimes \Omega(R)) = f \circ \bar{q} \circ s_{\Omega(R)} \circ (t\epsilon \otimes \Omega(R)) = \\
 &= f \circ s_{\Omega(X)} \circ (t\bar{q} \otimes \bar{q}) \circ (t\epsilon \otimes \Omega(R)) = f \circ s_{\Omega(X)} \circ (t\Omega(X) \otimes \bar{q}),
 \end{aligned}$$

or

$$[I, f] \circ \tilde{s}_{\Omega(X)} \circ \bar{p} = [I, f] \circ \tilde{s}_{\Omega(X)} \circ \bar{q}$$

by adjoint transformation. Therefore there exists a (unique) arrow

$$\delta: \Omega(R) \rightarrow E_f \text{ such that } \alpha \circ \delta = \bar{p}, \gamma \circ \delta = \bar{q}.$$

It is now clear that the morphism

$$h = \delta \circ j_R: R \rightarrow E_f$$

has the desired property, and this concludes the proof of the theorem.

REFERENCE.

A.A.M. ANDERSON B.D.O., ARBIB M. A. & MANES E.G., Foundations of system theory: Finitary and infinitary conditions, *Lecture Notes in Economics and Math. systems* 115, Springer (1974).

Department of Mathematics
University of Ioannina
IOANNINA, GRECE.