

# GROUPE DE TRAVAIL D'ANALYSE ULTRAMÉTRIQUE

WERNER LÜTKEBOHMERT

## **Families of Mumford curves**

*Groupe de travail d'analyse ultramétrique*, tome 9, n° 3 (1981-1982), exp. n° J12, p. J1-J5

[http://www.numdam.org/item?id=GAU\\_1981-1982\\_\\_9\\_3\\_A13\\_0](http://www.numdam.org/item?id=GAU_1981-1982__9_3_A13_0)

© Groupe de travail d'analyse ultramétrique  
(Secrétariat mathématique, Paris), 1981-1982, tous droits réservés.

L'accès aux archives de la collection « Groupe de travail d'analyse ultramétrique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

FAMILIES OF MUMFORD CURVES

by Werner LÜTKEBOHMERT (\*)  
 [Universität Münster]

In this lecture, we consider the following :

$k$  algebraically closed,  $p$ -adic field with residue field  $\tilde{k}$ .

$\pi : X \rightarrow S$  family of Mumford curves, i. e. a proper, flat,  $k$ -analytic morphism such that all fibres  $X_s := \pi^{-1}(s)$  are Mumford curves with genus  $g(X_s)$ .

0. General facts

0.1. The Euler-Poincaré characteristic  $EP(\mathcal{O}, \pi, s) = g(X_s)$  is locally constant on  $S$  in the sense of Grothendieck topologies. If  $S$  is connected, then  $g(X_s) = g$  is constant on  $S$ .

0.2. If  $g = g(X_s) \geq 2$ , the relative-tricanonical linebundle  $\omega_{X/S}^{\otimes 3}$  is very ample on all fibres. The direct image  $\xi := \pi_* \omega_{X/S}^{\otimes 3}$  is a holomorphic vectorbundle on  $S$  of rank  $5g - 5$  and gives an embedding of the family

$$\begin{array}{ccc} X & \xrightarrow{\quad\quad\quad} & P(\xi) \\ & \searrow & \swarrow \\ & S & \end{array}$$

If  $g = 1$ , one gets the same, if one takes  $\mathcal{O}(-3D)$  instead of  $\omega_{X/S}^{\otimes 3}$ , where  $D \subset X$  is a divisor, finite over  $S$ .

0.3. Now by the GAGA-theorem in the relative case one knows, that for all affinoid subdomains  $U = \text{Sp}A \subset S$  the restricted family  $\pi : X|U \rightarrow U$  is an algebraic morphism.

I. Uniformization of families

All fibres  $X_s$  in the family have an uniformization. Now one wants to get a

(\*) Werner LÜTKEBOHMERT, Mathematisches Institut der Universität, 62 Einsteinstrasse, D-4400 MÜNSTER (Allemagne fédérale).


simultaneous uniformization of the family locally on the base  $S$ . To explain what I mean, I will repeat the definition of Mumford curve : In our situation, we know

$$X_s \simeq \Omega_s / \Gamma_s ,$$

$$\Gamma_s = \langle \gamma_1^s , \dots , \gamma_g^s \rangle \subset \text{PGL}(2 , k) \text{ free, discontinuous group,}$$

$$\Omega_s \subset \mathbb{P}^1 \text{ the set of ordinary points.}$$

Moreover one can assume that  $\gamma^s$  is a geometric base, i. e. there are closed balls  $U_i^s$  which are pairwise disjoint such that  $\gamma_i^s(\mathbb{P}^1 - U_i^s) = \dot{U}_{-i}^s$  for  $i = 1 , \dots , g$ .



$$U_i^s = \{z \in \mathbb{P}^1 ; |z - a_i^s| \leq |r_i^s|\}$$

$$U_i^s = \{z \in \mathbb{P}^1 ; |z - a_i^s| < |r_i^s|\}$$

"Simultaneous Uniformization" means that all  $\gamma_i^s , a_i^s , r_i^s$  depend holomorphically on  $s \in S$ .

THEOREM I. - Let  $S$  be a reduced affinoid space,  $\pi : X \rightarrow S$  an analytic family of Mumford curves of genus  $g$  (with  $g + 1 < p = \text{char } \tilde{k}$  or  $0 = \text{char } \tilde{k}$ ), then there is a finite base extension  $S' \rightarrow S$  and a finite covering  $\{S'_1 , \dots , S'_r\}$  of  $S'$  by affinoids  $S'_i$ , such that  $X'_i := X \times_S S'_i$  have simultaneous uniformizations, i. e. :

$$\mathbb{P}_{S'_i}^1 \supset \Omega'_i \rightarrow X'_i \text{ } S'_i\text{-morphism,}$$

$$\gamma^i : S'_i \rightarrow (\text{PGL}(2 , k))^g \text{ holomorphic,}$$

such that

$\Gamma'_i := \langle \gamma^i \rangle$  act holomorphically, discontinuously on  $\Omega'_i$  by the canonical action  
 $\mathbb{P}_{S'_i}^1$  ,

$$\Omega'_i / \Gamma'_i \simeq X'_i \text{ isomorphic over } S'_i ,$$

$$\gamma^i(s) \text{ is a geometric base for } \Gamma'_i(s) .$$

To prove this theorem, one makes the following steps :

I.1. There is a finite base extension  $S' \rightarrow S$ , a finite covering  $\{S'_1 , \dots , S'_r\}$  of  $S'$  by affinoids  $S'_i$ , and a finite covering  $\{X'_{ij}\}_j$  von  $S'_i$  with

$$X'_{ik} \simeq \{(s, z) \in S'_i \times \tilde{P}^1; |z - a_{ik}^0(s)|^{m_0} \leq |r_{ik}^0(s)|,\$$

$$|z - a_{ik}^v(s)|^{m_v} \leq |r_{ik}^v(s)| \text{ for } 1 \leq v \leq n_{ik}\}$$

where  $m_v \geq 1$  are natural members, and  $a_{ik}^v, r_{ik}^v \in \mathcal{O}(S'_i)$  with

$$|a_{ik}^v(s) - a_{ik}^0(s)|^{m_0} < |r_{ik}^0(s)| \text{ for } 1 \leq v \leq n_{ik}$$

$$|a_{ik}^v(s) - a_{ik}^\mu(s)|^{m_v} > |r_{ik}^v(s)| \text{ for } 1 \leq v \neq \mu \leq n_{ik}$$


$$0 < |r_{ik}^v(s)|^{m_0} < |r_{ik}^0(s)|^{m_v} \text{ for } 1 \leq v \leq n_{ik}$$

For the components  $U_v$  of  $X_{ik} \cap X_{i\ell}$  one has :

$$U_v \simeq \{(s, z) \in S'_i \times \tilde{P}^1; |r_v^-(s)| \leq |z - a_v(s)|^{m_v} \leq |r_v^+(s)|\}$$

where  $r_v^+, r_v^-, a_v \in \mathcal{O}(S'_i)$  satisfy  $0 < |r_v^-(s)| < |r_v^+(s)|$  for all  $s \in S'_i$ .

Sketch of proof : After finite base extension  $S^* \rightarrow S$  one finds a section  $\sigma$  to  $\pi^* := \pi \times_S S^*$ . Then by the theorem of Riemann-Roch, one finds a finite morphism




with  $\deg \varphi \leq g + 1 < p$ .

Now after finite base extension  $S^+ \rightarrow S^*$  the ramification set of  $\varphi \times_{S^*} S^+$  "essentially" splits into divisors of degree 1. So there are holomorphic sections  $\tau_i : S^+ \rightarrow \tilde{P}^1_{S^+}$  such that

$$\text{Ram}(\varphi \times_{S^*} S^+) = \tau_1 S^+ \cup \dots \cup \tau_r S^+.$$

Then one constructs the coverings  $\{S'_i\}$  of  $S'$  and  $\{X'_{ij}\}$  of  $X'_i$  by using these sections  $\{\tau_\rho\}$  and by making a further finite base extension  $S' \rightarrow S^+$ . The main tool here is the method of constructing a stable reduction of an algebraic curve  $M$ , which admits a morphism  $\psi : M \rightarrow \tilde{P}^1$  with  $\deg \psi < p$ , by the ramification points of  $\psi$ .


I.2. By the covering  $\{X'_{ij}\}_j$  one constructs a  $k$ -analytic  $S'_i$ -morphism  $\Omega'_i \rightarrow X'_i$  such that  $\Omega'_i(s)$  is the universal covering of  $X'_i(s)$  for all  $s \in S'_i$ . Next one proves an embedding theorem



By this embedding the group  $\text{Deck}(\Omega'_i/X'_i)$  can be regarded as a subgroup of  $\text{PGL}(2, \mathcal{O}(S'_i))$ . From this, one easily gets the theorem.

COROLLARY. - Let  $\mathcal{B}_g$  be the space of geometric bases for Mumford curves,  $\pi_g$  the space of Mumford curves and  $g \geq 2$ .

For every family  $X \rightarrow S$  of Mumford curves of genus  $g \geq 2$ , the canonical map  $S \rightarrow \pi_g$  admits locally a holomorphic lifting after finite base extension




II. Rigidity of algebraic families

An algebraic family of Mumford curves  $\pi : X \rightarrow S$  is a proper, flat, algebraic morphism where  $S$  is a connected scheme of finite type over  $k$ , such that all fibres  $X_s$  are Mumford curves. Let  $g$  be the genus of the fibres.

THEOREM II. - Every algebraic family of Mumford curves is constant, i. e. the map  $S \rightarrow \pi_g, s \mapsto [X_s]$ , is constant (if  $g + 1 < p = \text{char } \tilde{k}$  or  $0 = \text{char } k$ ).

Idea of the proof. - It is enough to consider affine, irreducible, algebraic curves for  $S$ . Similar as in § I, one can show, that there is a finite extension  $S' \rightarrow S$  by an algebraic curve such that  $X' = X \times_S S'$  has locally on  $S'$  a simultaneous uniformization. If  $\Omega' \rightarrow S'$  is the universal covering, then the canonical map can be lifted



Now  $\mathcal{B}_g$  is bounded in some sense, then  $\Omega' \rightarrow \mathcal{B}_g$  has to be constant by the :

PROPOSITION. - Bounded holomorphic functions on the universal covering of an algebraic curve are constant.

COROLLARY. - Analytic families  $X \rightarrow S$  over an connected, complete, algebraic curve are constant (GAGA).

Remarks.

(1) All considerations are made only for  $g \geq 2$ , because the case  $g = 0$  is trivial and, in the case  $g = 1$ , one gets rigidity for analytic families over connected, affine, algebraic curves by considering the  $j$ -invariants of the fibres as holomorphic function on the base.

(2) The assumption concerning the genus " $g + 1 < \text{char } \tilde{k}$ " should not be necessary, like a new research by the author has shown.

(3) Theorem II should be true for analytic families of Mumford curves over connected schemes of finite type over  $k$  as in the case  $g = 1$ .

This lecture is a very short version of my paper "Ein globaler Starrheitssatz für Mumford Kurven" which will come out in "Journal für reine und angewandte Mathematik" in the next months.

-----