

GROUPE DE TRAVAIL D'ANALYSE ULTRAMÉTRIQUE

ALAIN ESCASSUT

Numbers with order ≤ 1

Groupe de travail d'analyse ultramétrique, tome 9, n° 3 (1981-1982), exp. n° J6, p. J1-J4

http://www.numdam.org/item?id=GAU_1981-1982__9_3_A7_0

© Groupe de travail d'analyse ultramétrique
(Secrétariat mathématique, Paris), 1981-1982, tous droits réservés.

L'accès aux archives de la collection « Groupe de travail d'analyse ultramétrique » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

NUMBERS WITH ORDER ≤ 1

by Alain ESCASSUT (*)

[Université de Bordeaux-I]

Let p be a prime integer, let \mathbb{Q}_p be the completion of \mathbb{Q} for the p -adic absolute value $|\cdot|$, let Ω_p be an algebraic closure of \mathbb{Q}_p provided with the unique extension of the absolute value $|\cdot|$, and let \mathbb{C}_p be the (algebraically closed) completion of Ω_p .

In \mathbb{C}_p , we propose to study a notion of "transcendence order $\leq \alpha$ over \mathbb{Q}_p " which looks a little like the transcendence type over \mathbb{Q} ([2], [3], [4]).

Let $\|\cdot\|$ be the Gauss norm on $\mathbb{Q}_p[X]$ defined by

$$\|\sum_{i=0}^n a_i X^i\| = \max_{0 \leq i \leq n} |a_i|.$$

Let K be a transcendence degree 1 extension of \mathbb{Q}_p . Let $|\cdot|$ be an extension to K of the absolute value defined on \mathbb{Q}_p , and let v be the valuation defined by $v(x) = -\log |x|$ where \log is the logarithm function in base p . An element $x \in K$, transcendental over \mathbb{Q}_p , is said to have order $\leq \alpha$ ($\alpha > 0$) if there exists a positive constant C_x such that every polynomial $P(X) \in \mathbb{Q}_p[X]$ satisfies

$$v(P(x)) \leq -\log \|P\| + C_x (\deg P)^\alpha.$$

No $x \in \mathbb{C}_p$ can have order $\leq \alpha$, if $\alpha < 1$, but with help of an example, we prove there exist some $x \in \mathbb{C}_p$ transcendental over \mathbb{Q}_p , with order ≤ 1 (theorem 1). On the other hand, we know that order $\leq \alpha$ is stable by algebraic extension.

At the Oberwolfach meeting, we proved the existence of some $x \in \mathbb{C}_p$ with order $\leq 1 + \epsilon$ whenever $\epsilon > 0$. Here, we give a new kind of number, with order ≤ 1 .

THEOREM. - There exist some numbers $t \in \mathbb{C}_p$, transcendental over \mathbb{Q}_p , of order ≤ 1 .

In the proof of theorem, we will use the following notations.

For any finite set S , let $\text{card}(S)$ be the number of elements of S .

Let H be an algebraic extension of \mathbb{Q}_p , and let $\text{Aut}(H)$ be the group of the \mathbb{Q}_p -automorphisms in H . We know that every algebraic extension of \mathbb{Q}_p is normal [1]

(*) Alain ESCASSUT, Laboratoire de Mathématiques et informatique associé au CNRS, Université de Bordeaux-I, 351 cours de la Libération, 33405 TALENCE CEDEX.

hence it follows the classical result.

LEMMA. - Let H be an algebraic extension of \mathbb{Q}_p . Then $[H : \mathbb{Q}_p] = \text{card}(\text{Aut}(H))$

Proof of theorem. - Let (u_i) , $i \in \mathbb{N}$, be a sequence in Ω_p such that $v(u_i) = 0$ and $\mathbb{Q}_p[u_i]$ is the unramified extension of degree 2^i over \mathbb{Q}_p . Then we have

$$(1) \quad v(u_i - \sigma(u_i)) = 0 \text{ for every } \mathbb{Q}_p\text{-automorphism } \sigma \text{ in } \Omega_p.$$

Let (f_i) , $i \in \mathbb{N}$, be a strictly increasing sequence of positive integers such that the series $\sum_{i=1}^{\infty} f_i/2^i$ converges.

Set $a_i = u_i p^{f_i}$, and let $t = \sum_{i=1}^{\infty} a_i \in \mathbb{C}_p$. We are going to show that t has order ≤ 1 .

Clearly, it is enough to prove the existence of some $C \in \mathbb{R}^+$ such that every irreducible polynomial $F(X) \in \mathbb{Q}_p[X]$, such that $\|F\| = 1$, satisfies

$$(2) \quad v(F(t)) \leq C \deg F.$$

Consider such a polynomial F and set

$$F(X) = \lambda \prod_{i=1}^d (X - b_i), \text{ where } b_1, \dots, b_d \text{ are conjugate over } \mathbb{Q}_p.$$

We know that the $v(b_i)$, $1 \leq i \leq d$, are equal [1]. If the $v(b_i)$ are < 0 , we have

$$v(F(t)) = v(\lambda) + d v(b_i) = -\log \|F\| = 0.$$

So we can assume $v(b_i) \geq 0$ to prove the relation (2).

Then, by classical results [1], we have $v(\lambda) = 0$ hence we can assume that F is monic.

Now, by reordering the b_i , we can obviously assume that

$$v(t - b_i) \geq v(t - b_j) \text{ for every } i = 2, \dots, d.$$

Set $b = b_1$ and let N be the unique integer such that

$$f_N < v(t - b) \leq f_{N+1}.$$

Write $b = a_1 + \dots + a_N + a'$; then $v(a') \geq v(t - b) > f_N$.

Let $L = \mathbb{Q}_p[a_1, \dots, a_N, a']$. Obviously, $\mathbb{Q}_p[b] \subseteq L$.

We are going to prove $\mathbb{Q}_p[b] = L$. Let $\sigma \in \text{Aut}(L)$ be different of the identical automorphism $\underline{1}$ and let us prove that $\sigma(b) \neq b$. First suppose there exist integers $h \leq N$ such that $\sigma(a_h) \neq a_h$, and let i be the lowest of these integers h . By hypothesis (1), we have $v(\sigma(a_i) - a_i) = v(a_i) = f_i$. It follows that $v(\sigma(b) - b) = f_i$ since the f_N are strictly increasing, and $v(a') > f_N$. Similarly, suppose now i does not exist; then

$$\sigma(b) - b = \sigma(a') - a'.$$

But necessarily $\sigma(a') \neq a'$ since $\sigma \neq 1$ and $\sigma(a_i) = a_i, \forall i = 1, \dots, N$.

Thus we see that the $\sigma(b)$ are all different, hence $\text{card}(\text{Aut } \mathbb{Q}_p[b]) \geq \text{card}(\text{Aut } L)$; then by lemma 1, $[\mathbb{Q}_p[b] : \mathbb{Q}_p] \geq [L : \mathbb{Q}_p]$ and finally $\mathbb{Q}_p[b] = L$.

Now, for each $j = 1, \dots, d$, b_j is some $\sigma(b)$ (with $\sigma \in \text{Aut } \Omega_p$).

Then $b_j = \sigma(a_1) + \dots + \sigma(a_N) + \sigma(a')$. Consider

$$b - b_j = \sum_{i=1}^N a_i - \sigma(a_i) + \sum_{i=N+1}^{\infty} a_i - \sigma(a').$$

By (1), we have $v(a_i - \sigma(a_i)) = v(a_i) = f_i$, and

$$v(\sum_{i=N+1}^{\infty} a_i - \sigma(a')) > f_N \text{ since } v(\sum_{i=N+1}^{\infty} a_i) \geq f_{N+1} \text{ and } \sigma(a') > f_N.$$

Thus the $v(b - b_j)$ are necessarily in the form f_ℓ with $\ell \leq N$, when they are not strictly superior to f_N . Then it is the same thing for the $v(t - b_j)$.

For each integer $\ell \leq N$, let

$$E_\ell = \{b_j ; v(t - b_j) = f_\ell\};$$

then it follows $\text{card } E_\ell$ is equal to the number A_ℓ of the $\sigma \in \text{Aut}(L)$ which fix $a_1, \dots, a_{\ell-1}$. Then by definition of the a_n (from u_n), we see clearly that every $\sigma \in \text{Aut}(L)$ which fixes a_i also fixes a_1, \dots, a_{i-1} , hence $A_\ell = d/2^\ell$ and $\text{card}(E_\ell) = d/2^\ell$. So, we have

$$\sum_{\substack{b_j \in E_\ell \\ 1 \leq \ell \leq N}} v(t - b_j) = \sum_{i=1}^N \frac{df_i}{2^i}$$

Similarly the set $\tilde{E}_N = \{b_j ; v(t - b_j) > f_N\}$ has $d/2^N$ elements. Since, for any $j \leq d$, we have $v(t - b_j) \leq v(t - b_1) \leq f_{N+1}$, we see that

$$\sum_{b_j \in \tilde{E}_N} v(t - b_j) \leq \frac{df_{N+1}}{2^N}.$$

Finally

$$v(F(t)) = \sum_{j=1}^d v(t - b_j) \leq (\sum_{i=1}^N \frac{df_i}{2^i}) + \frac{df_{N+1}}{2^N}.$$

Set $C = \sum_{i=1}^{\infty} f_i/2^{i-1}$; with greater reason, $v(F(t)) \leq C d$.

Thus, every irreducible polynomial F such that $\|F\| = 1$ satisfies $v(F(t)) \leq C \text{ deg } F$, hence by preliminary remarks, theorem 1 is proved.

BIBLIOGRAPHY

- [1] AMICE (Y.). - Les nombres p-adiques. - Paris, Presses universitaires de France, 1975 (Collection SUP, "Le Mathématicien", 14).

- [2] ESCASSUT (A.). - Type de transcendance p -adique, Groupe d'étude d'Analyse ultramétrique, 5e année, 1977/78, n° 8, 10 p.
 - [3] LANG (S.). - Introduction to transcendental numbers. - Reading, Addison-Wesley, 1966 (Addison-Wesley Series in Mathematics).
 - [4] WALDSCHMIDT (M.). - Nombres transcendants. - Berlin, Springer-Verlag, 1974 (Lecture Notes in Mathematics, 402).
-