

# JOURNÉES ÉQUATIONS AUX DÉRIVÉES PARTIELLES

CHARLES FEFFERMAN

D. H. PHONG

## Lower bounds for Schrödinger equations

*Journées Équations aux dérivées partielles* (1982), p. 1-7

[http://www.numdam.org/item?id=JEDP\\_1982\\_\\_\\_\\_A7\\_0](http://www.numdam.org/item?id=JEDP_1982____A7_0)

© Journées Équations aux dérivées partielles, 1982, tous droits réservés.

L'accès aux archives de la revue « Journées Équations aux dérivées partielles » (<http://www.math.sciences.univ-nantes.fr/edpa/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

LOWER BOUNDS FOR SCHRODINGER EQUATIONS

by C. FEFFERMAN and D. H. PHONG

The main purpose of this article is to establish the following theorem on the lowest eigenvalue of Schrödinger equations with no regularity assumptions on the potentials :

Theorem : Let  $V(x)$  be a negative function on  $\mathbb{R}^n$ , and set

$$E_{\text{small}} = \sup_Q \left[ \left( \frac{1}{|Q|} \int_Q |V(x)| dx \right) - c(\text{diam } Q)^{-2} \right]$$

$$E_{\text{big}} = \sup_Q \left[ \left( \frac{1}{|Q|} \int_Q |V(x)|^p dx \right)^{1/p} - c(\text{diam } Q)^{-2} \right] \quad p > 1$$

where  $Q$  ranges over cubes in  $\mathbb{R}^n$  with sides parallel to the axes,  $|Q|$  and  $\text{diam } Q$  are respectively the measure and the diameter of  $Q$ , and  $C, c$  denote throughout constants depending only on the dimension of  $\mathbb{R}^n$ .

Then the lowest bound  $-E$  of the operator  $-\Delta + V(x)$  satisfies the inequalities

$$cE_{\text{small}} \leq E \leq CE_{\text{big}} \tag{1}$$

Remarks (a) It may happen that  $E_{\text{big}} = 0$ , in which case  $-\Delta + V(x)$  has been proved to be positive. When  $n \geq 3$  we may thus apply the theorem with  $p < n/2$  and obtain as a corollary the Sobolev inequality, i.e.,  $\forall f \in L^2$  implies  $f \in L^{2n/(n-2)}$ . In  $\mathbb{R}^2$  the sharp form of the Sobolev inequality states that  $\forall f \in L^2$  implies  $e^{\lambda|f|^2} \in L^1_{\text{loc}}$  for small positive  $\lambda$ . This is sharper than our result.

(b) In the definition of  $E_{\text{big}}$ , one cannot replace  $p > 1$  simply by 1. In fact in  $\mathbb{R}^2$  this would imply  $\forall f \in L^2 \Rightarrow f \in L^\infty$ , which is false. To get counterexamples in  $\mathbb{R}^n$  ( $n > 2$ ) just consider potentials of the form  $V(x', x'') = V(x')$  with  $x' \in \mathbb{R}^2$ ,  $x'' \in \mathbb{R}^{n-2}$ .

(c) It is a natural guess that the sharp lower bound should involve  $L \text{Log}^+ L$  norms of  $|V|$  rather than  $L^p$  norms. This would agree with the Sobolev inequality on  $\mathbb{R}^2$ . If this were the case, however, a study of the function  $f^\#$  (see below) alone would not suffice, since  $f^\# \in L^\infty$  would merely imply  $f \in \exp(L)$ .

(d) Similar information can be obtained for higher eigenvalues, this development will be reported on in detail elsewhere.

The proof of the theorem is based on three lemmas, the first providing the key estimate in this work, while the other two are variants of results from harmonic analysis established in the last decade (see e. g. [1] [2] and the references therein). We begin by recalling some definitions :

(1) A positive measure  $\mu$  is said to satisfy condition  $(A_\infty)$  if for any  $\varepsilon > 0$  there exists a  $\delta > 0$  such that

$$E \subset Q \text{ and } |E| < \delta |Q| \text{ implies } \mu(E) < \varepsilon \mu(Q).$$

$$(2) \quad \begin{aligned} f_Q &= \frac{1}{|Q|} \int_Q f(x) dx \\ f^*(x) &= \sup_{x \in Q} \frac{1}{|Q|} \int_Q |f(y)| dy \\ f^\#(x) &= \sup_{x \in Q} \frac{1}{|Q|} \int_Q |f(y) - f_Q| dy \end{aligned}$$

Fix a cube  $Q^\circ$ .

Lemma 1 : Let  $\mu$  be a positive measure on  $Q^\circ$  such that

$$\mu(Q) \leq C |Q| (\text{diam } Q)^{-2} \quad \text{for all cubes } Q \subseteq Q^\circ. \quad (2)$$

Then

$$\int_{Q^\circ} |f^\#|^2 d\mu \leq c' \int_{Q^\circ} |\nabla f|^2 dx \quad (3)$$

Lemma 2 : If  $\mu$  satisfies  $(A_\infty)$  then

$$\int_{Q^\circ} |f(x) - f_{Q^\circ}|^2 d\mu(x) \leq c \int_{Q^\circ} |f^\#(x)|^2 d\mu(x) \quad (4)$$

Lemma 3 : Let  $d\mu = |V(x)| dx$ , and assume that  $(p > 1)$

$$\left( \frac{1}{|Q|} \int_Q |V(x)|^p dx \right)^{1/p} \leq c (\text{diam } Q)^{-2} \quad \text{for all cubes } Q \subset Q^\circ.$$

Then the measure  $d\mu^+$  defined by  $d\mu^+ = [(|V(x)|^p)^*]^{1/p} dx$  will satisfy both  $(A_\infty)$  and the condition (2).

The constants in the conclusions can evidently be taken to be small if the constants in the hypotheses are small.

Proof of the theorem : Applying  $-\Delta + V(x)$  to the dilates and translates of a fixed  $C_0^\infty$  function, we get the estimate involving  $E_{\text{small}}$  at once. To establish the estimate involving  $E_{\text{big}}$ , cut  $\mathbb{R}^n$  into a grid of cubes  $Q^0$  of equal size  $\text{diam } Q^0 = (cE_{\text{big}})^{-1/2}$ , and define  $d\mu, d\mu^+$  as in Lemma 3. Observe then that if the constants  $c$  appearing in the definition of  $E_{\text{big}}$  and  $Q^0$  are small, we have

$$\begin{aligned} \int_{Q^0} |f_{Q^0}|^2 d\mu &\leq |f_{Q^0}|^2 \mu(Q^0) \\ &\leq \left( \frac{1}{|Q^0|} \int_{Q^0} |f|^2 dx \right) [C|Q^0| (\text{diam } Q^0)^{-2}] \\ &\leq \frac{E_{\text{big}}}{2} \int_{Q^0} |f|^2 dx \end{aligned} \quad (5)$$

while

$$\begin{aligned} \int_{Q^0} |f(x) - f_{Q^0}|^2 d\mu(x) &\leq \int_{Q^0} |f(x) - f_{Q^0}|^2 d\mu^+(x) \\ &\leq c \int_{Q^0} |f^\#(x)|^2 d\mu^+(x) \\ &\leq \frac{1}{2} \int_{Q^0} |\nabla f(x)|^2 dx \end{aligned} \quad (6)$$

in view of Lemmas 1, 2, and 3. The inequalities (5) and (6) in turn yield

$$\int_{Q^0} |f(x)|^2 d\mu(x) \leq \int_{Q^0} |\nabla f(x)|^2 dx + E_{\text{big}} \int_{Q^0} |f(x)|^2 dx$$

and the desired bound follows by summing over all cubes  $Q^0$ .

Proof of Lemma 1 : Let  $\alpha_0 = \frac{1}{|Q^0|} \int_{Q^0} |f(x) - f_{Q^0}| dx$ . For any  $\alpha > \alpha_0$ ,  $K \gg 1$  we may apply the Calderón-Zygmund stopping process to obtain a decomposition of the sets  $\{x \in Q^0, f^\#(x) > \alpha\}$  and  $\{x \in Q^0; f^\#(x) > K\alpha\}$  into cubes with the following properties.

$$\{f^\# > \alpha\} = \bigcup_k Q_k \quad \{f^\# > K\alpha\} = \bigcup_{k,v} Q_{kv}$$

$$(i) \quad Q_{kv} \subseteq Q_k$$

$$(ii) \quad \frac{1}{|Q_k|} \int_{Q_k} |f(x) - f_{Q_k}| dx \sim \alpha$$

$$(iii) \quad \frac{1}{|Q_{kv}|} \int_{Q_{kv}} |f(x) - f_{Q_{kv}}| dx \sim K\alpha$$

Next observe that

$$(\nabla f)^* \geq \alpha (\text{diam } Q_k)^{-1} \text{ throughout } Q_k \quad (7)$$

$$|Q_k| \geq 2 \left| \bigcup_v Q_{kv} \right| \quad (8)$$

In fact (7) follows from

$$\frac{1}{|Q_k|} \int_{Q_k} |f(x) - f_{Q_k}| dx \leq (\text{diam } Q_k) \frac{1}{|Q_k|} \int_{Q_k} |\nabla f(x)| dx$$

while (8) is a consequence of

$$\begin{aligned} \alpha |Q_k| &\sim \int_{Q_k} |f(x) - f_{Q_k}| dx \geq \sum_v \int_{Q_{kv}} |f(x) - f_{Q_k}| dx \\ &\geq \frac{1}{2} \sum_v \int_{Q_{kv}} |f(x) - f_{Q_{kv}}| dx \sim \frac{1}{2} K\alpha \sum_v |Q_{kv}| \end{aligned}$$

and the fact that  $K$  is large. We may thus write

$$\begin{aligned} \int_{Q_k \setminus \bigcup_v Q_{kv}} |(\nabla f)^*(x)|^2 dx &\geq \alpha^2 (\text{diam } Q_k)^{-2} |Q_k \setminus \bigcup_v Q_{kv}| \\ &\geq \frac{1}{2} \alpha^2 (\text{diam } Q_k)^{-2} |Q_k| \\ &\geq \alpha^2 \mu(Q_k) / (2C). \end{aligned}$$

which implies in turn

$$\int_{\{\alpha < f^\# \leq K\alpha\}} |(\nabla f)^*(x)|^2 dx \geq \alpha^2 \mu(f^\# > \alpha) / C .$$

Let now  $\alpha$  range over  $2K^m \alpha_0$ ,  $m = 0, 1, 2, \dots$ , and sum over  $m$  to obtain

$$\int_{Q_0} |(\nabla f)^*(x)|^2 dx \geq \left( \int_{\{f^\# > 2\alpha_0\}} |f^\#(x)|^2 d\mu(x) \right) / C \quad (9)$$

On the other hand note that

$$\begin{aligned} \int_{\{f^\# \leq 2\alpha_0\}} |f^\#|^2 d\mu &\leq 4\alpha_0^2 \mu(Q_0) \\ &\leq 4[(\text{diam } Q_0)^2 \frac{1}{|Q_0|} \int_{Q_0} |\nabla f(x)|^2 dx] [C|Q_0| (\text{diam } Q_0)^{-2}] \\ &\leq 4C \int_{Q_0} |\nabla f(x)|^2 dx \end{aligned} \quad (10)$$

Combining (9), (10) and applying the maximal theorem establishes (3).

Proof of Lemma 2 : We may assume that  $f|_{Q_0} = 0$ , so it suffices to show that

$$\int_{Q_0} |f^*(x)|^2 d\mu(x) \leq C \int_{Q_0} |f^\#(x)|^2 d\mu(x) \quad (11)$$

Let  $\alpha_0 = \frac{1}{|Q_0|} \int_{Q_0} |f(x)| dx$ , and let  $K$  be a large positive number. A Calderon-

Zygmund decomposition yields for each  $\alpha > \alpha_0$  :

$$\{f^* > \alpha\} = \bigcup_k Q_k \quad \{f^* > K\alpha\} = \bigcup_{k,v} Q_{kv}$$

(i)  $Q_{kv} \subseteq Q_k$

(ii)  $\frac{1}{|Q_k|} \int_{Q_k} |f(x)| dx \sim \alpha$

(iii)  $\frac{1}{|Q_{kv}|} \int_{Q_{kv}} |f(x)| dx \sim K\alpha$

Obviously  $f^\# > \delta\alpha$  on cubes  $Q_k$  satisfying  $\frac{1}{|Q_k|} \int_{Q_k} |f(x) - f_{Q_k}| dx > \delta\alpha$ ; on the remaining cubes, for any  $\varepsilon > 0$  a choice of  $\delta$  small enough will guarantee that

$$\mu\left(\bigcup_{\nu} Q_{k\nu}\right) \leq \varepsilon \mu(Q_k) \quad (12)$$

Indeed we have in that case

$$\begin{aligned} \delta\alpha|Q_k| &\geq \int_{Q_k} |f(x) - f_{Q_k}| dx \geq \sum_{\nu} \int_{Q_{k\nu}} |f(x) - f_{Q_k}| dx \\ &\geq \frac{1}{2} \sum_{\nu} \int_{Q_{k\nu}} |f(x) - f_{Q_{k\nu}}| dx \sim \frac{1}{2} K\alpha \sum_{\nu} |Q_{k\nu}| \end{aligned}$$

and (12) follows from (A<sub>∞</sub>). Thus

$$\mu(f^* > K\alpha) \leq \varepsilon \mu(f^* > \alpha) + \mu(f^\# > \delta\alpha) \quad \text{for } \alpha > \alpha_0.$$

For  $\alpha \leq \alpha_0$  this estimate is trivial, since  $f^\# > \alpha_0$  throughout  $Q^0$ . To get (11), integrate with respect to  $\alpha d\alpha$  and choose  $\varepsilon = K^{-2}/2$ . Q.E.D.

Proof of Lemma 3 : We need only prove a reverse Hölder inequality for some  $r > 1$

$$\left(\frac{1}{|Q|} \int_Q (v^+(x))^r dx\right)^{1/r} \leq c \frac{1}{|Q|} \int_Q v^+(x) dx \quad \text{all } Q \subseteq Q^0 \quad (13)$$

since then

$$\mu(E) = \int_Q \chi_E v^+(x) dx \leq |E|^{1/r'} \left(\int_Q (v^+(x))^r dx\right)^{1/r} \leq \left(\frac{|E|}{|Q|}\right)^{1/r'} \mu(Q)$$

for  $E \subset Q$ . Now introduce

$$\begin{aligned} K(Q) &= \sup_{Q' \supseteq Q} \left[\frac{1}{|Q'|} \int_{Q'} |v(x)|^p dx\right]^{1/p} \\ v_Q(x) &= \sup_{x \in Q' \subseteq Q} \left[\frac{1}{|Q'|} \int_{Q'} |v(y)|^p dy\right]^{1/p} \end{aligned}$$

and observe that

$$v^+(x) = \max(K(Q), v_Q(x))$$

$$K(Q) \leq \min\{C(\text{diam } Q)^{-2}, \frac{1}{|Q|} \int_Q v^+(x) dx\}$$

Thus (13) and condition (2) for  $d\mu^+$  are simple consequences of the following inequality

$$\frac{1}{|Q|} \int_Q (v_Q(x))^r dx \leq c K(Q)^r \quad (14)$$

In view of the maximal theorem we may write

$$\begin{aligned} \frac{|\{x \in Q; v_Q(x) > \alpha\}|}{|Q|} &\leq \frac{c}{\alpha^p} \frac{1}{|Q|} \int_Q |v_Q(x)|^p dx \\ &\leq \frac{c}{\alpha^p} [K(Q)]^p \end{aligned}$$

Integrating with respect to  $\alpha$  for  $\alpha > K(Q)$ ,  $1 < r < p$  yields

$$\frac{1}{|Q|} \int_{v_Q(x) > K(Q)} (v_Q(x))^r dx \leq c [K(Q)]^r$$

The inequality (14) now follows since  $\frac{1}{|Q|} \int_{v_Q(x) \leq K(Q)} (v_Q(x))^r dx$  is also bounded by  $K(Q)^r$ . The proof of Lemma 3 and hence of the theorem is complete.

#### REFERENCES

- [1] R. Coifman, C. Fefferman : Weighted norm inequalities for maximal functions and singular integrals, *Studia Math* , 51 (1974), pp. 241-250.
- [2] C. Fefferman, E. M. Stein :  $H^p$  spaces in several variables, *Acta Math.* 129 (1972), pp. 137-193.

\*  
\*  
\*