

WENPENG ZHANG

On the mean values of Dedekind sums

Journal de Théorie des Nombres de Bordeaux, tome 8, n° 2 (1996),
p. 429-442

http://www.numdam.org/item?id=JTNB_1996__8_2_429_0

© Université Bordeaux 1, 1996, tous droits réservés.

L'accès aux archives de la revue « Journal de Théorie des Nombres de Bordeaux » (<http://jtnb.cedram.org/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

On the mean values of Dedekind Sums

par WENPENG ZHANG

RÉSUMÉ. On étudie dans ce papier le comportement asymptotique de valeurs moyennes de sommes de Dedekind. On donne en particulier une formule asymptotique améliorant un résultat antérieur.

ABSTRACT. In this paper we study the asymptotic behavior of the mean value of Dedekind sums, and give a sharper asymptotic formula.

1. Introduction

For a positive integer k and an arbitrary integer h , the Dedekind sum $S(h, k)$ is defined by

$$S(h, k) = \sum_{a=1}^k \left(\left(\frac{a}{k} \right) \right) \left(\left(\frac{ah}{k} \right) \right)$$

where

$$\left(\left(x \right) \right) = \begin{cases} x - [x] - \frac{1}{2} & \text{if } x \text{ is not an integer;} \\ 0 & \text{if } x \text{ is an integer.} \end{cases}$$

The various properties of $S(h, k)$ were investigated by many authors. Perhaps the most famous property of Dedekind sums is the reciprocity formula (see [3], [5], [6]):

$$(1) \quad S(h, k) + S(k, h) = \frac{h^2 + k^2 + 1}{12hk} - \frac{1}{4}$$

for all $(h, k) = 1$, $h > 0$, $k > 0$. A three term version of (1) was discovered by Rademacher [7]. Walum [8] has shown that for prime $p \geq 3$,

$$(2) \quad \sum_{\substack{\chi \pmod{p} \\ \chi(-1)=-1}} |L(1, \chi)|^4 = \frac{\pi^4(p-1)}{p^2} \sum_{h=1}^p |S(h, p)|^2$$

and

$$(3) \quad \sum_{\substack{\chi \pmod p \\ \chi(-1)=-1}} |L(1, \chi)|^2 = \frac{\pi^2(p-1)^2(p-2)}{12p^2}$$

Recently, J.B. Conrey et al. [4] studied the mean value distribution of $S(h, k)$, and proved that

$$(4) \quad \sum'_{h=1}^k |S(h, k)|^{2m} = f_m(k) \left(\frac{k}{12}\right)^{2m} + O\left(\left(k^{\frac{9}{5}} + k^{2m-1+\frac{1}{m+1}}\right) \log^3 k\right),$$

where \sum'_h denotes the summation over all h such that $(k, h) = 1$, and

$$\sum_{n=1}^{\infty} \frac{f_m(n)}{n^s} = 2 \frac{\zeta^2(2m)}{\zeta(4m)} \cdot \frac{\zeta(s+4m-1)}{\zeta^2(s+2m)} \zeta(s).$$

In the spirit of [4] and [8], we use the estimates for character sum and the properties of Dirichlet’s L -functions to give a sharper asymptotic formula for (4), where $m = 1$ and $k = p^n$, a power of a prime. That is, we shall prove the following two theorems.

THEOREM 1. *Let p be a prime and n be a positive integer. Then for $k = p^n$, we have the asymptotic formula*

$$\sum'_{h=1}^k |S(h, k)|^2 = \frac{5}{144} k^2 \frac{(p^2 - 1)^2}{p(p^3 - 1)} + O\left(k \exp\left(\frac{3 \ln k}{\ln \ln k}\right)\right),$$

where the constant implied by the O -symbol does not depend on any parameter, and $\exp(y) = e^y$.

THEOREM 2. *Let n be an integer ≥ 1 , p be a prime, and $k = p^n$. Then we have*

$$\sum'_{h=1}^k \frac{S(h, k)}{h} = \frac{\pi^2}{72} k \left(1 - \frac{1}{p^2}\right) + O\left(\sqrt{k}\right).$$

COROLLARY. *Let p be a prime. Then we have the asymptotic formulas*

$$\begin{aligned}
 1) \quad & \sum_{h=1}^p |S(h, p)|^2 = \frac{5}{144} p^2 + O\left(p \exp\left(\frac{3 \ln p}{\ln \ln p}\right)\right). \\
 2) \quad & \sum_{h=1}^p \frac{S(h, p)}{h} = \frac{\pi^2}{72} p + O(\sqrt{p}).
 \end{aligned}$$

2. Some Lemmas

To prove the theorems, we need the following lemmas.

LEMMA 1. *Let q be an integer ≥ 3 , and let χ be any Dirichlet character modulo q with $\chi(-1) = -1$. Then we have*

$$L(1, \chi) = \frac{\pi}{2q} \sum_{r=1}^q \chi(r) \cot \frac{\pi r}{q}$$

where $L(s, \chi)$ denotes Dirichlet L -function corresponding to χ modulo q .

Proof. Let $N \geq 3$ be an integer. Then we have

$$\begin{aligned}
 \sum_{1 \leq n \leq Nq} \frac{\chi(n)}{n} &= \sum_{a=1}^q \sum_{\ell=0}^{N-1} \frac{\chi(\ell q + a)}{\ell q + a} \\
 (5) \quad &= \frac{1}{q} \sum_{a=1}^q \chi(a) \sum_{\ell=0}^{N-1} \frac{1}{\ell + a/q}.
 \end{aligned}$$

Note that since $\chi(-1) = -1$, we also have

$$\begin{aligned}
 \sum_{1 \leq n \leq Nq} \frac{\chi(n)}{n} &= \frac{1}{q} \sum_{a=1}^q \chi(q - a) \sum_{\ell=0}^{N-1} \frac{1}{\ell + 1 - a/q} \\
 (6) \quad &= \frac{-1}{q} \sum_{a=1}^q \chi(a) \sum_{\ell=1}^N \frac{1}{\ell - a/q}.
 \end{aligned}$$

Thus from (5) and (6) we get

$$\begin{aligned}
 2 \sum_{1 \leq n \leq Nq} \frac{\chi(n)}{n} &= \frac{1}{q} \sum_{a=1}^q \chi(a) \left(\sum_{\ell=0}^{N-1} \frac{1}{\ell + a/q} - \sum_{\ell=1}^N \frac{1}{\ell - a/q} \right) \\
 &= \frac{1}{q} \sum_{a=1}^q \chi(a) \left(\frac{1}{a/q} + \sum_{\ell=1}^{N-1} \left(\frac{1}{\ell + a/q} - \frac{1}{\ell - a/q} \right) - \frac{1}{N - a/q} \right) \\
 &= \frac{1}{q} \sum_{a=1}^q \chi(a) \left(\frac{1}{a/q} + \sum_{\ell=1}^{N-1} \left(\frac{1}{\ell + a/q} - \frac{1}{\ell - a/q} \right) \right) + O\left(\frac{1}{N-1}\right).
 \end{aligned}$$

Taking $N \rightarrow \infty$ in the above, and noting that

$$\lim_{n \rightarrow \infty} \sum_{1 \leq n \leq Nq} \frac{\chi(n)}{n} = L(1, \chi)$$

and

$$\frac{a}{q} + \sum_{\ell=1}^{\infty} \left(\frac{1}{\ell + a/q} - \frac{1}{\ell + a/q} \right) = \pi \cot \frac{\pi a}{q},$$

the lemma follows.

LEMMA 2. *Let k be an integer ≥ 3 and $(h, k) = 1$. Then*

$$S(h, k) = \frac{1}{\pi^2 k} \sum_{d|k} \frac{d^2}{\phi(d)} \sum_{\substack{\chi \pmod{d} \\ \chi(-1)=-1}} \chi(h) |L(1, \chi)|^2,$$

where $\phi(k)$ is the Euler function.

Proof. Note that $\sum_{a=1}^k \chi(a) \cot \frac{\pi a}{k} = 0$ if $\chi(-1) = 1$. For $(c, k) = 1$, from Lemma 1 and the orthogonality of characters modulo k we have the identity

$$\begin{aligned} \sum_{\substack{\chi \pmod{k} \\ \chi(-1)=-1}} \bar{\chi}(c) L(1, \chi) &= \sum_{\substack{\chi \pmod{k} \\ \chi(-1)=-1}} \bar{\chi}(c) \left(\frac{\pi}{2k} \sum_{b=1}^k \chi(b) \cot \frac{\pi b}{k} \right) \\ &= \sum_{\chi \pmod{k}} \bar{\chi}(c) \left(\frac{\pi}{2k} \sum_{b=1}^k \chi(b) \cot \frac{\pi b}{k} \right) = \frac{\pi \phi(k)}{2k} \cot \frac{\pi c}{k}. \end{aligned}$$

Hence

$$(7) \quad \cot \frac{\pi c}{k} = \frac{2k}{\pi \phi(k)} \sum_{\substack{\chi \pmod{k} \\ \chi(-1)=-1}} \bar{\chi}(c) L(1, \chi)$$

From [2] we know that

$$(8) \quad \left(\left(\frac{uh}{k} \right) \right) = -\frac{1}{2k} \sum_{a=1}^{k-1} \sin \frac{2\pi ahu}{k} \cot \frac{\pi a}{k}.$$

Thus

$$\begin{aligned}
 (9) \quad \left(\left(\frac{uh}{k} \right) \right) &= -\frac{1}{2k} \sum_{a=1}^k \sin \frac{2\pi ahu}{k} \cot \frac{\pi a}{k} \\
 &= -\frac{1}{2k} \sum_{d|k} \sum_{a=1}^{k/d} \left(\frac{2k/d}{\pi\phi(k/d)} \sum_{\substack{\chi \pmod{k/d} \\ \chi(-1)=-1}} \bar{\chi}(a)L(1, \chi) \right) \frac{e\left(\frac{ahu}{k/d}\right) - e\left(\frac{-ahu}{k/d}\right)}{2i} \\
 &= \frac{i}{2\pi} \sum_{d|k} \frac{1}{d\phi(k/d)} \sum_{\substack{\chi \pmod{k/d} \\ \chi(-1)=-1}} L(1, \chi) \sum_{a=1}^{k/d} \bar{\chi}(a) \left(e\left(\frac{ahu}{k/d}\right) - e\left(\frac{-ahu}{k/d}\right) \right) \\
 &= \frac{i}{2\pi} \sum_{d|k} \frac{1}{d\phi(k/d)} \sum_{\substack{\chi \pmod{k/d} \\ \chi(-1)=-1}} \chi(h)L(1, \chi) \sum_{a=1}^{k/d} \bar{\chi}(a) \left(e\left(\frac{au}{k/d}\right) - e\left(\frac{-au}{k/d}\right) \right) \\
 &= \frac{i}{\pi} \sum_{d|k} \frac{1}{d\phi(k/d)} \sum_{\substack{\chi \pmod{k/d} \\ \chi(-1)=-1}} \chi(h)L(1, \chi) \sum_{a=1}^{k/d} \bar{\chi}(a) e\left(\frac{au}{k/d}\right),
 \end{aligned}$$

where $e(y) = e^{2\pi iy}$, $i^2 = -1$, $(h, k) = 1$.

Similarly, we also have

$$(10) \quad \left(\left(\frac{u}{k} \right) \right) = \frac{i}{\pi} \sum_{d|k} \frac{1}{d\phi(k/d)} \sum_{\substack{\chi \pmod{k/d} \\ \chi(-1)=-1}} L(1, \chi) \sum_{a=1}^{k/d} \bar{\chi}(a) e\left(\frac{au}{k/d}\right).$$

Note that from the definitions of $S(h, k)$, and from (9) and (10) we get

$$\begin{aligned}
 S(h, k) &= \sum_{c=1}^k \left(\left(\frac{c}{k} \right) \right) \left(\left(\frac{ch}{k} \right) \right) \\
 &= \frac{-1}{\pi^2} \sum_{u|k} \sum_{v|k} \frac{1}{u\phi(k/u)} \frac{1}{v\phi(k/v)} \\
 &\quad \times \sum_{\substack{\chi_1 \pmod{k/u} \\ \chi_1(-1)=-1}} \sum_{\substack{\chi_2 \pmod{k/v} \\ \chi_2(-1)=-1}} \chi_1(h)L(1, \chi_1)L(1, \chi_2) \\
 &\quad \times \sum_{c=1}^k \left(\sum_{a=1}^{k/u} \bar{\chi}_1(a) e\left(\frac{ac}{k/u}\right) \right) \left(\sum_{b=1}^{k/v} \bar{\chi}_2(b) e\left(\frac{bc}{k/v}\right) \right) \\
 &= \frac{k}{\pi^2} \sum_{d|k} \frac{1}{d^2\phi(k/d)} \sum_{\substack{\chi \pmod{k/d} \\ \chi(-1)=-1}} \chi(h) |L(1, \chi)|^2 \\
 &= \frac{1}{\pi^2 k} \sum_{d|k} \frac{d^2}{\phi(d)} \sum_{\substack{\chi \pmod{d} \\ \chi(-1)=-1}} \chi(h) |L(1, \chi)|^2,
 \end{aligned}$$

where we have used the identities

$$\begin{aligned}
 &\sum_{a=1}^{k/u} \sum_{b=1}^{k/v} \bar{\chi}_1(a) \bar{\chi}_2(b) \sum_{c=1}^k e\left(\frac{(au + bv)c}{k}\right) \\
 &= \begin{cases} k \sum_{a=1}^{k/u} \bar{\chi}_1(a) \bar{\chi}_2(k/u - a), & \text{if } u = v; \\ 0, & \text{if } u \neq v. \end{cases} \\
 &= \begin{cases} -k\phi(k/u), & \text{if } \bar{\chi}_1 = \chi_2 \text{ and } u = v; \\ 0, & \text{if } \bar{\chi}_1 \neq \chi_2 \text{ or } u \neq v. \end{cases}
 \end{aligned}$$

This completes the proof of Lemma 2.

LEMMA 3. *Let q be an integer ≥ 3 . Then we have the asymptotic formula*

$$\sum_{\substack{\chi \pmod{q} \\ \chi(-1)=-1}} |L(1, \chi)|^4 = \frac{5\pi^4}{144} \phi(q) \prod_{p|q} \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} + O\left(\frac{\phi(q)}{q} \exp\left(\frac{3 \ln q}{\ln \ln q}\right)\right).$$

Proof. For any parameter $N \geq q$, applying Abel's identity we have

$$\begin{aligned}
 L^2(1, \chi) &= \sum_{n=1}^{\infty} \frac{\chi(n)d(n)}{n} \\
 (11) \qquad &= \sum_{1 \leq n \leq N} \frac{\chi(n)d(n)}{n} + \int_N^{\infty} \frac{A(y, \chi)}{y^2} dy,
 \end{aligned}$$

where $d(n)$ is the divisor function. We have $A(y, \chi) = \sum_{N < n \leq y} \chi(n)d(n)$.

Note that the identities

$$\begin{aligned}
 A(y, \chi) &= 2 \sum_{n \leq \sqrt{y}} \chi(n) \sum_{m \leq y/n} \chi(m) - 2 \sum_{n \leq \sqrt{N}} \chi(n) \sum_{m \leq N/n} \chi(m) - \\
 &\quad - \left(\sum_{n \leq \sqrt{y}} \chi(n) \right)^2 + \left(\sum_{n \leq \sqrt{N}} \chi(n) \right)^2.
 \end{aligned}$$

Applying the Cauchy inequality and estimates for character sums we have

$$\begin{aligned}
 \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} |A(y, \chi)|^2 &\ll \sqrt{y} \sum_{n \leq \sqrt{y}} \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} \left| \sum_{m \leq y/n} \chi(m) \right|^2 + \\
 &\quad + \sqrt{N} \sum_{n \leq \sqrt{N}} \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} \left| \sum_{m \leq N/n} \chi(m) \right|^2 + \\
 &\quad + y \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} \left| \sum_{n \leq \sqrt{y}} \chi(n) \right|^2 + N \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} \left| \sum_{n \leq \sqrt{N}} \chi(n) \right|^2 \\
 (12) \qquad &\ll y\phi^2(q).
 \end{aligned}$$

Thus from (12) we get

$$\begin{aligned}
 & \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} \left| \int_N^\infty \frac{A(y, \chi)}{y^2} dy \right|^2 \\
 &= \int_N^\infty \int_N^\infty \frac{1}{y^2 z^2} \left(\sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} |A(y, \chi)| \cdot |A(z, \chi)| \right) dy dz \\
 &\ll \left(\int_N^\infty \frac{1}{y^2} \left(\sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} |A(y, \chi)|^2 \right)^{\frac{1}{2}} dy \right)^2 \\
 (13) \quad &\ll \left(\int_N^\infty \frac{1}{y^{3/2}} \phi(q) dy \right)^2 \ll \frac{\phi^2(q)}{N}.
 \end{aligned}$$

Note that for $(ab, q) = 1$, from the orthogonality relation for character sums modulo q we have

$$\sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} \chi(a)\overline{\chi}(b) = \begin{cases} \frac{1}{2}\phi(q), & \text{if } a \equiv b(q); \\ -\frac{1}{2}\phi(q), & \text{if } a \equiv -b(q); \\ 0, & \text{otherwise.} \end{cases}$$

Hence

$$(14) \quad \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} \left| \sum_{1 \leq n \leq N} \frac{\chi(n)d(n)}{n} \right|^2 =$$

$$\begin{aligned}
 &= \frac{1}{2}\phi(q) \sum_{\substack{1 \leq a, b \leq N \\ (ab, q) = 1 \\ a \equiv b(q)}} \frac{d(a)d(b)}{ab} - \frac{1}{2}\phi(q) \sum_{\substack{1 \leq a, b \leq N \\ (ab, q) = 1 \\ a \equiv -b(q)}} \frac{d(a)d(b)}{ab} \\
 &= \frac{1}{2}\phi(q) \sum_{\substack{1 \leq a \leq N \\ (a, q) = 1}} \frac{d^2(a)}{a^2} + O\left(\phi(q) \sum_{b=1}^N \sum_{\ell=1}^{[N/q]} \frac{d(b)d(\ell q + b)}{(\ell q + b)b}\right) + \\
 &\quad + O\left(\phi(q) \sum_{a=1}^{q-1} \frac{d(a)d(q-a)}{a(q-a)}\right) + \\
 &\quad + O\left(\phi(q) \sum_{1 \leq a \leq N} \sum_{(1+a/q) \leq \ell \leq N/q} \frac{d(a)d(\ell q - a)}{a(\ell q - a)}\right) \\
 &= \frac{1}{2}\phi(q) \sum_{\substack{n=1 \\ (n, q) = 1}}^{\infty} \frac{d^2(n)}{n^2} + O\left(\frac{\phi(q)}{q} \exp\left(\frac{\ln N}{\ln \ln N}\right)\right) \\
 &= \frac{1}{2}\phi(q) \frac{\zeta^4(2)}{\zeta(4)} \prod_{p|q} \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} + O\left(\frac{\phi(q)}{q} \exp\left(\frac{\ln N}{\ln \ln N}\right)\right) \\
 &= \frac{5}{144} \pi^4 \phi(q) \prod_{p|q} \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} + O\left(\frac{\phi(q)}{q} \exp\left(\frac{\ln N}{\ln \ln N}\right)\right),
 \end{aligned}$$

where we have used the estimate $d(n) \ll \exp\left(\frac{(1+\epsilon)\ln 2 \ln n}{\ln \ln n}\right)$, and $\zeta(s)$ is the Riemann zeta-function.

$$\begin{aligned}
 (15) \quad &\sum_{\substack{\chi \pmod q \\ \chi(-1) = -1}} \left(\sum_{1 \leq n \leq N} \frac{\chi(n)d(n)}{n} \right) \left(\int_N^\infty \frac{A(y, \bar{\chi})}{y^2} dy \right) \\
 &\ll (\ln N)^2 \int_N^\infty \frac{1}{y^2} \left(\sum_{\substack{\chi \pmod q \\ \chi(-1) = -1}} |A(y, \bar{\chi})| \right) dy \ll \phi^{\frac{3}{2}}(q) (\ln N)^2 N^{-\frac{1}{2}}.
 \end{aligned}$$

Taking the parameter $N = q^3$, from (11), (13), (14) and (15) we obtain

$$\sum_{\substack{\chi \pmod q \\ \chi(-1) = -1}} |L(1, \chi)|^4 = \frac{5}{144} \pi^4 \phi(q) \prod_{p|q} \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} + O\left(\frac{\phi(q)}{q} \exp\left(\frac{3 \ln q}{\ln \ln q}\right)\right).$$

This proves lemma 3.

3. Proof of the theorems

In this section, we shall complete the proof of the theorems. First we prove theorem 1. For $k = p^n$, applying Lemma 2 and Lemma 3 we have

$$\begin{aligned}
 \sum_{h=1}^k |S(h, k)|^2 &= \frac{1}{\pi^4 k^2} \sum_{h=1}^k \left| \sum_{d|k} \frac{d^2}{\phi(d)} \sum_{\substack{\chi \pmod d \\ \chi(-1)=-1}} \chi(h) |L(1, \chi)|^2 \right|^2 \\
 &= \frac{1}{\pi^4 k^2} \sum_{u|k} \sum_{v|k} \frac{u^2}{\phi(u)} \frac{v^2}{\phi(v)} \\
 &\quad \times \sum_{\substack{\chi_1 \pmod u \\ \chi_1(-1)=-1}} \sum_{\substack{\chi_2 \pmod v \\ \chi_2(-1)=-1}} |L(1, \chi_1)|^2 |L(1, \chi_2)|^2 \sum_{h=1}^k \bar{\chi}_1(h) \chi_2(h) \\
 &= \frac{\phi(k)}{\pi^4 k^2} \left[2 \sum_{u|k} \sum_{v|u} \frac{u^2}{\phi(u)} \frac{v^2}{\phi(v)} \sum_{\substack{\chi \pmod v \\ \chi(-1)=-1}} |L(1, \chi)|^4 \right. \\
 &\quad \left. - \sum_{u|k} \frac{u^4}{\phi^2(u)} \sum_{\substack{\chi \pmod u \\ \chi(-1)=-1}} |L(1, \chi)|^4 \right] \\
 &= \frac{\phi(k)}{\pi^4 k^2} \sum_{u|k} \sum_{v|u} \frac{u^2}{\phi(u)} \frac{v^2}{\phi(v)} \left[\frac{10}{144} \pi^4 \phi(v) \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} + O\left(\frac{\phi(v)}{v} \exp\left(\frac{3 \ln v}{\ln \ln v}\right)\right) \right] \\
 &\quad - \frac{\phi(k)}{\pi^4 k^2} \sum_{u|k} \frac{u^4}{\phi^2(u)} \left[\frac{5}{144} \pi^4 \phi(u) \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} + O\left(\frac{\phi(u)}{u} \exp\left(\frac{3 \ln u}{\ln \ln u}\right)\right) \right] \\
 &= \frac{5}{144} \frac{\phi(k)}{k^2} \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} \left[2 \sum_{u|k} \sum_{v|u} \frac{u^2 v^2}{\phi(u)} - \sum_{u|k} \frac{u^4}{\phi(u)} \right] + O\left(k \exp\left(\frac{3 \ln k}{\ln \ln k}\right)\right) \\
 &= \frac{5}{144} k^2 \frac{(p^2 - 1)^2}{p(p^3 - 1)} + O\left(k \exp\left(\frac{3 \ln k}{\ln \ln k}\right)\right).
 \end{aligned}$$

This completes the proof of Theorem 1.

Proof of Theorem 2. For real $y \geq k = p^n$, let $U(y, \chi) = \sum_{k < n \leq y} \chi(n)$. Note

that

$$\begin{aligned}
 \sum_{h=1}^k \frac{S(h, k)}{h} &= \frac{1}{\pi^2 k} \sum_{d|k} \frac{d^2}{\phi(d)} \sum_{\substack{\chi \pmod d \\ \chi(-1)=-1}} \left(\sum_{h=1}^k \frac{\chi(h)}{h} \right) |L(1, \chi)|^2 \\
 (16) \qquad &= \frac{1}{\pi^2 k} \sum_{d|k} \frac{d^2}{\phi(d)} \sum_{\substack{\chi \pmod d \\ \chi(-1)=-1}} \left(L^2(1, \chi)L(1, \bar{\chi}) - |L(1, \chi)|^2 \int_k^\infty \frac{U(y, \chi)}{y^2} dy \right).
 \end{aligned}$$

Using the method of proving Lemma 3 we get

$$\begin{aligned}
 (17) \qquad \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} L^2(1, \chi)L(1, \bar{\chi}) &= \frac{\pi^4}{72} \phi(q) \prod_{p|q} \left(1 - \frac{1}{p^2} \right)^2 + O \left(\exp \left(\frac{3 \ln q}{\ln \ln q} \right) \right).
 \end{aligned}$$

Note that the estimate

$$\sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} |U(y, \chi)|^2 \ll \phi^2(q)$$

holds. Now, using the Cauchy inequality and Lemma 3 we get

$$(18) \qquad \sum_{\substack{\chi \pmod d \\ \chi(-1)=-1}} |L(1, \chi)|^2 \int_k^\infty \frac{U(y, \chi)}{y^2} dy \ll \frac{\phi^{\frac{3}{2}}(d)}{k}.$$

From (16), (17) and (18) we obtain

$$\begin{aligned}
 \sum_{h=1}^k \frac{S(h, k)}{h} &= \frac{1}{\pi^2 k} \sum_{d|k} \frac{d^2}{\phi(d)} \left[\frac{\pi^4}{72} \phi(d) \left(1 - \frac{1}{p^2} \right)^2 + O \left(\frac{\phi^{\frac{3}{2}}(d)}{d} \right) \right] \\
 &= \frac{\pi^2}{72} k \left(1 - \frac{1}{p^2} \right) + O \left(\sqrt{\phi(k)} \right).
 \end{aligned}$$

REMARK. In [9] and [10], we proved a generalized version of (3). i.e.

THEOREM A. *Let q be an integer ≥ 3 . Then we have*

$$\sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} |L(1, \chi)|^2 = \frac{\pi^2 \phi^2(q)}{12 q^2} \left[q \prod_{p|q} \left(1 + \frac{1}{p} \right) - 3 \right],$$

where the summation is over all odd Dirichlet characters modulo q .

THEOREM B. *Let q be a square-full number (i.e. $q \geq 4$ and a prime $p|q$ implies $p^2|q$). Then we have*

$$\sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}}^* |L(1, \chi)|^2 = \frac{\pi^2 \phi^3(q)}{12 q^2} \prod_{p|q} \left(1 + \frac{1}{p} \right),$$

where the summation is over all odd primitive Dirichlet characters modulo q .

Using Lemma 2 we can give a simple proof of Theorem A and Theorem B. In fact, note that for $k \geq 2$, from the definition of $S(h, k)$ we have

$$S(1, k) = \sum_{a=1}^{k-1} \left(\frac{a}{k} - \frac{1}{2} \right)^2 = \frac{(k-1)(k-2)}{12k}$$

From this identity, Lemma 2 with $h = 1$ and the Möbius inversion formula we get

$$\begin{aligned} \frac{q^2}{\phi(q)} \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}} |L(1, \chi)|^2 &= \pi^2 \sum_{d|q} \mu(d) \frac{q}{d} S\left(1, \frac{q}{d}\right) \\ &= \pi^2 \sum_{d|q} \mu(d) \frac{\left(\frac{q}{d} - 1\right) \left(\frac{q}{d} - 2\right)}{12} = \frac{\pi^2}{12} \sum_{d|q} \mu(d) \left(\frac{q^2}{d^2} - 3\frac{q}{d} + 2 \right) \\ &= \frac{\pi^2}{12} \left[q^2 \sum_{d|q} \frac{\mu(d)}{d^2} - 3q \sum_{d|q} \frac{\mu(d)}{d} + 2 \sum_{d|q} \mu(d) \right] \\ &= \frac{\pi^2}{12} \phi(q) \left[q \prod_{p|q} \left(1 + \frac{1}{p} \right) - 3 \right]. \end{aligned}$$

This proves Theorem A.

If q be a square-full number, then using the Möbius inversion formula and Theorem A we get

$$\begin{aligned} \sum_{\substack{\chi \pmod q \\ \chi(-1)=-1}}^* |L(1, \chi)|^2 &= \sum_{d|q} \mu(d) \left(\sum_{\substack{\chi \pmod{q/d} \\ \chi(-1)=-1}} |L(1, \chi)|^2 \right) \\ &= \sum_{d|q} \mu(d) \left\{ \frac{\pi^2}{12} \frac{\phi^2(q/d)}{(q/d)^2} \left[\frac{q}{d} \prod_{p|q/d} \left(1 + \frac{1}{p} \right) - 3 \right] \right\} \\ &= \frac{\pi^2}{12} \frac{\phi^3(q)}{q^2} \prod_{p|q} \left(1 + \frac{1}{p} \right). \end{aligned}$$

This completes the proof of Theorem B.

ACKNOWLEDGEMENTS

The author expresses his gratitude to the referee for his helpful and detailed comments.

REFERENCES

1. Tom M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag, New York, 1976.
2. ———, *Modular Functions and Dirichlet Series in Number Theory*, Springer-Verlag, New York, 1976.
3. L. Carlitz, *The reciprocity theorem for Dedekind sums*, Pacific J. Math. **3** (1953), 523-527.
4. J. B. Conrey, Eric Fransen, Robert Klein and Clayton Scott, *Mean values of Dedekind sums*, J. Number Theory **56** (1996), 214-226.
5. L. J. Mordell, *The reciprocity formula for Dedekind sums*, Amer. J. Math. **73** (1951), 593-598.
6. H. Rademacher, *On the transformation of $\log \eta(\tau)$* , J. Indian Math. Soc. **19** (1955), 25-30.
7. H. Rademacher and E. Grosswald, *Dedekind Sums*, *The Carus Mathematical Monographs*, Math. Assoc. Amer., Washington D.C., 1972.
8. H. Walum, *An exact formula for an average of L -series*, Illinois J. Math. **26** (1982), 1-3.
9. Zhang Wenpeng, *Lecture Notes in Contemporary Mathematics*, Science Press, Beijing, China, 1989, pp. 173-179.

10. Zhang Wenpeng, *A note on a class mean square values of L-functions*, Journal of Northwest University **20** (1990), 9-12.
11. D. E. Knuth, *Notes on generalized Dedekind sums*, Acta Arith. **33** (1977), 297-325.

Wenpeng ZHANG
Department of Mathematics
Northwest University
Xi'an, Shaanxi
P. R. China.