

STEFAN MITITELU

**Extremum conditions for nonlinear
functions on convex sets**

Revue française d'automatique, informatique, recherche opérationnelle. Recherche opérationnelle, tome 8, n° V2 (1974), p. 113-118

http://www.numdam.org/item?id=RO_1974__8_2_113_0

© AFCET, 1974, tous droits réservés.

L'accès aux archives de la revue « Revue française d'automatique, informatique, recherche opérationnelle. Recherche opérationnelle » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

EXTREMUM CONDITIONS FOR NONLINEAR FUNCTIONS ON CONVEX SETS

by Stefan MITITELU (1)

Abstract. — *The paper establishes necessary and sufficient extremum conditions for a function f that is nonlinear, differentiable on a closed convex set generated by quasi-convex and quasi-concave differentiable functions. Conditions of the same type are also established for the case when nonlinear function f is quasi-convex.*

* * *

Let C be a convex set in the real Euclidean space R^n .

Definition 1. Any function $f : C \rightarrow R$ is called quasi-convex in C if for any points $x^1, x^2 \in C$ and any number $\lambda \in [0, 1]$ we have

$$(1) \quad f(\lambda x^1 + (1 - \lambda)x^2) \leq \max \{f(x^1), f(x^2)\}.$$

If for any points $x^1, x^2 \in C$, $x^1 \neq x^2$, and any number $\lambda \in (0, 1)$ relation (1) exists with the sign of strict inequality, then function f is called strictly quasi-convex on the set C .

Any strictly quasi-convex function is quasi-convex [7].

Definition 2. The function $f : X \subseteq R^n \rightarrow R$ is called positive definite in the point $x^0 \in X$ if its hessian matrix in x^0 ,

$$H_f(x^0) = \left(\frac{\partial^2 f}{\partial x_i^0 \partial x_j^0} \right)_{1 \leq i, j \leq n}$$

(1) Institute of Civil Engineering, Department of Mathematics, Bucarest.

is positive definite, i.e. if the following conditions are satisfied :

$$\frac{\partial^2 f}{\partial x_1^{02}} > 0, \left| \begin{array}{cc} \frac{\partial^2 f}{\partial x_1^{02}} & \frac{\partial^2 f}{\partial x_1^0 \partial x_2^0} \\ \frac{\partial^2 f}{\partial x_2^0 \partial x_1^0} & \frac{\partial^2 f}{\partial x_2^{02}} \end{array} \right| > 0, \dots, \left| \begin{array}{ccc} \frac{\partial^2 f}{\partial x_1^{02}} & \frac{\partial^2 f}{\partial x_1^0 \partial x_2^0} & \frac{\partial^2 f}{\partial x_1^0 \partial x_n^0} \\ \frac{\partial^2 f}{\partial x_2^0 \partial x_1^0} & \frac{\partial^2 f}{\partial x_2^{02}} & \frac{\partial^2 f}{\partial x_2^0 \partial x_n^0} \\ \dots & \dots & \dots \\ \frac{\partial^2 f}{\partial x_n^0 \partial x_1^0} & \frac{\partial^2 f}{\partial x_n^0 \partial x_2^0} & \frac{\partial^2 f}{\partial x_n^{02}} \end{array} \right| > 0.$$

Theorem 1. Let be function f quasi-convex and differentiable on the convex set $C \subseteq R^n$ and a point $x^0 \in C$ where $\nabla f(x^0) = 0$. If function f is positive definite in x^0 , then x^0 is the global minimum of the function f on C .

Proof. Since f is positively defined in $x^0 \in C$ it results that x^0 is a point of strict local minimum of the function f in C . However, for a quasi-convex function any strict local minimum is the global minimum of the function [2], [8].

Theorem 2. Let be functions $f, g = (g_1, \dots, g_m)$ and $h = (h_1, \dots, h_p)$ differentiable on the convex set $K \subseteq R^n$, the convex set $C = \{x \in K/g(x) \leq 0, h(x) = 0\}$, a point $x^0 \in C$ and the set $J^0 = \{i/g_i(x^0) = 0\}$.

We assume that function $g_i (i \in J^0)$ are quasi-convex on K , while h is quasi-convex and quasi-concave on K .

If $u_i^0 \geq 0 (i \in J^0)$ and $v^0 = (v_1^0, \dots, v_p^0) \in R^p$ exist, such that

$$(2) \quad (x - x^0)' \nabla_x L(x^0, u^0, v_0) \geq 0, (\forall) x \in C,$$

where

$$L(x, u, v) = \varepsilon f(x) + \sum_{i \in J^0} u_i^0 g_i(x) + v'h(x), \varepsilon = +1 [\varepsilon = -1]$$

and if

(a) for a $u_{i_0}^0 > 0 \quad (i_0 \in J^0)$ we have

$$(3) \quad (x - x^0)' \nabla g_{i_0}(x^0) < 0, (\forall) x \in C - \{x^0\},$$

or at least for a $v_{j_0}^0 > 0$ we have

$$(x - x^0)' \nabla h_{j_0}(x^0) < 0, \quad (\forall) x \in C - \{x^0\},$$

or

$$(x - x^0)' \nabla h_{j_0}(x^0) > 0, \quad (\forall) x \in C - \{x^0\},$$

then x^0 is a local minimum [maximum] point of the function f on C .

$$(4) \quad (b) \quad C \cap \{x \in R^n / (x - x^0)' \nabla f(x^0) = 0\} = \{x^0\},$$

then x^0 is a local minimum [maximum] of the function f on C .

(c) function f is quasi-convex continuously differentiable in C and $\nabla f(x^0) \neq 0$, then x^0 is the global minimum [maximum] of the function f on C .

Proof. (a) Since relation $h(x) = 0$ is equivalent to relations $h(x) \leq 0$ and $-h(x) \leq 0$ it is enough to prove the theorem for the case when we have only $g(x) \leq 0$. Then only rel. (3) is necessary for the proof.

Since functions g_i are quasi-convex on C , relations

$$g_i(x) \leq g_i(x^0) = 0 (i \in J^0), \quad (\forall)x \in C,$$

imply relations [7] :

$$(5) \quad (x - x^0)' \nabla g_i(x^0) \leq 0, \quad i \in J^0, \quad (\forall)x \in C,$$

where for $i = i_0$ there exists the sign of strict inequality, according to (3).

For $u_i^0 \geq 0, i \in J^0$, where $u_{i_0}^0 > 0$, from (5) it results

$$(6) \quad (x - x^0)' \sum_{i \in J^0} u_i^0 \nabla g_i(x^0) < 0, \quad (\forall)x \in C - \{x^0\}.$$

From rels. (2) and (6) it results

$$(x - x^0)' \nabla f(x^0) > 0, \quad (\forall)x \in C - \{x^0\},$$

and according to [3], P. 36, point x^0 is a local minimum [maximum] of the function f in the set C .

(b) From relations (2) and (5) where $u_i^0 \geq 0, i \in J^0$, it results

$$(7) \quad (x - x^0)' \nabla f(x^0) \geq 0, \quad (\forall)x \in C$$

and taking account of (4), from (7) it results

$$(x - x^0)' \nabla f(x^0) > 0, \quad (\forall)x \in C - \{x^0\}.$$

According to point (a) x^0 is a point of local minimum [maximum] of the function f in C .

(c) Relation (7) result, where f is quasi-convex on C . If $\nabla f(x^0) \neq 0$ from (7) :

$$(8) \quad x^0' \nabla f(x^0) = \min_{x \in C} x' \nabla f(x^0)$$

will result and according to proposition 5 in [5] rel. (8) is equivalent to

$$f(x^0) = \min_{x \in C} f(x)$$

Theorem 3. Let be functions f differentiable and $g = (g_1, \dots, g_m)$ differentiable and quasi-convex on the convex set $K \subseteq R^n$, point $x^0 \in C = \{x \in K/g(x) \leq 0, x \geq 0\}$ with nonempty interior and the set

$$J^0 = \{i/g_i(x^0) = 0\}.$$

(n) A necessary condition that point x^0 should be a local minimum [maximum] of the function f in C is that $u^0 \in R^m$ should exist, so that

$$(9) \quad \begin{aligned} x^0 \geq 0, y^0 &= \varepsilon \nabla f(x^0) + \sum_{i=1}^m u_i^0 \nabla g_i(x^0) \geq 0, x^{0'} y^0 = 0 \\ g(x^0) &\leq 0, \quad u^0 \geq 0, \quad u^{0'} g(x^0) = 0, \end{aligned}$$

where $\varepsilon = +1$ [$\varepsilon = -1$] or equivalently :

$$(9') \quad \begin{aligned} x^0 \geq 0, \quad \nabla_x L(x^0, u^0) &\geq 0, \quad x^{0'} \nabla_x L(x^0, u^0) = 0 \\ u^0 \geq 0, \quad \nabla_u L(x^0, u^0) &\leq 0, \quad u^{0'} \nabla_u L(x^0, u^0) = 0 \end{aligned}$$

where

$$L(x, u) = \varepsilon f(x) + u' g(x)$$

(S₁) If for a $u_{i_0}^0 > 0$ ($i_0 \in J^0$) we have

$$(10) \quad (x - x^0)' \nabla g_{i_0}(x^0) < 0, (\forall) x \in C - \{x^0\},$$

then relations (9) or (9') are also sufficient so that x^0 should be a point of local minimum [maximum] of the function f in C .

(S₂) If $C \cap \{x \in R^n / (x - x^0)' \nabla f(x^0) = 0\} = \{x^0\}$,

then relations (9) or (9') are also sufficient so that point x^0 should be a local minimum [maximum] of the function f in C .

(S₃) If function f is quasi-convex in C and $\nabla f(x^0) \neq 0$, then relations (9) or (9') are also sufficient so that point x^0 should be the global minimum [maximum] of the function f on C .

Proof. (n) If x^0 is a point of local minimum of the function f in C , then functions f , g and $h = -x$ verify the Kuhn-Tucker conditions in x^0 [9]: there are $u^0 \in R^m$ and $y^0 \in R^n$ such that

$$\begin{aligned}
 \varepsilon \nabla f(x^0) + u^0 \nabla g(x^0) + y^0 \nabla h(x^0) &= 0 \quad (1) \\
 u^0 g(x^0) &= 0 \\
 y^0 h(x^0) &= 0 \\
 u^0 &\geq 0 \\
 y^0 &\geq 0
 \end{aligned}
 \tag{11}$$

However $\nabla h(x^0) = \underbrace{(-1, \dots, -1)}_n$ and relations (11) lead immediately to relations (9).

(S₁) From relations (9') it results

$$(x - x^0)' \nabla_x L(x^0, u^0) \geq 0, \quad (\forall) x \in C$$

The hypotheses of theorem 2 (a) are verified for $h \equiv 0$ and hence x^0 is a point of local minimum [maximum] of the function f in C .

(S₂) The hypotheses of the theorem 2 (b) with $h \equiv 0$ are verified.

(S₃) The hypotheses of the theorem 2 (c) with $h \equiv 0$ are verified.

From theorem 3 the following corollary will result.

Corollary. Let be functions $f, g = (g_1, \dots, g_m)$ and $h = (h_1, \dots, h_p)$ differentiable on an open convex set $K \subseteq R^{n_1} \times R^{n_2}$.

Besides, we assume that function g is quasi-convex on K , while h is quasi-convex and quasi-concave on K .

(n) A necessary condition so that point $(x^0, y^0) \in K$ should be a point of local minimum [maximum] for the function f on the set

$$C = \{ (x, y) \in K / g(x, y) \leq 0, \quad h(x, y) = 0, \quad y \geq 0 \},$$

with nonempty interior is that $u^0 \in R^m$ and $v^0 \in R^p$ should exist so that

$$\begin{aligned}
 x^0 \in R^{n_1}, \quad \nabla_x L(x^0, y^0, u^0, v^0) &= 0 \\
 y^0 \geq 0, \quad \nabla_y L(x^0, y^0, u^0, v^0) \geq 0, \quad y^0 \nabla_y L(x^0, y^0, u^0, v^0) &= 0 \\
 u^0 \geq 0, \quad \nabla_u L(x^0, y^0, u^0, v^0) \leq 0, \quad u^0 \nabla_u L(x^0, y^0, u^0, v^0) &= 0 \\
 v^0 \in R^p, \quad \nabla_v L(x^0, y^0, u^0, v^0) &= 0,
 \end{aligned}
 \tag{12}$$

where

$$L(x, y, u, v) = \varepsilon f(x, y) + u'g(x, y) + v'h(x, y),$$

$$\begin{aligned}
 (1) \quad \nabla g(x^0) &= \begin{pmatrix} \nabla g_1(x^0) \\ \vdots \\ \nabla g_m(x^0) \end{pmatrix}
 \end{aligned}$$

while $\varepsilon = +1$ [$\varepsilon = -1$].

(s₁) If for a $u_{i_0}^0 > 0$ we have

$$(x - x^0)' \nabla_x g_{i_0}(x^0, y^0) + (y - y^0)' \nabla_y g_{i_0}(x^0, y^0) < 0, (\forall)(x, y) \in C - \{(x^0, y^0)\}$$

or at least for a $v_{j_0}^0 > 0$ we have

$$(x - x^0)' \nabla_x h_{j_0}(x^0, y^0) + (y - y^0)' \nabla_y h_{j_0}(x^0, y^0) < 0, (\forall)(x, y) \in C - \{(x^0, y^0)\}$$

or

$$(x - x^0)' \nabla_x h_{j_0}(x^0, y^0) + (y - y^0)' \nabla_y h_{j_0}(x^0, y^0) > 0, (\forall), (x, y) \in C - \{(x^0, y^0)\}$$

the relations (12) are also sufficient, such that point (x^0, y^0) should be a local minimum [maximum] of the function f on C .

(s₂) If

$$C \cap \{ (x, y) \in R^{n_1} \times R^{n_2} / (x - x^0)' \nabla_x f(x^0, y^0) + \\ + (y - y^0)' \nabla_y f(x^0, y^0) = 0 \} = \{ (x^0, y^0) \},$$

the relations (12) are also sufficient such that point (x^0, y^0) should be a local minimum [maximum] of the function f on C .

(s₃) If function f is quasi-convex in C and $\nabla f(x^0, y^0) \neq 0$ then relations (12) are also sufficient such that point (x^0, y^0) should be the global minimum [maximum] of the function f on C .

REMARK. If the vectorial function $g(x)$ is quasi-concave then there are the theorems dual to theorem 2 and 3.

REFERENCE

- [1] ARROW H. J. and ENTHOVEN A. C., *Quasi-concave Programming*, *Econometrica*, 29 (1961), 779-800.
- [2] BECTOR C. R., *Some aspects of quasi-convex programming*, *Z. Angew. Math., Mech.*, 50 (1970), 495-497.
- [3] DRAGOMIRESCU M. et MALITA M., *Programare neliniara*, Ed. st., Bucuresti, 1972.
- [4] FIACCO A. V. and McCORMICK G. P., *Nonlinear Programming : Sequential Unconstrained Minimization Techniques*, Wiley, New York, 1968.
- [5] KORTANEK K. O. and EVANS I. P., *Pseudo-concave Programming and Lagrange Regularity*, *Operat. Res.*, 15 (1967), 882-891.
- [6] MANGASARIAN O. L., *Nonlinear Programming*, McGraw-Hill, New York, 1969.
- [7] MANGASARIAN O. L., *Pseudo-convex Functions*, *SIAM Control*, 3 (1965), 281-290.
- [8] MITITELU St., *Generalized Quasi-Convex Programmes*, *Rev. Roum. Math. Pures Appl.*, 4, tom XVII, 1972.
- [9] MITITELU St., *Kuhn-Tucker, Conditions for Nonlinear Programmes with Quasi-convex Constraints*, *Rev. Roum. Math. Pures Appl.* (to appear).