

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

VITTORIO SCORNAZZANI

Pointwise estimates for minimizers of some non-uniformly degenerate functionals

Rendiconti del Seminario Matematico della Università di Padova,
tome 81 (1989), p. 151-169

http://www.numdam.org/item?id=RSMUP_1989__81__151_0

© Rendiconti del Seminario Matematico della Università di Padova, 1989, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Pointwise Estimates for Minimizers of Some Non-uniformly Degenerate Functionals.

VITTORIO SCORNAZZANI (*)

RIASSUNTO - In questo articolo si prova l'hölderianità e una disuguaglianza di Harnack per i minimi di funzionali che degenerano in maniera non uniforme. Si adattano le tecniche di Giaquinta-Giusti e di Di Benedetto-Trudinger introducendo su R^n una metrica « naturale », diversa da quella euclidea, per i funzionali considerati che tiene conto della non uniforme degenerazione dei funzionali stessi.

Introduction.

In this paper we prove the Hölder regularity and the Harnack inequality for the minimizers of following functional:

$$(1) \quad F(u, \Omega) = \int_{\Omega} F(x, Du) dx, \quad D = \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n} \right)$$

where Ω is an open subset of R^n . We assume that the following hypotheses are satisfied by function F : F is a Caratheodory function, and there exist two constants $M > 0$ and $m > 1$ such that:

$$(2) \quad M^{-1}w(x) \left(\sum_{j=1}^n \lambda_j^2 P_j^2 \right)^{m/2} \leq F(x, P) \leq Mw(x) \left(\sum_{j=1}^n \lambda_j^2 P_j^2 \right)^{m/2}.$$

(*) Indirizzo dell'A.: Università degli Studi di Bologna, Dipartimento di Matematica, Piazza di Porta San Donato 5, 40127 Bologna.

Here w is a nonnegative weight in the sense of Muckenhaupt (see later on) and λ_j , $j = 1, \dots, n$ are nonnegative functions defined on \mathbf{R}^n satisfying the same hypotheses of [FL1] and [FL2]. In the case $\lambda_j = 1$ and $w = 1$, Giaquinta and Giusti ([GG1], [GG2]) proved the Hölder regularity of minimizers of F while Di Benedetto and Trudinger ([BT]) proved the Harnack inequality for nonnegative minimizers of F . Afterwards Modica ([M]) extended these results to the case $\lambda_j = 1$ and w is a weight that satisfies the following assumptions: $w \geq 0$ and there exist $p > 1$, $c_w \equiv c(w, p) \geq 1$ such that

$$(3) \quad \left(\frac{1}{|S_R|} \int_{S_R} w \, dx \right) \left(\frac{1}{|S_R|} \int_{S_R} w^{-1/(p-1)} \, dx \right)^{p-1} < c_w$$

for any euclidean ball S_R of radius R . Here $|S_R|$ is Lebesgue measure of S_R . The results of the works listed above cannot apply to non-uniformly degenerating functionals like, for example, to the following one

$$F(u, \Omega) = \int_{\Omega} (|D_x u|^2 + |x|^{2\sigma} |D_y u|^2)^{m/2} \, dx \, dy, \quad (x, y) \in \mathbf{R}^2 \times \mathbf{R}^q, \quad \sigma > 0.$$

On the other hand it seems to be natural to handle such a functional by equipping \mathbf{R}^n by a metric d constructed in order to taking account of the special nonuniformly degeneration of F (and therefore of λ_j). A class of metrics of this kind has been studied by several authors, see, for example, [FL1], [FL2], [FL3], [NSW]. By using the metric d in the place of the euclidean metric one can adapt the technics of Giaquinta and Giusti, of Di Benedetto and Trudinger and of Modica to the study of functionals of the kind (1) (2). It is reasonable to require that the weight w satisfies the condition of Muckenhaupt with respect to the metric d ; hence we substitute the hypothesis (3) with the following one: $w \geq 0$, there exist $p > 1$, and $c_w \equiv c(w, p) \geq 1$ such that

$$(4) \quad \left(\frac{1}{|B_R|} \int_{B_R} w \, dx \right) \left(\frac{1}{|B_R|} \int_{B_R} w^{-1/(p-1)} \, dx \right)^{p-1} < c_w$$

for any d -ball of radius R , $B_R \subset \Omega$.

As a consequence, our results can be applied, for example, to the functional defined by:

$$F(x, y, D_x u, D_y u) = (|D_x u|^2 + |x|^{2\sigma} |D_y u|^2)^{m/2} \|(x, y)\|^\alpha,$$

$$(x, y) \in \mathbf{R}^p \times \mathbf{R}^q, \sigma > 0, \alpha \in \mathbf{R}.$$

In Section 1 we introduce the class $\mathcal{DS}_m(\Omega, \lambda, w)$ which is the natural De Giorgi's class modelled on metric d and weight w . Then we prove that the functions in De Giorgi's class $\mathcal{DS}_m(\Omega, \lambda, w)$ are Hölder continuous and then we prove that the minimizers are in $\mathcal{DS}_m(\Omega, \lambda, w)$ and therefore they are Hölder continuous.

In Section 2 we prove the Harnack inequality by using a Krylov-Safanov ([KS]) covering type Lemma, that we prove by using fundamentally the homogeneity of the space $(\mathbf{R}^n, d, w(x) dx)$.

Now we list explicitly the hypotheses on the function λ_i . We suppose:

i) $\lambda_j \geq 0, \lambda_1 \equiv 1, \lambda_j(x) = \lambda_j(x_1, \dots, x_{j-1}) \quad \forall x \in \mathbf{R}^n, j = 1, \dots, n.$

ii) Put $\Pi = \left\{ x \in \mathbf{R}^n : \prod_{k=1}^n x_k = 0 \right\}$, then

$$\lambda_j \in C(\mathbf{R}^n) \cap C^1(\mathbf{R}^n - \Pi), \quad 0 < \lambda_j(x) \leq 1 \quad \forall x \in \mathbf{R}^n - \Pi, j = 1, \dots, n$$

$$\lambda_j(x_1, \dots, x_i, \dots, x_{j-1}) = \lambda_j(x_1, \dots, -x_i, \dots, x_{j-1})$$

$$j = 2, \dots, n; \quad i = 1, \dots, j-1.$$

iii) There exists a family of nonnegative numbers ϱ_{ij} such that

$$0 \leq x_i \frac{\partial}{\partial x_i} (\lambda_j(x)) \leq \varrho_{ij} \lambda_j(x), \quad j = 2, \dots, n, \quad i = 1, \dots, j-1.$$

These hypotheses allow to construct a «natural» metric d for the functional, associated with the fields $X_j = \lambda_j(\partial/\partial x_j)$, $j = 1, \dots, n$ (see [FL1], [FL2], [FL3], [NSW]) in the following way.

A continuous curve $\gamma \subseteq \Omega$ is X -admissible if:

i) γ is piecewise C^1 ;

ii) each piece C^1 of γ is an integral curve of one of the vector fields $\pm X_j$, $j = 1, \dots, n$.

If $\gamma: [0, T] \rightarrow \Omega$, we put $1(\gamma) = T$. The hypotheses on λ allow to prove that for any $x, y \in \Omega$ there exists an X -admissible curve γ joining x and y . Hence we can define $d(x, y) = \inf \{1(\gamma): \gamma \text{ is } X\text{-admissible and joins } x \text{ and } y\}$. For every compact subset K of Ω there exists $c = c(K)$ and $\varepsilon > 0$ such that

$$(5) \quad c^{-1}|x - y| \leq d(x, y) \leq c|x - y|^\varepsilon$$

where $|\cdot|$ denotes the euclidean norm on \mathbb{R}^n (see [FL2], Prop. 2.9). In the following we shall write

$$w(E) = \int_E w \, dx$$

and, for $k \in \mathbb{R}$

$$B(k, R) = \{x \in B_R / u(x) > k\}.$$

It is well known that $|E| = 0$ iff $w(E) = 0$ (see, for example, Lemma 4 in [C]) and that for the measure $w(x) \, dx$ the following doubling property holds: there exists a constant $\beta = \beta(p, c(w, p), d)$ such that

$$(6) \quad w(B_{2R}) \leq \beta w(B_R)$$

for every d -ball of radius R (see [FS], Lemma 2.10).

Given $u \in \text{Lip}(\Omega)$, we denote

$$D_\lambda u = \left(\lambda_1 \frac{\partial u}{\partial x_1}, \dots, \lambda_n \frac{\partial u}{\partial x_n} \right) \quad \text{and} \quad |D_\lambda u| = \left(\sum_{j=1}^n \lambda_j^2 \left(\frac{\partial u}{\partial x_j} \right)^2 \right)^{\frac{1}{2}},$$

besides we denote by $W_m^1(\Omega)$ the closure of the space $\text{Lip}(\Omega)$ with respect to the norm

$$\|u; W_m^1(\Omega)\| = \left(\int_\Omega |u|^m w \, dx \right)^{1/m} + \left(\int_\Omega |D_\lambda u|^m w \, dx \right)^{1/m}.$$

DEFINITION 1. We call $u \in W_m^1(\Omega)$ a minimizer for $F(\cdot, \Omega)$ if for all $\Phi \in W_m^1(\Omega)$ with $\text{supp } \Phi \subset \Omega$

$$F(u, \text{supp } \Phi) \leq F(u + \Phi, \text{supp } \Phi).$$

DEFINITION 2. A function u is said to belong to De Giorgi's class, $\mathcal{DG}_m(\Omega, \lambda, w) \equiv \mathcal{DG}_m(\Omega)$ if:

- i) $u \in W_m^1(\Omega)$;
- ii) for any d -ball $B_R \subset \Omega$ and for any $\varrho > 0$, $\varrho < R$, $k \in \mathbf{R}$

$$(8) \quad \int_{B(k, \varrho)} |D_\lambda u|^m w \, dx = \frac{C}{(R - \varrho)^m} \int_{B(k, R)} |u - k|^m w \, dx \quad (1)$$

where B_ϱ is the d -ball concentric to B_R and radius ϱ .

REMARK. In the case $\lambda_j \equiv 1$, the class $\mathcal{DG}_m(\Omega)$ coincide with the usual De Giorgi's class (see [DG], [GG1], [M]).

§ 1. Here we prove that the functions in $\mathcal{DG}_m(\Omega)$ are Hölder continuous (Th. 5) and that the minimizers are in $\mathcal{DG}_m(\Omega)$. To this aim we make use of the following embedding Theorem whose proof is similar to the one of Th. 4.1 of [FS]:

THEOREM 3. Let w be a weight such that satisfies (4) with $p \leq m$, let B_R be the d -ball with centre \bar{x} and radius R and $u \in W_m^1(\Omega)$ such that for a $\beta > 0$ we have $w(\{x \in B_R / u(x) = 0\}) \geq \beta w(B_R)$. Then exist $l > 1$ and $C > 0$ such that:

$$(9) \quad \left(\int_{B_R} |u|^{lm} w \, dx \right)^{1/lm} \leq CR w(B_R)^{(1-l)/lm} \left(\int_{B_R} |D_\lambda u|^m w \, dx \right)^{1/m}.$$

Here C depends on \bar{x} and β and on constants of (4), l depends on m , ϱ_{ij} , and constants of (4).

PROOF. Put $E = \{x \in B_R / u(x) = 0\}$ and $u_R = (w(B_R))^{-1} \int_{B_R} u w \, dx$. Then, making use of Hölder inequality:

$$|u_R| \leq (w(E))^{-1} \int_E |u_R - u| w \, dx \leq \beta^{-1} \left((w(B_R))^{-1} \int_{B_R} |u_R - u|^{lm} w \, dx \right)^{1/lm}$$

(1) In the following we denote by C a positive constant which is not always the same at each occurrence.

by that, we have:

$$\begin{aligned} \left((w(B_R))^{-1} \int_{B_R} |u|^{lm} w \, dx \right)^{1/lm} &\leq C \left((w(B_R))^{-1} \int_{B_R} |u_R - u|^{lm} w \, dx \right)^{1/lm} \leq \\ &\leq CR \left((w(B_R))^{-1} \int_{B_R} |D_\lambda u|^m w \, dx \right)^{1/m} \end{aligned}$$

by Th. 4.5 of [FS].

By (9) it is possible to prove that, if u and $-u \in \mathcal{DS}_m(\Omega)$, then u is locally essentially bounded. In fact, we have:

THEOREM 4. Let $u \in \mathcal{DS}_m(\Omega)$, $k \in R$, $\sigma \in]0, 1[$, $0 < R < R_0$: Then for any d -ball $B_R \subset \Omega$ we have:

$$(10) \quad \sup_{B_{\sigma R}} u \leq k + \frac{C}{(1-\sigma)^{\Theta/(l-1)}} \cdot (w(B_R))^{-\Theta/m} (w(B(k, R)))^{(\Theta-1)/m} \left(\int_{B(k, R)} |u - k|^m w \, dx \right)^{1/m}$$

where $\Theta = \Theta(l) > 1$ and l is the constant in (9), and $C = C(\bar{x}) > 0$.

We first prove two Lemmas in which we use the following notation like those used by Giaquinta ([G])

$$\begin{aligned} u(h, \varrho) &:= \int_{B(h, \varrho)} |u - k|^m w \, dx, \\ b(h, \varrho) &:= w(B(h, \varrho)), \quad \Phi(h, \varrho) := u^{l\Theta/(l-1)}(h, \varrho) b(h, \varrho) \end{aligned}$$

where $\Theta = (1 + (1 + 4(l-1)/l)^{1/2})/2$.

LEMMA 4.1. Let $0 < \varrho < R < R_0$ and $h > k$. For any $u \in \mathcal{DS}_m(\Omega)$ we have:

$$(11) \quad \Phi(h, \varrho) = \frac{CR^{m\Theta/(l-1)} (w(B_R))^{-\Theta}}{(R-\varrho)^{m\Theta/(l-1)} |h-k|^m} \Phi^\Theta(k, R).$$

PROOF. Let B_R be a d -ball contained in Ω . Then there exists $\eta \in C_0^\infty(B_{(R+\varrho)/2})$, $\eta = 1$ on B_ϱ , $0 < \eta < 1$, and such that $|D_\lambda \eta| < C/(R-\varrho)$ (see [FL2], proof of Lemma 4.2). Let $u \in \mathcal{DS}_m(\Omega)$. In particular

$u \in W_m^1(\Omega)$, so by Corollary 5.2 in [FS], it results $\eta \max(u - k, 0) \in W_m^1(\Omega)$. From inequality (9) and doubling property for the measure $w(x) dx$, we have:

$$\begin{aligned} \left(\int_{B((R+\varrho)/2)} |\eta \max(u - k, 0)|^{lm} w dx \right)^{1/lm} &\leq \\ &\leq CR(w(B_R))^{(1-l)/lm} \left(\int_{B((R+\varrho)/2)} |D_\lambda(\eta \max(u - k, 0))|^m w dx \right)^{1/m} \end{aligned}$$

for some $l > 1$. By Hölder inequality, we get:

$$\int_{B_\varrho} |\max(u - k, 0)|^m w dx \leq \left(\int_{B_\varrho} |\max(u - k, 0)|^{lm} w dx \right)^{1/l} (w(B(k, \varrho)))^{(l-1)/l}.$$

Hence, because $\varrho < (R + \varrho)/2 < R$ and $u \in \mathcal{DS}_m(\Omega)$:

$$\begin{aligned} \int_{B(k, \varrho)} |u - k|^m w dx &\leq CR^m (w(B_R))^{(1-l)/l} \\ &\cdot \int_{B(k, (R+\varrho)/2)} |D_\lambda(\eta \max(u - k, 0))|^m w dx \cdot (w(B(k, \varrho)))^{(l-1)/l} \leq \\ &\leq \frac{CR^m (w(B_R))^{(1-l)/l}}{(R - \varrho)^m} \int_{B(k, R)} |u - k|^m w dx \cdot (w(B(k, \varrho)))^{(l-1)/l}. \end{aligned}$$

Now, if $h > k$, and $0 < \varrho < R$, we have

$$|h - k|^m w(B(h, \varrho)) \leq \int_{B(h, \varrho)} |u - k|^m w dx \leq \int_{B(k, R)} |u - k|^m w dx$$

hence, because $u(h, \varrho) \leq u(k, \varrho)$, using our notation we get

$$\begin{cases} u(h, \varrho) \leq \frac{CR^m (w(B_R))^{(1-l)/l}}{(R - \varrho)^m} u(k, R) b(k, R)^{(l-1)/l}, \\ b(h, \varrho) = \frac{1}{|h - k|^m} u(k, R). \end{cases}$$

For any positive numbers ξ and ζ we find

$$u^\xi(h, \varrho) b^\zeta(h, \varrho) \leq \frac{CR^{m\xi}(w(B_R))^{(1-\xi)/l}}{(R-\varrho)^{m\xi}|h-k|^{m\xi}} u^{\xi+\zeta}(k, R) b(k, R)^{(1-\xi)/l}.$$

Now we choose ξ and ζ in such way that for some $\Theta > 0$, we have $\xi + \zeta = \Theta$, $(l-1)\xi/l = \Theta\zeta$, (then Θ must be the positive solution of $\Theta^2 - \Theta - (l-1)/l = 0$ i.e. $\Theta = (1 + (1 + 4(l-1)/l)^{1/2})/2 > 1$. We can choose $\zeta = 1$ and $\xi = l\Theta/(l-1)$ and consequently the Lemma follows.

LEMMA 4.2. For any $k \in \mathbb{R}$, $0 < R < R_0$, $\sigma \in]0, 1[$, it holds:

$$\Phi(k + d, \sigma R) = 0$$

where

$$d = \frac{2^{(l\Theta/(l-1)+1)\Theta/(\Theta-1)} C^{1/m} (w(B_R))^{-\Theta/m}}{(1-\sigma)^{l\Theta/(l-1)}} \Phi^{(\Theta-1)/m}(k, R)$$

and C is the constant that appears in Lemma 4.1.

PROOF. We set $h_n = k + d(1 - 2^{-n})$, $\varrho_n = \sigma R + (1 - \sigma)R2^{-n}$; $n \in \mathbb{N}$ then $h_n \nearrow k + d$, $\varrho_n \searrow \sigma R$. For every $n \in \mathbb{N}$ we have

$$\Phi(h_n, \varrho_n) \leq \frac{\Phi(k, R)}{2^{m/(\Theta-1)(l\Theta/(l-1)+1)n}}.$$

This easily follows by Lemma 4.1 and by induction. Now it is enough take the limit $n \rightarrow +\infty$ to prove the Lemma.

PROOF OF THEOREM 4. By Lemma 4.2, we have $\Phi(k + d, \sigma R) = 0$, thus either $u(k + d, \sigma R) = 0$ or $b(k + d, \sigma R) = 0$. Hence, since $(\Theta - 1)\Theta/l = 1$, the thesis follows.

This holds also for $-u$ and hence u is locally essentially bounded.

THEOREM 5. If u and $-u \in \mathcal{DS}(\Omega)$, then u is Hölder continuous. We first prove the following two Lemmas:

LEMMA 5.1. Let $h > k > k_0$, if $w(B(k_0, R)) < \gamma w(B_R)$, $0 < \gamma < 1$, then:

$$(12) \quad |h - k|^{(m+1)/2} (w(B(h, R)))^{1/l} \leq C(\gamma) R^{(m+1)/2} (w(B_R))^{(1-l)/l} \left(\int_{B(k, R)} |D_\lambda u|^m w \, dx \right)^{(m+1)/2m} \cdot (w(B(k, R) - B(h, R)))^{(m-1)/2m}.$$

PROOF. Put $v = \min(u, h) - \min(u, k)$. Then:

$$\begin{aligned} w(\{x \in B_R / v(x) = 0\}) &= w(B_R - B(k, R)) = \\ &= w(\{x \in B_R / u(x) \leq k\}) \geq w(\{x \in B_R / u(x) \leq k_0\}) \geq (1 - \gamma) w(B_R). \end{aligned}$$

So we can apply (9). There exists $l > 1$, such that

$$\begin{aligned} |h - k|^{(m+1)l/2} w(B(h, R)) &= \int_{B(h, R)} |v|^{(m+1)l/2} w \, dx \leq \\ &\leq \int_{B(k, R)} |v|^{(m+1)l/2} w \, dx \leq \int_{B_R} |v|^{(m+1)l/2} w \, dx \leq \\ &\leq C(\gamma) R^{(m+1)l/2} (w(B_R))^{1-l} \left(\int_{B_R} |D_\lambda v|^{(m+1)/2} w \, dx \right)^l = \\ &= C(\gamma) R^{(m+1)l/2} (w(B_R))^{1-l} \left(\int_{B(k, R) - B(h, R)} |D_\lambda u|^{(m+1)/2} w \, dx \right)^l. \end{aligned}$$

By Hölder inequality (12) follows.

LEMMA 5.2. Let $u \in \mathcal{DS}_m(\Omega)$; setting

$$\begin{aligned} M(2R) &= \sup_{B_{2R}} u, & m(2R) &= \inf_{B_{2R}} u; \\ k &= \frac{1}{2}(M(2R) + m(2R)), & k_\nu &= M(2R) - \frac{M(2R) - k_0}{2^\nu}. \end{aligned}$$

Then, if $w(B(k_0, R)) \leq \gamma w(B_R)$, $0 < \gamma < 1$, we have:

$$(13) \quad w(B(k_\nu, R)) \leq \frac{Cw(B_R)}{\gamma^{l(m-1)/2m}}.$$

PROOF. In inequality (12) we put $k \equiv k_{i-1}$ and $h \equiv k_i$: Hence, by (8) and by doubling property of measure $w(x) \, dx$:

$$\begin{aligned} (w(B(k_i, R)))^{2ml(m-1)} &\leq \\ &\leq C(w(B_R))^{(2m(1-l) + l(m+1))l(m-1)} (w(B(k_{i-1}, R)) - w(B(k_i, R))). \end{aligned}$$

Summing up for $i = 1, 2, \dots, \nu$ and using the inclusion $B(k_\nu, R) \subset \subset B(k_i, R)$, we have

$$\nu(w(B(k_\nu, R)))^{2m/l(m-1)} \leq C(w(B_R))^{(2m(1-1)+l(m+1))/l(m-1)} w(B(k_0, R)).$$

So the assertion follows.

PROOF OF THEOREM 5. We use the notations of Lemma 5.2. We may assume $w(B(k_0, R)) < \frac{1}{2}w(B_R)$. Otherwise, since

$$\{x \in B_R - u(x) > -k_0(-u)\} = \{x \in B_R \mid u(x) < k_0(u)\},$$

we can work with $-u$. Applying Theorem 4 and replacing k by k_ν , we have

$$\begin{aligned} M(R/2) &\leq k_\nu + C(w(B_R))^{-\Theta/m} \left(\int_{B(k_\nu, R)} |u - k_\nu|^m w \, dx \right)^{1/m} (w(B(k_\nu, R)))^{(\Theta-1)/m} \leq \\ &\leq k_\nu + C(w(B_R))^{-\Theta/m} |M(2R) - k_\nu| (w(B(k_\nu, R)))^{\Theta/m}. \end{aligned}$$

By Lemma 5.2 we can choose a $\bar{\nu}$ independent on R , such that $C(w(B_R))^{-\Theta/m} (w(B(k_\nu, R)))^{\Theta/m} < \frac{1}{2}$. Then

$$\begin{aligned} M(R/2) &\leq M(2R) - 2^{-(\bar{\nu}+1)}(M(2R) - m(2R)) + \frac{1}{2}(M(2R) - k_\nu) = \\ &= M(2R) - 2^{-(\bar{\nu}+2)}(M(2R) - m(2R)). \end{aligned}$$

We now subtract $m(R/2)$:

$$M(R/2) - m(R/2) \leq (M(2R) - m(2R))(1 - 2^{-(\bar{\nu}+2)})$$

i.e. $\omega(R/2) = M(R/2) - m(R/2) \leq \beta\omega(2R)$, with $\beta \in]0, 1[$ and thus there exists a constant α such that $\omega(\rho) \leq C(\rho/R)^\alpha \omega(R)$ and u is Hölder continuous by (5).

Now we prove that the minimizers are Hölder continuous. This result follows from Theorem 5 and the following

THEOREM 6. Let $u \in W_m^1(\Omega)$ be a minimizer for the functional F defined in (1), (2). Then u and $-u \in \mathcal{DS}_m(\Omega)$.

PROOF. Let B_R be the d -ball with centre $\bar{x} \in \Omega$, such that $B_R \subset \subset \Omega$. Moreover let u a minimizer (see (7)). Let now $\eta \in C_0^\infty(B_R)$, $\eta = 1$ in

B_ϱ , $0 < \varrho < R$ and such that $|D_\lambda u| \leq C/(R - \varrho)$ (see [FL2], proof of Lemma 4.2). For a fixed $k \in \mathbf{R}$, we put $\Phi = -\eta \max(u - k, 0)$. Since $B(k, \varrho) = \{x \in B_\varrho / u(x) > k\} \subseteq \text{supp } \Phi$, we have

$$\begin{aligned} M^{-1} \int_{B(k, \varrho)} |D_\lambda u|^m w \, dx &\leq M^{-1} \int_{\text{supp } \Phi} |D_\lambda u|^m w \, dx \leq \int_{\text{supp } \Phi} F(x, Du) \, dx \leq \\ &\leq \int_{\text{supp } \Phi} F(x, D(u + \Phi)) \, dx \leq M \int_{\text{supp } \Phi} |D_\lambda(u + \Phi)|^m w \, dx \leq \\ &\leq C \int_{\text{supp } \Phi} (1 - \eta)^m |D_\lambda u|^m w \, dx + C \int_{\text{supp } \Phi} |u - k|^m |D_\lambda \eta|^m w \, dx \leq \\ &\leq C \int_{B(k, R) - B(k, \varrho)} |D_\lambda u|^m w \, dx + \frac{C}{(R - \varrho)^m} \int_{B(k, R)} |u - k|^m w \, dx. \end{aligned}$$

In the last step we used $\text{supp } \Phi \subseteq \{x \in B_R / u(x) > k\}$ and $\eta = 1$ in B_ϱ . From this inequality adding $C \int_{B(k, \varrho)} |D_\lambda u|^m w \, dx$, we obtain

$$(M^{-1} + C) \int_{B(k, \varrho)} |D_\lambda u|^m w \, dx \leq C \int_{B(k, R)} |D_\lambda u|^m w \, dx + \frac{C}{(R - \varrho)^m} \int_{B(k, R)} |u - k|^m w \, dx.$$

This, by Lemma 1.1 in [GG1], implies

$$\int_{B(k, \varrho)} |D_\lambda u|^m w \, dx \leq \frac{C}{(R - \varrho)^m} \int_{B(k, R)} |u - k|^m w \, dx;$$

that is $u \in \mathcal{DS}_m(\Omega)$.

At this point, to prove that $-u \in \mathcal{DS}_m(\Omega)$, it is enough to observe that $-u$ is a minimizer of the functional $\tilde{F}(v, \Omega) = \int_\Omega \tilde{F}(x, Dv) \, dx$ where $\tilde{F}(x, Dv) = F(x, D(-v))$. It is evident that \tilde{F} satisfies the same hypotheses as F .

§ 2. Here we prove the Harnack inequality for the nonnegative minimizers of F . To this aim we prove a Krylow-Safanov covering type Lemma:

LEMMA 7. Let B_R be a d -ball, $E \subseteq B_R$, E measurable, $\delta \in]0, 1[$,

$$B = \{B_{\rho}(x) \cap B_R / x \in B_R, \rho > 0, w(E \cap B_{\rho}(x)) \geq \delta w(B_{\rho}(x) \cap B_R)\}$$

($B_{\rho}(x)$ is the d -ball of centre x and radius ρ). We consider

$$E_{\delta} = \bigcup_{B \in \mathcal{B}} B.$$

Then either i) $E_{\delta} = B_R$, or ii) $w(E) < C\delta w(E_{\delta})$.

Here $C > 1$ is dependent on the doubling constant for the measure $w(x) dx$.

PROOF. If $w(E) \geq \delta w(B_R)$ then, since $E \subseteq B_R$, we have $w(E) = w(E \cap B_R) \geq \delta w(B_R)$ and hence $B_R \in B$ and hence i) is valid.

Let now $w(E) < \delta w(B_R)$. If $w(E) = 0$ then ii) is valid. Let so $w(E) > 0$. We say that a d -ball of centre x and radius ρ intersects substantially E (B i.s. E) if $w(E \cap B_{\rho}(x)) \geq \delta w(B_{\rho}(x) \cap B_R)$. We suppose that every point of E let be a Lebesgue point ([C]) that is

$$(14) \quad \lim_{r \rightarrow 0^+} \frac{w(E \cap B_r(x))}{w(B_r(x))} = 1.$$

Afterwards, we cover B_R by means of d -balls $B_{R/2}(x_i^{(1)})$ of radius $R/2$ which are centred at the points of a maximal set $\{x_1^{(1)}, \dots, x_{I_1}^{(1)}\} \subset B_R$ with $d(x_i^{(1)}, x_j^{(1)}) > R/2$, $\forall i \neq j$ ([CW]). Moreover we cover B_R by means of d -ball of radius $R/4$ centred at the points of a maximal set $\{x_1^{(2)}, \dots, x_{I_2}^{(2)}\} \subset B_R$ with $d(x_i^{(2)}, x_j^{(2)}) > R/4$ and so on; in this way we obtain a class of families

$$R^{(p)} = \{B_{2^{-(p-1)}R}(x_i^{(p)}) / i = 1, \dots, I_p\} \quad \text{of covering of } B_R.$$

Then we consider

$$\mathcal{S}_p = \left\{ B_{R/2^{p-1}}(x_i^{(p)}) \ i = 1, \dots, I_p, B \text{ i.s. } E, \right. \\ \left. x_i \notin \bigcup_{h=1}^{p-1} \bigcup_{B \in R^p} B, B_{R/2^{h-1}}(x_j^{(h)}) \text{ i.s. } E \right\}$$

It results that for every $x \in E$, $\exists p \in \mathbb{N}$, $i \in \mathbb{N}$ such that $x \in B_{R/2^p}(x_i^{(p)})$ and $w(E \cap B_{R/2^{p-1}}(x_i^{(p)})) \geq \delta w(B_{R/2^{p-1}}(x_i^{(p)}))$. In fact, by (14), taken

$\sigma > \delta/A^2$ (A is the doubling constant for the measure $w(x) dx$), $\sigma < 1$, it exists $r_x \in \mathbf{R}_+$ such that $\forall r < r_x$ it holds $w(E \cap B_r(x)) \geq \sigma w(B_r(x))$. Let be $x \in E$, there exist $p, i \in \mathbf{N}$ such that $x \in B_{R/2^p}(x_i^{(p)})$ with $R/2^p < r_x$, then

$$\begin{aligned} w(E \cap B_{R/2^{p-1}}(x_i^{(p)})) &\geq w(E \cap B_{R/2^p}(x)) \geq \sigma w(B_{R/2^p}(x)) \geq A\sigma w(B_{R/2^{p-1}}(x)) \geq \\ &\geq A\sigma w(B_{R/2^p}(x_i^{(p)})) \geq A^2 \sigma w(B_{R/2^{p-1}}(x_i^{(p-1)})) \geq \delta w(B_{R/2^{p-1}}(x_i^{(p)})) . \end{aligned}$$

To any $B \in \bigcup_{p \in \mathbf{N}} S_p$, we associate \tilde{B}_B as following manner: let $B \in \bigcup_{p \in \mathbf{N}} S_p$ then $\exists p \in \mathbf{N}, \exists i \in \{1, \dots, I_p\}$ such that $B \equiv B_{R/2^{p-1}}(x_i^{(p)}) \in S_p$ with $x_i^{(p)} \in B_{R/2^{p-1}}(x_j^{(p-1)})$, $j \in \{1, \dots, I_{p-1}\}$. We put $\tilde{B}_B = B_{R/2^{p-1}}(x_j^{(p-1)})$. It results $B \subset \tilde{B}_B$ and \tilde{B}_B not intersects substantially E . Then we put

$$\tilde{E}_\delta = \bigcup_{p \in \mathbf{N}} \left(\bigcup_{B \in S_p} (\tilde{B}_B \cap B_R) \right).$$

It results $\tilde{E}_\delta \subseteq E_\delta$. In fact: let $y \in \tilde{E}_\delta$, then $\exists p \in \mathbf{N}, \exists B_{R/2^{p-1}}(x_i^{(p)}) \in S_p$ that intersects substantially E , such that $y \in \tilde{B}_B = B_{R/2^{p-1}}(x_j^{(p-1)})$ and $B_{R/2^{p-1}}(x_j^{(p-1)}) \subset B_{4R/2^{p-1}}(x_i^{(p)})$ (2), and $B_{4R/2^{p-1}}(x_i^{(p)}) \cap B_R \subset E_\delta$.

It results $w(\tilde{E}_\delta \cap E) = w(E)$. In fact $E \subset \tilde{E}_\delta$ because every point of E is Lebesgue point.

Let now $0 < \alpha < \frac{1}{4}$. We can prove that

$$\begin{aligned} B_{\alpha R/2^{p-1}}(x_i^{(p)}) \cap B_{\alpha R/2^{q-1}}(x_j^{(q)}) &= \emptyset, \\ \forall p, q \in \mathbf{N}, \forall i &= 1, \dots, I_p, \forall j = 1, \dots, I_q . \end{aligned}$$

In fact, if (let $q \geq p$) $B_{\alpha R/2^{p-1}}(x_i^{(p)}) \cap B_{\alpha R/2^{q-1}}(x_j^{(q)}) \neq \emptyset$, because $B_{R/2^p}(x_i^{(p)}) \in R^{(p)}$, $B_{R/2^q}(x_j^{(q)}) \in R^{(q)}$, if $p = q$, then $d(x_i^{(p)}, x_j^{(q)}) > R/2^p$. If $q > p$ then $B_{R/2^{p-1}}(x_i^{(p)}) \in S_p$, $B_{R/2^{q-1}}(x_j^{(q)}) \in S_q$ and $x_j^{(q)} \notin B_{R/2^p}(x_i^{(p)})$ then $d(x_i^{(p)}, x_j^{(q)}) > R/2^p$.

In every case $d(x_i^{(p)}, x_j^{(q)}) > R/2^p$. Let $z \in B_{\alpha R/2^{p-1}}(x_i^{(p)}) \cap B_{\alpha R/2^{q-1}}(x_j^{(q)})$ then

$$R/2^p < d(x_i^{(p)}, x_j^{(q)}) \leq d(x_i^{(p)}, z) + d(z, x_j^{(q)}) \leq \alpha R/2^{p-1} + \alpha R/2^{q-1} \Leftrightarrow \alpha \geq \frac{1}{4} .$$

Then, by taking $0 < \alpha < \frac{1}{4}$ and by recalling that the doubling prop-

(2) In fact, let $z \in B_{R/2^{p-1}}(x_j^{(p-1)})$, then

$$d(z, x_i^{(p)}) \leq d(z, x_j^{(p-1)}) + d(x_j^{(p-1)}, x_i^{(p)}) < R/2^{p-2} + R/2^{p-1} = 3R/2^{p-1} .$$

erty holds also for $B \cap B_R$ (see Prop. 2.10 of [FL2] and Lemma 4 of [C]).

$$\begin{aligned}
 w(E) &= w(\tilde{E}_\delta \cap E) \leq \sum_{B \in \mathcal{U}_{S_p}} w(\tilde{B}_B \cap E) < \\
 &< \delta \sum_{B \in \mathcal{U}_{S_p}} w(\tilde{B}_B \cap B_R) \stackrel{(*)}{\leq} \delta D \sum_{B \in \mathcal{U}_{S_p}} w(B \cap B_R) \stackrel{(*)}{\leq} \\
 &\leq C\delta \sum_{B \in \mathcal{U}_{S_p}} w((\alpha B) \cap B_R) = C\delta w\left(\bigcup_{B \in \mathcal{U}_{S_p}} (\alpha B) \cap B_R\right) \leq C\delta w\left(\bigcup_{B \in \mathcal{U}_{S_p}} B \cap B_R\right) \leq \\
 &= C\delta w\left(\bigcup_{B \in \mathcal{U}_{S_p}} (\tilde{B}_B \cap B_R)\right) = C\delta w(\tilde{E}_\delta) \leq C\delta w(E_\delta).
 \end{aligned}$$

Now we prove the Harnack inequality. We have, in fact:

THEOREM 8. Let u and $-u \in \mathcal{D}\mathcal{G}_m(\Omega)$, $u \geq 0$. Then it exists a constant C dependent on the functional and on \bar{x} such that for any d -ball $B_R(\bar{x}) \equiv B_R$, $B_{\delta R} \subset \Omega$, we have:

$$\sup_{B_{R/2}} u \leq C \inf_{B_{R/2}} u.$$

The proof follows by following propositions:

PROPOSITION 8.1. Let $u \in \mathcal{D}\mathcal{G}_m(\Omega)$, $u \geq 0$. Then for any $q > 0$, it exists $C = C(\bar{x}, q)$ such that

$$(15) \quad \sup_{B_{R/2}} u \leq C \left(\frac{1}{w(B_R)} \int_{B_R} u^q w \, dx \right)^{1/q}.$$

PROOF. Let $q \geq m$. The (15) follows by (10) and Hölder inequality. Let now $0 < q < m$. By (10) for $\sigma \in]0, 1[$ fixed and $k = 0$

$$\begin{aligned}
 \sup_{B_{\sigma R}} u &\leq \frac{C}{(1-\sigma)^{i\Theta/(l-1)}} \left(\frac{1}{w(B_R)} \int_{B_R} u^m w \, dx \right)^{1/m} \leq \\
 &\leq \frac{C}{(1-\sigma)^{i\Theta/(l-1)}} \left(\frac{1}{w(B_R)} \int_{B_R} u^q (\sup_{B_R} u)^{m-q} w \, dx \right)^{1/m} \leq \\
 &\leq \varepsilon \sup_{B_R} u + \frac{C(\varepsilon)}{(1-\sigma)^{i\Theta m/q(l-1)}} \left(\frac{1}{w(B_R)} \int_{B_R} u^q w \, dx \right)^{1/q}
 \end{aligned}$$

(*) By doubling property.

with $0 < \varepsilon < 1$ fixed. (Here we have utilized the inequality $ab \leq \varepsilon a^{m/(m-a)} + C(\varepsilon)b^{m/a}$, $a, b \geq 0$). Now by Lemma 1.1 in [GG1] we have

$$\sup_{B_{\sigma R}} u \leq \frac{C}{(1-\sigma)^{l\Theta m/a(l-1)}} \left(\frac{1}{w(B_R)} \int_{B_R} u^q w \, dx \right)^{1/q}$$

by which, for $\sigma = \frac{1}{2}$, the thesis follows.

PROPOSITION 8.2. Let $u \geq 0$, $-u \in \mathcal{DS}_m(\Omega)$, $\tau > 0$, $\gamma \in]0, 1[$. If $w(\{x \in B_R / u < \tau\}) \leq \gamma w(B_R)$, then:

(16) i) $w(\{x \in B_R / u < \tau/2^{\nu+1}\}) \leq C(\gamma)(\gamma/\nu)^{l(m-1)/2m} w(B_R)$, $\nu \in \mathbb{N}$,

(17) ii) $\inf_{B_{4R}} u \geq \lambda(\gamma)\tau$, with $0 < \lambda(\gamma) < 1$.

PROOF. We observe that

$$w(\{x \in B_R / -u > -\tau\}) = w(\{x \in B_R / u < \tau\}) \leq \gamma w(B_R).$$

So we can apply the (12) of Lemma 5.1 to the function $-u$, for $h > k > -\tau$:

$$\begin{aligned} |h-k|^{m(m+1)/(m-1)} w(B(h, R))^{2m/l(m-1)} &\leq \\ &\leq C(w(B_R))^{2m(1-l)/l(m-1)} \left(\int_{B(k, 2R)} |-u-k|^m w \, dx \right)^{(m+1)/(m-1)} \cdot \\ &\quad \cdot (w(B(k, R)) - w(B(h, R))). \end{aligned}$$

We put $h = -\tau/2^{s+1}$, $k = -\tau/2^s$, $s \in \mathbb{N}$ in the last inequality. We observe that on $B(-\tau/2^s, 2R)$ it is $0 \leq -\tau + \tau/2^s \leq \tau/2^s$. By doubling property for the measure $w(x) \, dx$, we obtain:

$$\begin{aligned} \left(w \left(\left\{ x \in B_R / u < \frac{\tau}{2^{s+1}} \right\} \right) \right)^{2m/l(m-1)} &\leq \\ &\leq C(\gamma)(w(B_R))^{(2m(1-l)+l(m+1))/l(m-1)} \left(w \left(B \left(\frac{-\tau}{2^s}, R \right) \right) - w \left(B \left(\frac{-\tau}{2^{s+1}}, R \right) \right) \right). \end{aligned}$$

Summing up for $s = 0, \dots, \nu$, we have

$$\begin{aligned} \nu \left(w \left(\left\{ x \in B_R / u < \frac{\tau}{2^{\nu+1}} \right\} \right) \right)^{2m/l(m-1)} &< \\ &\leq C(\gamma) (w(B_R))^{(2m(1-l)+l(m+1))/l(m-1)} w(\{x \in B_R \mid u < \tau\}) \leq \\ &\leq C(\gamma) (w(B_R))^{(2m(1-l)+l(m+1))/l(m-1)} \gamma w(B_R). \end{aligned}$$

From which it follows that

$$w \left(\left\{ x \in B_R / u < \frac{\tau}{2^{\nu+1}} \right\} \right) < C(\gamma) \left(\frac{\gamma}{\nu} \right)^{l(m-1)/2m} w(B_R)$$

that is (16).

At last we prove the ii). By (10) applied to $-u$, with $k = -\tau$, $\sigma = \frac{1}{2}$, we have

$$\begin{aligned} \inf_{B_{R/2}} u &\geq \tau - C(w(B_R))^{-\Theta/m} \left(\int_{B(-\tau, R)} |-u + \tau|^m w \, dx \right)^{1/m} (w(B(-\tau, R)))^{(\Theta-1)/m} \geq \\ &\geq \tau \left(1 - C \left(\frac{w(B(-\tau, R))}{w(B_R)} \right)^{\Theta/m} \right) \geq \tau (1 - C\gamma^{\Theta/m}) \end{aligned}$$

(we observe that on $B(-\tau, R)$ it is $0 \leq -u + \tau \leq \tau$).

Put $\gamma_0 = (1/2C)^{m/\Theta}$. Then

$$(18) \quad \inf_{B_{R/2}} u \geq \frac{1}{2} \tau \quad \gamma \in]0, \gamma_0[,$$

Let now $\nu \in \mathbb{N}$ fixed such that $C(\gamma)(\gamma/\nu)^{l(m-1)/2m} < \gamma_0$. $C(\gamma)$ is the constant that appears in (16). Then by (16) and (18) it is

$$(19) \quad \inf_{B_{R/2}} u \geq \frac{1}{2} \frac{\tau}{2^{\nu+1}} = \lambda(\gamma) \tau .$$

We note that $\lambda(\gamma) < 1$. The conclusion of proof of ii) follows making use of the doubling property of the measure $w(x) \, dx$. It exists $\alpha > 0$ (see Lemma 2.10 in [FS]) such that

$$w(\{x \in B_{8R} / u \geq \tau\}) \geq (1 - \gamma)(1/8)^\alpha w(B_{8R}) ,$$

that is equivalent to $w(\{x \in B_{8R}/u < \tau\}) \leq \gamma' \cdot w(B_{8R})$. So by (19) follows (17).

PROPOSITION 8.3. Let $u \geq 0$, $-u \in \mathcal{DS}_m(\Omega)$. Then $\forall q > 0$, $q < 1/C_0$ (see proof), $\exists C > 0$ dependent on q such that

$$(20) \quad \left(\frac{1}{w(B_R)} \int_{B_R} u^q w \, dx \right)^{1/q} \leq C \inf_{B_{4R}} u .$$

PROOF. If $u = 0$ the (20) is evident. Now let $u \neq 0$. Then it exists $t > 0$ such that $w(\{x \in B_R/u(x) \geq t\}) > 0$. Let $\delta \in]0, 1[$ fixed, we consider $E = \{x \in B_R/u(x) \geq \lambda^{i-1} t\} \equiv A_i^{i-1}$ ($\lambda = \lambda(\gamma)$ is the constant that appears in (17)). We can apply the Lemma 7 to the set E . There exist $z \in B_R$ and $\rho > 0$ such that $w(E \cap B(z, \rho)) \geq \delta w(B(z, \rho))$. Then by (17) replacing $\tau = \lambda^{i-1} t$, it follows $u(x) = \lambda^i t$, $\forall x \in B(z, 4\rho)$. Hence or

$$w(\{x \in B_R/u(x) \geq \lambda^i t\}) \geq \frac{1}{C\delta} w(\{x \in B_R/u(x) \geq \lambda^{i-1} t\})$$

or

$$w(\{x \in B_R/u(x) \geq \lambda^i t\}) = w(B_R) .$$

In both case we conclude: if $w(A_i^0) \geq C^{s-1} \delta^s w(B_R)$ for some s (we can suppose $C\delta < 1$) then

$$w(A_i^{s-1}) \geq \frac{1}{C\delta} w(A_i^{s-2}) \geq \dots \geq \frac{1}{(C\delta)^{s-1}} w(A_i^0) \geq \delta w(B_R) ,$$

and hence by (17) we have $\inf_{B_{4R}} u \geq \lambda^s t$.

Choose s so that $w(A_i^0) \geq w(\{x \in B_R/u(x) \geq t\}) \geq C^{s-1} \delta^s$ i.e.

$$s \geq \frac{\log(Cw(A_i^0)/w(B_R))}{\log(C\delta)} .$$

And so $\inf_{B_{4R}} u \geq C_1 t (w(A_i^0)/w(B_R))^{C_0}$ i.e.

$$w(A_i^0)/w(B_R) \leq C_1^{-1} t^{-1/C_0} \left(\inf_{B_{4R}} u \right)^{1/C_0} .$$

Now, since

$$\frac{1}{w(B_R)} \int_{B_R} u^q w \, dx = \frac{1}{w(B_R)} \left(q \int_{\xi}^{+\infty} t^{q-1} w(A_t^0) \, dt + q \int_0^{\xi} t^{q-1} w(A_t^0) \, dt \right),$$

setting $\xi = \inf_{B_{4R}} u$ and $q < 1/C_0$, it holds

$$\begin{aligned} \frac{1}{w(B_R)} \int_{B_R} u^q w \, dx &= \frac{q}{w(B_R)} \int_{\xi}^{+\infty} t^{q-1} w(A_t^0) \, dt + \xi^q \leq \\ &\leq C_1^{-1} q \int_{\xi}^{+\infty} t^{q-1} t^{-1/C_0} (\inf_{B_{4R}} u)^{1/C_0} \, dt + (\inf_{B_{4R}} u)^q = \frac{C_1^{-1} q}{(1/C_0) - q} (\inf_{B_{4R}} u)^q + (\inf_{B_{4R}} u)^q. \end{aligned}$$

From this it follows

$$\left(\frac{1}{w(B_R)} \int_{B_R} u^q w \, dx \right)^{1/q} \leq C \inf_{B_{4R}} u \leq C \inf_{B_{R/2}} u.$$

The conclusion of Theorem 8 now follows by (15).

REMARK. The previous results are true for *quasi-minima* (see [G], [GG2], [BT]). Recall that $u \in W_m^1(\Omega)$ is a quasi-minimum for the functional (1) with constant Q if, for any $\Phi \in W_m^1(\Omega)$, with $\text{supp } \Phi \subset \Omega$

$$(7') \quad F(u, \text{supp } \Phi) \leq Q F(u + \Phi, \text{supp } \Phi).$$

REFERENCES

- [BT] E. DI BENEDETTO - N. S. TRUDINGER, *Harnack inequalities for quasi-minima of variational integrals*, Ann. Inst. Henri Poincaré, **1**, n. 4 (1984).
- [C] A. P. CALDERON, *Inequalities for the maximal function relative to a metric*, Studia Math., **57** (1976).
- [CW] R. R. COIFMAN - G. WEISS, *Analyse harmonique non-commutative sur certain spaces homogenes*, Lecture Notes in Mathematics, **242**, Springer-Verlag, Berlin.

- [DG] E. DE GIORGI, *Sulla differenziabilità e l'analiticità delle estremali degli integrali multipli regolari*, Mem. Accad. Sci. Torino Cl. Fis. Mat. Nat., (3) **3** (1957).
- [FL1] B. FRANCHI - E. LANCONELLI, *Une metrique associée à une classe de operateurs elliptiques dégénérés*, in Proceedings of the Meeting « Linear Partial and Pseudo-Differential Operators », Torino 1982, Rend. Sem. Mat. Univ. e Politec. Torino.
- [FL2] B. FRANCHI - E. LANCONELLI, *Hölder regularity theorem for a class of linear non uniformly elliptic operators with measurable coefficients*, Ann. Scuola Norm. Sup. Pisa, Serie IV, **10**, n. 4 (1983).
- [FL3] B. FRANCHI - E. LANCONELLI, *An embedding theorem for Sobolev spaces related to non smooth vector fields and Harnack inequality*, Comm. Part. Diff. Equat., **9** (13) (1984).
- [FS] B. FRANCHI - R. SERAPIONI, *Pointwise estimates for a class of strongly degenerate elliptic operators: a geometrical approach*, Ann. Scuola Norm. Sup. Pisa, to appear.
- [G] M. GIAQUINTA, *An introduction to the regularity for non linear elliptic systems*, Mathematik Departement eth Zurich und Forschungsinstitut fur Mathematik eth Zurich 1983/84.
- [GG1] M. GIAQUINTA - E. GIUSTI, *On the regularity of the minima of variational integrals*, Acta Math., **148** (1982).
- [GG2] M. GIAQUINTA - E. GIUSTI, *Quasi-minima*, Ann. Inst. Henri Poincaré, **1**, n. 2 (1984).
- [M] G. MODICA, *Quasi minimi di alcuni funzionali degeneri*, Ann. Mat. Pura Appl., **142** (1985).
- [NSW] A. NAGEL - E. M. STEIN - S. WAINGER, *Balls and metrics defined by vector fields - I: Basic properties*, Acta Math., **155** (1985).
- [KS] N. V. KRYLOV - M. V. SAFANOV, *Certain properties of solutions of parabolic equations with measurable coefficients*, Izvestia Akad. Nauk. SSSR, **40** (1980); English transl. Math. USSR Izv., **16** (1981).

Manoscritto pervenuto in redazione l'11 marzo 1988.