

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

MARIO MARINO

ANTONIO MAUGERI

**A remark on the note : “Partial Hölder continuity of
the spatial derivatives of the solutions to nonlinear
parabolic systems with quadratic growth”**

Rendiconti del Seminario Matematico della Università di Padova,
tome 95 (1996), p. 23-28

http://www.numdam.org/item?id=RSMUP_1996__95__23_0

© Rendiconti del Seminario Matematico della Università di Padova, 1996, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques*

<http://www.numdam.org/>

**A Remark on the Note: «Partial Hölder Continuity
of the Spatial Derivatives of the Solutions
to Nonlinear Parabolic Systems with Quadratic Growth».**

MARIO MARINO - ANTONIO MAUGERI(*)

*Dedicated to Professor Sergio Campanato
with our deepest esteem on his 65th birthday*

SUNTO - In questa nota si dimostra che le soluzioni di classe

$$L^2(-T, 0, H^2(\Omega, \mathbb{R}^N)) \cap C^{0,\gamma}(\bar{Q}, \mathbb{R}^N), \quad 0 < \gamma < 1,$$

del sistema (1.1) hanno derivate spaziali parzialmente hölderiane in Q . Si rimuove quindi la condizione richiesta in [1] che la soluzione sia anche di classe $H^1(-T, 0, L^2(\Omega, \mathbb{R}^N))$.

1. - Let Ω be a bounded open subset of \mathbb{R}^n ($n \geq 2$), with sufficiently smooth boundary $\partial\Omega$, for instance of class C^3 , and Q the cylinder $\Omega \times (-T, 0)$ ($T > 0$). In [1] we were concerned with the following second order nonlinear parabolic system of variational type⁽¹⁾:

$$(1.1) \quad - \sum_{i=1}^n D_i a^i(X, u, Du) + \frac{\partial u}{\partial t} = B^0(X, u, Du),$$

(*) Authors' address: Dipartimento di Matematica, Viale A. Doria 6, 95125 Catania.

Research partially supported by M.U.R.S.T. and G.N.A.F.A. of C.N.R.

⁽¹⁾ We follow the notations used in [1].

where $a^i(X, u, p)$, $i = 1, 2, \dots, n$, and $B^0(X, u, p)$ are vectors of \mathbb{R}^N (N integer ≥ 1) measurable in X and continuous in (u, p) . Under the assumptions:

$$(1.2) \quad a^i(X, u, p) \in C^1(\overline{Q} \times \mathbb{R}^N \times \mathbb{R}^{nN}), \quad i = 1, 2, \dots, n,$$

(1.3) *there exists a constant $\nu > 0$ such that*

$$\sum_{h,k=1}^N \sum_{i,j=1}^n \frac{\partial a_h^i(X, u, p)}{\partial p_k^j} \xi_h^i \xi_k^j \geq \nu \sum_{i=1}^n \|\xi^i\|^2$$

for every system $\{\xi^i\}_{i=1,2,\dots,n}$ of vectors of \mathbb{R}^N and for every $(X, u, p) \in \overline{Q} \times \mathbb{R}^N \times \mathbb{R}^{nN}$,

(1.4) *the vectors $\partial a^i / \partial p_k^j$, $i, j = 1, 2, \dots, n$, $k = 1, 2, \dots, N$, are uniformly continuous in $\overline{Q} \times \mathbb{R}^N \times \mathbb{R}^{nN}$,*

(1.5) $\forall (X, u, p) \in \overline{Q} \times \mathbb{R}^N \times \mathbb{R}^{nN}$ *we have:*

$$\sum_{k=1}^N \sum_{j=1}^n \left\| \frac{\partial a^i}{\partial p_k^j} \right\| \leq M, \quad i = 1, 2, \dots, n,$$

$$\|a^i\| + \sum_{s=1}^N \left\| \frac{\partial a^i}{\partial x_s} \right\| + \sum_{k=1}^N \left\| \frac{\partial a^i}{\partial u_k} \right\| \leq M \left(1 + \sum_{j=1}^n \|p^j\| \right),$$

$$i = 1, 2, \dots, n,$$

(1.6) $\forall (X, u, p) \in \overline{Q} \times \mathbb{R}^N \times \mathbb{R}^{nN}$ *we have:*

$$\|B^0(X, u, p)\| \leq g^0(X) + M \sum_{j=1}^n \|p^j\|^2,$$

$$g^0(X) \in L^q(Q), \quad q > n + 2,$$

we established the partial Hölder continuity in Q of the spatial gradient of the weak solutions u of class

$$L^2(-T, 0, H^2(\Omega, \mathbb{R}^N)) \cap$$

$$\cap H^1(-T, 0, L^2(\Omega, \mathbb{R}^N)) \cap C^{0,\gamma}(\overline{Q}, \mathbb{R}^N), \quad 0 < \gamma < 1,$$

to the system (1.1) (see [1], Theorem 4.1).

The aim of this work is to find again the partial Hölder continuity of the spatial derivatives of the weak solutions to the system (1.1), under the assumption that these solutions belong merely to $L^2(-T, 0, H^2(\Omega, \mathbb{R}^N)) \cap C^{0,\gamma}(\bar{Q}, \mathbb{R}^N)$. In fact it is possible to remove the condition $u \in H^1(-T, 0, L^2(\Omega, \mathbb{R}^N))$ thanks to an interpolation theorem due to Nirenberg [3] (see also Miranda [2]), which ensures the result $D_i u \in L^s(Q, \mathbb{R}^N)$ under the mere assumption $u \in L^2(-T, 0, H^2(\Omega, \mathbb{R}^N)) \cap C^{0,\gamma}(\bar{Q}, \mathbb{R}^N)$.

2. - It is well known the following interpolation result

LEMMA 2.1. *If $u \in H^2(\Omega, \mathbb{R}^N) \cap C^{0,\gamma}(\bar{\Omega}, \mathbb{R}^N)$, $0 < \gamma < 1$, then*

$$(2.1) \quad D_i u \in L^s(\Omega, \mathbb{R}^N), \quad i = 1, 2, \dots, n,$$

and

$$(2.2) \quad \int_{\Omega} \|D_i u\|^s dx \leq c_1 [u]_{\gamma, \bar{\Omega}}^{s-\frac{2}{\gamma}} \int_{\Omega} \sum_{i,j=1}^n \|D_{ij} u\|^2 dx + c_2 [u]_{\gamma, \bar{\Omega}}^s,$$

where $s = (2(2 - \gamma))/(1 - \gamma)$, c_1 and c_2 are constants depending on Ω , γ , n (2).

See [3], Theorem 1' (with $\beta = \gamma$, $m = r = 2$, $j = 1$, $a = (1 - \gamma)/(2 - \gamma)$) (3).

LEMMA 2.2. *If $u \in L^2(-T, 0, H^2(\Omega, \mathbb{R}^N)) \cap C^{0,\gamma}(Q, \mathbb{R}^N)$, $0 < \gamma < 1$, then*

$$(2.3) \quad D_i u \in L^s(Q, \mathbb{R}^N), \quad i = 1, 2, \dots, n,$$

and

$$(2.4) \quad \int_Q \|D_i u\|^s dX \leq c_1 [u]_{\gamma, \bar{Q}}^{s-\frac{2}{\gamma}} \int_Q \sum_{i,j=1}^n \|D_{ij} u\|^2 dX + c_2 T [u]_{\gamma, \bar{Q}}^s,$$

$$(2) [u]_{\gamma, \bar{\Omega}} = \sup_{\substack{x, y \in \bar{\Omega} \\ x \neq y}} \frac{\|u(x) - u(y)\|}{\|x - y\|^\gamma}.$$

(3) We would like to thank warmly Prof. Francesco Guglielmino for pointing to us this result.

where $s = (2(2 - \gamma))/(1 - \gamma)$, c_1 and c_2 are the constants (depending on Ω , γ , n) that appear in the (2.2) (4).

PROOF. If $u \in L^2(-T, 0, H^2(\Omega, \mathbb{R}^N)) \cap C^{0,\gamma}(\bar{Q}, \mathbb{R}^N)$, $0 < \gamma < 1$, for a.e. $t \in (-T, 0)$ it results:

$$u(x, t) \in H^2(\Omega, \mathbb{R}^N) \cap C^{0,\gamma}(\bar{\Omega}, \mathbb{R}^N),$$

then, by Lemma 2.1, we have:

$$D_i u(x, t) \in L^s(\Omega, \mathbb{R}^N), \quad s = \frac{2(2 - \gamma)}{1 - \gamma}, \quad i = 1, 2, \dots, n,$$

and

$$(2.5) \quad \int_{\Omega} \|D_i u(x, t)\|^s dx \leq c_1 [u]_{\gamma, \bar{Q}}^{s-\frac{2}{\gamma}} \int_{\Omega} \sum_{i,j=1}^n \|D_{ij} u(x, t)\|^2 dx + c_2 [u]_{\gamma, \bar{Q}}^s.$$

Now, by integrating with respect to t both the sides of (2.5) in the interval $(-T, 0)$, we achieve the conclusion.

From the lemma above the partial Hölder continuity of the spatial derivatives of the weak solutions to system (1.1) easily follows.

THEOREM 2.1. *If $u \in L^2(-T, 0, H^2(\Omega, \mathbb{R}^N)) \cap C^{0,\gamma}(\bar{Q}, \mathbb{R}^N)$, $0 < \gamma < 1$, is a weak solution in Q to system (1.1) (5) and if conditions (1.2)-(1.6) are fulfilled, then there exists a set $Q_0 \subset Q$, closed in Q , such that*

$$D_i u \in C^{0,\mu}(Q \setminus Q_0, \mathbb{R}^N), \quad \forall \mu < 1 - \frac{n+2}{q}, \quad i = 1, 2, \dots, n,$$

$$(4) \quad [u]_{\gamma, \bar{Q}} = \sup_{\substack{X, Y \in \bar{Q} \\ X \neq Y}} \frac{\|u(X) - u(Y)\|}{d^\gamma(X, Y)}, \quad d(X, Y) = \max\{\|x - y\|, |t - \tau|^{1/2}\},$$

$$X = (x, t), \quad Y = (y, \tau).$$

(5) In the sense that it results:

$$\int \left\{ \sum_{i=1}^n (a^i(X, u, Du) | D_i \varphi) - \left(u \mid \frac{\partial \varphi}{\partial t} \right) \right\} dX = \int (B^0(X, u, Du) | \varphi) dX,$$

$$\forall \varphi \in C_0^\infty(Q, \mathbb{R}^N).$$

and

$$\mathcal{H}_{n+2-r}(Q_0) = 0,$$

for every $r \in (2, (2/\gamma_0) \wedge (2(n+2))/(n+2-2\gamma))$ ⁽⁶⁾.

PROOF. From the assumption

$$u \in L^2(-T, 0, H^2(\Omega, \mathbb{R}^N)) \cap C^{0,\gamma}(\bar{Q}, \mathbb{R}^N)$$

and from the estimate $(2(2-\gamma))/(1-\gamma) > 4$, it follows, by Lemma 2.2:

$$D_i u \in L^4(Q, \mathbb{R}^N), \quad i = 1, 2, \dots, n;$$

then, taking into account that $B^0(X, u, p)$ satisfies (1.6), we have:

$$(2.6) \quad B^0(X, u, Du) \in L^2(Q, \mathbb{R}^N).$$

On the other hand from assumption (1.5) on $a^i(X, u, p)$ we obtain

$$\begin{aligned} \|D_i a^i(X, u, Du)\| &\leq \left\| \frac{\partial a^i}{\partial x_i} \right\| + \sum_{k=1}^N \left\| \frac{\partial a^i}{\partial u_k} \right\| \|D_i u_k\| + \\ &+ \sum_{j=1}^n \sum_{k=1}^N \left\| \frac{\partial a^i}{\partial p_k^j} \right\| \|D_{ij} u_k\| \leq c \left(1 + \sum_{j=1}^n \|D_j u\|^2 + \sum_{i,j=1}^n \|D_{ij} u\| \right) \end{aligned}$$

from which

$$(2.7) \quad D_i a^i(X, u, Du) \in L^2(Q, \mathbb{R}^N), \quad i = 1, 2, \dots, n.$$

Now let us recall that u is a solution in Q of system (1.1); then (see footnote ⁽⁵⁾), $\forall \varphi \in C_0^\infty(Q, \mathbb{R}^N)$, it results:

$$\int_Q \left(u \mid \frac{\partial \varphi}{\partial t} \right) dX = - \int_Q \left(\sum_{i=1}^n D_i a^i(X, u, Du) + B^0(X, u, Du) \mid \varphi \right) dX,$$

⁽⁶⁾ \mathcal{H}_{n+2-r} is the $(n+2-r)$ -dimensional Hausdorff measure with respect to the parabolic metric:

$$d(X, Y) = \max \{ \|x - y\|, |t - \tau|^{1/2} \}, \quad X = (x, t), \quad Y = (y, \tau).$$

γ_0 is the real number in the interval $((n-2)/n, 1)$ that appears in the statement of Theorem 3.1 in [1].

from which, by means of (2.6) and (2.7), we reach

$$\exists \frac{\partial u}{\partial t} \in L^2(Q, \mathbb{R}^N).$$

Then u verifies all the assumptions of Theorem 4.1 in [1] and therefore the conclusion follows by this theorem.

REMARK 2.1. The Theorem 2.1 can be proved in a «direct way», following the technique used in [1], that is without applying the Theorem 4.1 of [1]. We preferred to make use of Theorem 4.1 in [1] for the sake of shortness.

REFERENCES

- [1] M. MARINO - A. MAUGERI, *Partial Hölder continuity of the spatial derivatives of the solutions to nonlinear parabolic systems with quadratic growth*, Rend. Sem. Mat. Padova, **76** (1986), pp. 219-245.
- [2] C. MIRANDA, *Su alcuni teoremi di inclusione*, Annales Polonici Math., **16** (1965), pp. 305-315.
- [3] L. NIRENBERG, *An extended interpolation inequality*, Ann. Sc. Norm. Sup. Pisa (3), **20** (1966), pp. 733-737.

Manoscritto pervenuto in redazione il 25 agosto 1994.