

RENDICONTI
del
SEMINARIO MATEMATICO
della
UNIVERSITÀ DI PADOVA

V. ALEXANDRU

A. POPESCU

N. POPESCU

Completion of r.t. extensions of local fields (II)

Rendiconti del Seminario Matematico della Università di Padova,
tome 100 (1998), p. 57-66

http://www.numdam.org/item?id=RSMUP_1998__100__57_0

© Rendiconti del Seminario Matematico della Università di Padova, 1998, tous droits réservés.

L'accès aux archives de la revue « Rendiconti del Seminario Matematico della Università di Padova » (<http://rendiconti.math.unipd.it/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>

Completion of r.t. Extensions of Local Fields (II).

V. ALEXANDRU (*) - A. POPESCU (**) - N. POPESCU (***) (*)

Let (K, v) be a local field and let $(K(X), w)$ be a residual transcendental extension of it. The following question arises: describe the completion of $(K(X), w)$ (see [5]. ch VI Sect. 6) using (K, v) and related concepts. In the first part of this work (see [1]) had been considered the particular, but important case, when $w = w_0$ is the Gauss r.t. extension of v to $K(X)$. Thus $(\widehat{K(X)}, \tilde{w}_0)$ is described as the subfield of $(K\{\{X\}\}, u)$ consisting of all Laurent power series whose positive parts are almost periodic (see [1], Theorem 2.4).

In this paper we consider the general case when $(K(X), w)$ is any r.t. extension of (K, v) . In Section 2 we try to give a «combinatorial» description of elements from $(\widehat{K(X)}, \tilde{w})$, using a natural inclusion $\widehat{K(X)} \rightarrow K\{\{X - a, \delta\}\}$. For that we use some elementary remarks in section 1 and a characterization of r.t. extensions of a valuation given in [3]. In Section 3 we describe $(\widehat{K(X)}, \tilde{w})$ as a finite extension of $(\widehat{K(r)}, \tilde{w}_0)$, where r is a suitable element of $K(X)$ and w_0 is the Gauss r.t. extension of v to $K(r)$. Finally, in Section 4 we study the conditions when $(\widehat{K(X)}, \tilde{w})$ is coincident to the completion of $(K'(X), w')$, where K' is a suitable finite extension of K and w' an extension of w to $K'(X)$ and moreover w' is a slightly modified Gauss r.t. extension of v' to $K'(X)$ (here v' is the unique extension of v to K').

(*) Indirizzo dell'A.: University of Bucharest, Department of Mathematics, Str. Academiei 14, 70109, Bucharest, Romania.

(**) Civil Engineering University of Bucharest, 124, B-dul Lacul Tei, Department of Mathematics, R-72302, Bucharest, Romania.

(***) Institut of Mathematics of the Romanian Academy, P.O. Box I-764, Ro-70700 Bucharest, Romania.

1. – Taylor's type embedding of rational function field.

1) Let K be a field and let $K[X]$ be the polynomial ring of one variable over K . Let \bar{K} be an algebraic closure of K , a an element in \bar{K} and let $f(X)$ be the monic minimal polynomial of a with respect to K . Denote $K' = K(a)$. For any $g(X) \in K[X]$, we consider the Taylor's expansion $T(g) = \sum_{i=0}^n (g^{(i)}/i!)(a)(X-a)^i$ in $K'[X]$, where $n = \text{deg} g$. In this way we obtain a ring embedding $T: K[X] \rightarrow K'[X-a] = K'[X]$. Extend T in an usual manner to a field embedding $T: K(X) \rightarrow K'(X-a) = K'(X)$ and call a rational function from the range of T a K -rational function in $K'(X)$ with respect to a . One calls T the Taylor's embedding of $K(X)$ in $K'(X)$.

2) For any natural number $k \geq 0$ one consider the following upper triangular $(k+1) \times (k+1)$ -matrix:

$$\mathcal{A}_k = (a_{ij}) = \begin{cases} 0, & \text{if } i > j, \\ 1, & \text{if } i = j, \\ \binom{i}{j-1} a^{j-i}, & \text{if } i < j, \end{cases}$$

where $i, j \in \{0, 1, \dots, k\}$ and $\binom{n}{m} = \frac{n!}{m!(n-m)!}$, $n, m \in \mathbb{N}$, $m \leq n$.

PROPOSITION. 1.1. a) A polynomial in $K'[X]$ of the form $G(X) = \sum_{i=0}^n A_i (X-a)^i$ is a K -polynomial in $K'[X]$ if and only if the column vector $\mathcal{A}_n (A_0, A_1, \dots, A_n)^t = (a_0, a_1, \dots, a_n)^t$ has its components in K , i.e. $a_0, \dots, a_n \in K$. (Here " t " means the transposed). One has $T\left(\sum_{i=0}^n a_i X^i\right) = g(X)$.

b) A rational function in $K'(X)$,

$$\frac{P(X)}{Q(X)} = \frac{\sum_{i=0}^m P_i (X-a)^i}{\sum_{j=0}^n Q_j (X-a)^j}$$

is a K -function in $K'(X)$ if and only if P and Q are K -polynomials in $K'(X)$.

The proof is based on the Newton's binomial formula for $(X - a)^i$, $i = 0, 1, \dots$

3) A sequence Q_n, Q_{n-1}, \dots, Q_0 of elements of K' not all zero, is called a K -sequence if there exists a K -rational function P/Q in $K'(X)$ such that $Q(X) = \sum_{i=0}^n Q_i(X - a)^i$.

PROPOSITION 1.2. *Let P be an element of $K'(X)$ and let $\sum_{i \geq i_0} A_i(X - a)^i$ be the associated power series in an usual manner. This power series represent a rational function in $K(X)$ if and only if there exists a natural number m and a K -sequence Q_n, \dots, Q_0 in K' such that:*

- 1) $\sum_{i=0}^n Q_i A_{k-i} = 0$, for every $k > m$, and
- 2) if one denote $P_j = \sum_{i=0}^n Q_i A_{j-i}$, $j = 0, 1, \dots, m$, the components of the matrix $\mathfrak{A}_m(P_0, \dots, P_m)^t$ are in K .

PROOF. It is enough to identify the coefficients in the equality $P(X) = Q(X) \sum_{i \geq i_0} A_i(X - a)^i$ and apply Proposition 1.1.

2. - $w_{(a, \delta)}$ -completion of $K(X)$.

1) Let (K, v) be a local field (see [9] i.e a valued field K complete relative to a rank one valuation v). Denote by \overline{K} a fixed algebraic closure of K and denote by \overline{v} the unique extension of v to \overline{K} . If $a \in \overline{K}$ denote $\deg a = [K(a) : K]$, the *degree* of a with respect to K . Let $a \in \overline{K}$ and let $\delta \in \mathbf{Q}$, the additive group of rational numbers. We say that (a, δ) is *minimal pair* if for any $b \in \overline{K}$, the condition $\overline{v}(a - b) \geq \delta$ implies $\deg b \geq \deg a$. According to [8], the number $\delta(a) = \sup_b (\overline{v}(a - b), b \in \overline{K}, \deg b < \deg a)$ is called the *main invariant* of a . Then (a, δ) is a minimal pair if and only if $\delta > \delta(a)$.

An extension w of v to $K(X)$ is called residual transcendental (r.t.- extension) if the residue field of w is a transcendental extension of the residue field of v .

Let (a, δ) be a minimal pair. Denote $w = w_{(a, \delta)}$ the extension of v to $K(X)$ defined as follows: if $P(X) \in K[X]$, then

$$w(P(X)) = \inf_{i=0}^m \left\{ \bar{v} \left(\frac{P^{(i)}(a)}{i!} \right) + i\delta \right\}$$

where $m = \deg P(X)$, According to [3], $w_{(a, \delta)}$ (extended in an usual manner to $K(X)$) is a r.t. extension of v to $K(X)$ and by ([3], Theorem 2.1) any r.t. extension of u to $K(X)$ is the form $w_{(a, \delta)}$ for a suitable minimal pair (a, δ) .

2) Let $a \in \bar{K}$ and $K' = K(a)$. Denote v' the restriction of \bar{v} to K' . According to [1] denote by $K' \{ \{X - a, \delta\} \}$ the valued field of δ -formal Laurent series over the local field (K', v') , where $\delta > \delta(a)$ is a rational number. According to ([1], § 2) we can also consider the embedding $\omega: K'(X) = K'(X - a) \rightarrow K' \{ \{X - a, \delta\} \}$. In this way $K'(X - a)$ can be viewed as a subfield of $K' \{ \{X - a, \delta\} \}$ and by the composition $\omega \circ T$ we can view $K(X)$ as a subfield of $K' \{ \{X - a, \delta\} \}$. As usual (see [1]) denote u the natural valuation of $K' \{ \{X - a, \delta\} \}$. Now if $w = w_{(a, \delta)}$ is the valuation on $K(X)$ defined in the previous point, then it is easy to see that for any $\alpha \in K(X)$ one has $w(\alpha) = u((T \circ \omega)(\alpha))$. Then, through $\omega \circ T$, $K' \{ \{X - a, \delta\} \}$ becomes an extension of the valued field $(K(X), w)$. In what follows we consider $(K(X), w)$ as a subfield of $K' \{ \{X - a, \delta\} \}$ and the mapping $\omega \circ T$ does not appear explicitly.

Because $K' \{ \{X - a, \delta\} \}$ is complet, the topological adherence of $K(X)$ in it is exactly the completion of $(K(X), w)$. Denote by $(\bar{K}(X), \tilde{w})$ this completion.

3) Using now the results of [1], we try to describe the elements of $K \{ \{X - a, \delta\} \}$ which lie in $\bar{K}(\bar{X})$.

If $S = \sum_{i \in \mathbb{Z}} b_i (X - a)^i$ is a Laurent series in $K' \{ \{X - a, \delta\} \}$ and n is a naturd number, the partial sum: $S_{i(n)} = \sum_{i \geq i(n)} b_i (X - a)^i$ is called a n -section of S if $u(S - S_{i(n)}) > n$. To every series $S_{i_0} = \sum_{i \geq i_0} a_i (X - a)^i$ we associate the sequence $S_{i_0}^* = \{a_i^*\}_{i \in \mathbb{Z}}$ such that $a_i^* = a_i$ if $i \geq i_0$, and $a_j^* = 0$ if $j < i_0$.

If $D = \{d_0, d_1, \dots, d_q\}$ is a finite sequence of elements in K' and $A = \{a_i\}_{i \in \mathbb{Z}}$ is an infinite set of elements in K' , we define a «multiplication» between D and A in the following way:

$$[D, A] = C = \{c_k\}_{k \in \mathbb{Z}}, \quad \text{where } c_k = \sum_{i=0}^q d_i a_{k-i}.$$

If $C = \{c_k\}_{k \in \mathbb{Z}}, B = \{b_k\}_{k \in \mathbb{Z}}$ are subsets of K' , denote $u(C, B) = \inf_{k \in \mathbb{Z}} \{v(c_k - b_k) + k\delta\}$. It is possible that $u(C, B) = \infty$.

A Laurent series $S = \sum_{i \in \mathbb{Z}} a_i (X - a)^i$ in $K' \{\{X - a, \delta\}\}$ is called *K-local periodic series* if for any $n \in \mathbb{N}$, there exists a n -section $S_{i(n)} = \sum_{i \geq i(n)} a_i (X - a)^i$ of it and two sequences in $K', B = \{b_0, b_1, \dots, b_p\}$ and $D = \{d_0, d_1, \dots, d_q\}, D \neq 0$ such that $\alpha_p(b_0, b_1, \dots, b_p)^t$ and $\alpha_q(d_0, \dots, d_q)^t$ are vectors with components in K (for the definition of α_p see § 1) and that $u([D, S_{i(n)}^*], B) > n + u(D)$, where $u(D) = \inf_i \{v(d_i) + i\delta\}$.

THEOREM 2.1. *An element $S = \sum_{i \in \mathbb{Z}} a_i (X - a)^i$ of $K' \{\{X - a, \delta\}\}$ belongs to $\widetilde{K(X)}$ if and only if it is a K -local periodical series.*

PROOF. Let S be an element in $\widetilde{K(X)}$, n a natural number and $S_{i(n)}$ a n -section of S . Let

$$\alpha = \frac{P}{Q} = \frac{b_0 + b_1(x - a) + \dots + b_p(x - a)^p}{d_0 + d_1(x - a) + \dots + d_q(x - a)^q}$$

be an element in $K(X)$ such that $u(S - \alpha) > n$. Since $u(S - S_{i(n)}) > n$, then $u(S_{i(n)} - \alpha) > n$. If $B = (b_k)_k, D = (d_k)_k$, then it is easy to see that S is K -local periodic.

Conversely, let us assume that S is a K -local periodic element in $K' \{\{X - a, \delta\}\}$. If $B = (b_0, \dots, b_p)$ and $D = (d_0, \dots, d_q)$ are as in the definition of a local periodic series S , for a natural number n , then

$$\alpha = \frac{P}{Q} = \frac{a_0 + a_1(X - a) + \dots + b_p(X - a)^p}{d_0 + d_1(X - a) + \dots + d_q(X - a)^q}$$

belongs to $K(X)$ and $u(S - \alpha) > n$ etc.

REMARK 2.2. The above results try to give a «combinatorial» description of elements of $K' \{ \{X' - a, \delta\} \}$ which lie in $\overline{K(X)}$. We remark that even in the case $a = 0$ this result is not as powerful as the one given in ([1], Theorem 2.3).

3. - Another type of characterization of the completion of $(K(X), w_{(a, \delta)})$.

1) The notation and the hypotheses are as in the section 2. Let f be the minimal and monic polynomial of a with respect to K . According to ([3], Theorem 2.1) and [7], there exists a natural number e , and a polynomial $h(X)$ in $K[X]$, with $\deg h < n = \deg f$ such that:

i) The rational function $r = r(X) = (f^e/h) \in K(X)$ is such that $w(r) = 0$ and t , the image of r in the residue field of w , is transcendental over the residue field of v , and

ii) r is of minimal degree with these properties:

Let us denote by w_0 the restriction of w to $K(r)$ and by $e(w|w_0)$, $f(w|w_0)$ the ramification index and respectively the inertia degree of w over w_0 . Using ([3], Theorem 2.1, Corollaries 2.4, 2.5, and 2.6) and ([4]), Theorem 4.5) we conclude that $[K(X) : K(r)] = ne = e(w|w_0) \cdot f(w|w_0)$, so w is the unique extension of w_0 to $K(X)$. The valuation w_0 on $K(r)$ is exactly the Gauss extension of v to $K(r)$ and from [1] we know a characterization for the completion $\overline{K(r)}$. It is easy to see that $K(X)$ and $\overline{K(r)}$ are linear disjoint over $K(r)$, i.e. $f^e(Y) - h(Y) r(X)$, the minimal polynomial of X over $K(r)$ is also irreducible over $\overline{K(r)}$, so $\overline{K(X)}$, the completion of $K(X)$ with respect to w , is exactly $\overline{K(r)}(X)$.

It is clear that any element in $K(X)$ is of the form $\sum_{i=0}^{ne-1} a_i(r) X^i$, where $a_i(r) \in K(r)$. Let us consider now the embedding ω (see [1]) of $K(r)$ in $K\{\{r\}\}$, the local field of 0-formal Laurent series in the variable r over K , (see [1]), and describe any element from $\overline{K(X)}$ as being of the form $\sum_{i=0}^{ne-1} b_i(r) X^i$, where $b_i(r) \in \overline{K(r)}$.

Using these last remarks above it is not difficult to see that any element in $\overline{K(X)}$ may be described as being a Laurent series of the type $\sum_{i \in \mathbb{Z}} s_i(X) r^i$, where $s_i(X) \in K[X]$, $\deg s_i < ne$, the values of coefficients of $s_i(X)$ are inferior bounded (for all $i \in \mathbb{Z}$) and when $i \rightarrow -\infty$ the values of them tendes to ∞ . Even the sequence $\{s_i(X)\}_i$ is almost periodic and we

have the equality

$$w\left(\sum_{i \in \mathbb{Z}} s_i(X) r^i\right) = \inf_{i \in \mathbb{Z}} w(s_i(X)).$$

4. – Some properties of the algebraic extension $K(X) \hookrightarrow K'(X)$.

The notations and definitions are as in section 1. In this section we are interested in some properties of the extension $\overline{K'(X)}/\overline{K(X)}$. Particularly we derive some conditions such that $\overline{K(X)}$ is coincident to $\overline{K'(X)}$.

1) Let $a \in \overline{K}$ and f the monic minimal polynomial of a with respect to K . Let $S = \{a = a_1, \dots, a_n\}$ be all roots of f in \overline{K} . Let δ be a rational number such that (a, δ) is a minimal pair, i.e. $\delta > \delta(a)$. Denote $\gamma = \sum_{i=1}^n \inf(\delta, \bar{v}(a - a_i))$ and let e be the smallest non-zero natural number such that $e\gamma \in G_{v'}$, the value group of v' and by e' denote the smallest non-zero natural number such that $e' \delta \in G_{v'}$.

Denote $w = w_{(a, \delta)}$ the r.t. extension of v to $K(X)$ defined by the minimal pair (a, δ) . According to ([3], Theorem 2.1) relative to the value group G_w of w one has:

(1)
$$\gamma = w(f); \quad G_w = G_{v'} + Z\gamma.$$

In what follows we assume that a is separable over k . We remark, according to ([4], Theorem 3.1) that any r.t. extension w of v to $K(X)$ is of the form $w = w_{(a, \delta)}$ where a is separable over K .

Denote s the number of all conjugates a' of a over K such that $\bar{v}(a - a') \geq \delta$. As usual by $e(v' | v)$ and $f(v' | v)$ we denote the ramification index and the residual degree of v' with respect to v (see [10]).

THEOREM 4.1. *With the above notations and hypothesis one has:*

- 1) s divides $n = \deg f = [K' : K]$,
- 2) There exists exactly n/s distinct extensions of w to $K'(X)$.
- 3) If w' is an extension of w to $K'(x)$ and $d = e(w' | w)$ then d is the greatest common divisor between s and e' , and $f(w' | w) = s/d$.

PROOF. Let $K'' = K(a_1, \dots, a_n)$. Then K''/K is a Galois extension and denote by $G = \text{Gal}(K''/K)$. Let $H = \{g \in G/\bar{v}(a - g(a)) \geq \delta\}$ and $S(H) = \{g(a), g \in H\}$. Then H is a subgroup of G and $S(H)$ contains exactly s elements. Since G acts transitively on $S(H)$ then s divides n , as claimed.

2) and 3). Let $t = n/s$ and let $H = H_1, \dots, H_t$ be all conjugates of H in G . Denote by $S(H_i) = \{g(a) | g \in H_i\}$, $1 \leq i \leq t$ and $S(H_i) = \{a_i^{(1)}, \dots, a_i^{(s)}\}$. Denote \bar{w}_i , $1 \leq i \leq t$, the r.t. extension of \bar{v} to $\bar{K}(X)$, defined by the minimal pair $(a_i^{(1)}, \delta)$. Then according to ([4], Theorem 2.2), the extensions \bar{w}_i , $1 \leq i \leq t$ are pair distinct and are all extensions of w to $\bar{K}(X)$. Denote w_i the restriction of \bar{w}_i to $K'(X)$. It is clear that the set of w_i , $1 \leq i \leq t$ contains all possible non - equivalent extension of w to $K'(X)$. We assert that w_i , $1 \leq i \leq t$ are all pair distinct. Using the above notations and relations it will be enough to show the following equalities

$$e(w_i | w) = d, \quad f(w_i | w) = s/d, \quad 1 \leq i \leq t.$$

Because of symmetry it is enough to show the above equalities for $w_1 = w'$. Now we denote $\varepsilon = \gamma - s\delta$. It is clear that $\varepsilon = \sum v(a - a_j)$, where a_j runs all conjugates a_j of a such that $v(a - a_j) < \delta$. By a permutation of a_i , $1 \leq i \leq n$ we can assume that $\bar{v}(a - a_i) \geq \delta$ if $1 \leq i \leq s$ and $v(a - a_i) < \delta$ if $s < i \leq n$. We assert that $\varepsilon \in G_{v'}$. Indeed, one has $(f^{(s)}/s!)(a) \in K''$. But $(f^{(s)}/s!)(a) = \sum_{2 \leq i_1 < i_2 < \dots < i_{n-s} \leq n} (a - a_{i_1}) \dots (a - a_{i_{n-s}})$. Then according to definition of s one has $v'((f^{(s)}/s!)(a)) = \bar{v}((a - a_{s+1}) \dots (a - a_n)) = \varepsilon \in G_{v'}$, as claimed.

Now we use the definitions of e and e' and the equality $e' \gamma = se' \delta + e' \varepsilon$ in order to derive that $e' \gamma \in G_{w'}$. It follows that $e' = ed$ for a natural number d . Since by (1), $G_w = G_{v'} + Z\gamma$ and $G_{w'} = G_v + Z\delta$, it follows that $e(w' | w) = d$. But d is a divisor of s since $sed = e\gamma - e\varepsilon \in G_{v'}$ and so se is a multiple of $e' = ed$. It is easy to see that d is exactly the greatest common divisor between s and e' (use the definition of e').

Now let us compute $f(w' | w)$. Denote by b an element in \bar{K} such that $\bar{v}(b) = \delta$ and by $k_{\bar{w}}$, $k_{\bar{v}}$, etc. the residue field of \bar{w} ; \bar{v} , etc. Then θ , the image of $(X - a)/b$ in $k_{\bar{w}}$, is transcendental over $k_{\bar{v}}$ and $k_{\bar{w}} = k_{\bar{v}}(\theta)$ (see [3], Theorem 2.1). Now consider the element $r \in k(X)$ defined in section 3. Then $k_w = k_{v'}(t) \subset k_{w'} \subset k_{\bar{w}}$. It is not difficult to see that t is a polynomial of θ of degree se . If $l(X) \in K'[X]$ is such that $w'(l(X)) = e' \delta$ then t' , the image in $k_{w'}$ of $(X - a)^{e'}/l(X)$, is transcendental over $k_{v'}$. The degree of t'

as polynomial of θ is exactly e' . Then $[k_w : k_w] = f(w|w') = se/e' = s/d$. The proof of Theorem 4.1 is complete.

COROLLARY 4.2. *The extension $(K(X), w) \subset (K'(X), w')$ is immediate (see [11]) if and only if $s = 1$. In this case $X - a \in (\overline{K(X)}, \tilde{w}) = (\overline{K(X)}, \tilde{w}')$. This is true if and only if $\delta > \omega(a) = \sup(\bar{v}(a - a'))$, where a' runs all conjugates of a distinct to a .*

Now we indicate some cases when the conditions of Corollary 4.2 are verified

COROLLARY 4.3. *Let $p = \text{char } k_u > 0$. If $n = [K' : K]$ is relative prime to p then $s = 1$.*

PROOF. Indeed, in this case $\delta(a) = \omega(a)$. For that let $S(a) = \{a' \in S, \bar{v}(a - a') \geq \omega(a)\}$ and let h be the cardinal of $S(a)$. Let H be the subgroup of $G = \text{Gal}(K''|K)$ defined by: $H = \{g \in G | g(a) \in S(a)\}$. If $m = |G/H|$ then m divides n and let $H = H_1, \dots, H_m$ be all conjugates of H in G . If $H_i = g_i H g_i^{-1}$, let $a_i = g_i(a)$ and let $S(a_i)$ defined as $S(a)$. Then the $\{S(a_i)\}_i$ gives a partition of S . Denote $b_i = \sum a'$, where a' runs over $S(a_i)$, $1 \leq i \leq m$. It is clear that G acts transitively on the set $\{b_1, \dots, b_m\}$. It follows that the coefficients of the polynomial $\varphi(X) = (X - b_1) \dots (X - b_m)$ belongs to K . Hence the element b_1 is such that $[K(b_1) : K] \leq m \leq n$. Since one has $\bar{v}(a - b_1/h) \geq \omega(a)$ then, according to definition of $\delta(a)$, one has necessarily $\delta(a) = \omega(a)$.

REMARK 4.4. It can be proved that the polynomial φ is irreducible and the pair (a, b_1) is admissible (see [8]).

COROLLARY 4.5. *If a is an uniformising element of K' and $e(v/v')$ is relatively prime to p , or $f(v|v') = n = [k' : k]$, $\bar{v}(a) = 0$ and $k_v = k_v(\alpha)$, where α is the image of a in k_v , then $s = 1$.*

The proof follows by [9], since in both cases one has $\omega(a) = \delta(a)$.

REFERENCES

- [1] V. ALEXANDRU - A. POPESCU - N. POPESCU, *Completion of r.t. extensions of local fields (I)*, Math. Z., **221** (1996), pp. 675-682.
- [2] V. ALEXANDRU - N. POPESCU, *Some elementary remarks about n-local fields*, Rend. Sem. Mat. Univ. Padova, **92** (1994), pp. 17-28.
- [3] V. ALEXANDRU - N. POPESCU - A. ZAHARESCU, *A theorem of characterization of residual transcendental extensions of a valuation*, J. Math. Kyoto Univ., **28**, No. 4 (1988), pp. 579-592.
- [4] V. ALEXANDRU - N. POPESCU - A. ZAHARESCU, *Minimal pairs of definition of a residual transcendental extension of a valuation*, J. Math. Kyoto Univ., **30**, No. 2 (1990), pp. 207-225.
- [5] N. BOURBAKI, *Algèbre Commutative*, Ch. V, VI, Herman, Paris (1964).
- [6] A. N. PARSHIN, *Abelian covering of arithmetic schemes (russian)*, Dokl. Acad. Nauk, SSSR, **273**, No. 4 (1978), pp. 855-858.
- [7] L. POPESCU - N. POPESCU, *Sur la définition des prolongements résiduels transcendants d'une valuation sur un corps K à $K(X)$* , Bull. Math. Soc. Sci. Math. R. S. Romania, **33** (81), No. 3 (1989), pp. 257-264.
- [8] N. POPESCU - A. ZAHARESCU, *On the structure of the irreducible polynomials over local fields*, J. Number Theory, **52**, No. 1 (1995), pp. 98-118.
- [9] N. POPESCU - A. ZAHARESCU, *On the main invariant of an element over a local field*, Portugaliae Mathematica, **54**, Fasc. 1 (1997), pp. 73-83.
- [10] I. P. SERRE, *Corps locaux*, Hermann, Paris (1962).
- [11] O. F. G. SHILLING, *The theory of valuations*, Amer. Math. Soc., Mathematical Surveys, Number IV (1950).

Manoscritto pervenuto in redazione il 6 settembre 1996.