

SÉMINAIRE N. BOURBAKI

JOHN G. THOMPSON

Sylow 2-subgroups of simple groups

Séminaire N. Bourbaki, 1968, exp. n° 345, p. 543-545

http://www.numdam.org/item?id=SB_1966-1968__10__543_0

© Association des collaborateurs de Nicolas Bourbaki, 1968, tous droits réservés.

L'accès aux archives du séminaire Bourbaki (<http://www.bourbaki.ens.fr/>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

SYLOW 2-SUBGROUPS OF SIMPLE GROUPS

by John G. THOMPSON

I will primarily limit this lecture to a discussion of results obtained by two students, Goldschmidt and MacWilliams.

For each group X , let $m(X)$ be the minimal number of generators of X and let $d(X) = \max\{m(A)\}$, where A ranges over all the normal abelian subgroups of X .

Suppose G is simple and T is a Sylow 2-subgroup of G . In studying the minimal simple groups, it became clear that the case $d(T) \leq 2$ was anomalous. I handled the problem by first determining all the possibilities for a Sylow 2-subgroup and then using techniques available in any minimal simple group.

For further work in simple groups, it is desirable to classify all simple G such that $d(T) \leq 2$. The case $d(T) = 1$ is non trivial, but seems well on the way to a solution, so we assume $d(T) = 2$.

The most naive way to tackle this problem is first to classify all 2-groups T with $d(T) = 2$. This is difficult, but one result about 2-groups is helpful.

LEMMA 1.- If T is a 2-group with $d(T) \leq 2$, then every subgroup of T is generated by 4 elements.

This result then leads fairly rapidly to

THEOREM 1 (MacWilliams).- Suppose T is a Sylow 2-subgroup of the simple group G ,
 $d(T) = 2$ and $T.C(T) \subset N(T)$. Then $|T| = 4, 64$ or 128 and T is determined
by $|T|$.

If $|T| = 64$, T is isomorphic to a Sylow 2-subgroup of $U_3(4)$ and if
 $|T| = 128$, T is isomorphic to a Sylow 2-subgroup of the new simple groups of
 Janko of orders 604800 and 50232960 .

The structure of T in case $T.C(T) = N(T)$ is not yet determined. Several of
 the families of known simple groups satisfy these hypotheses.

Goldschmidt's work had a different origin. Initially, he studied simple groups
 with a Sylow 2-subgroup whose class of nilpotency is 2 . One of the results obtai-
 ned is that a Sylow 2-subgroup has exponent 4 . However, this emerges from a more
 general set up, the starting point being

LEMMA 2.- Suppose p is a prime and P is a Sylow p-subgroup of a group G .
Let n be the smallest integer such that $n(p-1) \geq c-1$, where c is the
class of nilpotency of P . Let $W = \langle x^p \mid x \in Z(P) \rangle$, where $Z(P)$ is the center
of P . Then W is weakly closed in P (that is, $g \in G$ and $W^g \subseteq P$ imply
 $W = W^g$).

This is elementary, but clever. The crucial result is

THEOREM 2 (Goldschmidt).- Suppose T is a Sylow 2-subgroup of G , W is a
weakly closed subgroup of T , $1 \subset W \subseteq Z(T)$ and t is an involution of $T - W$.
If $W \subseteq O_{2',2}(C(x))$ for every involution x of Wt , then G is not simple.

SYLOW 2-SUBGROUPS OF SIMPLE GROUPS

Here $O_{2',2}(X)$ is the largest normal subgroup of X with a normal 2-complement. The proof is character-theoretic.

If one couples this result with work of Gorenstein, we get

THEOREM 3 (Goldschmidt).- If W is weakly closed in T and $W \subseteq Z(T)^2$ where T is a Sylow 2-subgroup of G , then $W \subseteq O_{2',2}(G)$.

All these results are fragmentary, but given the state of finite group theory, this is not surprising.