

# SÉMINAIRE ÉQUATIONS AUX DÉRIVÉES PARTIELLES – ÉCOLE POLYTECHNIQUE

A. MENIKOFF

## **Hypoelliptic operators with double characteristics**

*Séminaire Équations aux dérivées partielles (Polytechnique)* (1976-1977), exp. n° 10,  
p. 1-8

[http://www.numdam.org/item?id=SEDP\\_1976-1977\\_\\_\\_\\_A9\\_0](http://www.numdam.org/item?id=SEDP_1976-1977____A9_0)

© Séminaire Équations aux dérivées partielles (Polytechnique)  
(École Polytechnique), 1976-1977, tous droits réservés.

L'accès aux archives du séminaire Équations aux dérivées partielles (<http://sedp.cedram.org>) implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme  
Numérisation de documents anciens mathématiques  
<http://www.numdam.org/>

ÉCOLE POLYTECHNIQUE

CENTRE DE MATHÉMATIQUES

PLATEAU DE PALAISEAU 91128 PALAISEAU CEDEX

Téléphone : 941.82 00 - Poste N°

Télex : ECOLEX 691 596 F

S E M I N A I R E   G O U L A O U I C - S C H W A R T Z   1 9 7 6 - 1 9 7 7

HYPSELLIPTIC OPERATORS WITH DOUBLE CHARACTERISTICS

par A. MENIKOFF

Exposé n° X

11 Janvier 1977



## X.1

I want to discuss pseudo-differential operators of the form  $P = P_m(x,D) + P_{m-1}(x,D) + \dots$  such that in a conic neighborhood  $\Gamma$  of  $(x_0, \xi_0) \in \mathbb{R}^{2n}$ ,  $P_m(x, \xi) \geq 0$  and vanishes to exactly second order on a submanifold  $\Sigma$  of codimension 1 transverse to the fiber axis.  $P$  is a classical pseudo-differential operator of order  $m$ . These assumptions imply that  $P_m = QU^2$  where  $Q \neq 0$  and  $d_\xi U \neq 0$  near  $(x_0, \xi_0)$ ,  $Q$  and  $U$  are real and homogeneous of order  $m-2$  and  $1$  respectively. Recall that the definition of subprincipal symbol of  $P$  is

$$P_{m-1}^S(x, \xi) = P_{m-1}(x, \xi) - \frac{1}{2i} \sum_{j=1}^n \frac{\partial^2 P_m}{\partial x_j \partial \xi_j} \quad .$$

The first two results I want to talk about concern necessary and sufficient conditions for  $P$  to be locally solvable and hypoelliptic.

**Theorem 1** : Suppose that  $\operatorname{Re} P_{m-1}^S \neq 0$  in  $\Gamma$  and that whenever  $\operatorname{Re} P_{m-1}^S < 0$  at a point of  $\Sigma$  then along the null bicharacteristic of  $U$  through that point  $\operatorname{Im} P_{m-1}^S$  has only zeros of even order  $\leq k$ .

(A) Then given any  $\varphi \in S^0(\Gamma)$ ,  $\varphi(x_0, \xi_0) \neq 0$  there exist operators  $E_i$ , and  $R_i$ ,  $i = 1, 2$  such that

- (1)  $E_i : H_S(\Gamma) \rightarrow H_{S+m-2+(k+2)/2(k+1)}(\Gamma)$  and
- (2)  $R_i : H_S(\Gamma) \rightarrow H_{S+m-2+1/k+1}(\Gamma)$  are bounded and
- (3)  $PE_1 = \varphi(x_1 D)I + R_1$
- (4)  $E_2 P = \varphi(x_1 D)I + R_2$

(B)  $P$  is hypoelliptic and locally solvable in  $\Gamma$

In the converse direction there is

**Theorem 2** : If at  $(x_0, \xi_0) \in \Sigma$ ,  $\operatorname{Re} P_{m-1}^S(x_0, \xi_0) < 0$  and  $\operatorname{Im} P_{m-1}^S$  changes sign and has a zero of finite order on the null bicharacteristic of  $U$  through  $(x_0, \xi_0)$  then  $P(x,D)$  is not locally solvable at  $x_0$ .

Note that since the hypotheses of theorem 2 also hold for  $P^*$ ,  $P$  is also not hypoelliptic.

Theorem 2 is due to P. Weston [5] for partial differential operators. Theorems 1 and 2 are due to P. Popivanov [2] and myself [1].

By means of Fourier integral operators the problem of studying  $P$  may be reduced to considering

$$(5) \quad Q = D_t^2 + a(t, x, D_x) + b(t, x, D_x, D_t)D_t$$

where  $a \in S^1$ ,  $b \in S^0$ . This is a consequence of the invariance of the assumptions and conclusions of theorem 2 under multiplication by real elliptic factors or conjugation by Fourier integral operators. Noting that  $a(t, x, \xi)$  is the sub-principal symbol of  $Q$  the assumptions of theorem 1 become that if  $\operatorname{Re} a < 0$  then  $\operatorname{Im} a(t, x, \xi)$  has only zeros of even order  $\leq k$  as a function of  $t$ . This, of course, implies that  $a$  has constant sign.

To understand the hypotheses of theorems 1 and 2 consider the ordinary differential operator

$$(6) \quad L = D_t^2 + a(t, x, \xi)$$

depending on  $(x, \xi) \in \mathbb{R}^{2n}$  as parameters. Considering functions which are oscillatory in  $x$ , the local solvability of  $Q^*$  is related to whether or not an estimate of the form

$$(7) \quad \| \| u \| \| \leq C \| Lu \|$$

holds for some pair of norms on  $C_0^\infty(\mathbb{R})$ . Using the Green-Liouville approximation,  $Lu \sim 0$  has solutions

$$(8) \quad u_{\pm}(t, x, \xi) = \frac{1}{a^{1/4}(t, x, \xi)} e^{\pm \int_{T(x, \xi)}^t \sqrt{a(t', x, \xi)} dt'}$$

If  $\operatorname{Re} a < 0$  and  $\operatorname{Im} a(t, x, \xi)$  changes signs at  $T(x, \xi)$  then  $\operatorname{Re} \sqrt{a}$  will also have a change of sign. This means that  $Lu \sim 0$  has a solution in  $\mathcal{S}$  and consequently an estimate of form (7) cannot hold.

To construct parametrices under the assumptions of theorem 1, I will use the notion of vector-valued pseudo-differential operators. The idea of using pseudo-differential operators whose symbols are operators between Hilbert space originated in Trèves [4]. I shall use an extension of this idea due to Sjöstrand [3] which allows the norms on the Hilbert spaces to vary.

Let  $H_1$  and  $H_2$  be a pair of Hilbert spaces whose norms depend on  $\xi \in \mathbf{R}^n$  such that

$$(9) \quad c \|u\|_{H_i} \leq \|u\|_{H_i(\xi)} \leq C(1 + |\xi|)^{m_i} \|u\|_{H_i}$$

for  $i = 1, 2$ . Let  $\mathcal{L}(H_1(\xi), H_2(\xi))$  be the space of bounded operators from  $H_1 \rightarrow H_2$  with the uniform operator norm, also varying with  $\xi$ . I will define the symbol in class  $S_{\rho, \delta}^m(\Omega \times \mathbf{R}^n; H_1(\xi), H_2(\xi))$  to be the class of functions  $A(x, \xi) : \Omega \times \mathbf{R}^n \rightarrow \mathcal{L}(H_1(\xi), H_2(\xi))$  such that

$$(10) \quad \left\| \partial_x^\alpha \partial_\xi^B A(x, \xi) \right\|_{\mathcal{L}(H_1(\xi), H_2(\xi))} \leq C(1 + |\xi|)^{m + \delta|\alpha| - \rho|\beta|}$$

The corresponding class of operators given by

$$A(x, D)u(x) = (2\pi)^{-n} \int e^{ix\xi} A(x, \xi) \hat{u}(\xi) d\xi$$

will be denoted by  $L_{\rho, \delta}^m(\Omega, H_1(D), H_2(D))$ , where  $\Omega \subset \mathbf{R}^n$ . These operators are at least maps

$$A(x, D) : C_0^\infty(\Omega, H_1) \rightarrow C_0^\infty(\Omega, H_2)$$

Furthermore, the standard calculus of pseudo-differential operators still holds for these operators.

The special case of such operators I will use is obtained by taking  $H = L^2(\mathbf{R}, dt)$  and  $B(\xi)$  to be the completion of  $C_0^\infty(\mathbf{R})$  in the norm

$$\|u\|_{B(\xi)} = \left\| (1 + |D_t|^2 + |D_x|)^{(k+2)/2(k+1)} u \right\|_{L^2}$$

An example of the boundlessness of results for this class of operators is that if  $A \in L_{\rho, \delta}^0(\mathbf{R}^n; B(D), H)$  and  $0 \leq \delta < 1/2 < \rho \leq 1$  then for any compact subset  $K$  of  $\Omega$

$$\|Au\|_{L^2(\mathbf{R}^{n+1})} \leq C \left\| (1 + |D_t|^2 + |D_x|)^{(k+2)/2(k+1)} u \right\|$$

for all  $u \in C_0^\infty(K \times \mathbf{R})$ .

To construct a parametrix of  $Q = D_t^2 + a(t, x, D_x)$  define

$$(11) \quad e(x, \xi, t, s) = \begin{cases} \frac{1}{2(a(t, x, \xi)a(s, x, \xi))^{1/4}} e^{-\int_t^s \sqrt{a(t', x, \xi)} dt'} & \text{if } t \leq s \\ e(x, \xi, s, t) & \text{if } s \leq t \end{cases}$$

Suppose  $\operatorname{Re} a < 0$ , and  $\operatorname{Im} a \geq 0$ , then  $a^{1/2}$  is a smooth function of all its variables. Choose the square root in (11) so that  $\operatorname{Re} a^{1/2} \geq 0$ . Define the corresponding integral operator with kernel  $g$  as

$$(12) \quad E_g(x, \xi)f(t) = \int_{-\infty}^{\infty} g e(x, \xi, t, s)f(s)ds$$

where  $g \in S^0(\mathbb{R}^{2n+2})$ . My candidate for a parametrix of  $Q$  is  $E_1(x, D)$ .

A calculation will show that

$$(13) \quad L(x, t, D_t) E(x, \xi) = I + E_g$$

where

$$(14) \quad g = \left( \frac{a''}{4a} - \frac{5}{16} \frac{a'^2}{a} \right) \in S^0$$

Equation (13) may be thought of as a statement about the multiplication of the symbols of  $Q$  and  $E(x, D)$ . The crucial step in proving theorem 1 is

**Lemma 3** : With the above assumptions if  $g \in S^0(\mathbb{R}^{2n+2})$  then  $E_g(x, \xi) \in S_{\rho, \delta}^0(\Omega; H, B(\xi))_s$  where

$$\frac{1}{2} - \frac{1}{2(k+1)} = \delta < 1/2 < \rho = \frac{1}{2} + \frac{1}{2(k+1)}$$

To prove lemma 3, I must show that

$$(15) \quad \left\| (1 + (D_t)^2 + |\xi|^2)^{(k+2)/2(k+1)} E u \right\|_{L^2} \leq C \|u\|_{L^2}$$

and similar estimates for  $\partial_x^\alpha \partial_\xi^\beta E(x, \xi)$ . I will need the following lemmas.

**Lemma 4** : Suppose  $k(t, s)$  is a measurable function on  $\mathbb{R}^2$  such that

$$\int |k(t,s)| ds \leq B \quad \text{and} \quad \int |k(t,s)| dt \leq B .$$

Then  $K f(t) = \int k(t,s)f(s)ds$  is a bounded operator on  $L^2(\mathbb{R})$  and  $\|K\| \leq B$ .

**Lemma 5** : If  $0 \leq a(t,x,\xi) \in S^0(\mathbb{R}^{n+1} \times \mathbb{R}^n)$  and has zeros in  $t$  of order  $\leq k$ , then there is a constant  $c > 0$  such that

$$(16) \quad c|t-s|^{k+1}|\xi| \leq \int_t^s a(t',x,\xi) dt'$$

For a proof see the appendix of [4].

**Lemma 6** : Given  $C > 0$  there is a constant  $C' > 0$  such that for any complex number  $z$   $|\operatorname{Im} z| \leq C|\operatorname{Re} z|$  implies  $|\operatorname{Re} z|^{1/2} \geq C'|\operatorname{Im} z|/|\operatorname{Re} z|^{1/2}$ .

To prove (15), I will estimate

$$I = \int_t^\infty |e(x,\xi,t,s)| ds \quad \text{and} \quad \int_{-\infty}^t |e(x,\xi,t,s)| ds .$$

Using the definition of  $E$ ,

$$(17) \quad I \leq \int_t^\infty \frac{1}{|a(t)a(s)|^{1/4}} e^{-\int_s^t \sqrt{a} dt} ds .$$

Since  $\operatorname{Re} a \neq 0$   $|a(s)a(t)|^{1/4} \geq C|\xi|^{1/2}$ . Combining, lemmas 5 and 6 give

$$\operatorname{Re} \int_t^s \sqrt{a(t')} dt' \geq C|\xi|^{1/2}|t-s|^{k+1}$$

Using the last two inequalities in (17), I get

$$I \leq C|\xi|^{-1/2} \int_t^\infty e^{-c|\xi|^{1/2}|t-s|^{k+1}} ds = C|\xi|^{-\frac{1}{2} - \frac{1}{2(k+1)}} .$$

Lemma 4, then yields the bound

$$\| |\xi|^{(k+1)/2(k+2)} E(x,\xi)u \| \leq C \|u\| .$$

Since  $D_t^2 E = -a(t,x,\xi)E +$  bounded operators and  $a \in S^1$  we also may



estimate twice as many  $t$ -derivatives as  $x$ -derivatives. This gives (15).

I am now in a position to complete the proof of theorem 1 by applying the calculus of pseudo-differential operators. Since  $D_t^2$  is independent of  $t$  the symbol of  $D_t^2 \circ E(x, D_x)$  is  $D_t^2$  applied to the symbol of  $E$ . To compute the composition of  $a(t, x, D)$  and  $E$ , consider  $a$  as being in the class  $S^1(\Omega \times \mathbb{R}^n; B(\xi), B(\xi))$ . I then have

$$a(t, x, D_x) \circ E(x, D_x) = (aE)(x, D) \text{ mod } L^{-\min(\rho, 1-\delta)}(H, B(\xi)).$$

Combining the above observations with (13) gives

$$Q(t, x, D_t, D_x) \circ E(x, D_x) = I + R,$$

where  $R \in L^{k/2(k+1)}(\Omega; H, B(D))$ .

For  $R$ , there will be the estimate

$$\|(1 + |D_t|^2 + |D_x|^2)^{1/k+1} Ru\| \leq C \|u\|, \quad u \in C_0^\infty(\mathbb{R}^{n+1}).$$

This completes the proof of theorem 1.

An extension of the argument used for theorem 1 will give

**Theorem 7 :** Let  $L = D_t^2 + a(t, x, \xi) + b(t, x, D_t, D_x)D_t + \dots$  where  $a \in S^1$ ,  $b \in S^0$  and suppose that  $\operatorname{Re} a \neq 0$  and if  $\operatorname{Re} a < 0$  then  $\operatorname{Im} a$  has a constant sign as a function of  $t$  and  $\operatorname{Im} a$  never vanishes on an open  $t$ -interval for fixed  $x$  and  $\xi$ .

Then,  $L$  is locally solvable and for any  $\varepsilon > 0$  there is a neighborhood  $\omega$  of sufficiently small diameter such that

$$\|u\|_{S^{m-3/2}} \leq \varepsilon \|PLu\|_S + C \|u\|_{S^{m-2}}, \quad u \in C_0^\infty(\omega).$$

Operators whose sub-principal symbol vanishes on the characteristic variety may also be treated. Now let

$$Q = D_t^2 + b(t, x, D_x, D_t)$$

where  $b(0, x, \xi) = t^k a(t, x, \xi)$ ,  $a \neq 0$ .

A parametrix for  $L$  may be attempted to be constructed in the same way as the one in theorem 1 was. Consider the ordinary differential operator  $L = D_t^2 + t^k a(t, x, \xi)$ . A solution of  $Lu \sim 0$  may be sought of the form

$$u = g(t, x, \xi) V(\phi(t, x, \xi))$$

where  $V$  is a solution of  $-V'' + s^k V(s) = 0$ . This will force that

$$\phi = (k+2/2 \int_0^t \sqrt{t^k a} dt)^{2/k+2},$$

and  $g = \phi^{-1/2}$ . Then  $Lu = \left(-\frac{d^2 g}{dt^2}\right) V(\phi)$ .

If  $Lu \sim 0$  has two independent solutions which increase and decrease exponentially on opposite sides of the  $t$ -axis, a parametrix may again be constructed. For instance, I have shown

Theorem 8 : If either  $a(0, x_0, \xi_0)$  is not real,  $\phi$  or  $k$  is even and  $a(0, x_0, \xi_0) > 0$  then  $L$  is hypoelliptic and locally solvable and has left and right parametrices.

Theorem 9 : If  $a(0, x_0, \xi_0)$  is real and  $\text{Im} a$  has a zero of first order then  $Q$  is hypoelliptic.

Theorem 10 : If  $a(0, x, \xi)$  is real and  $\text{Im} b(0, x, \xi) \neq 0$ , then

$$Q = D_t^2 + t^k (a(t, x, D_x) + t^\ell b(t, x, D_x))$$

is locally solvable when  $\ell < k+2$ .

I don't know what the situation is for the operators of theorem 10 if  $\ell \geq k+2$ , nor do I have non-solvability results if  $k > 1$  in theorem 8 etc...

---

## REFERENCES

- [1] A. Menikoff : On hypoelliptic operators with double characteristics  
An. Sc. N. Sup. di Pisa, to appear.

- [2] P. Popivanov : Proc. Bulgarian Acad. Sc. (1975).
  - [3] J. Sjöstrand : Parametrices for pseudo-differential operators with multiple characteristics. Ark. för Mat. 12, 85-130 (1974).
  - [4] F. Trèves : A new method of proof of the subelliptic estimates. Comm. Pure and Appl. Math. 24, 71-115 (1971).
  - [5] P. Weston : A necessary condition for the local solvability of  $P_m^2(x,D) + P_{2m-1}(x,D)$  ; J. Diff. Eq. ,to appear.
-