

SÉMINAIRE DE THÉORIE DES NOMBRES DE GRENOBLE

BENEDICT H. GROSS

Topological Invariants of Real Algebraic Curves

Séminaire de théorie des nombres de Grenoble, tome 9 (1980-1981), exp. n° 2, p. 1-2

<http://www.numdam.org/item?id=STNG_1980-1981__9__A2_0>

© Institut Fourier – Université de Grenoble, 1980-1981, tous droits réservés.

L'accès aux archives du séminaire de théorie des nombres de Grenoble implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/conditions>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

*Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques
<http://www.numdam.org/>*

Séminaire de Théorie des Nombres
21 janvier 1981

Grenoble

TOPOLOGICAL INVARIANTS OF REAL ALGEBRAIC CURVES

par Benedict H. GROSS

Let X be an absolutely irreducible projective smooth curve over the field \mathbb{R} of real numbers. One can associate to X three topological invariants :

$$(1) \quad \begin{cases} g = \text{genus of the Riemann surface } X(\mathbb{C}) \\ n = \text{number of components of the real locus } X(\mathbb{R}) \\ a = \begin{cases} 0 & \text{if } X(\mathbb{R}) \text{ disconnects } X(\mathbb{C}) \\ 1 & \text{if } X(\mathbb{C}) - X(\mathbb{R}) \text{ is connected.} \end{cases} \end{cases}$$

These invariants satisfy the following relations, which follow from the classification of compact 2-manifolds :

$$(2) \quad \begin{cases} g \geq 0 . \\ 0 \leq n \leq g+1 . \\ \text{If } n = 0 \text{ then } a = 1 . \\ \text{If } n = g+1 \text{ then } a = 0 . \\ \text{If } a = 0 \text{ then } n \equiv g+1 \pmod{2} . \end{cases}$$

Klein showed that any triple (g, n, a) satisfying the relations (2) actually occurs as the invariants of a real algebraic curve X , and that the moduli of all real curves with a fixed triple of invariants is a connected real analytic space of dimension g for $g \leq 1$ and $3g - 3$ for $g \geq 2$.

The invariant g can also be described as the algebraic genus of the function field $\mathbb{R}(X)$. Here we give an algebraic description of the invariants n

and a in the case where $n > 0$ (so X has a real point). Consider the finite set S of real theta-characteristics on X ; this is partitioned into even and odd characteristics, and one can show that :

$$(3) \quad \begin{cases} \text{Card } S = 2^{g+n-1} \\ \text{Card } S_{\text{odd}} = 2^{g-1}(2^{n-1} + a - 1) . \end{cases}$$

For example, any smooth quintic $X \subset \mathbb{P}^2$ with one real component has $a = 1$, as S contains an odd characteristic which is cut out by lines.

For more details, see the forthcoming article by the author and J. Harris in the Annals of the Ecole Normale Supérieure.

Benedict H. GROSS
Dept. of mathematics
Princeton University
PRINCETON, NEW JERSEY 08540
U.S.A.